



Seasonal variation of cladocera in a perennial pond, Ambajogai (Maharashtra), India

Vishwas Balasaheb Sakhare¹  • Ashwini Dnyandeo Chalak² 

¹ Post Graduate Department of Zoology, Yogeshwari Mahavidyalaya, Ambajogai – 431517, India

² Department of Zoology, Late Shankarrao Gutte Gramin Arts, Commerce and Science College, Dharmapuri – 431515, India

Correspondence

Vishwas Balasaheb Sakhare; Post Graduate Department of Zoology, Yogeshwari Mahavidyalaya, Ambajogai – 431517, India

 vbsakhare@gmail.com

Manuscript history

Received 12 December 2019 | Revised 7 April 2020 | Accepted 8 April 2020 | Published online 29 May 2020

Citation

Sakhare VB, Chalak AD (2020) Seasonal variation of cladocera in a perennial pond, Ambajogai (Maharashtra), India. *Journal of Fisheries*. <http://journal.bdfish.org/index.php/fisheries/article/view/JFish20181>

Abstract

A systematic investigation on seasonal variations of cladocera in Udayan Sarovar, a perennial pond near Swami Ramanand Teerth Mahavidyalaya at Ambajogai (Maharashtra, India) was carried out for a period of one year from June 2018 to May 2019. A total of 9 species of cladocera were identified. The cladoceran diversity fluctuated seasonally and the maximum number of 240 organisms L⁻¹ was recorded during month of April and minimum number of 22 organisms L⁻¹ during the month of September.

Keywords: Cladoceran diversity; zooplankton; seasonal variations; perennial pond; Maharashtra

1 | INTRODUCTION

Cladocera, commonly known as ‘water fleas’, are planktonic crustaceans ranging in size from 0.2 to 5.00 mm and belong to the class Branchiopoda. Globally about 4000 species of cladocera have been described (Covich and Tharp 1991). About 187 species of freshwater cladocera have been reported from India (Raghunathan and Kumar 2003). A range of studies on cladocera have been conducted from different ponds of India (e.g. Biswas 1966, 1971; Nayar 1971; Michael and Sharma 1988; Raghunathan 1989; Sharma and Sharma 1991; Gupta 2002; Raghunathan and Suresh 2002; Raghunathan and Kumar 2003; Sharma *et al.* 2005; Siraj *et al.* 2007; Chanderkiran 2008; Sharma and Chanderkiran 2011; Sharma and Kotwal 2011; Sharma *et al.* 2012). However, sufficient investigations have not been done on the cladoceran diversity of freshwater bodies in Maharashtra, India. Con-

sidering the lack of knowledge of cladoceran diversity of this region, a systematic investigation on seasonal variations on cladoceran diversity from Udayan Sarovar, a perennial pond of Ambajogai, Maharashtra was carried out for one year.

2 | METHODOLOGY

With aim to investigate the seasonal diversity of the cladocerans, a perennial pond near Swami Ramanand Teerth Mahavidyalaya at Ambajogai (18043'37.6860"N, 76022'51.9168"E; at an elevation of 633 m above sea level) in Indian state of Maharashtra was surveyed monthly between June 2018 and May 2019. Approximately 50 litres of water was filtered through plankton net of standard bolting cloth number 25 (0.03 – 0.04 μ mesh size). The filtrate was taken in plastic vials and preserved in 5% formaldehyde solution. These fixed samples were brought to the laboratory for taxonomical studies (Ward and Whip-

ple 1959; Edmondson and Winberg 1971; Pennak 1978). Drop count method of Trivedy and Goel (1992) was followed for enumeration of cladocerans and expressed as organisms per litre.

3 | RESULTS AND DISCUSSION

Recorded cladocerans were represented by nine species including *Diaphanosoma sarsi*, *D. excisum*, *Ceriodaphnia cornuta*, *Moina micrura*, *Macrothrix spinosa*, *Alona rectangular*, *A. nana*, *Indialona ganapati* and *Daphnia pulex* (Table 1). The cladoceran diversity fluctuated seasonally and the maximum number of 240 organisms L⁻¹ was recorded during month of April and minimum number of 22 organisms L⁻¹ during the month of September (Figure 1).

Ceriodaphnia cornuta and *Alona nana* were absent for three months in 2018 i.e. July, August and September.

Diaphanosoma sarsi and *A. rectangular* were absent in August and September, while *Macrothrix spinosa* was not seen in September. The cladocerans such as *Diaphanosoma excisum*, *Moina micrura*, *Indialona ganapati* and *Daphnia pulex* were present throughout the study period. All the cladoceran species showed minima in monsoon and maxima in summer. The decline in cladoceran population during monsoon coincided with the inflow of allochthonous nutrients, along with surface run off (Mishra et al. 2010). Besides, cladocerans produce resting eggs in autumn or winter months of the year which hatch from February onwards (Dokulil et al. 1990) culminating in peak population during summer. Sarmila Sree and Shameem (2017) and Chalak and Sakhare (2018) also reported higher cladoceran density in summer and lower in monsoon.

TABLE 1 Species composition of cladocera during year 2018–19.

Species	Cladocera density (organisms L ⁻¹) in different months											
	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May
<i>Diaphanosoma sarsi</i>	02	04	00	00	08	10	08	06	19	30	42	17
<i>Diaphanosoma excisum</i>	06	06	01	08	10	10	14	10	17	24	38	20
<i>Ceriodaphnia cornuta</i>	03	00	00	00	12	14	10	14	22	16	35	15
<i>Moina micrura</i>	10	10	05	04	08	10	12	10	12	20	30	24
<i>Macrothrix spinosa</i>	10	05	07	00	07	10	10	19	20	24	28	25
<i>Alona rectangular</i>	05	02	00	00	05	05	05	08	15	18	18	20
<i>Alona nana</i>	08	00	00	00	05	05	05	08	08	10	12	14
<i>Indialona ganapati</i>	07	09	07	05	08	08	12	10	12	15	12	16
<i>Daphnia pulex</i>	10	10	12	05	18	15	15	15	20	20	25	30
Total	61	46	32	22	81	87	91	100	145	177	240	181




FIGURE 1 Seasonal variation of cladocera in the study pond.

Hutchinson (1967) pointed out that water temperature increases the development of cladocera while Patalas (1972) stated that temperature is the primary factor influencing the zooplankton abundance. Quadri and Yousuf (1980) also recorded the significance of temperature in occurrence and the distribution of cladocera in Kashmir lake. The present investigation corroborate with the findings of Hutchinson (1967).

The overall view of the present investigation reveals good diversity of cladocera with maxima and minima in the month of April and September respectively. This investigation will be useful to monitor the health (water quality) and wealth (fish population) of this pond in near future.

ACKNOWLEDGEMENTS

The authors are grateful to Principal, Yogeshwari Mahavidyalaya, Ambajogai for providing laboratory facilities. Much appreciation is also given to the reviewers of this research paper.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES


Biswas S (1966) Five species of Daphnidae (Crustacea: Cladocera) from Shimla Hills in India with a new record of *Alona costata* Sars from Kameng Division; NEFA. Journal of the Zoological Society of India 16(1–2): 92–98.

- Biswas S (1971) Fauna of Rajasthan, part II (Crustacea: Cladocera). Records of the Zoological Survey of India 63: 95–141.
- Chalak AD, Sakhare VB (2018) Reservoir ecology. Discovery Publishing House Pvt. Ltd., New Delhi, India.
- Chanderkiran (2008) Diversity, dynamics and cyclomorphosis in cladocera inhabiting freshwater ponds of Jammu. MPhil dissertation, University of Jammu, Jammu.
- Covich AP and Tharp JH (1991) Crustacea, introduction to Percaridae. In: Tharp JH, Covich AP (Eds) Ecology and classification of North American freshwater invertebrates. Academic Press, San Diego. pp. 665–690.
- Dokulil M, Herzig A, Jagsch, A (1990) Trophic relationships in the pelagic zone of Mondsee, Australia. Hydrobiologia 191: 199–212.
- Edmondson WT, Winberg CG (1971) A manual on methods for assessment of secondary production in freshwater. IBP handbook 17: 358.
- Gupta S (2002) Studies on the diversity of cladocerans in lake Mansar, Jammu. MPhil dissertation, University of Jammu, Jammu.
- Hutchinson GE (1967) A treatise on limnology. Introduction to lake biology and the limnoplankton. John Wiley and Sons, New York, London.
- Michael RG, Sharma BK (1988) Indian cladocera. Fauna of India and adjacent countries. Zoological Survey of India.
- Mishra A, Chakraborty SK, Jaiswar AK, Sharma AP, Deshmukhe G, Mohan M (2010) Plankton diversity in Dhaura and Baigul reservoirs of Uttarakhand. Indian Journal of Fisheries 57(3): 19–27.
- Nayar CKG (1971) Cladocera of Rajasthan. Hydrobiologia 37: 509–519.
- Patalas K (1972) Crustacea plankton and eutrophication of St. Lawrence Great Lakes. Journal of the Fisheries Research Board of Canada 29: 1451–1462.
- Pennak RW (1978) Freshwater invertebrates of United States, 2nd edition. Wiley Inter Science Publication. John Wiley and Sons, New York.
- Quadri MY, Yousuf AR (1980) Influence of the physico-chemical factors on the seasonality of cladoceras in Lake Manasbal. Geobios 7: 273–276.
- Raghunathan MB (1989) Indian cladocera (Crustacea). Indian Review of Life Sciences 9: 137–152.
- Raghunathan MB, Kumar S (2003) A checklist of Indian Cladocera (Crustacea). Zoos' Print Journal 18: 1180–1182.
- Raghunathan MB, Suresh R (2002) Checklist of Indian cladocera (Crustacea). Zoos' Print Journal 18(8): 1180–1182.
- Sharma BK, Sharma S (1991) Freshwater cladocerans (Crustacea: Branchiopoda: Cladocera), Zoological Survey of India, Kolkata, State Fauna Series: fauna of Meghalaya, 4(9): 469–550.
- Sharma KK, Chanderkiran (2011) Comparative analysis of cladoceran communities from three subtropical freshwater ponds of Jammu: present composition and diversity. The Bioscan 6(2): 233–237.
- Sharma KK, Kaur S, Gupta S (2005) Distributional pattern of cladoceran in Subtropical lake, Mansoor, J. & K. National Academy of Sciences 76(B): 434–350.
- Sharma KK, Kotwal S (2011) Studies on diversity and dynamics of Cladocera in a subtropical Sungal pond, Akhnour (J & K). The Bioscan 6(4): 623–625.
- Sharma V, Verma BK, Sharma R, Sharma MS, Gaur KS (2012) A report on the freshwater cladocera (Crustacea: Branchiopoda) of South Rajasthan (India). International Journal of Environmental Sciences 3(1): 275–296.
- Sharmila Sree J, Shameem U (2017) Zooplankton diversity indices and seasonal variations in Meghadrigedda reservoir, Visakhapatnam, Andhra Pradesh, India. European Journal of Biotechnology and Bioscience 5(1): 4–11.
- Siraj S, Yousuf, AR, Parveen M (2007) Cladoceran community in Sheshbug wetland, Kashmir. Journal of Research and Development 7: 67–74.
- Trivedy RK, Goel PK (1992) Chemical and biological methods for water pollution studies. Environmental Publication, Karad (India).
- Ward HB, Whipple GC (1959) Freshwater biology, 2nd edition, John Wiley and Sons, New York.

CONTRIBUTION OF THE AUTHORS

VBS research design, sampling, laboratory works and manuscript writing;
ADC sampling and laboratory works.



VB Sakhare  <https://orcid.org/0000-0002-0871-4257>
AD Chalak  <https://orcid.org/0000-0002-3343-2227>