

NANOSTRUCTURED LIPID CARRIER LOADED WITH ZINGIBER
OFFICINALE AND ZINGIBER ZERUMBET OIL FOR INDUCTION OF
LIPOLYSIS IN SUBCUTANEOUS SKIN LAYER

NUR AYSHAH BINTI ROSLI

A thesis submitted in fulfilment of the
requirements for the award of the degree of
Doctor of Philosophy (*Bioprocess Engineering*)

School of Chemical and Energy Engineering
Faculty of Engineering
Universiti Teknologi Malaysia

MAY 2019

To my beloved husband, sons and family

ACKNOWLEDGEMENT

In preparing this thesis, I was in contact with many people, researchers, academicians, and practitioners. They have contributed towards my understanding and thoughts. In particular, I wish to express my sincere appreciation to my main supervisor, Assoc. Prof. Dr. Azila Abdul Aziz, for encouragement, guidance, critics and friendship. I am also very thankful to my co-supervisors Dr. Rosnani Hasham for her guidance, advices and motivation. Without their continued support and interest, this thesis would not have been the same as presented here.

I am also indebted to Universiti Sains Malaysia (USM) and Ministry of Education Malaysia for funding my Ph.D. study through Rancangan Latihan Kakitangan Akademik Scheme. Librarians at UTM and especially Encik Abas also deserve special thanks for their assistance in supplying the relevant literatures and information. Not to forget, all staffs and technicians in IBD, UTM, UPM, UKM and UiTM whom assisted me during sample analyses.

My fellow postgraduate colleague should also be recognised for their support. My sincere appreciation also extends to all my colleagues and others who have helped at various occasions. Their views and tips are useful indeed.

Unfortunately, it is not possible to list all of them in this limited space. I am grateful to all my family members whom patiently wait for me to complete this journey. Alhamdulillah and thank you all.

ABSTRACT

Body weight loss strategies include combination of nutrition consultation, regular exercise, drug prescription, invasive intervention and/or non-invasive intervention. A non-invasive intervention such as transdermal administration of active ingredients helps in reducing localized subcutaneous adipose tissue through lipolysis. *Zingiber officinale* (ZO) and *Zingiber zerumbet* (ZZ) belong to the Zingiberaceae family and both have been discovered to possess lipolysis activity. An efficient drug delivery system is required to encapsulate ZO and ZZ, and deliver these substances up to the subcutaneous skin layer. In this study, ZO and ZZ oil were successfully encapsulated in nanostructured lipid carrier (NLC) using hot homogenization technique followed by ultrasonication. D-optimal mixture design was used to optimize the NLC in which the composition of ZO and ZZ oil, solid lipid, and liquid lipid were the independent variables, while particle size, polydispersity index (PDI), zeta potential, and encapsulation efficiency were the dependent variables. From the study, the optimum formulation for NLC-ZO was 3.6% ZO oil, 1.4% glyceryl monostearate, and 5.0% virgin coconut oil, and the NLC-ZO obtained had the following properties: 90.7 nm particle size, 0.15 PDI, -45.7 mV zeta potential and 88.8% encapsulation efficiency. The optimum NLC-ZZ formulation was 3.7% ZZ oil, 1.3% glyceryl monostearate, and 5.0% virgin coconut oil, and the NLC-ZZ obtained had the following properties: 91.0 nm particle size, 0.17 PDI, -40.9 mV zeta potential and 94.4% encapsulation efficiency. Morphology study revealed the spherical shape of NLC-ZO and NLC-ZZ, and the particle size obtained through transmission electron microscope conformed with the size measured through dynamic light scattering technique. Thermogram profile of NLC showed incorporation of ZO and ZZ into NLC lowered the melting temperature of lipid mixture, thus producing a less ordered crystalline structure of NLC to accommodate the active ingredients. Fourier transform infrared spectroscopy analysis demonstrated an interaction existed between ZO/ZZ and the NLC system. *In vitro* penetration study using Strat-M® membrane and freshly excised rat skin showed NLC-ZO and NLC-ZZ had higher penetration flux compared to free ZO and ZZ. The release of ZO from NLC-ZO followed Korsmeyer-Peppas model, and the release of ZZ from NLC-ZZ followed zero order kinetic model. 90-day storage stability study showed no significant changes for both NLC-ZO and NLC-ZZ in terms of particle size, PDI and zeta potential. *In vitro* lipolysis study revealed the potential of NLC-ZO and NLC-ZZ as anti-obesity agents as they stimulated the release of glycerol in 3T3-L1 adipocytes cells. *In vivo* NLC distribution in female Sprague-Dawley rats showed penetration deep into the dermis layer after 12 h of application. Free fatty acid release was detected when the rats were treated with NLC-ZO and NLC-ZZ for 3 h. This study demonstrated that ZO and ZZ were successfully encapsulated in NLC and the nanoparticles were effectively utilized to enhance dermis penetration and induce lipolysis activity via *in vitro* and *in vivo*.

ABSTRAK

Strategi penurunan berat badan meliputi kombinasi perundingan pemakanan, kekerapan bersenam, preskripsi ubat-ubatan, campur tangan invasif dan/atau campur tangan tidak-invasif. Campur tangan tidak invasif seperti pengurusan bahan aktif secara transdermal membantu mengurangkan tisu adiposa subkutanus setempat menerusi lipolisis. *Zingiber officinale* (ZO) dan *Zingiber zerumbet* (ZZ) adalah daripada keluarga *Zingiberaceae* dan kedua-duanya ditemui mempunyai aktiviti lipolisis. Sistem penyampaian ubat yang cekap diperlukan untuk merangkum ZO dan ZZ dan menghantar bahan ini sehingga ke lapisan subkutanus kulit. Dalam kajian ini, minyak ZO dan minyak ZZ telah berjaya dimuatkan dalam pembawa lipid berstruktur nano (NLC) menggunakan teknik homogenisasi panas diikuti ultrasonik. Reka bentuk campuran D-optimal telah digunakan untuk mengoptimumkan NLC di mana komposisi minyak ZO dan ZZ, lipid pepejal dan lipid cecair sebagai pembolehubah tidak bersandar, manakala saiz zarah, indeks poliserakan (PDI), potensi zeta, dan kecekapan enkapsulasi sebagai pembolehubah bersandar. Dari kajian ini, rumusan yang optimum untuk NLC-ZO adalah 3.6% minyak ZO, 1.4% gliseril monostearat, dan 5.0% minyak kelapa dara, dan NLC-ZO yang dihasilkan mempunyai sifat-sifat berikut: saiz zarah 90.7 nm, PDI 0.15, keupayaan zeta -45.7 mV, dan kecekapan pengkapsulan 88.8%. Rumusan NLC-ZZ yang optimum adalah 3.7% minyak ZZ, 1.3% gliseril monostearat, dan 5.0% minyak kelapa dara, dan NLC-ZZ yang dihasilkan mempunyai sifat berikut: saiz zarah 91.0 nm, PDI 0.17, keupayaan zeta -40.9 mV dan kecekapan pengkapsulan 94.4%. Kajian morfologi menunjukkan yang NLC-ZO dan NLC-ZZ berbentuk sfera, dan saiz zarah yang diperolehi melalui analisis mikroskop transmisi elektron mengesahkan saiz yang diukur melalui teknik penyelerakan cahaya dinamik. Profil termogram NLC menunjukkan penggabungan ZO dan ZZ ke dalam NLC telah menurunkan suhu lebur campuran lipid, sehingga menghasilkan struktur kristal NLC yang kurang teratur untuk menampung bahan-bahan aktif. Analisis spektroskopi inframerah transformasi Fourier menunjukkan interaksi antara sistem ZO/ZZ dan NLC. Kajian penembusan secara *in vitro* menggunakan membran Strat-M® dan kulit tikus yang disiat segar menunjukkan NLC-ZO dan NLC-ZZ mempunyai fluks penembusan yang lebih tinggi berbanding ZO dan ZZ. Pelepasan ZO daripada NLC-ZO mengikuti model Korsmeyer-Peppas dan pelepasan ZZ daripada NLC-ZZ mengikuti model kinetik tertib sifar. Kestabilan penyimpanan selama 90 hari menunjukkan tiada perubahan ketara bagi NLC-ZO dan NLC-ZZ dari segi saiz zarah, PDI dan keupayaan zeta. Kajian lipolisis secara *in vitro* telah mendedahkan potensi NLC-ZO dan NLC-ZZ sebagai agen anti-obesiti yang merangsang pembebasan gliserol dalam sel-sel adiposit 3T3-L1. Kajian penyebaran NLC secara *in vivo* pada tikus Sprague-Dawley betina telah menunjukkan penembusan sehingga ke lapisan dermis selepas 12 jam rawatan. Pembebasan asid lemak bebas telah dikesan apabila tikus dirawat dengan NLC-ZO dan NLC-ZZ selama 3 jam. Kajian ini menunjukkan bahawa ZO dan ZZ telah berjaya dirangkumkan dalam NLC dan zarah nano ini berkesan untuk meningkatkan penembusan sehingga ke lapisan dermis dan mendorong aktiviti lipolisis secara *in vitro* dan *in vivo*.

TABLE OF CONTENTS

	TITLE	PAGE
	DECLARATION	ii
	DEDICATION	iii
	ACKNOWLEDGEMENT	iv
	ABSTRACT	v
	ABSTRAK	vi
	TABLE OF CONTENTS	vii
	LIST OF TABLES	xiv
	LIST OF FIGURES	xvii
	LIST OF ABBREVIATIONS	xxii
	LIST OF SYMBOLS	xxviii
	LIST OF APPENDICES	xxix
CHAPTER 1	INTRODUCTION	1
1.1	Research Background	1
1.2	Problem Statement	4
1.3	Objective of the Research	6
1.4	Scope of the Research	6
1.5	Significance and Original Contributions of the Research	7
1.6	Thesis Structure and Organization	8
CHAPTER 2	LITERATURE REVIEW	11
2.1	Obesity: The Overview	11
2.1.1	Current Progress for Obesity Treatment	13
2.1.2	Lipolysis	15
2.1.3	Topical Formulation for Fat Reduction	19

2.2	Essential Oil	26
2.2.1	<i>Zingiber Officinale</i>	28
2.2.2	<i>Zingiber Zerumbet</i>	30
2.2.3	Role of <i>Zingiber officinale</i> and <i>Zingiber zerumbet</i> in Obesity Management	32
2.3	Skin	35
2.3.1	Subcutaneous Adipose Tissue – Drug Administration	38
2.4	Drug Delivery	40
2.4.1	Transdermal Drug Delivery	43
2.4.2	Lipid Nanoparticles	48
2.4.3	Nanostructured Lipid Carrier (NLC)	51
2.4.3.1	Method Preparation of NLC	57
2.4.3.2	Selection of Lipid	61
2.4.3.3	Selection of Emulsifier	63
2.5	Characterization Techniques of Lipid Nanoparticles	65
2.5.1	Particle Size, Polydispersity Index and Zeta Potential Analysis	65
2.5.2	Encapsulation Efficiency and Drug Loading Capacity	68
2.5.3	Nanoparticle Morphology	70
2.5.4	Differential Scanning Calorimetry (DSC)	71
2.5.5	Fourier-Transform Infrared Spectroscopy (FTIR)	73
2.5.6	Penetration Study	74
2.5.7	In Vitro Release study	76
2.5.8	Physical Stability Study	80
2.5	Summary	81
CHAPTER 3	RESEARCH METHODOLOGY	83
3.1	Introduction	83
3.2	Materials	85
3.2	Preparation of ZO and ZZ Loaded NLC	85

3.2.1	Addition of Dye-Marker	86
3.2.2	Lipid Screening	87
3.3	Optimization of Preparation Condition for NLC Encapsulated ZO and ZZ	87
3.4	Characterization of Optimized NLC Encapsulates ZO and ZZ	89
3.4.1	Particle Size, Zeta Potential and PDI Analysis	89
3.4.2	Encapsulation Efficiency Study	90
3.4.3	Transmission Electron Microscopy (TEM)	89
3.4.4	Differential Scanning Calorimetry (DSC)	91
3.4.5	Fourier-Transform Infrared Spectroscopy (FTIR)	91
3.4.6	Skin Penetration Study Using Franz Diffusion Cell	92
3.4.6.1	Skin Preparation	95
3.4.7	In Vitro Release Study	95
3.4.8	Stability Study	96
3.5	Efficacy Study of NLCs system	97
3.5.1	<i>In Vitro</i> Adipose 3T3-L1 Cell Lines	97
3.5.1.1	3T3-L1 Preparations and Differentiations	97
3.5.1.2	MTT Assay	98
3.5.1.3	Lipolysis Assay	99
3.5.2	Efficacy of NLCs <i>in vivo</i>	100
3.5.2.1	<i>In Vivo</i> Skin Deposition Study	100
3.5.2.2	Observation under the Skin	101
3.5.2.3	Lipolysis Marker Measurement	102
3.6	Statistical Analysis	102

CHAPTER 4 RESULTS AND DISCUSSION 103

4.1	Introduction	103
4.2	Preliminary Study	104
4.2.1	Lipid Screening	104
4.2.2	Screening of Ultrasonication Parameters	106
4.3	Optimization of NLC Composition Mixture using D-Optimal Mixture Design	109
4.3.1	Optimization of Preparation Condition for NLCs Encapsulated ZO	110
4.3.1.1	Influence of Material Composition on Particle Size	111
4.3.1.2	Influence of Material Composition on PDI	116
4.3.1.3	Influence of Material Composition on Zeta-Potential	120
4.3.1.4	Influence of Material Composition on Encapsulation Efficiency	124
4.3.1.5	Optimization of NLC-ZO Formulation with The Mixture Design	128
4.3.2	Optimization of Preparation Condition of NLC Encapsulated ZZ	128
4.3.2.1	Influence of Material Composition on Particle Size	131
4.3.2.2	Influence of Material Composition on PDI	134
4.3.2.3	Influence of Material Composition on Zeta-Potential	138
4.3.2.4	Influence of Material Composition on Encapsulation Efficiency	141
4.3.2.5	Optimization of NLC-ZZ Formulation with The Mixture Design	145
4.3.3	Summary	145
4.4	Characterization of Optimized NLC	146

4.4.1	Morphology Analysis	146
4.4.2	Crystallization Behaviour of Lipids	148
4.4.3	Functional Group Analysis for Crude ZO, ZZ and NLCs	150
4.4.4	Skin Penetration Study Using Franz Diffusion Cell	154
4.4.4.1	Penetration of NLC-ZO and NLC-ZZ Through Strat-M®	154
4.4.4.2	Penetration of NLC-ZO and NLC-ZZ Through Rat Skin	157
4.4.4.3	Summary	160
4.4.5	Release of Encapsulated ZO and ZZ	160
4.4.6	Storage Stability Study	166
4.4.7	Summary	168
4.5	Efficacy of NLC: <i>In Vitro</i> and <i>In Vivo</i> Study	169
4.5.1	<i>In Vitro</i> 3T3-L1 Cell Lines	170
4.5.1.1	<i>In-Vitro</i> Cytotoxicity Assay	170
4.5.1.2	Lipolysis Assay	175
4.5.2	<i>In Vivo</i> Efficacy Study of NLC	178
4.5.2.1	Skin Deposition Study	179
4.5.2.2	Observation Under the Skin	180
4.5.2.3	Lipolysis Marker Measurement	182
4.5.3	Summary	183
CHAPTER 5	CONCLUSIONS AND RECOMMENDATIONS	185
5.1	Conclusions	185
5.2	Recommendations	187
	REFERENCES	189

LIST OF TABLES

TABLE NO.	TITLE	PAGE
Table 2.1	Summary of plant extracts with anti-obesity activity and their hypothetical mechanism of anti-obesity action	20
Table 2.2	List of topical formulation containing plant extracts/ compounds for anti-obesity application	25
Table 2.3	Summary of various route of administration of drug	42
Table 2.4	Transdermal therapeutic system technologies for controlled release of drug	44
Table 2.5	Various NLC formulations for various applications	54
Table 2.6	Method of NLC preparation	58
Table 2.7	Common surfactant in NLC formulation with respective HLB value	64
Table 2.8	The ranges of infrared (IR) radiation	73
Table 3.1	Experimental design matrixes	88
Table 4.1	Lipid screening for selection of suitable solid lipid for designing NLC-ZO and NLC-ZZ	105
Table 4.2	Liquid lipid selection for designing NLC-ZO and NLC-ZZ after mixing with glyceryl monostearate	106
Table 4.3	NLC-ZO preparation matrix	110

Table 4.4	Estimation of model coefficients for NLC-ZO optimization	110
Table 4.5	NLC-ZO particle size	112
Table 4.6	ANOVA analysis for particle size analysis for NLC-ZO	112
Table 4.7	NLC-ZO PDI	116
Table 4.8	ANOVA analysis for PDI analysis of NLC-ZO	117
Table 4.9	NLC-ZO zeta potential	120
Table 4.10	ANOVA analysis for zeta potential analysis for NLC-ZO	121
Table 4.11	NLC-ZO encapsulation efficiency	124
Table 4.12	ANOVA analysis for encapsulation efficiency analysis for NLC-ZO	125
Table 4.13	The compression of the mixture results and the responses gave by mixture design (n=3)	128
Table 4.14	NLC-ZZ preparation matrix	129
Table 4.15	Estimation of model coefficients for NLC-ZZ optimization	130
Table 4.16	NLC-ZZ particle size	131
Table 4.17	ANOVA analysis for particle size analysis for NLC-ZZ	132
Table 4.18	NLC-ZZ PDI	135
Table 4.19	ANOVA analysis for PDI analysis for NLC-ZZ	135
Table 4.20	NLC-ZZ zeta potential	138

Table 4.21	ANOVA analysis for zeta potential analysis for NLC-ZZ	139
Table 4.22	NLC-ZZ encapsulation efficiency	142
Table 4.23	ANOVA analysis for encapsulation efficiency analysis for NLC-ZZ	143
Table 4.24	The compression of the mixture results and the responses gave by mixture design (n=3) for NLC-ZZ optimization	145
Table 4.25	Strats-M [®] diffusion parameters for NLC-ZO, ZO, NLC-ZZ, and ZZ.	156
Table 4.26	Permeation parameters of ZO and ZO-NLC, ZZ and NLC-ZZ through excised Sprague-Dawley rat skin	159
Table 4.27	Correlation coefficients ($R^2_{adjusted}$) of different models to describe ZO oil and ZZ released from NLC	165
Table 4.28	IC ₅₀ determination using Graphpad Prism 7.0.2 software using nonlinear regression analysis	174

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE
Figure 2.1	Subcutaneous fat distribution in males and females (Sparavigna <i>et al.</i> , 2012).	12
Figure 2.2	Summary of possible intracellular signalling pathways involved in the regulation of lipolysis in adipocytes. (Adapted from Chaves <i>et al.</i> , 2011).	17
Figure 2.3	Chemical structures of gingerols	29
Figure 2.4	Chemical structure of zerumbone	31
Figure 2.5	Beneficial effects of ginger and its active compounds on metabolic responses (Adapted from Wang <i>et al.</i> , 2017).	33
Figure 2.6	Drug permeation route across the skin (Adapted from Herman and Herman., 2015)	37
Figure 2.7	The schematic representation of nanodelivery system over the past 19 th century to today.	41
Figure 2.8	Type of carrier for transdermal delivery system using passive strategies for penetration enhancement	47
Figure 2.9	Published item under 'lipid nanoparticles' from 1995 to 2014 (source from Web of Science)	48
Figure 2.10	The lipid matrix of SLN and NLC	52

Figure 3.1	Flow of study	84
Figure 3.2	Preparation step of NLC using ultrasonication method (Bose <i>et al.</i> , 2013).	86
Figure 3.3	Nile Red (Sigma Aldrich)	87
Figure 3.4	Vertical Franz-style diffusion cell (adapted from Yang <i>et al.</i> , 2017)	93
Figure 3.5	Schematic set up for <i>in vitro</i> release study	96
Figure 4.1	Effect of sonication time on particle size and PDI of NLC. Data presented in mean \pm SD (n=3).	107
Figure 4.2	Effect of sonication power on particle size and PDI of NLC. Data presented in mean \pm SD (n=3).	108
Figure 4.3	Trace plot for NLC-ZO particle size with varying compositions of X_1 , X_2 , and X_3 .	113
Figure 4.4	Contour plot (a) and 3D diagram (b) of changes in particle size of NLC-ZO with varying compositions of X_1 , X_2 , and X_3 .	115
Figure 4.5	Trace plot for NLC-ZO PDI with varying compositions of X_1 , X_2 , and X_3 .	117
Figure 4.6	Contour plot (a) and 3D diagram (b) of changes in PDI of NLC-ZO with varying compositions of X_1 , X_2 , and X_3 .	118
Figure 4.7	Trace plot for NLC-ZO zeta potential	121
Figure 4.8	Contour plot (a) and 3D diagram (b) of changes in zeta potential NLC-ZO with varying compositions of X_1 , X_2 , and X_3 .	123
Figure 4.9	Trace plot for NLC-ZO encapsulation efficiency	125

Figure 4.10	Contour plot (a) and 3D diagram (b) of changes in encapsulation efficiency NLC-ZO with varying compositions of X_1 , X_2 , and X_3 .	126
Figure 4.11	Trace plot for NLC-ZZ particle size	132
Figure 4.12	Contour plots illustrating the effect of (a) ZZ oil concentration, (b) liquid lipid concentration and (c) solid lipid concentration on particle size of NLC-ZZ	133
Figure 4.13	Trace plot for NLC-ZZ PDI	136
Figure 4.14	Contour plots illustrate the effect of (a) ZZ oil concentration, (b) liquid lipid concentration and (c) solid lipid concentration on PDI of NLC-ZZ	137
Figure 4.15	Trace plot for NLC-ZZ zeta potential	139
Figure 4.16	Contour plots illustrate the effect of (a) ZZ oil concentration, (b) liquid lipid concentration and (c) solid lipid concentration on zeta potential of NLC-ZZ	140
Figure 4.17	Trace plot for NLC-ZZ encapsulation efficiency	143
Figure 4.18	Contour plots illustrate the effect of (a) ZZ oil concentration, (b) liquid lipid concentration and (c) solid lipid concentration on encapsulation efficiency of NLC-ZZ	144
Figure 4.19	Transmission electron micrographs of optimized NLC-ZO prepared by ultrasonication technique (magnification of 50 000x and 150 000x).	147
Figure 4.20	Morphology of NLC-ZZ formulation observed under transmission electron (magnification of 150 000x).	147
Figure 4.21	The differential scanning calorimetry thermograms profile of ZO, NLC, NLC-ZO, SLN-ZO and GMS.	148

Figure 4.22	The DSC thermograms profile of ZZ, NLC, NLC-ZZ, SLN-ZZ and GMS.	148
Figure 4.23	Comparison of FTIR peaks between (a) NLC-ZO with (b) ZO oil and (c) NLC	152
Figure 4.24	Comparison of FTIR peaks between (a) NLC-ZZ with (b) ZZ oil and (c) NLC	153
Figure 4.25	Cumulative amount of actives permeated through Strat-M®, comparison between NLC-ZO, ZO, NLC-ZZ, and ZZ. Values are mean \pm SEM (n=3) and are expressed as cumulative amount permeated through Strat-M®.	155
Figure 4.26	Cumulative amount of actives permeated through rat skin. Comparison among NLC-ZO, ZO, NLC-ZZ, and ZZ. Permeation profile of ZO and NLC-ZO. Values are mean \pm SEM (n=3) and are expressed as cumulative amount permeated through rat skin.	158
Figure 4.27	ZO and ZZ release profiles from NLC. Values are mean \pm SD (n=3) and are expressed as percentage of release to medium	162
Figure 4.28	Optimized NLC-ZO formulation stability based on particle size observed for 90 days in storage at room temperature, 10°C, and -2°C. Data presented in mean \pm standard deviation (SD) (n=3).	167
Figure 4.29	Optimized NLC-ZZ formulation stability based on particle size observed for 90 days in storage at room temperature, 10°C, and -2°C. Data presented in mean \pm standard deviation (SD) (n=3).	167
Figure 4.30	Cytotoxic effects of ZO and NLC-ZO at dose concentration on 3T3-L1 preadipocytes through MTT assay. Values are mean \pm SEM (n=3) and are expressed as percentage viability	172

Figure 4.31	Cytotoxic effects of ZZ and NLC-ZZ at dose concentration on 3T3-L1 preadipocytes through MTT assay. Values are mean \pm SEM (n=3) and are expressed as percentage viability	172
Figure 4.32	Cytotoxic effects of ZO/ZZ and NLC-ZOZZ at dose concentration on 3T3-L1 preadipocytes through MTT assay. Values are mean \pm SEM (n=3) and are expressed as percentage viability.	173
Figure 4.33	Lipolytic effect of ZO and NLC-ZO in 3T3-L1 differentiated adipocytes compared with positive control, isoproterenol and IBMX. Values are mean \pm SEM (n=3) represent as glycerol released and significantly different (p<0.05) from control.	176
Figure 4.34	Lipolytic effect of ZZ and NLC-ZZ in 3T3-L1 differentiated adipocytes compared with positive control, isoproterenol and IBMX. Values are mean \pm SEM (n=3) represent as glycerol released and significantly different (p<0.05) from control.	177
Figure 4.35	Lipolytic effect of ZO/ZZ and NLC-ZO/ZZ in 3T3-L1 differentiated adipocytes compared with positive control, isoproterenol and IBMX. Values are mean \pm SEM (n=3) represent as glycerol released and significantly different (p<0.05) from control.	178
Figure 4.36	Amount of NLC deposited in rat skin for 4 h, 8 h and 12 h in vivo. Data presented in mean \pm SD (n = 4)	179
Figure 4.37	Images revealing the penetration and distribution of Nile red within rat skin when treated with; (a) NLC-Nile red and (b) NLC-ethanol for 4 h; (c) NLC-Nile red and (d) NLC-ethanol for 8 h; and (e) NLC-Nile red and (f) NLC-ethanol for 12 h	181
Figure 4.38	Changes in FFA plasma concentration after 3 h treatment with NLC-ZO, NLC-ZZ and water (control) compared to basal level of plasma FFA. Data presented in mean \pm SD (n = 4).	183

LIST OF ABBREVIATIONS

11HSD1	-	11 β -hydroxysteroids dehydrogenase
ABTS, 2,2'-azinobis	-	3-ethylbenzothiazoline-6-sulfonic acid
AC	-	adenyl cyclase
ACC	-	acetyl-CoA carboxylase
ACE	-	Angiotensin-1-converting enzyme
ADD1/SREBP-1c	-	adipocyte determination and differentiation factor 1
ALT	-	alanine aminotransferase
AOX	-	anti-oxidant capacity
aP2	-	fatty acid-binding protein
AST	-	aspartate aminotransferase
ATGL	-	Adipose triglyceride lipase
ATP	-	adenosine and three phosphate
ATR	-	attenuated total reflectance
BMI	-	body mass index
C/EBP α , C/EBP β and C/EBP δ	-	CCAAT/enhancer binding proteins
cAMP	-	Cyclic adenosine monophosphate

CAT	-	catalase
cGMP	-	cyclic guanosine monophosphate
CMT-3	-	chemically modified tetracyclines
DAG	-	diacylglycerol
DEX	-	1 μ M dexamethasone
DI	-	deionized water
DL	-	drug loading
DLS	-	dynamic light scattering
DLX	-	duloxetine
DMEM	-	Dulbecco's modified Eagle's medium
DSC	-	differential scanning methods
EE	-	Encapsulation efficiency
EMA	-	European Medicines Agency
ETH	-	Swiss Federal Institute of Technology
FAOSTAT	-	Food and Agricultural Organizations of the United Nations
FAS	-	fatty-acid synthase
F-C	-	Folin-Ciocalteu
FCS	-	fetal calf serum
FDA	-	Food and Drug Administration
FESEM	-	field-emission scanning electron microscopy
FFA	-	free fatty acids

FoxO1	-	forkhead box class O 1
FTIR	-	Fourier-transform infrared spectroscopy
GA	-	glycyrrhetic acid
GAE	-	gallic acid equivalents
GC	-	guanylyl cyclase
GCI-58	-	comparative gene identification 58
GI	-	gastrointestinal
GMS	-	Glyceryl monostearate
GPx	-	glutathione peroxidase
GSH	-	glutathione
GTP	-	guanosine triphosphate
HIFU	-	high intensity focused ultrasound
HLB	-	hydrophile-lipophile balance
HMG-CoA	-	3-hydroxy-3-methylglutaryl-coenzyme A reductase
HOMA-IR	-	homeostasis model assessment of insulin resistance
HPH	-	High pressure homogenisation
HSL	-	hormone sensitive lipase
IBMX	-	isobutylmethylxanthine
IL-1 β	-	interleukin-1 β
IL-6	-	interleukin-6
IL-8	-	interleukin-8
iNOS	-	inducible nitric oxide

IP3	-	inositol triphosphate
IR	-	Infrared
IR	-	Infrared
IRS-1	-	insulin receptor substrate 1
JAK	-	Janus kinase
JNK	-	c-Jun N-terminal kinase
LCAT	-	lecithin choline acyl transferase
LDL	-	low density lipoprotein
LLLT	-	low-level laser therapy
LPAAT	-	lysophosphatidic acid acyltransferase
LPL	-	lipoprotein lipase
MAG	-	monoacylglycerol
MALS	-	multi-angle light scattering
MaPk	-	mitogen-activated protein kinase
MCP-1	-	monocyte chemoattractant protein-1
MDA	-	malondialdehyde
MGL	-	monoacylglycerol lipase
Mo	-	molybdenum,
MTT	-	3-(4,5-Dimethylthiazol-2-yl, 5-diphenyltetrazolium bromide
MWCO	-	Molecular weight cut-off
NIH	-	National Institutes of Health

NLC	-	Nanostructured lipid carrier
NLC-ZO	-	Nanostructured lipid carrier loaded with <i>Zingiber officinale</i>
NLC-ZZ	-	Nanostructured lipid carrier loaded with <i>Zingiber zerumbet</i>
NMR	-	Nuclear magnetic resonance
NO	-	nitric oxide
o/w	-	oil in water
OAT 2	-	Organic Anion Transporter 2
OECD	-	Organisation for Economic Co-operation and Development
PCS	-	correlation spectroscopy
PDE	-	phosphodiesterase
PDI	-	polydispersity index
PDK	-	phosphoinositide-dependent kinase
PES	-	polyethersulfone
PKA	-	protein kinase A
PKB	-	protein kinase B
PKC	-	Protein kinase C
PKG	-	cGMP-dependent protein kinase
PLC	-	phospholipase C
PLIN	-	perilipin
PPAR- γ	-	peroxisome proliferator-activated receptor
QLS	-	quasi-light scattering
RF	-	radio frequency

SC	-	stratum corneum
SD	-	Standard deviation
SEM	-	scanning electron microscopy
SET	-	single electron transfer
SLN	-	solid lipid nanoparticles
SOD	-	superoxide dismutase
SREBP-1c	-	sterol regulatory element-binding protein-1c
STAT	-	signal transducers and activators of transcription
TAG	-	triacylglycerol
TBARS	-	thiobarbituric acid–reactive substances
TC	-	total cholesterol
TEM	-	transmission electron microscopy
TG	-	triglycerides
TNF- α	-	tumor necrosis factor alpha
VCO	-	Virgin coconut oil
w/o	-	water in oil
WHO	-	World Health Organization
ZO	-	Zingiber officinale
ZZ	-	Zingiber zerumbet

LIST OF SYMBOLS

cm	-	Centimeter
µg/ml	-	Microgram per mililitre
%	-	Percent
°C	-	Degree celcius
mg/ml	-	Miligram per mililitre
µL	-	Microlitre
nm	-	Nanometer
mM	-	Milimolar
µM	-	Micromolar
nmole	-	Nanomole
mm	-	Milimetre
ml	-	Mililitre
rpm	-	Revolutions per minute
g	-	Gram

LIST OF APPENDICES

APPENDIX	TITLE	PAGE
Appendix A	Supplementary Figures	227
Appendix B	Supplementary Tables	230

CHAPTER 1

INTRODUCTION

1.1 Research Background

Obesity has reached epidemic proportions and is a significant contributor to the global burden of chronic disease and disability. Currently, nearly two billion adults worldwide are overweight, and at least 600 million of them are clinically obese (WHO, 2016). Obesity often related to the appearance of excess fat or adiposities which may generate social distress, physiological embarrassment, affecting the quality of life and highlighting possible undetected metabolic disturbance. The modern lifestyle does take its toll on current generations, for instance, long working hours and less physical activities, as well as exposure to imbalanced high caloric diet. The imbalance of diet intake and energy expenditure are generating more obesity-prone individual. The excess of calories is naturally stored within adipocytes.

White adipose tissues provide insulation and serve as useful energy depot. It also plays a vital role in maintaining an excellent endocrine equilibrium and actively contributes to the control of lipid and glucose metabolism. However, excessive adipose tissue distribution in human bodies does affect negatively to appearance and health state of an individual. The adiposities have a number of side effect including skin barrier alterations, sebaceous gland hyperactivity, apocrine and eccrine sweat glands hyperactivity, impaired lymphatic drainage leading to inflammation and fibrosis, altered collagen structure, impaired wound healing, and decreased microvascular homeostatic efficiency (Sparagvina *et al.*, 2012).

To date, there are two types of obesity-treatment drugs available in the market which is orlistat and sibutramine. Both drugs treat obesity through the gastrointestinal route in which they reduce energy intake. However, due to the dissatisfaction of high costs and potentially hazardous side effects of these drugs, numerous researches have reported potential plant extracts which may be an excellent alternative strategy for the anti-obesity drug (Yun, 2010). Alternatively, various non-invasive techniques in reducing localized subcutaneous adipose tissue were reported safe and effective, for example, low-level laser therapy (LLLT), cryolipolysis, radio frequency (RF) and high intensity focused ultrasound (HIFU) (Kennedy *et al.*, 2015). On the other hand, liposuction surgery has been reported as extremely effective in reducing adiposities although the post-procedural pain may cause discomfort (Hunstad *et al.*, 2018). Reports by Greenway *et al.*, (1995) and Sparagvina *et al.*, (2012) have proved a reduction of weight through transdermal administration of drug which promotes lipolysis of fat cell under the skin.

Subcutaneous adipose tissue constitutes the largest fat depot and altered distribution of subcutaneous fat may affect skin elasticity and cosmetically undesirable. Inefficient subcutaneous fat cell lipolysis also independently linked to long-term weight gain and disturbed glucose metabolism (Arner *et al.*, 2018). It is well established that obesity is characterized by increased spontaneous (basal) but decreased hormone stimulated fat cell lipolysis. Major endogenous stimulators of lipolysis are catecholamines, which accelerates lipolysis after binding to beta-adrenergic receptors and stimulates the production of cyclic AMP (cAMP). The cAMP production activates lipases that hydrolyze fat cell triglycerides into glycerol and fatty acids. The energy-rich fatty acids are oxidized for energy consumption in skeletal muscle, heart, and liver. Therefore, it is possible that variation in basal and/or stimulated adipocyte lipolysis may impact weight development. On the other hand, reports by Roure *et al.*, (2011) showed the potential of caffeine and forskolin as active ingredients which stimulated lipolysis through adenylate cyclase activation and inhibit phosphodiesterase. The lipolysis activity was measured from glycerol release after treated with active ingredients.

The treatment of obesity is commonly done together with regulation of diet and regular exercise. There are rising needs to manage obesity with low cost, high efficacy and give very fewer side effects. Oral drug delivery poses multiple challenges such as relatively low uptake and low bioavailability which is due to the absorption of the drug in digestive system and enters the hepatic portal system. As the liver metabolizes the drug, only a small amount of active drug emerges from the liver to the rest of the circulatory system. Therefore, transdermal drug delivery is a preferred candidate due to reduced systemic effect as compared to oral and parenteral drug delivery. However, very scarce data available on anti-obesity treatment strategy which utilizing local plants for transdermal drug delivery.

Zingiber officinale (ginger), ZO and *Zingiber zerumbet* (pinecone ginger), ZZ are common spices used around the world and long been used in traditional medicine as a cure for ailments. A recent study has shown the anti-obesity activity of ZO and its constituents through *in vitro* and *in vivo* studies. Consumption of ginger has significantly reduced lipid profile and reduce body weight of male Albino rats after four weeks of treatment (Mahmoud and Elnour, 2013). Similarly, Misawa *et al.*, (2015) have reported the role of ginger extracts in preventing obesity via activation of peroxisome proliferator-activated receptor, PPAR- γ pathway in C57BL/6J mice treated with dietary ginger extract. In this study, the compound 6-gingerol and 6-shogaol acted as specific PPAR- γ ligands and stimulated PPAR- γ -dependent gene expression in cultured human skeletal muscle myotubes, therefore stimulating fat utilization. In another report, 6-gingerol reduced the expression of PPAR- γ associated genes and reduced adipogenesis in 3T3-L1 adipocytes (Isa *et al.*, 2008). Obese rats treated orally with gingerol for 30 days have shown to reduce body weight and tissue lipid suggested ginger supplementation could suppress obesity induced by high-fat diet (Saravanan *et al.*, 2014). Ethanol extract of ZZ has reported suppressing body weight gain and body fat accumulation by increasing fatty acid oxidation which likely mediated via up-regulation of PPAR- α in the liver of high fed diet male Wistar rats (Chang *et al.*, 2012). The *in vivo* and *in vitro* data does support the potential of ZO and ZZ to induce the anti-obesity effect. Therefore, this study was designed to load ZO and ZZ extracts into a specific carrier to penetrate into the skin in an attempt to reduced localized subcutaneous adipose tissue.

The skin barrier is a major limitation for transdermal drug administration. The barriers generally designed to keep foreign material out and, in some cases, to allow highly regulated transfer of specific desired molecules. Different skin levels such as skin surface, epidermis, dermis, and hypodermis lower the chance for penetration of drug. Therefore, in drug delivery strategies, various types of carrier for transdermal drug delivery have been developed and colloidal vesicle carriers are one of them. The modern development of colloidal carriers started with the discovery of liposome. These vesicles act as drug reservoir and have the ability to control release rate for drug. They also assist the drug to reach the desired pharmacological site of action and have continuous duration of action. The other type of colloidal-based carriers is solid lipid nanoparticles (SLN) and nanostructured lipid particles (NLC). Prof R. Muller discovered SLN in 1995 and the successor of SLN was NLC which developed later in 1990s (Muller *et al.*, 2007). SLN were developed to improve the stability issues arise from liposome and at the same time retain high absorption and biocompatibility properties. NLC is the improvised version of SLN which offers more stable nanoparticle and high performance of drug-controlled release. The lipid nanoparticle, NLC has nano-size range and its formulation is suitable candidate to encapsulate active ingredients and assists in drug delivery to target site. The preparation method of NLC significantly influence the physical properties of NLC. The NLC must possess good loading capacity and long-term stability to ensure no premature release of drug before reaching the target tissue. In another aspect, a smaller size of NLC ensure close contact, specifically skin, and therefore increase the amount of drug penetrated.

1.2 Problem Statement

There are plenty of products on the shelf claimed to have anti-adiposity effect or in a simple word, slimming effect (Stickel, 2007; Phattanawasin *et al.*, 2012). The products contain active ingredients such as essential oil and herbs extracts which potentially have anti-adiposity influence on human body. However, slimming requires reduction in unwanted adipose in subcutaneous tissue under the skin layer. The

efficacy studies of topical application for slimming or anti-adiposity product rarely focus on the delivery mode of active ingredient into the targeted layer. The data only considers the reduction of adipose tissue in subcutaneous layer and theoretically assumed that the active ingredient induced lipolysis in fat layer (Kennedy *et al.*, 2015; Grosskreutz *et al.*, 2017). The mechanism of carrier penetrating the subcutaneous skin layer for lipolysis needs further explanation and scientific proof as many topical slimming products claimed possibility to reduce adiposities (Sparavigna *et al.*, 2012). The delivery of substance into subcutaneous layer of skin itself is challenging enough (Anissimov *et al.*, 2013). The skin is an ultimate barrier against external mechanical, chemical, microbial and physical effects. Nevertheless, the large surface area and easy accessibility is the main reason for potential application in topical drug delivery. In order for a substance to reach into subcutaneous layer of skin, suitable carrier must be developed with desirable properties specifically to enhance deeper skin delivery.

NLC has been established since 2000 in encapsulation of active ingredients for various applications (Pardeike *et al.*, 2009). The nano-scale size, stability, and high bioavailability as drug delivery system have made NLC are among preferred carrier especially in carrying lipophilic/hydrophobic type of active ingredient (O'Driscoll and Griffin, 2008; Das *et al.*, 2012; Khan *et al.*, 2013; Ranpise *et al.*, 2014). Common size for nanoparticle for transdermal delivery system is in the range of 50-500 nm. Penetration through the skin depends on the particle size of nanoparticles, in which larger nanoparticles (>300 nm) will stay on stratum corneum, while smaller nanoparticles (<300 nm) will penetrate epidermis and dermis layer of the skin (Iannuccelli *et al.*, 2014). However, smaller particle size (20 nm) possessed toxicity to the skin compared to larger particle size (Park *et al.*, 2013). Additionally, it may cause interaction of nanoparticle with cells, body fluids, and protein. The interaction causes biological changes and ability to distribute throughout the body, hence generate complexes that are more mobile and can enter tissue sites that normally inaccessible (Nel *et al.*, 2006). For instance, smaller particle allows penetration up to the dermis layer and causes particle trafficking to regional lymph nodes. Such trafficking can deliver particles to paracortical areas in the lymph nodes where macrophage and dendritic cells specialized in the uptake of particulate matter. This could lead to the effects on the immune system. Although the reticuloendothelial system is able to clear

or sequester nanoparticles, self-protein interactions with particles may change their antigenicity and initiate autoimmune responses. Therefore, the NLC must be able to penetrate into skin by having size of less than 300 nm and must not impose toxicity towards cells and tissue.

In the search for novel anti-obesity strategies via transdermal penetration for ZO and ZZ, the present study was designed to establish the formulation of NLC with better skin targeting efficiency, high stability, prolonged release to achieve maximum efficacy. The penetration of NLC encapsulated ZO and ZZ into subcutaneous layer was further enlightened through several series of experiments mimicking real skin condition.

1.3 Objective of the Research

The objective of this research is to formulate *Zingiber officinale* and *Zingiber zerumbet* oil loaded NLC system which having desirable physical characteristics and good stability to penetrate subcutaneous layer of skin hence activate lipolysis in adipose tissue.

1.4 Scopes of the Research

In order to achieve the objective, studies have been narrowed down and identified in this research. The scopes of this research listed as below:

- 1) Development of an optimized formulation of NLC encapsulated ZO and ZZ using hot homogenization and ultrasonication.
- 2) Characterization of NLC encapsulated ZO and ZZ by morphology, crystalline behavior, functional group, and skin penetration analysis, and storage stability study. Particle characteristics were measured including particle size, zeta potential, and polydispersity index analysis.
- 3) Determination of kinetic mechanism of active ingredients release from NLC using dialysis method.
- 4) Investigation of the efficacy of NLC encapsulated ZO and ZZ into subcutaneous layer of skin to induce lipolysis in adipose tissue, utilizing *in vitro* and *in vivo* analysis. *In vitro* study was conducted using 3T3-L1 cells to measure lipolysis activity. *In vivo* studies included *in vivo* skin deposition and penetration study as well as lipolysis marker measurement after treatment with NLC.

1.5 Significance and Original Contributions of the Research

This investigation offers several contributions in the area of essential oil encapsulation using NLC and its application in transdermal drug delivery which focused on subcutaneous skin layer. The contributions are as follows:

- 1) Transdermal delivery of active ingredients can circumvent first pass metabolism and maintains the concentration of drug at site of action for longer periods. Therefore, an investigation on permeation and release of active ingredients onto the targeted side may give useful insight into the understanding of NLC's mechanism of action *in vitro* and *in vivo*. *In vitro* study

focuses on the extent of active ingredient diffusion across the skin and the ability of lipolytic activity induction on adipocyte cell. The release mechanism of active ingredient from nanoparticle is identified according to kinetic release profile in the designated experimental condition. *In vivo* study highlights the extent of NLC penetration across rat skin in different treatment duration and lipolysis activity detected in plasma level.

- 2) ZO and ZZ belonged to Zingiberaceae family. It has a wide spectrum of traditional uses, as well as biological and pharmacological properties. New emerging *in vitro* data suggest both ZO and ZZ have demonstrated potential anti-obesity activity through various mechanisms *in vitro* and *in vivo*. NLC has been a good candidate for encapsulation ZO and ZZ due to the lipophilicity properties and capability to permeate challenging physiological barriers such as human and animal skins.

1.6 Thesis Structure and Organization

This thesis is divided into five chapters. Chapter 1 covers a brief overview of the research background, problems statement, a central objective, scopes of analyses, originality and significant contributions of the study.

Chapter 2 offer an overview of obesity and topical formulation for fat reduction. The literature also highlights the current drug delivery technology available for transdermal administration as well as current methodologies for nanoparticles characterization.

Chapter 3 covers the overall methodologies used for optimization of NLC encapsulated *Zingiber officinale* and *Zingiber zerumbet*, the physical characterization

of NLC and *in vitro* release study of active ingredients. The methodologies of *in vitro* and *in vivo* penetration of NLC through skin were described in this chapter.

Chapter 4 presents the comprehensive results and discussions on the optimization of NLC formulation using D-optimal mixture design and the characterization including particle size measurement, zeta potential, morphology, thermal properties, functional group analysis, *in vitro* permeation and kinetic of active ingredient release from NLC. The *in vitro* and *in vivo* analysis were also presented.

Chapter 5 provides the overall summary of the research findings and specific future recommendations for the upcoming works.

REFERENCES

- Abdul Rahman, H., Saari, N., Abas, F., Ismail, A., Mumtaz, M.W., & Abdul Hamid, A. (2017). Anti-obesity and antioxidant activities of selected medicinal plants and phytochemical profiling of bioactive compounds. *International Journal of Food Properties*, 20(11), 2616-2629.
- Acheson, K. J., Gremaud, G., Meirim, I., Montigon, F., Krebs, Y., Fay, L. B., & Tappy, L. (2004). Metabolic effects of caffeine in humans: lipid oxidation or futile cycling?. *The American Journal of Clinical Nutrition*, 79(1), 40-46.
- Ado, M.A., Abas, F., Mohammed, A.S., & Ghazali, H.M. (2013). Anti-and pro-lipase activity of selected medicinal, herbal and aquatic plants, and structure elucidation of an anti-lipase compound. *Molecules*, 18(12), 14651-14669.
- Adorjan, B., & Buchbauer, G. (2010). Biological properties of essential oils: an updated review. *Flavour and Fragrance Journal*, 25(6), 407-426.
- Ahmadian, M., Duncan, R.E., Varady, K.A., Frasson, D., Hellerstein, M.K., Birkenfeld, A.L., Samuel, V.T., Shulman, G.I., Wang, Y., Kang, C., & Sul, H.S. (2009). Adipose overexpression of desnutrin promotes fatty acid use and attenuates diet-induced obesity. *Diabetes*, 58(4), 855-866.
- Ahmadian, M., Wang, Y., & Sul, H.S. (2010). Lipolysis in adipocytes. *The International Journal of Biochemistry & Cell Biology*, 42(5), 555-559.
- Ahmed, T.A., & Khalid, M. (2014). Development of alginate-reinforced chitosan nanoparticles utilizing W/O nanoemulsification/internal crosslinking technique for transdermal delivery of rabeprazole. *Life sciences*, 110(1), 35-43.
- Aisha, A.F., Majid, A.M.S.A., & Ismail, Z. (2014). Preparation and characterization of nano liposomes of Orthosiphon stamineus ethanolic extract in soybean phospholipids. *BMC Biotechnology*, 14(23), 1-11.
- Akalin, O., Akay, K.U., Sennaroglu, B., & Tez, M. (2010). Optimization of chemical admixture for concrete on mortar performance tests using mixture experiments. *Chemometrics and Intelligent Laboratory Systems*, 104, 233-242.

- Alexander, A., Dwivedi, S., Giri, T.K., Saraf, S., Saraf, S., & Tripathi, D.K. (2012). Approaches for breaking the barriers of drug permeation through transdermal drug delivery. *Journal of Controlled Release*, 164(1), 26-40.
- Al-Haj, N.A., Shamsudin, M.N., Alipiah, N.M., Zamri, H.F., Ahmad, B., Siddig, I., & Rasedee, A. (2010). Characterization of *Nigella sativa* L. essential oil-loaded solid lipid nanoparticles. *American Journal of Pharmacology and Toxicology*, 5(1), 52-57.
- Ali, B. H., Blunden, G., Tanira, M. O., & Nemmar, A. (2008). Some phytochemical, pharmacological and toxicological properties of ginger (*Zingiber officinale* Roscoe): a review of recent research. *Food and Chemical Toxicology*, 46, 409-402.
- Allaf, T., & Allaf, K. (2016). Instant controlled pressure drop (DIC) in food processing. Springer-Verlag New York.
- Allen, T. M., & Cullis, P. R. (2013). Liposomal drug delivery systems: from concept to clinical applications. *Advanced Drug Delivery Reviews*, 65(1), 36-48.
- Anandhakumar, S., Gokul, P., & Raichur, A. M. (2016). Stimuli-responsive weak polyelectrolyte multilayer films: A thin film platform for self triggered multi-drug delivery. *Materials Science and Engineering: C*, 58, 622-628.
- Andrade, J.M., Faustino, C., Garcia, C., Ladeiras, D., Reis, C.P., & Rijo, P. (2018). *Rosmarinus officinalis* L.: An update review of its phytochemistry and biological activity. *Future Science OA*, 4(4), FSO283.
- Ansari, M. N., Bhandari, U., & Pillai K. K. (2006). Ethanolic *Zingiber officinale* R. extract pretreatment alleviates isoproterenol-induced oxidative myocardial necrosis in rats. *Indian Journal of Experimental Biology*, 44(11), 892-897.
- Aqil, M., Ahad, A., Sultana, Y., & Ali, A. (2007). Status of terpenes as skin penetration enhancers. *Drug Discovery Today*, 12(23-24), 1061-1067.
- Araújo, J., Gonzalez-Mira, E., Egea, M. A., Garcia, M. L., & Souto, E. B. (2010). Optimization and physicochemical characterization of a triamcinolone acetone-loaded NLC for ocular antiangiogenic applications. *International Journal of Pharmaceutics*, 393(1-2), 168-176.
- Armanini, D., Nacamulli, D., Francini-Pesenti, F., Battagin, G., Ragazzi, E., & Fiore, C. (2005). Glycyrrhetic acid, the active principle of licorice, can reduce the thickness of subcutaneous thigh fat through topical application. *Steroids*, 70(8), 538-542.

- Arner, P., Andersson, D. P., Bäckdahl, J., Dahlman, I., & Rydén, M. (2018). Weight gain and impaired glucose metabolism in women are predicted by inefficient subcutaneous fat cell lipolysis. *Cell metabolism*, 28(1), 45-54.
- Arranz, E., Jaime, L., de las Hazas, M. L., Reglero, G., & Santoyo, S. (2015). Supercritical fluid extraction as an alternative process to obtain essential oils with anti-inflammatory properties from marjoram and sweet basil. *Industrial Crops and Products*, 67, 121-129.
- Attari, V. E., Ostadrahimi, A., Jafarabadi, M. A., Mehralizadeh, S., & Mahluji, S. (2016). Changes of serum adipocytokines and body weight following Zingiber officinale supplementation in obese women: a RCT. *European Journal of Nutrition*, 55(6), 2129-2136.
- Aulton, M.E. (Ed.). (1995). *Pharmaceutics: The Science of Dosage Form Design* (pp. 290–292). Churchill Livingstone, New York.
- Azagury, A., Khoury, L., Enden, G., & Kost, J. (2014). Ultrasound mediated transdermal drug delivery. *Advanced Drug Delivery Reviews*, 72, 127-143.
- Baharvand-Ahmadi, B., Bahmani, M., Tajeddini, P., Naghdi, N., & Rafieian-Kopaei, M. (2016). An ethno-medicinal study of medicinal plants used for the treatment of diabetes. *Journal of Nephropathology*, 5(1), 44-50.
- Bajerski, L., Michels, L.R., Colomé, L.M., Bender, E.A., Freddo, R.J., Bruxel, F., & Haas, S.E. (2016). The use of Brazilian vegetable oils in nanoemulsions: an update on preparation and biological applications. *Brazilian Journal of Pharmaceutical Sciences*, 52(3), 347-363.
- Baker R.W., & Lonsdale H.S. (1974). *Controlled release of biologically active agents*, Tanquary A.C., Lacey R.E. Eds., Plenum Press, New York.
- Bakkali, F., Averbeck, S., Averbeck, D., & Idaomar, M. (2008). Biological effects of essential oils—a review. *Food and Chemical Toxicology*, 46(2), 446-475.
- Banerjee, S., Chattopadhyay, P., Ghosh, A., Datta, P., & Veer, V. (2014). Aspect of adhesives in transdermal drug delivery systems. *International Journal of Adhesion and Adhesives*, 50, 70-84.
- Bangham, A.D., Standish, M.M., & Watkins, J.C. (1965). Diffusion of univalent ions across the lamellae of swollen phospholipids. *Journal of Molecular Biology*, 13(1), 238-IN27.

- Belfrage, P., Fredrikson, G., Nilsson, N.O., & Strålfors, P. (1981). Regulation of adipose-tissue lipolysis by phosphorylation of hormone-sensitive lipase. *International Journal of Obesity*, 5(6), 635-641.
- Bellas, E., Seiberg, M., Garlick, J., & Kaplan, D. L. (2012). *In vitro* 3D full-thickness skin-equivalent tissue model using silk and collagen biomaterials. *Macromolecular Bioscience*, 12(12), 1627-1636.
- Berardesca, E., Abril, E., Rona, C., Vesnaver, R., Cenni, A. & Oliva, M. (2012). An effective night slimming topical treatment. *International Journal of Cosmetic Science*, 34(3), 263-272.
- Berg, L. R. (1997). *Introductory Botany: Plants, People and the Environment*, Ft. Worth: Saunders College Publication.
- Beyki, M., Zhavah, Z., Khalili, S.D., Rahmani-Cherati, T., Abollahi, A., Bayat, M., Tabatabaei, M., & Mohsenifar, A. (2014). Encapsulation of Mentha piperita essential oils in chitosan–cinnamic acid nanogel with enhanced antimicrobial activity against Aspergillus flavus, *Industrial Crops and Products*. 54, 310-319.
- Bilia, A. R., Guccione, C., Isacchi, B., Righeschi, C., Firenzuoli, F., & Bergonzi, M. C. (2014). Essential oils loaded in nanosystems: a developing strategy for a successful therapeutic approach. *Evidence-Based Complementary and Alternative Medicine*, 2014.
- Binks, B. P., & Horozov, T. S. (Eds.). (2006). Colloidal particles at liquid interfaces. Cambridge University Press.
- Boakye, C. H., Patel, K., & Singh, M. (2015). Doxorubicin liposomes as an investigative model to study the skin permeation of nanocarriers. *International Journal of Pharmaceutics*, 489(1), 106-116.
- Bolzinger, M.A., Briançon, S., Pelletier, J., & Chevalier, Y. (2012). Penetration of drugs through skin, a complex rate-controlling membrane. *Current Opinion in Colloid & Interface Science*, 17(3), 156-165.
- Boon, N., Goossens, G. H., Blaak, E. E., & Saris, W. H. (2007). The effects of hydralazine on lipolysis in subcutaneous adipose tissue in humans. *Metabolism*, 56(12), 1742-1748.

- Boqué, N., Campión, J., de la Iglesia, R., de la Garza, A.L., Milagro, F.I., San Román, B., Bañuelos, Ó., & Martínez, J.A. (2013). Screening of polyphenolic plant extracts for anti-obesity properties in Wistar rats. *Journal of the Science of Food and Agriculture*, 93(5), 1226-1232.
- Boqué, N., de la Iglesia, R., de la Garza, A. L., Milagro, F. I., Olivares, M., Bañuelos, Ó., & Campión, J. (2013). Prevention of diet-induced obesity by apple polyphenols in Wistar rats through regulation of adipocyte gene expression and DNA methylation patterns. *Molecular Nutrition & Food Research*, 57(8), 1473-1478.
- Bos, J. D., & Meinardi, M. M. (2000). The 500 Dalton rule for the skin penetration of chemical compounds and drugs. *Experimental Dermatology*, 9(3), 165-169.
- Bose, S., & Michniak-Kohn, B. (2012). Preparation and characterization of lipid based nanosystems for topical delivery of quercetin. *European Journal of Pharmaceutical Sciences*, 48(3), 442-452.
- Bose, S., Du, Y., Takhistov, P., & Michniak-Kohn, B. (2013). Formulation optimization and topical delivery of quercetin from solid lipid based nanosystems. *International Journal of Pharmaceutics*, 441(1-2), 56-66.
- Bray, G. A., Kim, K. K., Wilding, J. P. H., & World Obesity Federation. (2017). Obesity: a chronic relapsing progressive disease process. A position statement of the World Obesity Federation. *Obesity Reviews*, 18(7), 715-723.
- Braydich-Stolle, L., Hussain, S., Schlager, J. J., & Hofmann, M. C. (2005). In vitro cytotoxicity of nanoparticles in mammalian germline stem cells. *Toxicological Sciences*, 88(2), 412-419.
- Bujalska, I. J., Gathercole, L. L., Tomlinson, J. W., Darimont, C., Ermolieff, J., Fanjul, A. N., Rejto, P. A., & Stewart, P. M. (2008). A novel selective 11 β -hydroxysteroid dehydrogenase type 1 inhibitor prevents human adipogenesis. *The Journal of Endocrinology*, 197(2), 297-307.
- Bunjes, H., & Unruh, T. (2007). Characterization of Lipid Nanoparticles by Differential Scanning Calorimetry, X-Ray and Neutron Scattering. *Advanced Drug Delivery Reviews*, 59(6), 379-402.
- Bunjes, H., Westesen, K., & Koch, M. H. (1996). Crystallization tendency and polymorphic transitions in triglyceride nanoparticles. *International Journal of Pharmaceutics*, 129(1-2), 159-173.

- Campos, P. P., Fraceto, L. F., & Ferreira, M. (2018). Layer-by-layer films containing emodin or emodin encapsulated in liposomes for transdermal applications. *Colloids and Surfaces B: Biointerfaces*, 162, 69-75.
- Carrasco, F. R., Schmidt, G., Romero, A. L., Sartoretto, J. L., Caparroz-Assef, S. M., Bersani-Amado, C. A., & Cuman, R. K. N. (2009). Immunomodulatory activity of *Zingiber officinale* Roscoe, *Salvia officinalis* L. and *Syzygium aromaticum* L. essential oils: evidence for humor-and cell-mediated responses. *Journal of Pharmacy and Pharmacology*, 61(7), 961-967.
- Caruso, M. K., Pekarovic, S., Raum, W. J., & Greenway, F. (2007). Topical fat reduction from the waist. *Diabetes, Obesity and Metabolism*, 9(3), 300-303.
- Chaaban, A., Santos, V. M. C. S., Gomes, E. N., Martins, C. E. N., Amaral, W. D., Deschamps, C., & Molento, M. B. (2018). Chemical composition of *Piper gaudichaudianum* essential oil and its bioactivity against *Lucilia cuprina* (Diptera: Calliphoridae). *Journal of Essential Oil Research*, 30(3), 159-166.
- Chan, E. S., Yim, Z. H., Phan, S. H., Mansa, R. F., & Ravindra, P. (2010). Encapsulation of herbal aqueous extract through absorption with ca-alginate hydrogel beads. *Food and Bioproducts Processing*, 88(2-3), 195-201.
- Chandan, C., & Maheshwari, R. K. (2014). Mixed solvency concept in reducing surfactant concentration of self-emulsifying drug delivery systems of candesartan cilexetil using D-optimal mixture design. *Asian Journal of Pharmaceutics*, 7(2), 83-91.
- Chang, C. J., Tzeng, T. F., Liou, S. S., Chang, Y. S., & Liu, I. M. (2012). Regulation of lipid disorders by ethanol extracts from *Zingiber zerumbet* in high-fat diet-induced rats. *Food chemistry*, 132(1), 460-467.
- Changez, M., Varshney, M., Chander, J., & Dinda, A. K. (2006). Effect of the composition of lecithin/n-propanol/isopropyl myristate/water microemulsions on barrier properties of mice skin for transdermal permeation of tetracaine hydrochloride: in vitro. *Colloids and Surfaces B: Biointerfaces*, 50(1), 18-25.
- Changhyun R., & Uhee J. (2012). Screening of crude plant extracts with anti-obesity activity. *International Journal of Molecular Sciences*, 13, 1710-1719.
- Chaput, J. P., Després, J. P., Bouchard, C., & Tremblay, A. (2007). Short sleep duration is associated with reduced leptin levels and increased adiposity: results from the Quebec family study. *Obesity*, 15(1), 253-261.

- Charcosset, C., & Fessi, H. (2005). Preparation of nanoparticles with a membrane contactor. *Journal of Membrane Science*, 266(1-2), 115-120.
- Chaves, V. E., Frasson, D., & Kawashita, N. H. (2011). Several agents and pathways regulate lipolysis in adipocytes. *Biochimie*, 93(10), 1631-1640.
- Chen, F., Shi, Z., Neoh, K. G., & Kang, E. T. (2009). Antioxidant and antibacterial activities of eugenol and carvacrol-grafted chitosan nanoparticles. *Biotechnology and Bioengineering*, 104(1), 30-39.
- Chen, M., Liu, X., & Fahr, A. (2011). Skin penetration and deposition of carboxyfluorescein and temoporfin from different lipid vesicular systems: in vitro study with finite and infinite dosage application. *International Journal of Pharmaceutics*, 408(1-2), 223-234.
- Cirri, M., Bragagni, M., Mennini, N., & Mura, P. (2012). Development of a new delivery system consisting in "drug--in cyclodextrin--in nanostructured lipid carriers" for ketoprofen topical delivery. *European Journal of Pharmaceutics and Biopharmaceutics*, 80(1), 46-53.
- Cosmetic Surgery National Data Bank Statistics: American Society for Aesthetic Plastic Surgery; 2014. URL <http://www.surgery.org/media/statistics> (last accessed: 21 July 2014).
- Craft, B. D., Kerrihard, A. L., Amarowicz, R., & Pegg, R. B. (2012). Phenol-based antioxidants and the *in vitro* methods used for their assessment. *Comprehensive Reviews in Food Science and Food Safety*, 11(2), 148-173.
- Curri, S. B. (1993). Cellulite and fatty tissue microcirculation. *Cosmetic and Toiletries*, 108, 51-58.
- da Silva, J. A., Becceneri, A. B., Mutti, H. S., Martin, A. C. B. M., Fernandes, J. B., Vieira, P. C., & Cominetti, M. R. (2012). Purification and differential biological effects of ginger-derived substances on normal and tumor cell lines. *Journal of Chromatography B*, 903, 157-162.
- Dagon, Y., Avraham, Y., & Berry, E. M. (2006). AMPK activation regulates apoptosis, adipogenesis, and lipolysis by eIF2 α in adipocytes. *Biochemical and Biophysical Research Communications*, 340(1), 43-47.
- Damodaran, N. P., & Dev, S. (1968). Studies in sesquiterpenes—XXXVII: Sesquiterpenoids from the essential oil of *Zingiber zerumbet* smith. *Tetrahedron*, 24(11), 4113-4122.

- Dan, N. (2014). Nanostructured Lipid Carriers: Effect of Solid Phase Fraction and Distribution on the Release of Encapsulated Materials. *Langmuir*, 30(46), 13809-13814.
- Danaei, M., Dehghankhold, M., Ataei, S., Hasanzadeh Davarani, F., Javanmard, R., Dokhani, A., Khorasani, S., & Mozafari, M. R. (2018). Impact of particle size and polydispersity index on the clinical applications of lipidic nanocarrier systems. *Pharmaceutics*, 10(2), 57.
- Das, S., & Chaudhury, A. (2011). Recent advances in lipid nanoparticle formulations with solid matrix for oral drug delivery. *AAPS Pharmscitech*, 12(1), 62-76.
- Das, S., Ng, W. K., & Tan, R. B. (2012). Are nanostructured lipid carriers (NLCs) better than solid lipid nanoparticles (SLNs): development, characterizations and comparative evaluations of clotrimazole-loaded SLNs and NLCs?" *European Journal of Pharmaceutical Sciences*, 47(1), 139-151.
- Dash, S., Murthy, P. N., Nath, L., & Chowdhury, P. (2010). Kinetic modeling on drug release from controlled drug delivery systems. *Acta Poloniae Pharmaceutica*, 67(3), 217-23.
- Date, A. A., Desai, N., Dixit, R., & Nagarsenker, M. (2010). Self-nanoemulsifying drug delivery systems: formulation insights, applications and advances. *Nanomedicine*, 5(10), 1595-1616.
- de Barros Fernandes, R.V., Botrel, D.A., Silva, E.K., Borges, S.V., de Oliveira, C.R., Yoshida, M.I., de Andrade Feitosa, J.P., & de Paula, R.C.M. (2016). Cashew gum and inulin: New alternative for ginger essential oil microencapsulation. *Carbohydrate Polymers*, 153, 133-142.
- Deepak Prashar, S. S., Yogesh, B., Mohit S., Sushant, M., & Soni, A. (2012). Pharma-Economical World of Herbal Antitussive- An Overview. *Asian Journal of Research in Pharmaceutical Sciences*, 2(2), 48-51.
- Derjaguin, B. V., & Landau, L. D. (1941). *Acta Physicochim URSS*, 14, 733.
- Desai, P. P., Date, A. A., & Patravale, V. B. (2012). Overcoming poor oral bioavailability using nanoparticle formulations—opportunities and limitations. *Drug Discovery Today: Technologies*, 9(2), e87-e95.
- Despina H. (2012). *Antiobesity and antidiabetic activity of P. balsamifera, its active Salicortin, and L. laricina, medicinal plants from the traditional pharmacopoeia of the James Bay Cree*. PhD Theses. Université de Montréal. Canada.

- Dima, C., Cotarlet, M., Alexe, P., & Dima S. (2014). Reprint of “Microencapsulation of essential oil of pimento [*Pimenta dioica* (L) Merr.] by chitosan/k-carrageenan complex coacervation method”, *Innovative Food Science & Emerging Technologies*, 25, 97-105.
- Djuris, J., Vasiljevic, D., Jokic, S., & Ibric, S. (2014). Application of D-optimal experimental design method to optimize the formulation of O/W cosmetic emulsions, *International Journal of Cosmetic Science*, 36, 79-87.
- Doktorovová, S., Araújo, J., Garcia, M. L., Rakovský, E., & Souto, E. B. (2010). Formulating fluticasone propionate in novel PEG-containing nanostructured lipid carriers (PEG-NLC). *Colloids and Surfaces B: Biointerfaces*, 75(2), 538-542.
- Doktorovova, S., Souto, E. B., & Silva, A. M. (2014). Nanotoxicology applied to solid lipid nanoparticles and nanostructured lipid carriers—a systematic review of *in vitro* data. *European Journal of Pharmaceutics and Biopharmaceutics*, 87(1), 1-18.
- Dolhaji, N. H. (2010). *Effect of Ginger (zingiber officinale) Extracts on Triglyceride Content in 3T3-L1 Adipocytes*. Proceedings of the Regional Seminar on Science, Technology and Social Sciences 2010.
- Draelos, Z. D., & Marenus K. D. (1997). Cellulite. Etiology and purported treatment. *Dermatology Surgical*, 23(12), 1177-1181.
- Draize, J. H., Woodard, G., & Galvery, H. O. (1944). Methods for the study of irritation and toxicity of substances applied topically to the skin and mucous membranes, *Journal of Pharmacology and Experimental Therapeutics*, 82, 377-390.
- Dubey, A., Prabhu, P., & Kamath, J.V. (2012). Nano Structured lipid carriers a novel topical drug delivery system, *International Journal of PharmTech Research*, 4(2), 705-714.
- Dubey, B. K., Katare, O. P., Singh, R., & Jain, S. K. (1995). Lyophilized aqueous based polymer matrices for transdermal delivery of captopril. *Journal of Dermatological Science*, 10(3), 191-195.
- Ebrahimzadeh Attari, V., Malek Mahdavi, A., Javadivala, Z., Mahluji, S., Zununi Vahed, S., & Ostadrahimi, A. (2018). A systematic review of the anti-obesity and weight lowering effect of ginger (*Zingiber officinale* Roscoe) and its mechanisms of action. *Phytotherapy Research*, 32(4), 577-585.

- Edris, A. E., Wawrzyniak, P., & Kalembe, D. (2018). Subcritical CO₂ extraction of a volatile oil-rich fraction from the seeds of *Nigella sativa* for potential pharmaceutical and nutraceutical applications. *Journal of Essential Oil Research*, 30(2), 84-91.
- El Asbahani, A., Miladi, K., Badri, W., Sala, M., Addi, E. A., Casabianca, H. & Elaissari, A. (2015). Essential oils: From extraction to encapsulation. *International Journal of Pharmaceutics*, 483(1), 220-243.
- Eleazu, C. O., Iwo, G. A. C., Nwosu, P., & Ironua, C. F. (2013). Chemical composition and free radical scavenging activities of 10 elite accessions of ginger (*Zingiber officinale* Roscoe). *Journal of Nutrition and Food Sciences*, 3(2), 1-5.
- El-Menshaweh, S. F., Ali, A. A., Rabeh, M. A., & Khalil, N. M. (2018). Nanosized soy phytosome-based thermogel as topical anti-obesity formulation: an approach for acceptable level of evidence of an effective novel herbal weight loss product. *International Journal of Nanomedicine*, 13, 307-318.
- Elnaggar, Y. S., El-Refaie, W. M., El-Massik, M. A., & Abdallah, O. Y. (2014). Lecithin-based nanostructured gels for skin delivery: an update on state of art and recent applications. *Journal of Controlled Release*, 180, 10-24.
- Emami, J., Rezazadeh, M., Varshosaz, J., Tabbakhian, M., & Aslani, A. 2012. Formulation of LDL targeted nanostructured lipid carriers loaded with paclitaxel: a detailed study of preparation, freeze drying condition, and *In vitro* cytotoxicity. *Journal of Nanomaterials*, 2012, 1-10.
- Escudier, B., Fanchon, C., Labrousse, E., & Pellae, M. (2011). Benefit of a topical slimming cream in conjunction with dietary advice. *International Journal of Cosmetic Science*, 33(4), 334-337.
- Everette, J. D., Bryant, Q. M., Green, A. M., Abbey, Y. A., Wangila, G. W., & Walker, R. B. (2010). Thorough study of reactivity of various compound classes toward the Folin– Ciocalteu reagent. *Journal of Agricultural and Food Chemistry*, 58(14), 8139-8144.
- Fabroni, S., Ballistreri, G., Amenta, M., Romeo, F.V., & Rapisarda, P. (2016). Screening of the anthocyanin profile and *in vitro* pancreatic lipase inhibition by anthocyanin-containing extracts of fruits, vegetables, legumes and cereals. *Journal of the Science of Food and Agriculture*, 96(14), 4713-4723.
- Farooqi, I. S., & O’Rahilly, S. (2007). Genetic factors in human obesity. *Obesity Reviews*, 8, 37-40.

- Fathi, M., & Varshosaz, J. (2013). Novel hesperetin loaded nanocarriers for food fortification: Production and characterization. *Journal of Functional Foods*, 5(3), 1382-1391.
- Fiel, L. A., Paese, K., Rizzi, M., Guterres, S. S., & Pohlmann, A. R. (2014). A strategy to estimate the intrinsic flux of a poorly water soluble substance for comparison with its release from lipid-core nanocapsules. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 441, 716-724.
- Flynn, G. L., Yalkowsky, S. H., Roseman, T. J. (1974). Mass transport phenomena and models: theoretical concepts. *Journal of Pharmaceutical Sciences*, 63(4), 479-510.
- Fox, L. T., Gerber, M., Plessis, J. D., & Hamman, J. H. (2011). Transdermal drug delivery enhancement by compounds of natural origin. *Molecules*, 16(12), 10507-10540.
- Franz, C. M. (2010). Essential oil research: past, present and future. *Flavour and Fragrance Journal*, 25(3), 112-113.
- Freitas, C., & Müller, R. H. (1998). Effect of light and temperature on zeta potential and physical stability in solid lipid nanoparticle (SLNTM) dispersions. *International Journal of Pharmaceutics*, 168(2), 221-229.
- Fry, D. W., White, J. C., & Goldman, I. D. (1978). Rapid separation of low molecular weight solutes from liposomes without dilution. *Analytical Biochemistry*, 90(2), 809-815.
- Gasco, M. R. (1993). *U.S. Patent No. 5,250,236*. Washington, DC: U.S. Patent and Trademark Office.
- Gaumet, M., Vargas, A., Gurny, R., & Delie, F. (2008). Nanoparticles for drug delivery: the need for precision in reporting particle size parameters. *European Journal of Pharmaceutics and Biopharmaceutics*, 69(1), 1-9.
- Ghate, V. M., Lewis, S. A., Prabhu, P., Dubey, A., & Patel, N. (2016). Nanostructured lipid carriers for the topical delivery of tretinoin. *European Journal of Pharmaceutics and Biopharmaceutics*, 108, 253-261.
- Gomes, M. J., Martins, S., Ferreira, D., Segundo, M. A., & Reis, S. (2014). Lipid nanoparticles for topical and transdermal application for alopecia treatment: development, physicochemical characterization, and in vitro release and penetration studies. *International Journal of Nanomedicine*, 9, 1231.

- Greenberg, A. S., Shen, W. J., Muliro, K., Patel, S., Souza, S. C., Roth, R. A., & Kraemer, F. B. (2001). Stimulation of lipolysis and hormone-sensitive lipase via the extracellular signal-regulated kinase pathway. *Journal of Biological Chemistry*, 276(48), 45456-45461.
- Greenway, F. L., Bray, G. A., & Heber, D. (1995). Topical fat reduction. *Obesity Research*, 3, 561-568.
- Griffin, W.C. (1954). Calculation of HLB values of non-ionic surfactants. *Journal of the Society of Cosmetic Chemists*, 5, 249-256.
- Grosskreutz, C. L., Pasquale, L. R., Singer, M. S., & Kalayoglu, M. V. (2017). *U.S. Patent No. 9,795,614*. Washington, DC: U.S. Patent and Trademark Office.
- Gursoy, R. N., & Benita, S. (2004). Self-emulsifying drug delivery systems (SEDDS) for improved oral delivery of lipophilic drugs. *Biomedicine & Pharmacotherapy*, 58(3), 173-182.
- Guy, R. H. (2010). Transdermal drug delivery. In *Drug delivery* (pp. 399-410). Springer, Berlin, Heidelberg.
- Hadgraft J., & Guy R. H. (1989). *Transdermal Drug Delivery*. New York, Marcel Dekker.
- Hadgraft, J. (2001). Modulation of the barrier function of the skin. *Skin Pharmacology and Applied Skin Physiology*, 14 (1), 72-81.
- Haemmerle, G., Zimmermann, R., Hayn, M., Theussl, C., Waeg, G., Wagner, E., Sattler, W., Magin, T. M., Wagner, E. F., & Zechner, R. (2002). Hormone-sensitive lipase deficiency in mice causes diglyceride accumulation in adipose tissue, muscle, and testis. *Journal of Biological Chemistry*, 277(7), 4806-4815.
- Hales, C. M., Carroll, M. D., Fryar, C. D., & Ogden, C. L. (2017). Prevalence of obesity among adults and youth: United States, 2015–2016.
- Halpern, B. N., Benacerraf, B., & Biozzi, G. (1953). Quantitative study of the granulopoietic activity of the reticulo-endothelial system I: The effect of the ingredients present in India ink and of substances affecting blood clotting *in vivo* on the fate of carbon particles administered intravenously in rats, mice and rabbits. *British Journal of Experimental Pathology*, 34, 426–440.

- Hamidpour, R., Hamidpour, S., & Elias, G. (2017). Rosmarinus Officinalis (Rosemary): A Novel Therapeutic Agent for Antioxidant, Antimicrobial, Anticancer, Antidiabetic, Antidepressant, Neuroprotective, Anti-Inflammatory, and Anti-Obesity Treatment. *Biomedical Journal of Scientific & Technical Research*, 1(4), 1-6.
- Han, F., Li, S., Yin, R., Liu, H., & Xu, L. (2008). Effect of surfactants on the formation and characterization of a new type of colloidal drug delivery system: nanostructured lipid carriers. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 315(1-3), 210-216.
- Haniadka, R., Saldanha, E., Sunita, V., Palatty, P. L., Fayad R., & Baliga, M. S. (2013). A review of the gastroprotective effects of ginger (*Zingiber officinale* Roscoe). *Food & Function*, 4(6), 845-855.
- Harimurti, N., Nhestricia, N., Subardjo, S. Y., & Yuliani, S. (2011). Effect of oleoresin concentration and composition of encapsulating materials on properties of the microencapsulated ginger oleoresin using spray drying method. *Indonesian Journal of Agriculture*, 4(1), 33-39.
- Heal, D. J., Gosden, J., & Smith, S. L. (2012). What is the prognosis for new centrally-acting anti-obesity drugs?. *Neuropharmacology*, 63(1), 132-146.
- Herman, A., & Herman, A.P. (2015). Essential oils and their constituents as skin penetration enhancer for transdermal drug delivery: a review. *Journal of Pharmacy and Pharmacology*, 67(4), pp.473-485.
- Heurtault, B., Saulnier, P., Pech, B., Proust, J. E., & Benoit, J. P. (2002). A novel phase inversion-based process for the preparation of lipid nanocarriers. *Pharmaceutical Research*, 19(6), 875-880.
- Higuchi, T. (1963). Mechanism of sustained-action medication. Theoretical analysis of rate of release of solid drugs dispersed in solid matrices. *Journal of Pharmaceutical Sciences*, 52(12), 1145-1149.
- Ho, S. C., Chang, K. S., & Lin, C. C. (2013). Anti-neuroinflammatory capacity of fresh ginger is attributed mainly to 10-gingerol. *Food Chemistry*, 141(3), 3183-3191.
- Horne, D. S. (1995). Light scattering studies of colloid stability and gelation. *New Physicochemical Techniques for the Characterization of Complex Food Systems*. Blackie Academic and Professional, London, UK.

- Hosseini, S. F., Zandi, M., Rezaei, M., & Farahmandghavi, F. (2013). Two-step method for encapsulation of oregano essential oil in chitosan nanoparticles: preparation, characterization and *in vitro* release study. *Carbohydrate Polymers*, 95(1), 50-56.
- How, C. W., Rasedee, A., Manickam, S., & Rosli, R. (2013). Tamoxifen-loaded nanostructured lipid carrier as a drug delivery system: characterization, stability assessment and cytotoxicity. *Colloids and Surfaces B: Biointerfaces*, 112, 393-399.
- Hsu, Y. W., Chu, D. C., Ku, P. W., Liou, T. H., & Chou, P. (2010). Pharmacotherapy for obesity: past, present and future. *Journal of Experimental & Clinical Medicine*, 2(3), 118-123.
- Hu, F. Q., Jiang, S. P., Du, Y. Z., Yuan, H., Ye, Y. Q., & Zeng, S. (2005). Preparation and characterization of stearic acid nanostructured lipid carriers by solvent diffusion method in an aqueous system. *Colloids and Surfaces B: Biointerfaces*, 45(3-4), 167-173.
- Hu, F. Q., Yuan, H., Zhang, H. H., & Fang, M. (2002). Preparation of solid lipid nanoparticles with clobetasol propionate by a novel solvent diffusion method in aqueous system and physicochemical characterization. *International Journal of Pharmaceutics*, 239(1-2), 121-128.
- Huang, D., Ou, B., & Prior, R. L. (2005). The chemistry behind antioxidant capacity assays. *Journal of Agricultural and Food Chemistry*, 53(6), 1841-1856.
- Huang, M., Khor, E., & Lim, L. Y. (2004). Uptake and cytotoxicity of chitosan molecules and nanoparticles: effects of molecular weight and degree of deacetylation. *Pharmaceutical Research*, 21(2), 344-353.
- Hung, L. C., Basri, M., Tejo, B. A., Ismail, R., Nang, H. L. L., Hassan, H. A., & May, C. Y. (2011). An improved method for the preparations of nanostructured lipid carriers containing heat-sensitive bioactives. *Colloids and Surfaces B: Biointerfaces*, 87(1), 180-186.
- Hunstad, J. P., Godek, C. P., Van Natta, B. W., Kortesis, B. G., Bharti, G., Crantford, J. C., Daniels, M. A., & Andrew, M. S. (2018). A multicenter, prospective, randomized, contralateral study of tissue liquefaction liposuction vs suction-assisted liposuction. *Aesthetic Surgery Journal*, 38(9), 980-989.

- Husky, M. M., Mazure, C. M., Ruffault, A., Flahault, C., & Kovess-Masfety, V. (2018). Differential associations between excess body weight and psychiatric disorders in men and women. *Journal of Women's Health*, 27(2), 183-190.
- Iannuccelli, V., Bertelli, D., Romagnoli, M., Scalia, S., Maretti, E., Sacchetti, F., & Leo, E. (2014). In vivo penetration of bare and lipid-coated silica nanoparticles across the human stratum corneum. *Colloids and Surfaces B: Biointerfaces*, 122, 653-661.
- Ibrahim, M.M. (2010). Subcutaneous and visceral adipose tissue: structural and functional differences. *Obesity Reviews*, 11(1), 11-18.
- Iqbal, M. A., Md, S., Sahni, J. K., Baboota, S., Dang, S. and Ali, J. (2012). Nanostructured lipid carriers system: recent advances in drug delivery. *J Drug Target*, 20(10), 813-830.
- Isa, Y., Miyakawa, Y., Yanagisawa, M., Goto, T., Kang, M.S., Kawada, T., Morimitsu, Y., Kubota, K., & Tsuda, T. (2008). 6-Shogaol and 6-gingerol, the pungent of ginger, inhibit TNF- α mediated downregulation of adiponectin expression via different mechanisms in 3T3-L1 adipocytes. *Biochemical and Biophysical Research Communications*, 373(3), 429-434.
- Ito, K., Kato, Y., Tsuji, H., Nguyen, H.T., Kubo, Y., & Tsuji, A. (2007). Involvement of organic anion transport system in transdermal absorption of flurbiprofen. *Journal of Controlled Release*, 124(1-2), 60-68.
- Ito, K., Nguyen, H.T., Kato, Y., Wakayama, T., Kubo, Y., Iseki, S., & Tsuji, A. (2008). P-glycoprotein (Abcb1) is involved in absorptive drug transport in skin. *Journal of Controlled Release*, 131(3), 198-204.
- Jaworski, K., Ahmadian, M., Duncan, R. E., Sarkadi-Nagy, E., Varady, K. A., Hellerstein, M. K., Lee, H. Y., Samuel, V. T., Shulman, G. I., Kim, K. H., & De Val, S. (2009). AdPLA ablation increases lipolysis and prevents obesity induced by high-fat feeding or leptin deficiency. *Nature Medicine*, 15(2), 159-168.
- Jayarathne, S., Koboziev, I., Park, O. H., Oldewage-Theron, W., Shen, C. L., & Moustaid-Moussa, N. (2017). Anti-inflammatory and anti-obesity properties of food bioactive components: Effects on adipose tissue. *Preventive Nutrition and Food Science*, 22(4), 251.
- Jee, W., & Weiss, L. (1983). Histology: Cell and tissue biology. *Histology Cell and Tissue Biology*, 5(9), 200-255.

- Jemaa, M.B., Falleh, H., Serairi, R., Neves, M.A., Snoussi, M., Isoda, H., Nakajima, M., & Ksouri, R. (2018). Nanoencapsulated *Thymus capitatus* essential oil as natural preservative. *Innovative Food Science & Emerging Technologies*, 45, 92-97.
- Jeon, S., Yoo, C. Y., & Park, S. N. (2015). Improved stability and skin permeability of sodium hyaluronate-chitosan multilayered liposomes by Layer-by-Layer electrostatic deposition for quercetin delivery. *Colloids and Surfaces B: Biointerfaces*, 129, 7-14.
- Jeong, Y. S., Jung, H. K., Cho, K. H., Youn, K. S., & Hong, J. H. (2011). Anti-obesity effect of grape skin extract in 3T3-L1 adipocytes. *Food Science and Biotechnology*, 20(3), 635-642.
- Jepps, O. G., Dancik, Y., Anissimov, Y. G., & Roberts, M. S. (2013). Modeling the human skin barrier—Towards a better understanding of dermal absorption. *Advanced Drug Delivery Reviews*, 65(2), 152-168.
- Jiang, J., Oberdörster, G., & Biswas, P. (2009). Characterization of size, surface charge, and agglomeration state of nanoparticle dispersions for toxicological studies. *Journal of Nanoparticle Research*, 11(1), 77-89.
- Johji, Y., Keizo, M., Takeshi, C., Tokunosuke, S., Hajime, F., Toshiaki, T., Kimiko, N., & Toshihiro, N. (1985). Cholagogic effect of ginger and its active constituents. *Journal of Ethnopharmacology*, 13(2), 217-225.
- Joo, H., Kim, C. T., Kim, I. H., & Kim, Y. (2013). Anti-obesity effects of hot water extract and high hydrostatic pressure extract of garlic in rats fed a high-fat diet. *Food and Chemical Toxicology*, 55, 100-105.
- Jores, K., Mehnert, W., Drechsler, M., Bunjes, H., Johann, C., & Mäder, K. (2004). Investigations on the structure of solid lipid nanoparticles (SLN) and oil-loaded solid lipid nanoparticles by photon correlation spectroscopy, field-flow fractionation and transmission electron microscopy. *Journal of Controlled Release*, 95(2), 217-227.
- Joshi, M., & Patravale, V. (2008). Nanostructured lipid carrier (NLC) based gel of celecoxib. *International Journal of Pharmaceutics*, 346(1-2), 124-132.
- Jung, E. C., & Maibach, H. I. (2014). Animal models for percutaneous absorption. In *Topical drug bioavailability, bioequivalence, and Penetration* (pp. 21-40). Springer, New York, NY.

- Junyaprasert, V. B., Teeranachaideekul, V., Souto, E. B., Boonme, P., & Müller, R. H. (2009). Q10-loaded NLC versus nanoemulsions: stability, rheology and in vitro skin permeation. *International Journal of Pharmaceutics*, 377(1-2), 207-214.
- Kano, S., Todo, H., Sugie, K., Fujimoto, H., Nakada, K., Tokudome, Y., Hashimoto, F., & Sugibayashi, K. (2010). Utilization of reconstructed cultured human skin models as an alternative skin for permeation studies of chemical compounds. *Alternatives to Animal Testing and Experimentation*, 15(2), 61-70.
- Karadzovska, D., Brooks, J. D., Monteiro-Riviere, N. A., & Riviere, J. E. (2013). Predicting skin permeability from complex vehicles. *Advanced Drug Delivery Reviews*, 65(2), 265-277.
- Karamadoukis, L., Shivashankar, G. H., Ludeman, L., & Williams, A. J. (2009). An unusual complication of treatment with orlistat. *Clinical Nephrology*, 71(4), 430-432.
- Kasongo, W. A., Pardeike, J., Müller, R. H., & Walker, R. B. (2011). Selection and characterization of suitable lipid excipients for use in the manufacture of didanosine-loaded solid lipid nanoparticles and nanostructured lipid carriers. *Journal of Pharmaceutical Sciences*, 100(12), 5185-5196.
- Kassem, A. A., El-Alim, S. H. A., & Asfour, M. H. (2017). Enhancement of 8-methoxypsoralen topical delivery via nanosized niosomal vesicles: Formulation development, in vitro and in vivo evaluation of skin deposition. *International Journal of Pharmaceutics*, 517(1-2), 256-268.
- Kaur, S. & Mondal, P. (2014). Study of total phenolic and flavonoid content, antioxidant activity and antimicrobial properties of medicinal plants. *Journal of Microbiology and Experimentation*, 1(1), 1-6.
- Kazeem, M. I., Bankole, H. A., & Fatai, A. A. (2011). Protective effect of ginger in normal and carbon-tetrachloride induced hepatotoxic rats. *Annals of Biological Research*, 2(1), 1-8.
- Kelly, C. Z., Marcia, O. M. M., Ademir, J. P., & Meireles, M. A. A. (2002). Extraction of ginger (*Zingiber officinale* Roscoe) oleoresin with CO₂ and co-solvents a study of the antioxidant action of the extracts. *Journal of Supercritical Fluids*, 24, 57-76.

- Kennedy, J., Verne, S., Griffith, R., Falto-Aizpurua, L., & Nouri, K. (2015). Non-invasive subcutaneous fat reduction: a review. *Journal of the European Academy of Dermatology and Venereology*, 29(9), 1679-1688.
- Kesarwani, A., Yadav, A. K., Singh, S., Gautam, H., Singh, H. N., Sharma, A., & Yadav, C. (2013). Theoretical aspects of transdermal drug delivery system. *Bulletin of Pharmaceutical Research*, 3(2), 78-89.
- Khan, I., Pandotra, P., Gupta, A. P., Sharma, R., Gupta, B. D., Dhar, J. K., Ram, G., Bedi, Y., & Gupta, S. (2010). RP-thin layer chromatographic method for the quantification of three gingerol homologs of ultrasonic-assisted fresh rhizome extracts of *Zingiber officinale* collected for North Western Himalayas. *Journal of Separation Science*, 33, 558-563.
- Khater, D. S. (2010). The influence of ginger as a chemopreventive agent on proliferation and apoptosis in chemically induced oral carcinogenesis. *Nature Science*, 8, 44-51.
- Kim Y.S., Lee Y.M., Kim H., Kim J., Jang D.S., Kim J.H., & Kim J.S. (2010). Anti-obesity effect of morus bombycis root extract: anti-lipase activity and lipolytic effect. *Journal of Ethnopharmacology*, 120(3) 621-624.
- Kim, E. C., Min, J. K., Kim, T. Y., Lee, S.J., Yang, H.O., Han, S., Kim, Y. M., & Kwon, Y. G. (2005). [6]-Gingerol, a pungent ingredient of ginger, inhibits angiogenesis *in vitro* and *in vivo*. *Biochemical and Biophysical Research Communications*, 335(2), 300-308.
- Kim, G. W., Lin, J. E., Blomain, E. S., & Waldman, S. A. (2014). Antiobesity pharmacotherapy: new drugs and emerging targets. *Clinical Pharmacology & Therapeutics*, 95(1), 53-66.
- Kirjavainen, M., Mönkkönen, J., Saukkosaari, M., Valjakka-Koskela, R., Kiesvaara, J., & Urtti, A. (1999). Phospholipids affect stratum corneum lipid bilayer fluidity and drug partitioning into the bilayers. *Journal of Controlled Release*, 58(2), 207-214.
- Kong, B.J., Kim, A., & Park, S.N. (2016). Properties and in vitro drug release of hyaluronic acid-hydroxyethyl cellulose hydrogels for transdermal delivery of isoliquiritigenin. *Carbohydrate polymers*, 147, 473-481.
- Korsmeyer, R. W., Gurny, R., Doelker, E., Buri, P., & Peppas, N. A. (1983). Mechanisms of solute release from porous hydrophilic polymers. *International Journal of Pharmaceutics*, 15(1), 25-35.

- Kost, J., & Langer, R. (2012). Responsive polymeric delivery systems. *Advanced Drug Delivery Reviews*, 64, 327-341.
- Kreuter, J. (2007). Nanoparticles—a historical perspective. *International Journal of Pharmaceutics*, 331(1), 1-10.
- Kuipers, E., Dam, A., Held, N., Mol, I., Houtkooper, R., Rensen, P., & Boon, M. (2018). Quercetin lowers plasma triglycerides accompanied by white adipose tissue Browning in diet-induced obese mice. *International Journal of Molecular Sciences*, 19(6), 1786.
- Lacatusu, I., Mitrea, E., Badea, N., Stan, R., Oprea, O., & Meghea, A. (2013). Lipid nanoparticles based on omega-3 fatty acids as effective carriers for lutein delivery. Preparation and in vitro characterization studies. *Journal of Functional Foods*, 5(3), 1260-1269.
- Lademann, J., Knorr, F., Richter, H., Blume-Peytavi, U., Vogt, A., Antoniou, C., Sterry, W., & Patzelt, A. (2008). Hair follicles—an efficient storage and penetration pathway for topically applied substances. *Skin Pharmacology and Physiology*, 21(3), 150-155.
- Lakshmi, P., & Kumar, G. A. (2010). Nanosuspension technology: A review. *International Journal of Pharmacy and Pharmaceutical Sciences*, 2(4), 35-40.
- Lass, A., Zimmermann, R., Oberer, M., & Zechner, R. (2011). Lipolysis—a highly regulated multi-enzyme complex mediates the catabolism of cellular fat stores. *Progress in Lipid Research*, 50(1), 14-27.
- Lee, G. R., Shin, M. K., Yoon, D. J., Kim, A. R., Yu, R., Park, N. H., & Han, I. S. (2013). Topical application of capsaicin reduces visceral adipose fat by affecting adipokine levels in high-fat diet-induced obese mice. *Obesity*, 21(1), 115-122.
- Lee, H., Kang, R., & Yoon, Y. (2010). SH21B, an anti-obesity herbal composition, inhibits fat accumulation in 3T3-L1 adipocytes and high fat diet-induced obese mice through the modulation of the adipogenesis pathway. *Journal of Ethnopharmacology*, 127(3), 709-717.
- Lee, J. H., Moon, M. H., Jeong, J. K., Park, Y. G., Lee, Y. J., Seol, J. W., & Park, S. Y. (2012). Sulforaphane induced adipolysis via hormone sensitive lipase activation, regulated by AMPK signaling pathway. *Biochemical and Biophysical Research Communications*, 426(4), 492-497.

- Lee, M. K., Lim, S. J., & Kim, C. K. (2007). Preparation, characterization and in vitro cytotoxicity of paclitaxel-loaded sterically stabilized solid lipid nanoparticles. *Biomaterials*, 28(12), 2137-2146.
- Lee, S. G., Jeong, J. H., Kim, S. R., Lee, K. M., Ahn, B. K., Kang, M. H., & Choi, Y. W. (2012). Topical formulation of retinyl retinoate employing nanostructured lipid carriers. *Journal of Pharmaceutical Investigation*, 42(5), 243-250.
- Lewis, D. R., & Liu, D. J. (2012). Direct measurement of lipase inhibition by orlistat using a dissolution linked in vitro assay. *Clinical Pharmacology & Biopharmaceutics*, 1(103).
- Li, C., & Zhou, L. (2015). Inhibitory effect 6-gingerol on adipogenesis through activation of the Wnt/ β -catenin signaling pathway in 3T3-L1 adipocytes. *Toxicology in Vitro*, 30(1), 394-401.
- Li, F., Wang, Y., Parkin, K. L., Nitteranon, V., Liang, J., Yang, W., Li, Y., Zhang, G., & Hu, Q. (2011). Isolation of quinone reductase (QR) inducing agents from ginger rhizome and their *in vitro* anti-inflammatory activity. *Food Research International*, 44(6), 1597-1603.
- Li, M., Zheng, Y., Shan, F. Y., Zhou, J., Gong, T., & Zhang, Z. R. (2013). Development of ionic-complex-based nanostructured lipid carriers to improve the pharmacokinetic profiles of breviscapine. *Acta Pharmacologica Sinica*, 34(8), 1108-1115.
- Li, Y., Taulier, N., Rauth, A. M., & Wu, X. Y. (2006). Screening of lipid carriers and characterization of drug-polymer-lipid interactions for the rational design of polymer-lipid hybrid nanoparticles (PLN). *Pharmaceutical Research*, 23(8), 1877-1887.
- Liu, I.M., & Chang, C. J. (2013). U.S. Patent 8,501,249,B2. Washington D.C: U.S Patent and Trademark Office.
- Loo, C. H., Basri, M., Ismail, R., Lau, H. L. N., Tejo, B. A., Kanthimathi, M. S., Hassan, H. A., & Choo, Y. M. (2013). Effect of compositions in nanostructured lipid carriers (NLC) on skin hydration and occlusion. *International Journal of Nanomedicine*, 8, 13.
- Lopes, D. B., Speranza, P., & Macedo, G. A. (2016). A new approach for flavor and aroma encapsulation. In *Novel Approaches of Nanotechnology in Food* (pp. 623-661). Academic Press.

- Lundberg, G. D. (2008). Fat reduction by topical waist applications may actually work. *The Medscape Journal of Medicine*, 10(2), 43.
- Maherani, B., Arab-Tehrany, E., Kheirrolomoom, A., Reshetov, V., Stebe, M. J., & Linder, M. (2012). Optimization and characterization of liposome formulation by mixture design. *Analyst*, 137(3), 773-786.
- Mahmoud, R. H., & Elnour, W. A. (2013). Comparative evaluation of the efficacy of ginger and orlistat on obesity management, pancreatic lipase and liver peroxisomal catalase enzyme in male albino rats. *European Review for Medical and Pharmacological Sciences*, 17(1), 75-83.
- Mali, N., Darandale, S., & Vavia, P. (2013). Niosomes as a vesicular carrier for topical administration of minoxidil: formulation and in vitro assessment. *Drug Delivery and Translational Research*, 3(6), 587-592.
- Malik, Z. A., & Sharma, P. L. (2011). Anomalies after Chronic Administration of Ginger (*Zingiber officinale*) in Wistar Rats. *International Journal of Pharmacology*, 7(8), 801-812.
- Manea, A. M., Vasile, B. S., & Meghea, A. (2014). Antioxidant and antimicrobial activities of green tea extract loaded into nanostructured lipid carriers. *Comptes Rendus Chimie*, 17(4), 331-341.
- Margalit, R., Alon, R., Linenberg, M., Rubin, I., Roseman, T. J., & Wood, R. W. (1991). Liposomal drug delivery: thermodynamic and chemical kinetic considerations. *Journal of Controlled Release*, 17(3), 285-296.
- Martins, I. M., Barreiro, M. F., Coelho, M., & Rodrigues, A. E. (2014). Microencapsulation of essential oils with biodegradable polymeric carriers for cosmetic applications. *Chemical Engineering Journal*, 245, 191-200.
- Mashmoul, M., Azlan, A., Khaza'ai, H., Yusof, B., & Noor, S. (2013). Saffron: a natural potent antioxidant as a promising anti-obesity drug. *Antioxidants*, 2(4), 293-308.
- McClements, D. J. (2015). *Food emulsions: principles, practices, and techniques*. CRC press.
- Medeiros, L. B. (2003). *Lipogistrofia ginóide: tratamento clínico*. Rio de Janeiro, Atheneu.

- Mendes, A. I., Silva, A. C., Catita, J. A. M., Cerqueira, F., Gabriel, C., & Lopes, C. M. (2013). Miconazole-loaded nanostructured lipid carriers (NLC) for local delivery to the oral mucosa: improving antifungal activity. *Colloids and Surfaces B: Biointerfaces*, 111, 755-763.
- Mesomo, M. C., Corazza, M. L., Ndiaye, P. M., Dalla Santa, O. R., Cardozo, L., & de Paula Scheer, A. (2013). Supercritical CO₂ extracts and essential oil of ginger (*Zingiber officinale* R.): Chemical composition and antibacterial activity. *The Journal of Supercritical Fluids*, 80, 44-49.
- Michaels, A. S., Chandrasekaran, S. K., & Shaw, J. E. (1975). Drug permeation through human skin: theory and in vitro experimental measurement. *AIChE Journal*, 21(5), 985-996.
- Misawa, K., Hashizume, K., Yamamoto, M., Minegishi, Y., Hase, T., & Shimotoyodome, A. (2015). Ginger extract prevents high-fat diet-induced obesity in mice via activation of the peroxisome proliferator-activated receptor δ pathway. *The Journal of Nutritional Biochemistry*, 26(10), 1058-1067.
- Modi, S., & Anderson, B. D. (2013). Determination of drug release kinetics from nanoparticles: overcoming pitfalls of the dynamic dialysis method. *Molecular Pharmaceutics*, 10(8), 3076-3089.
- Mohamed, Y. I., Siddig, I. A. W., Manal, M. E., Adel, S. A., & Syam, M. M. (2009). Attenuation of cisplatin-induced hepatotoxicity in rats using zerumbone. *Research Journal of Biological Sciences*, 4(7), 777-784.
- Mojahedian, M. M., Daneshamouz, S., Samani, S. M., & Zargaran, A. (2013). A novel method to produce solid lipid nanoparticles using n-butanol as an additional co-surfactant according to the o/w microemulsion quenching technique. *Chemistry and Physics of Lipids*, 174, 32-38.
- Montenegro, L., Pasquinucci, L., Zappalà, A., Chiechio, S., Turnaturi, R., & Parenti, C. (2017). Rosemary essential oil-loaded lipid nanoparticles: In vivo topical activity from gel vehicles. *Pharmaceutics*, 9(4), 48.
- Moreno-Sastre, M., Pastor, M., Esquisabel, A., Sans, E., Viñas, M., Bachiller, D., & Pedraz, J. L. (2016). Stability study of sodium colistimethate-loaded lipid nanoparticles. *Journal of Microencapsulation*, 33(7), 636-645.
- Moreno-Sastre, M., Pastor, M., Esquisabel, A., Sans, E., Viñas, M., Bachiller, D., & Pedraz, J. L. (2016). Stability study of sodium colistimethate-loaded lipid nanoparticles. *Journal of Microencapsulation*, 33(7), 636-645.

- Muhammad Nadzri, N., Abdul, A. B., Sukari, M. A., Abdelwahab, S. I., Eid, E. E., Mohan, S., ... & Arbab, I. A. (2013). Inclusion complex of zerumbone with hydroxypropyl- β -cyclodextrin induces apoptosis in liver hepatocellular HepG2 cells via caspase 8/bid cleavage switch and modulating Bcl2/Bax ratio. *Evidence-Based Complementary and Alternative Medicine*, 2013.
- Müller, R. H., Petersen, R. D., Hommoss, A., & Pardeike, J. (2007). Nanostructured lipid carriers (NLC) in cosmetic dermal products. *Advanced Drug Delivery Reviews*, 59(6), 522-530.
- Müller, R. H., Radtke, M., & Wissing, S. A. (2002). Solid lipid nanoparticles (SLN) and nanostructured lipid carriers (NLC) in cosmetic and dermatological preparations. *Advanced Drug Delivery Reviews*, 54, S131-S155.
- Müller, R. H., Rühl, D., Runge, S., Schulze-Forster, K., & Mehnert, W. (1997). Cytotoxicity of solid lipid nanoparticles as a function of the lipid matrix and the surfactant. *Pharmaceutical Research*, 14(4), 458-462.
- Murdock, R. C., Braydich-Stolle, L., Schrand, A. M., Schlager, J. J., & Hussain, S. M. (2008). Characterization of nanomaterial dispersion in solution prior to in vitro exposure using dynamic light scattering technique. *Toxicological Sciences*, 101(2), 239-253.
- Nagoya Declaration 2015. The 8th Asia-Oceania Conference on Obesity (AOCO 2015). <http://www.jasso.or.jp/data/data/pdf/nagoya2015.pdf>
- Nahr, F. K., Ghanbarzadeh, B., Hamishehkar, H., & Kafil, H. S. (2018). Food grade nanostructured lipid carrier for cardamom essential oil: Preparation, characterization and antimicrobial activity. *Journal of Functional Foods*, 40, 1-8.
- Naik, A., Kalia, Y. N., & Guy, R. H. (2000). Transdermal drug delivery: overcoming the skin's barrier function. *Pharmaceutical Science & Technology Today*, 3(9), 318-326.
- Nakajima, I., Aso, H., Yamaguchi, T., & Ozutsumi, K. (1998). Adipose tissue extracellular matrix: newly organized by adipocytes during differentiation. *Differentiation*, 63(4), 193-200.
- Nastruzzi, C. (2004). *Lipospheres in drug targets and delivery: approaches, methods, and applications*. CRC Press.

- Nayak, A. P., Tiyafoonchai, W., Patankar, S., Madhusudhan, B., & Souto, E. B. (2010). Curcuminoids-loaded lipid nanoparticles: novel approach towards malaria treatment. *Colloids and Surfaces B: Biointerfaces*, 81(1), 263-273.
- Nel, A., Xia, T., Mädler, L., & Li, N. (2006). Toxic potential of materials at the nanolevel. *Science*, 311(5761), 622-627.
- Ng, S. F., Rouse, J. J., Sanderson, F. D., Meidan, V., & Eccleston, G. M. (2010). Validation of a static Franz diffusion cell system for in vitro permeation studies. *AAPS Pharmscitech*, 11(3), 1432-1441.
- Nguyen, H. M., Hwang, I. C., Park, J. W., & Park, H. J. (2012). Enhanced payload and photo-protection for pesticides using nanostructured lipid carriers with corn oil as liquid lipid. *Journal of Microencapsulation*, 29(6), 596-604.
- Nguyen, H. X., Puri, A., Bhattacharjee, S. A., & Banga, A. K. (2018). Qualitative and quantitative analysis of lateral diffusion of drugs in human skin. *International Journal of Pharmaceutics*, 544(1), 62-74.
- Ni, S., Sun, R., Zhao, G., & Xia, Q. (2015). Quercetin loaded nanostructured lipid carrier for food fortification: preparation, characterization and in vitro study. *Journal of Food Process Engineering*, 38(1), 93-106.
- Nikolić, S., Keck, C. M., Anselmi, C., & Müller, R. H. (2011). Skin photoprotection improvement: synergistic interaction between lipid nanoparticles and organic UV filters. *International Journal of Pharmaceutics*, 414(1-2), 276-284.
- Niwano, Y., Beppu, F., Shimada, T., Kyan, R., Yasura, K., Tamaki, M., Nishino, M., Midorikawa, Y., & Hamada, H. (2009). Extensive screening for plant foodstuffs in Okinawa, Japan with anti-obese activity on adipocytes in vitro. *Plant Foods for Human Nutrition*, 64(1), 6.
- Nobbmann, U., & Morfesis, A. (2009). Light scattering and nanoparticles. *Materials Today*, 12(5), 52-54.
- Nokhodchi, A., Sharabiani, K., Rashidi, M. R., & Ghafourian, T. (2007). The effect of terpene concentrations on the skin penetration of diclofenac sodium. *International Journal of Pharmaceutics*, 335(1-2), 97-105.
- Noor, N. M., Sheikh, K., Somavarapu, S., & Taylor, K. M. (2017). Preparation and characterization of dutasteride-loaded nanostructured lipid carriers coated with stearic acid-chitosan oligomer for topical delivery. *European Journal of Pharmaceutics and Biopharmaceutics*, 117, 372-384.

- Noor, N.M., Aziz, A. A., Sarmidi, M. R., & Aziz, R. (2013). The effect of virgin coconut oil loaded solid lipid particles (VCO-SLPs) on skin hydration and skin elasticity, *Jurnal Teknologi*, 62(1), 39-43.
- Nootheti, P. K., Magpantay, A., Yosowitz, G., Calderon, S., & Goldman, M. P. (2006). A single center, randomized, comparative, prospective clinical study to determine the efficacy of the VelaSmooth system versus the Triactive system for the treatment of cellulite. *Lasers in Surgery and Medicine*, 38(10), 908-912.
- Obeidat, W. M., Schwabe, K., Müller, R. H., & Keck, C. M. (2010). Preservation of nanostructured lipid carriers (NLC). *European Journal of Pharmaceutics and Biopharmaceutics*, 76(1), 56-67.
- Obesity and overweight. (2018, February 16). Retrieved February 11, 2019, from <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>
- Oliveira, C. P., Venturini, C. G., Donida, B., Poletto, F. S., Guterres, S. S., & Pohlmann, A. R. (2013). An algorithm to determine the mechanism of drug distribution in lipid-core nanocapsule formulations. *Soft Matter*, 9(4), 1141-1150.
- Orafidiya, L. O., & Oladimeji, F. A. (2002). Determination of the required HLB values of some essential oils. *International Journal of Pharmaceutics*, 237(1-2), 241-249.
- Palani, S. (2006). *Increased fat oxidation in 3T3-L1 adipocytes through forced expression of UCP 1* (Doctoral dissertation, Texas A&M University).
- Pandit, J., Aqil, M., & Sultana, Y. (2016). Nanoencapsulation technology to control release and enhance bioactivity of essential oils. In *Encapsulations* (pp. 597-640). Academic Press.
- Paolino, D., Ventura, C. A., Nistico, S., Puglisi, G., & Fresta, M. (2002). Lecithin microemulsions for the topical administration of ketoprofen: percutaneous adsorption through human skin and in vivo human skin tolerability. *International Journal of Pharmaceutics*, 244(1-2), 21-31.
- Pardeike, J., Hommoss, A., & Müller, R. H. (2009). Lipid nanoparticles (SLN, NLC) in cosmetic and pharmaceutical dermal products. *International Journal of Pharmaceutics*, 366(1-2), 170-184.

- Pardeike, J., Weber, S., Haber, T., Wagner, J., Zarfl, H. P., Plank, H., & Zimmer, A. (2011). Development of an itraconazole-loaded nanostructured lipid carrier (NLC) formulation for pulmonary application. *International Journal of Pharmaceutics*, 419(1-2), 329-338.
- Parhi, R., & Suresh, P. (2016). Transdermal delivery of Diltiazem HCl from matrix film: Effect of penetration enhancers and study of antihypertensive activity in rabbit model. *Journal of advanced research*, 7(3), 539-550.
- Park, M. J., Balakrishnan, P., & Yang, S. G. (2013). Polymeric nanocapsules with SEDDS oil-core for the controlled and enhanced oral absorption of cyclosporine. *International Journal of Pharmaceutics*, 441(1-2), 757-764.
- Park, Y. H., Bae, H. C., Jang, Y., Jeong, S. H., Lee, H. N., Ryu, W. I., Yoo, M. G., Kim, Y. R., Kim, M. K. Lee, J. K., Jeong, J. & Son, S. W. (2013). Effect of the size and surface charge of silica nanoparticles on cutaneous toxicity. *Molecular & Cellular Toxicology*, 9(1), 67-74.
- Patel, K., Padhye, S., & Nagarsenker, M. (2012). Duloxetine HCl lipid nanoparticles: preparation, characterization, and dosage form design. *AAPS PharmSciTech*, 13(1), 125-133.
- Pellegrino, J., Schulte, L. R., De la Cruz, J., & Stoldt, C. (2017). Membrane processes in nanoparticle production. *Journal of Membrane Science*, 522, 245-256.
- Peppas, N. A. (1985). Analysis of Fickian and non-Fickian drug release from polymers. *Pharmaceutica Acta Helvetiae*, 60(4), 110.
- Petersen, S., Steiniger, F., Fischer, D., Fahr, A., & Bunjes, H. (2011). The physical state of lipid nanoparticles influences their effect on in vitro cell viability. *European Journal of Pharmaceutics and Biopharmaceutics*, 79(1), 150-161.
- Phaechamud, T., Koizumi, T., & Ritthidej, G. C. (2000). Chitosan citrate as film former: compatibility with water-soluble anionic dyes and drug dissolution from coated tablet. *International Journal of Pharmaceutics*, 198(1), 97-111.
- Phattanawasin, P., Sotanaphun, U., Sukwattanasinit, T., Akkarawarathorn, J., & Kitchaiya, S. (2012). Quantitative determination of sibutramine in adulterated herbal slimming formulations by TLC-image analysis method. *Forensic Science International*, 219(1-3), 96-100.

- Pichayakorn, W., Suksaeree, J., Boonme, P., Amnuaikit, T., Taweepreda, W., & Ritthidej, G. C. (2012). Nicotine transdermal patches using polymeric natural rubber as the matrix controlling system: effect of polymer and plasticizer blends. *Journal of Membrane Science*, 411, 81-90.
- Pokharkar, V. B., Shekhawat, P. B., Dhapte, V. V., & Mandpe, L. P. (2011). Development and optimization of eugenol loaded nanostructured lipid carriers for periodontal delivery. *International Journal of Pharmacy and Pharmaceutical Sciences*, 3(4), 138-143.
- Poulos, S. P., Dodson, M. V., & Hausman, G. J. (2010). Cell line models for differentiation: preadipocytes and adipocytes. *Experimental Biology and Medicine*, 235(10), 1185-1193.
- Pouton, C. W., & Porter, C. J. (2008). Formulation of lipid-based delivery systems for oral administration: materials, methods and strategies. *Advanced Drug Delivery Reviews*, 60(6), 625-637.
- Prakash, B., Kujur, A., Yadav, A., Kumar, A., Singh, P. P., & Dubey, N. K. (2018). Nanoencapsulation: An efficient technology to boost the antimicrobial potential of plant essential oils in food system. *Food control*, 89, 1-11.
- Prasannan, R., Kalesh, K. A., Shanmugam, M. K., Nachiyappan, A., Ramachandran, L., Nguyen, A. H., Kumar, A. P., Lakshmanan, M., Ahn, K. S., & Sethi, G. (2012). Key cell signaling pathways modulated by zerumbone: Role in the prevention and treatment of cancer. *Biochemical Pharmacology*, 84(10), 1268-1276.
- Prior, R. L., Wu, X., & Schaich, K. (2005). Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements. *Journal of Agricultural and Food Chemistry*, 53(10), 4290-4302.
- Prow, T. W., Grice, J. E., Lin, L. L., Faye, R., Butler, M., Becker, W., Wurm, E. M., Yoong, C., Robertson, T. A., Soyer, H. P., & Roberts, M. S. (2011). Nanoparticles and microparticles for skin drug delivery. *Advanced Drug Delivery Reviews*, 63(6), 470-491.
- Rabanel, J. M., Adibnia, V., Tehrani, S. F., Sanche, S., Hildgen, P., Banquy, X., & Ramassamy, C. (2019). Nanoparticle heterogeneity: an emerging structural parameter influencing particle fate in biological media?. *Nanoscale*, 11(2), 383-406.

- Radhakrishnan, R., Kulhari, H., Pooja, D., Gudem, S., Bhargava, S., Shukla, R., & Sistla, R. (2016). Encapsulation of biophenolic phytochemical EGCG within lipid nanoparticles enhances its stability and cytotoxicity against cancer. *Chemistry and Physics of Lipids*, 198, 51-60.
- Rahimpour, Y., & Hamishehkar, H. (2012). Niosomes as carrier in dermal drug delivery. In *Recent Advances in Novel Drug Carrier Systems*. IntechOpen.
- Rahman, H. S., Rasedee, A., How, C. W., Abdul, A. B., Zeenathul, N. A., Othman, H. H., Saeed, M. I., & Yeap, S. K. (2013). Zerumbone-loaded nanostructured lipid carriers: preparation, characterization, and antileukemic effect. *International Journal of Nanomedicine*, 8, 2769.
- Ranpise, N. S., Korabu, S. S., & Ghodake, V. N. (2014). Second generation lipid nanoparticles (NLC) as an oral drug carrier for delivery of lercanidipine hydrochloride. *Colloids and Surfaces B: Biointerfaces*, 116, 81-87.
- Rao, P., & Rathod, V. (2017). Phytochemicals: An insight to modern extraction technologies and their applications. In *Ingredients extraction by physicochemical methods in food* (pp. 495-521). Academic Press.
- Rashed, A. A., Mohd Nawi, M. N., & Sulaiman, K. (2017). Assessment of essential oil as a potential anti-obesity agent: a narrative review. *Journal of Essential oil Research*, 29(1), 1-10.
- Rasmussen, K., Rauscher, H., Mech, A., Fadeel, B., Pietroiusti, A., & Shvedova, A.A. (Eds.). (2017). *Physicochemical Characterization, Adverse Effects of Engineered Nanomaterials: Exposure, Toxicology and Impact on Human Health* (pp. 15 –49) .Academic Press.
- Reid, G. (2005). ThermoTRP channels and cold sensing: what are they really up to?. *Pflügers Archiv*, 451(1), 250-263.
- Reilly, S. M., & Lee, C. H. (2008). PPAR δ as a therapeutic target in metabolic disease. *FEBS Letters*, 582(1), 26-31.
- Reis, P., Watzke, H., Leser, M., Holmberg, K., & Miller, R. (2010). Interfacial mechanism of lipolysis as self-regulated process. *Biophysical Chemistry*, 147(3), 93-103.
- Reithmeier, H., Herrmann, J., & Göpferich, A. (2001). Lipid microparticles as a parenteral controlled release device for peptides. *Journal of Controlled Release*, 73(2), 339-350.

- Relkin, P., Shukat, R., & Moulin, G. (2014). Encapsulation of labile compounds in heat-and high-pressure treated protein and lipid nanoparticles. *Food Research International*, 63, 9-15.
- Ribier, A., Simonnet, J. T., Fanchon, C., Segot, E., & Cantin, H. (1997). *U.S. Patent No. 5,637,316*. Washington, DC: U.S. Patent and Trademark Office.
- Riss, T. L., Moravec, R. A., Niles, A. L., Duellman, S., Benink, H. A., Worzella, T. J., & Minor, L. (2016). Cell viability assays.
- Rodriguez, J., Martín, M. J., Ruiz, M. A., & Clares, B. (2016). Current encapsulation strategies for bioactive oils: From alimentary to pharmaceutical perspectives. *Food Research International*, 83, 41-59.
- Roh, C., & Jung, U. (2012). Screening of crude plant extracts with anti-obesity activity. *International Journal of Molecular Sciences*, 13(2), 1710-1719.
- Ronsard, N. C. (1973). Those lumps, bumps and bulges, you couldn't lose before. *Beauty and Health Publishing Co., New York*.
- Roure, R., Oddos, T., Rossi, A., Vial, F., & Bertin, C. (2011). Evaluation of the efficacy of a topical cosmetic slimming product combining tetrahydroxypropyl ethylenediamine, caffeine, carnitine, forskolin and retinol, in vitro, ex vivo and in vivo studies. *International Journal of Cosmetic Science*, 33(6), 519-526.
- Ruiz-Ojeda, F., Rupérez, A., Gomez-Llorente, C., Gil, A., & Aguilera, C. (2016). Cell models and their application for studying adipogenic differentiation in relation to obesity: A review. *International Journal of Molecular Sciences*, 17(7), 1040.
- Saatchi, K., Tod, S. E., Leung, D., Nicholson, K. E., Andreu, I., Buchwalder, C., Schmitt, V., Hafeli, U. O., & Gray, S. L. (2017). Characterization of alendronic-and undecylenic acid coated magnetic nanoparticles for the targeted delivery of rosiglitazone to subcutaneous adipose tissue. *Nanomedicine: Nanotechnology, Biology and Medicine*, 13(2), 559-568.
- Sabu, M. (2003). Revision of the genus *Zingiber* in South India. *Folia Malesiana*, 4, 25-52.
- Salleh, N., Jais, U. S., & Sarijo, S. H. (2015). Characterization of crosslinked zerumbone loaded gelatin–zeolite Y hybrid composites with glutaraldehyde for controlled release of natural anticancer drug. *Malaysian Journal of Analytical Sciences*, 19(6), 1290-1296.

- Sanap, G. S., & Mohanta, G. P. (2014). Investigation of the factors influencing the incorporation of miconazole in SLN and NLC dispersion. *Indo American Journal of Pharmaceutical Research*, 4, 1378-90.
- Santosh, K. S. C., Srinivas, P., Negi, P. S., & Bettadaiah, B. K. (2013). Antibacterial and antimutagenic activities of novel zerumbone analogues. *Food Chemistry*, 141(2), 1097-1103.
- São Pedro, A., Santo, I., Silva, C., Detoni, C., & Albuquerque, E. (2013). The use of nanotechnology as an approach for essential oil-based formulations with antimicrobial activity. *Microbial Pathogens and Strategies for Combating Them (Méndez-Vilas, A., ed.) Formatex Research Center Publisher*, 2, 1364-1374.
- Saravanan, G., Ponmurugan, P., Deepa, M. A., & Senthilkumar, B. (2014). Anti-obesity action of gingerol: effect on lipid profile, insulin, leptin, amylase and lipase in male obese rats induced by a high-fat diet. *Journal of the Science of Food and Agriculture*, 94(14), 2972-2977.
- Schubert, M. A., & Müller-Goymann, C. C. (2003). Solvent injection as a new approach for manufacturing lipid nanoparticles—evaluation of the method and process parameters. *European Journal of Pharmaceutics and Biopharmaceutics*, 55(1), 125-131.
- Schwarz, J. C., Baisaeng, N., Hoppel, M., Löw, M., Keck, C. M., & Valenta, C. (2013). Ultra-small NLC for improved dermal delivery of coenzyme Q10. *International Journal of Pharmaceutics*, 447(1-2), 213-217.
- Schwarz, J. C., Weixelbaum, A., Pagitsch, E., Löw, M., Resch, G. P., & Valenta, C. (2012). Nanocarriers for dermal drug delivery: influence of preparation method, carrier type and rheological properties. *International Journal of Pharmaceutics*, 437(1-2), 83-88.
- Seiller, M., Orecchioni, A. M., & Vaution, C. (1994). Vesicular systems and multiple emulsions in cosmetology. *Cosmetic dermatology. Martin Dunitz, Baltimore*, 27-36.
- Severino, P., Andreani, T., Macedo, A. S., Fanguero, J. F., Santana, M. H. A., Silva, A. M., & Souto, E. B. (2012). Current state-of-art and new trends on lipid nanoparticles (SLN and NLC) for oral drug delivery. *Journal of Drug Delivery*, 2012.

- Severino, P., Santana, M. H. A., & Souto, E. B. (2012). Optimizing SLN and NLC by 22 full factorial design: effect of homogenization technique. *Materials Science and Engineering: C*, 32(6), 1375-1379.
- Seymour, E. M., Lewis, S. K., Urcuyo-Llanes, D. E., Tanone, I. I., Kirakosyan, A., Kaufman, P. B., & Bolling, S. F. (2009). Regular tart cherry intake alters abdominal adiposity, adipose gene transcription, and inflammation in obesity-prone rats fed a high fat diet. *Journal of Medicinal Food*, 12(5), 935-942.
- Shah, P. P., Desai, P. R., Channer, D., & Singh, M. (2012). Enhanced skin permeation using polyarginine modified nanostructured lipid carriers. *Journal of Controlled Release*, 161(3), 735-745.
- Shah, P. P., Desai, P. R., Channer, D., & Singh, M. (2012). Enhanced skin permeation using polyarginine modified nanostructured lipid carriers. *Journal of Controlled Release*, 161(3), 735-745.
- Shemesh, R., Krepker, M., Goldman, D., Danin-Poleg, Y., Kashi, Y., Nitzan, N., Vaxman, A., & Segal, E. (2015). Antibacterial and antifungal LDPE films for active packaging. *Polymers for Advanced Technologies*, 26(1), 110-116.
- Shete, H., & Patravale, V. (2013). Long chain lipid based tamoxifen NLC. Part I: preformulation studies, formulation development and physicochemical characterization. *International Journal of Pharmaceutics*, 454(1), 573-583.
- Sikka, S. C., & Bartolome, A. R. (2018). Perfumery, Essential Oils, and Household Chemicals Affecting Reproductive and Sexual Health. In *Bioenvironmental Issues Affecting Men's Reproductive and Sexual Health* (pp. 557-569). Academic Press.
- Simon, J. A., & ESTRASORB Study Group. (2006). Estradiol in micellar nanoparticles: the efficacy and safety of a novel transdermal drug-delivery technology in the management of moderate to severe vasomotor symptoms. *Menopause*, 13(2), 222-231.
- Singh, A. B., Singh, N., Maurya, R., & Srivastava, A. K. (2009). Anti-hyperglycaemic, lipid lowering and anti-oxidant properties of [6]-gingerol in db/db mice. *International Journal of Medicine and Medical Sciences*, 1(12), 536-544.
- Singh, S., Dobhal, A. K., Jain, A., Pandit, J. K., & Chakraborty, S. (2010). Formulation and evaluation of solid lipid nanoparticles of a water soluble drug: zidovudine. *Chemical and Pharmaceutical Bulletin*, 58(5), 650-655.

- Smith, R. J., Lotya, M., & Coleman, J. N. (2010). The importance of repulsive potential barriers for the dispersion of graphene using surfactants. *New Journal of Physics*, 12(12), 125008.
- Söhle, J., Knott, A., Holtzmann, U., Siegner, R., Grönniger, E., Schepky, A., Gallinat, S., Wenck, H., Stab, F., & Winnefeld, M. (2009). White Tea extract induces lipolytic activity and inhibits adipogenesis in human subcutaneous (pre)-adipocytes. *Nutrition & Metabolism*, 6(1), 20.
- Soltani, S., & Madadlou, A. (2015). Gelation characteristics of the sugar beet pectin solution charged with fish oil-loaded zein nanoparticles. *Food Hydrocolloids*, 43, 664-669.
- Somchit, M. N., Bustamam, A. A., & Zuraini, A. (2005). Anti-pyretic and Analgesic Activity of *Zingiber zerumbet*. *International Journal of Pharmacology*, 1(3), 277-280.
- Souto, E. B., & Müller, R. H. (2005). SLN and NLC for topical delivery of ketoconazole. *Journal of Microencapsulation*, 22(5), 501-510.
- Sparavigna, A., Vesnaver, R., Cenni, A., & Oliva, M. (2012). Cosmeceutical slimming cream to treat localised subcutaneous adiposities.
- Sternby, B., Hartmann, D., BORGSTROÖM, B., & Nilsson, Å. (2002). Degree of in vivo inhibition of human gastric and pancreatic lipases by Orlistat (Tetrahydrolipstatin, THL) in the stomach and small intestine. *Clinical Nutrition*, 21(5), 395-402.
- Suttiponparnit, W., & Morgan, J. J. (2012). *Aquatic chemistry: chemical equilibria and rates in natural waters* (Vol. 126). John Wiley & Sons.
- Su, R., Fan, W., Yu, Q., Dong, X., Qi, J., Zhu, Q., Zhao, W., Wu, W., Chen, Z., Li, Y., & Lu, Y. (2017). Size-dependent penetration of nanoemulsions into epidermis and hair follicles: implications for transdermal delivery and immunization. *Oncotarget*, 8(24), 38214.
- Suk, S., Seo, S. G., Yu, J. G., Yang, H., Jeong, E., Jang, Y. J., Yaghmoor, S. S., Ahmed, Y., Yousef, J. M., Abualnaja, K. O., & Al-Malki, A. L. (2016). A Bioactive Constituent of Ginger, 6-Shogaol, Prevents Adipogenesis and Stimulates Lipolysis in 3T3-L1 Adipocytes. *Journal of Food Biochemistry*, 40(1), 84-90.

- Sun, L., Cun, D., Yuan, B., Cui, H., Xi, H., Mu, L., Chen, Y., Liu, C., Wang, Z., & Fang, L. (2012). Formulation and in vitro/in vivo correlation of a drug-in-adhesive transdermal patch containing azasetron. *Journal of Pharmaceutical Sciences*, 101(12), 4540-4548.
- Sun, M., Nie, S., Pan, X., Zhang, R., Fan, Z., & Wang, S. (2014). Quercetin-nanostructured lipid carriers: characteristics and anti-breast cancer activities *in vitro*. *Colloids and Surfaces B: Biointerfaces*, 113, 15-24.
- Suttiponparnit, K., Jiang, J., Sahu, M., Suvachittanont, S., Charinpanitkul, T., & Biswas, P. (2011). Role of surface area, primary particle size, and crystal phase on titanium dioxide nanoparticle dispersion properties. *Nanoscale Research Letters*, 6(1), 27.
- Tabatt, K., Sameti, M., Olbrich, C., Müller, R. H., & Lehr, C. M. (2004). Effect of cationic lipid and matrix lipid composition on solid lipid nanoparticle-mediated gene transfer. *European Journal of Pharmaceutics and Biopharmaceutics*, 57(2), 155-162.
- Takeuchi, H., Ishida, M., Furuya, A., Todo, H., Urano, H., & Sugibayashi, K. (2012). Influence of skin thickness on the in vitro permeabilities of drugs through Sprague-Dawley rat or Yucatan micropig skin. *Biological and Pharmaceutical Bulletin*, 35(2), 192-202.
- Takeuchi, H., Mano, Y., Terasaka, S., Sakurai, T., Furuya, A., Urano, H., & Sugibayashi, K. (2011). Usefulness of rat skin as a substitute for human skin in the *in vitro* skin permeation study. *Experimental Animals*, 60(4), 373-384.
- Tamjidi, F., Shahedi, M., Varshosaz, J., & Nasirpour, A. (2013). Nanostructured lipid carriers (NLC): A potential delivery system for bioactive food molecules. *Innovative Food Science & Emerging Technologies*, 19, 29-43.
- Tan, Y. J., Lee, C. S., Er, H. M., Lim, W. H., & Wong, S. F. (2016). In-vitro evaluation of griseofulvin loaded lipid nanoparticles for topical delivery. *Journal of Drug Delivery Science and Technology*, 31, 1-10.
- Tansey, J. T., Sztalryd, C., Gruia-Gray, J., Roush, D. L., Zee, J. V., Gavrilova, O., Reitman, M. L., Deng, C. X., Li, C., Kimmel, A. R., & Londos, C. (2001). Perilipin ablation results in a lean mouse with aberrant adipocyte lipolysis, enhanced leptin production, and resistance to diet-induced obesity. *Proceedings of the National Academy of Sciences*, 98(11), 6494-6499.

- Tardugno, R., Spagnoletti, A., Grandini, A., Maresca, I., Sacchetti, G., Pellati, F., & Benvenuti, S. (2018). Chemical profile and biological activities of Cedrelopsis grevei H. Baillon bark essential oil. *Plant Biosystems-An International Journal Dealing with all Aspects of Plant Biology*, 152(1), 120-129.
- Ternullo, S., de Weerd, L., Holsæter, A. M., Flaten, G. E., & Škalko-Basnet, N. (2017). Going skin deep: a direct comparison of penetration potential of lipid-based nanovesicles on the isolated perfused human skin flap model. *European Journal of Pharmaceutics and Biopharmaceutics*, 121, 14-23.
- Tiwari, B. K., Valdramidis, V. P., O'Donnell, C. P., Muthukumarappan, K., Bourke, P., & Cullen, P. J. (2009). Application of natural antimicrobials for food preservation. *Journal of Agricultural and Food Chemistry*, 57(14), 5987-6000.
- Tremblay, A., & Pérusse, L. (2017). Obesity, genes, and sleep habits. *The American Journal of Clinical Nutrition*, 105 (4), 779–780.
- Trigueros, L., Peña, S., Ugidos, A. V., Sayas-Barberá, E., Pérez-Álvarez, J. A., & Sendra, E. (2013). Food ingredients as anti-obesity agents: a review. *Critical Reviews in Food Science and Nutrition*, 53(9), 929-942.
- Turek, C., & Stintzing, F. C. (2013). Stability of essential oils: a review. *Comprehensive Reviews in Food Science and Food Safety*, 12(1), 40-53.
- Tzeng, T. F., & Liu, I. M. (2013). 6-Gingerol prevents adipogenesis and the accumulation of cytoplasmic lipid droplets in 3T3-L1 cells. *Phytomedicine*, 20(6), 481-487.
- U.S. Environmental Protection Agency (EPA), 1998. OPPTS 870.2500 and OPPTS 870.2600, Acute Dermal Irritation and Skin Sensitization.
- Uchida, T., Kadhum, W. R., Kanai, S., Todo, H., Oshizaka, T., & Sugibayashi, K. (2015). Prediction of skin permeation by chemical compounds using the artificial membrane, Strat-M™. *European Journal of Pharmaceutical Sciences*, 67, 113-118.
- Üstündağ-Okur, N., Gökçe, E. H., Bozbıyık, D. I., Eğrilmez, S., Özer, Ö., & Ertan, G. (2014). Preparation and in vitro–in vivo evaluation of ofloxacin loaded ophthalmic nano structured lipid carriers modified with chitosan oligosaccharide lactate for the treatment of bacterial keratitis. *European Journal of Pharmaceutical Sciences*, 63, 204-215.

- Vanić, Ž., & Škalco-Basnet, N. (2014). Mucosal nanosystems for improved topical drug delivery: Vaginal route of administration. *Journal of Drug Delivery Science and Technology*, 24(5), 435-444.
- Verma, D. D., Verma, S., Blume, G., & Fahr, A. (2003). Particle size of liposomes influences dermal delivery of substances into skin. *International Journal of Pharmaceutics*, 258(1-2), 141-151.
- Vignati, E., Piazza, R., & Lockhart, T. P. (2003). Pickering emulsions: interfacial tension, colloidal layer morphology, and trapped-particle motion. *Langmuir*, 19(17), 6650-6656.
- Villena, J. A., Roy, S., Sarkadi-Nagy, E., Kim, K. H. and Sul, H. S. (2004). Desnutrin, an adipocyte gene encoding a novel patatin domain-containing protein, is induced by fasting and glucocorticoids ectopic expression of desnutrin increases triglyceride hydrolysis. *Journal of Biological Chemistry*, 279(45), 47066-47075.
- Vitorino, C., Almeida, A., Sousa, J., Lamarche, I., Gobin, P., Marchand, S., Couet, W., Olivier, J. C., & Pais, A. (2014). Passive and active strategies for transdermal delivery using co-encapsulating nanostructured lipid carriers: in vitro vs. in vivo studies. *European Journal of Pharmaceutics and Biopharmaceutics*, 86(2), 133-144.
- Wallace, S. J., Li, J., Nation, R. L., & Boyd, B. J. (2012). Drug release from nanomedicines: selection of appropriate encapsulation and release methodology. *Drug Delivery and Translational Research*, 2(4), 284-292.
- Wang, H., & Ng, T. B. (2005). An antifungal protein from ginger rhizomes. *Biochemical and Biophysical Research Communications*, 336(1), 100-104.
- Wang, J., Ke, W., Bao, R., Hu, X., & Chen, F. (2017). Beneficial effects of ginger *Zingiber officinale* Roscoe on obesity and metabolic syndrome: a review. *Annals of the New York Academy of Sciences*, 1398(1), 83-98.
- Wang, W., Li, C. Y., Wen, X. D., Li, P., & Qi, L. W. (2009). Simultaneous determination of 6-gingerol, 8-gingerol, 10-gingerol and 6-shogaol in rat plasma by liquid chromatography–mass spectrometry: Application to pharmacokinetics. *Journal of Chromatography B*, 877(8-9), 671-679.
- Weisberg, S. P., Leibel, R., & Tortoriello, D. V. (2008). Dietary curcumin significantly improves obesity-associated inflammation and diabetes in mouse models of diabetes. *Endocrinology*, 149(7), 3549-3558.

- Wiedersberg, S., & Guy, R. H. (2014). Transdermal drug delivery: 30+ years of war and still fighting!. *Journal of Controlled Release*, 190, 150-156.
- Wikantyasning, E. R., Andasari, S. D., Da'i, M., & Nabila, A. (2017, October). Nanoencapsulation of Zerumbone in Oleic Acid-Modified Chitosan Nanoparticles. In *Proceedings of the 2017 International Conference on Computational Biology and Bioinformatics* (pp. 105-109). ACM.
- Williams, A. C., & Barry, B. W. (2012). Penetration enhancers. *Advanced Drug Delivery Reviews*, 64, 128-137.
- Wirth, A., & Steinmetz, B. (1998). Gender differences in changes in subcutaneous and intra-abdominal fat during weight reduction: an ultrasound study. *Obesity Research*, 6(6), 393-399.
- Wissing, S. A., & Müller, R. H. (2003). Cosmetic applications for solid lipid nanoparticles (SLN). *International Journal of Pharmaceutics*, 254(1), 65-68.
- Wissing, S. A., Kayser, O., & Müller, R. H. (2004). Solid lipid nanoparticles for parenteral drug delivery. *Advanced Drug Delivery Reviews*, 56(9), 1257-1272.
- Wokovich, A. M., Prodduturi, S., Doub, W. H., Hussain, A. S., & Buhse, L. F. (2006). Transdermal drug delivery system (TDDS) adhesion as a critical safety, efficacy and quality attribute. *European Journal of Pharmaceutics and Biopharmaceutics*, 64(1), 1-8.
- Wong, T. W. (2014). Electrical, magnetic, photomechanical and cavitation waves to overcome skin barrier for transdermal drug delivery. *Journal of Controlled Release*, 193, 257-269.
- Xavier, A. C. F., de Moraes, M. L., & Ferreira, M. (2013). Immobilization of aloin encapsulated into liposomes in layer-by-layer films for transdermal drug delivery. *Materials Science and Engineering: C*, 33(3), 1193-1196.
- Xia, Q., Hao, X., Lu, Y., Xu, W., Wei, H., Ma, Q., & Gu, N. (2008). Production of drug-loaded lipid nanoparticles based on phase behaviors of special hot microemulsions. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 313, 27-30.
- Yang, T. S., Liu, T. T., & Liu, H. I. (2017). Effects of aroma compounds and lipid composition on release of functional substances encapsulated in nanostructured lipid carriers lipolyzed by lipase. *Food Hydrocolloids*, 62, 280-287.

- Yang, X., Zhao, L., Almasy, L., Garamus, V. M., Zou, A., Willumeit, R., & Fan, S. (2013). Preparation and characterization of 4-dedimethylamino sancycline (CMT-3) loaded nanostructured lipid carrier (CMT-3/NLC) formulations. *International Journal of Pharmaceutics*, 450(1-2), 225-234.
- Yeh, H. Y., Chuang, C. H., Chen, H. C., Wan, C. J., Chen, T. L., & Lin L. Y. (2014). Bioactive components analysis of two various gingers (*Zingiber officinale* Roscoe) and antioxidant effect of ginger extracts. *Food Science and Technology*, 55(1), 329-334.
- Yen, H. F., Hsieh, C. T., Hsieh, T. J., Chang, F. R., & Wang, C. K. (2015). In vitro anti-diabetic effect and chemical component analysis of 29 essential oils products. *Journal of Food and Drug Analysis*, 23(1), 124-129.
- Yob, N. J., Jofrry, S. M., Affandi, M. M. R., Teh, L. K., Salleh, M. Z., & Zakaria, Z. A. (2011). *Zingiber zerumbet* (L.) Smith: a review of its ethnomedicinal, chemical, and pharmacological uses. *Evidence-Based Complementary and Alternative Medicine*, 2011.
- Yulvianti, M., Barleany, D. R., & Ernayati, W. (2015). Encapsulation Red Ginger Oleoresin (" *Zingiber officinale*" var. *Rubrum*) with Chitosan-alginate as Wall Material Using Spray Drying. *Research Journal of Applied Sciences, Engineering and Technology*, 10(12), 1370-1378.
- Yun, J. W. (2010). Possible anti-obesity therapeutics from nature—A review. *Phytochemistry*, 71(14-15), 1625-1641.
- Zadini, F., & Ratiner, B. (2015). *U.S. Patent No. 8,952,067*. Washington, DC: U.S. Patent and Trademark Office.
- Zanqui, A. B., de Moraes, D. R., da Silva, C. M., Santos, J. M., Gomes, S. T. M., Visentainer, J. V., Eberlin, M. N., Cardozo-Filho, L., & Matsushita, M. (2015). Subcritical extraction of flaxseed oil with n-propane: composition and purity. *Food Chemistry*, 188, 452-458.
- Zhai, H., & Maibach, H. I. (2001). Effects of skin occlusion on percutaneous absorption: an overview. *Skin Pharmacology and Physiology*, 14(1), 1-10.
- Zhang, H. L., Zhong, X. B., Tao, Y., Wu, S. H., & Su, Z. Q. (2012). Effects of chitosan and water-soluble chitosan micro-and nanoparticles in obese rats fed a high-fat diet. *International Journal of Nanomedicine*, 7, 4069.
- Zhang, J., Fan, Y., & Smith, E. (2009). Experimental design for the optimization of lipid nanoparticles. *Journal of Pharmaceutical Sciences*, 98(5), 1813-1819.

- Zhang, J., Xiao, X., Dong, Y., Shi, L., Xu, T., & Wu, F. (2017). The anti-obesity effect of fermented barley extracts with *Lactobacillus plantarum* dy-1 and *Saccharomyces cerevisiae* in diet-induced obese rats. *Food & Function*, 8(3), 1132-1143.
- Zhang, W. L., Zhu, L., & Jiang, J. G. (2014). Active ingredients from natural botanicals in the treatment of obesity. *Obesity Reviews*, 15(12), 957-967.
- Zhang, W., Li, X., Ye, T., Chen, F., Sun, X., Kong, J., Yang, X., Pan, W., & Li, S. (2013). Design, characterization, and in vitro cellular inhibition and uptake of optimized genistein-loaded NLC for the prevention of posterior capsular opacification using response surface methodology. *International Journal of Pharmaceutics*, 454(1), 354-366.
- Zhang, Y. T., Shen, L. N., Wu, Z. H., Zhao, J. H., & Feng, N. P. (2014). Evaluation of skin viability effect on ethosome and liposome-mediated psoralen delivery via cell uptake. *Journal of Pharmaceutical Sciences*, 103(10), 3120-3126.
- Zheng, M., Falkeborg, M., Zheng, Y., Yang, T., & Xu, X. (2013). Formulation and characterization of nanostructured lipid carriers containing a mixed lipids core. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 430, 76-84.
- Zimmermann, R., Strauss, J.G., Haemmerle, G., Schoiswohl, G., Birner-Gruenberger, R., Riederer, M., Lass, A., Neuberger, G., Eisenhaber, F., Hermetter, A., & Zechner, R. (2004). Fat mobilization in adipose tissue is promoted by adipose triglyceride lipase. *Science*, 306(5700), 1383-1386.
- zur Mühlen, A., Schwarz, C., & Mehnert, W. (1998). Solid lipid nanoparticles (SLN) for controlled drug delivery—drug release and release mechanism. *European Journal of Pharmaceutics and Biopharmaceutics*, 45(2), 149-155.