



## Article (refereed) - postprint

---

Peacock, M.; Jones, T.G.; Airey, B.; Johncock, A.; Evans, C.D.; Lebron, I; Fenner, N.; Freeman, C. 2015. **The effect of peatland drainage and rewetting (ditch blocking) on extracellular enzyme activities and water chemistry**. *Soil Use and Management*, 31 (1). 67-76. [10.1111/sum.12138](https://doi.org/10.1111/sum.12138)

© 2014 British Society of Soil Science

This version available <http://nora.nerc.ac.uk/512937/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

**This document is the author's final manuscript version of the journal article, incorporating any revisions agreed during the peer review process. There may be differences between this and the publisher's version. You are advised to consult the publisher's version if you wish to cite from this article.**

The definitive version is available at <http://onlinelibrary.wiley.com/>

Contact CEH NORA team at  
[noraceh@ceh.ac.uk](mailto:noraceh@ceh.ac.uk)

1            **The effect of peatland drainage and rewetting (ditch blocking) on**  
2            **extracellular enzyme activities and water chemistry**

3 Mike Peacock<sup>ac</sup>, Timothy G. Jones<sup>a</sup>, Belinda Airey<sup>a</sup>, Anna Johncock<sup>a</sup>, Chris D. Evans<sup>b</sup>, Inma  
4 Lebron<sup>b</sup>, Nathalie Fenner<sup>a</sup>, Chris Freeman<sup>a</sup>.

5  
6 <sup>a</sup>Wolfson Carbon Capture Laboratory, School of Biological Sciences, Bangor University,  
7 Deiniol Road, LL57 2UW, UK.

8 <sup>b</sup>Centre for Ecology and Hydrology, Environment Centre Wales, Deiniol Road, LL57 2UW,  
9 UK.

10 <sup>c</sup>Corresponding author. Centre for Earth, Planetary, Space and Astronomical Research  
11 (CEPSAR), Department of Environment, Earth and Ecosystems, The Open University,  
12 Walton Hall, Milton Keynes, MK7 6AA. mikepeacocknin@yahoo.co.uk

13  
14 **Abstract**

15 Extensive areas of European peatlands have been drained by digging ditches in an attempt to  
16 improve the land, resulting in increased carbon dioxide fluxes to the atmosphere and  
17 enhanced fluvial dissolved organic carbon (DOC) concentrations. Numerous peatland  
18 restoration projects have been initiated which aim to raise water tables by ditch blocking, thus  
19 reversing drainage-induced carbon losses. It has been suggested that extracellular hydrolase  
20 and phenol oxidase enzymes are partly responsible for controlling peatland carbon dynamics,  
21 and that these enzymes are affected by environmental change. The aim of this study was to  
22 investigate how drainage and ditch blocking affect enzyme activities and water chemistry in a  
23 Welsh blanket bog, and to study the relationship between enzyme activity and water  
24 chemistry. A comparison of a drained and undrained site showed that the drained site had  
25 higher phenol oxidase and hydrolase activities, and lower concentrations of phenolic

26 compounds which inhibit hydrolase enzymes. Ditch blocking had little impact upon enzyme  
27 activities; although hydrolase activities were lowered 4-9 months after restoration, the only  
28 significant difference was for arylsulphatase activity. Finally, we noted a negative  
29 correlation between  $\beta$ -glucosidase activity and DOC concentrations, and a positive  
30 correlation between arylsulphatase activity and sulphate concentration. Phenol oxidase  
31 activity was negatively correlated with DOC concentrations in pore water, but for ditch water  
32 phenol oxidase correlated negatively with the ratio of phenolics to DOC. Our results imply  
33 that drainage could exacerbate gaseous and fluvial carbon losses, and that peatland  
34 restoration may not reverse the effects, at least in the short term.

35

36 Key words: ditch blocking, peatland restoration, phenol oxidase,  $\beta$ -glucosidase, dissolved  
37 organic carbon, phenolics,

38

### 39 **1. Introduction**

40 Northern peatlands are important carbon stores, but many have been drained for  
41 forestry, agriculture, and peat harvesting. In the UK drainage ditches were predominantly  
42 dug during the 19<sup>th</sup> and 20<sup>th</sup> centuries. The size and spacing of ditches varies but in UK  
43 blanket bogs they are typically around 0.5 m deep, with 7-20 m spacing (Stewart & Lance,  
44 1991). It has been suggested that blanket bogs are somewhat resistant to drainage, with water  
45 table drawdown occurring only in the immediate vicinity of ditches (Stewart & Lance, 1991),  
46 and the magnitude of drawdown will depend on ditch spacing and the hydraulic conductivity  
47 of the peat (Armstrong, 2000). Nevertheless, long-term drainage can lead to the  
48 establishment of deeper water tables (Holden *et al.*, 2011), and even slight changes in water  
49 tables can have ecological effects (Price *et al.*, 2003).

50 Blanket bogs are largely ombrotrophic, and often found at the headwaters of river  
51 catchments, making them sources of potable water as well as sources of dissolved organic  
52 carbon (DOC) (Hope *et al.*, 1999). The quality of water draining these systems thus has  
53 relevance for aquatic ecosystems (Karlsson *et al.*, 2009), water treatment (McDonald *et al.*,  
54 1991), and human health issues (Chow *et al.*, 2003). DOC is a natural export from  
55 peatlands, but there is evidence that DOC concentrations are higher in drained bogs (Glatzel  
56 *et al.*, 2003, Wallage *et al.*, 2006). The drainage of ombrotrophic bogs generally leads to an  
57 increase in carbon dioxide (CO<sub>2</sub>) emissions and a decrease in methane (CH<sub>4</sub>) emissions  
58 (Bussell *et al.*, 2010).

59 In an attempt to reverse these drainage-induced biogeochemical changes, numerous  
60 peatland restoration projects have been initiated. Sites that have been ditched are restored by  
61 blocking the ditches with dams. The aim is to return the water table to pre-drainage levels.  
62 Some success has been observed on blanket bog; 6-7 years after rewetting, Holden *et al.*  
63 (2011) observed that a ditch-blocked site had hydrological functioning intermediate between  
64 an undrained site and drained site. Similarly, Wilson *et al.* (2011a) and Worrall *et al.* (2007)  
65 both noted increases in the water table after blocking

66 One aspect of drainage that has received little attention is the activity of soil  
67 extracellular enzymes. Extracellular enzymes are involved in peatland carbon cycling  
68 (Freeman *et al.*, 1997) but their activities are constrained by the conditions that exist in peat  
69 soils. Recalcitrant phenolic compounds are released by plants (Wetzel, 1992) and degraded  
70 by phenol oxidase, which has limited activity in northern peatlands due to the acidic pH, low  
71 temperatures and low oxygen content (Pind *et al.*, 1994, Freeman *et al.*, 2001a, Tahvanainen  
72 & Haraguchi, 2013). The build-up of phenolics in turn inhibits the activity of hydrolase  
73 enzymes (Freeman *et al.*, 1990, Wetzel, 1992); resulting in low rates of decomposition.  
74 Conversely, increased peat aeration stimulates phenol oxidase activity, lowers phenolic

75 concentrations, and removes the inhibitory effect on hydrolase enzymes (Freeman *et al.*,  
76 2001a). It can therefore be hypothesised that long-term drainage would lead to increased  
77 phenol oxidase activity, reduced phenolic concentrations and increased hydrolase activity,  
78 thereby resulting in greater overall soil decomposition rates and contributing to carbon loss  
79 (*hypothesis 1*). Theoretically, ditch blocking would reverse this by raising the water table,  
80 and leading to suppressed phenol oxidase activity, increased phenolic concentrations and  
81 reduced hydrolase enzyme activity (*hypothesis 2*). The aim of this study was to test these  
82 hypotheses using two sites located within a large peatland. A further aim was to examine  
83 enzyme activities and to determine if they were related to DOC or phenolic concentrations, as  
84 past studies have shown contradictory results.

85

## 86 **2. Materials and Methods**

### 87 *2.1. Study sites*

88 The study was carried out on the Migneint blanket bog, North Wales (UK).  
89 According to the JNCC National Vegetation Classification (NVC), it includes areas of mire  
90 habitat of classes M18, M19 and M20. Mean annual rainfall is 2.2 m and mean annual  
91 temperature 5.6 °C (Billett *et al.*, 2010).

92 The primary field site was the Afon Ddu catchment (latitude 52.99 N, longitude 3.82  
93 W, 490 m above sea level) which was drained during the 1970s and 1980s. The ditches run  
94 downslope and were blocked in February 2011. A replicated experiment was established in  
95 August 2010 which comprises four ditches that have been left open as controls, and eight that  
96 have been blocked using two different methods. Four have been blocked using peat dams,  
97 for which the peat is extracted from ‘borrow pits’ adjacent to each ditch. The other four have  
98 been blocked using a reprofiling technique, which involves the ditch vegetation being  
99 removed, and the peat bottom being compressed to destroy any natural pipes that may be

100 present. The ditch is then infilled with peat from borrow pits and the vegetation is replaced.  
101 As in the previous treatment peat dams are also constructed along the ditch.

102 A second nearby field site, was used to provide a comparison with undrained  
103 conditions; the Bryn Du site (latitude 52.97 N, longitude 3.82 W, 460 m above sea level)  
104 includes four control plots on intact blanket bog that has not been drained.

105

## 106 *2.2. Soil Sampling*

107 At the Afon Ddu soil samples were taken from each of the twelve ditches in June,  
108 July, August, September and November 2011. These samples were used to test the effect of  
109 ditch blocking on enzyme activities. Additional soil samples were taken from areas of bog  
110 between ditches to examine the effects of enzyme activities on DOC and phenolic  
111 concentrations. At Bryn Du, soil samples were taken from each of the four control plots in  
112 June and September 2011. All soil samples were taken to 10 cm depth. Each soil sample  
113 comprised 2-4 sub-samples of soil (taken from an area of approximately 1 m<sup>2</sup>) to minimise  
114 the influence of small-scale spatial variation in enzyme activity. Samples were stored in the  
115 dark at 4°C. Soil water content was determined by weighing 1 g of sample, drying for 24  
116 hours at 105°C and re-weighing.

117

## 118 *2.3. Water sampling and water tables*

119 Water samples were taken from the ditches at the Afon Ddu and from piezometers 2-3  
120 m adjacent to ditches (i.e. water and soil samples were taken from approximately the same  
121 locations for 'ditch' and 'bog' samples). Piezometers were constructed from PVC pipe with  
122 intakes at 10-15 cm depth. Water samples at Bryn Du were extracted using Rhizon samplers  
123 (Rhizosphere Research Products) at a depth of 10 cm. Water samples were collected in 60 ml  
124 Nalgene ® bottles and were stored in the dark at 4°C.

125 Water tables were measured using dipwells constructed from PVC pipe; for each  
126 ditch, a dipwell was positioned 2 m either side of the ditch. Water tables were manually  
127 recorded on an approximately monthly basis from April to November 2011. Dipwell length  
128 was 1000 mm. Every 100 mm, four drilled holes of 8 mm diameter were evenly spaced  
129 around the pipe to allow water entry.

130

#### 131 *2.4. Laboratory analysis*

132 Phenol oxidase activity was measured using a method modified from Pind *et al.*  
133 (1994), using 1 cm<sup>3</sup> of soil. Analysis of hydrolase activity was measured using a method  
134 modified from Freeman *et al.* (1995), using 1 cm<sup>3</sup> of soil. Further information concerning the  
135 enzyme assays can be found in Dunn *et al.* (2013).

136 Water samples were filtered at 0.45 µm. Ion concentrations were determined using  
137 either a DX-120 Ion Chromatograph (Dionex), or an 850 Professional IC (Metrohm). DOC  
138 concentrations were analysed using a Thermalox Total Carbon analyser (Analytical  
139 Sciences). Phenolic concentrations were determined using a method adapted from Box  
140 (1983), and were derived from a standard curve using phenol standards.

141

#### 142 *2.5. Statistical analysis*

143 Statistical analysis was carried out using SPSS v16.0.1 (IBM Corporation). The  
144 Shapiro-Wilk test was used to test the normality of data, and log 10 or square root  
145 transformations were attempted on any data that failed this. For the comparisons of the  
146 drained and undrained site, t-tests were used, or the non-parametric Mann-Whitney test (for  
147 any data that could not be transformed to normality). To compare unblocked ditches to the  
148 two ditch blocking treatments, repeated-measures ANOVAs with Tukey HSD post-hoc tests  
149 were carried out. If transformations failed to produce normal data, then the non-parametric

150 Kruskal-Wallis test was used. Linear regression was used to test for relationships between  
151 variables

152

### 153 **3. Results**

#### 154 *3.1. Site comparison – effect of long term drainage*

155 A comparison of the Bryn Du data with that from the open ditches at the Afon Ddu  
156 shows that the drained site had higher hydrolase (driven by arylsulphatase and  $\beta$ -glucosidase)  
157 and phenol oxidase activity (Figure 1 and 2). Additionally, Bryn Du displays a significantly  
158 higher phenolic concentration;  $5.6 \text{ mg L}^{-1}$  compared with  $4.8 \text{ mg L}^{-1}$  at the Afon Ddu (one-  
159 tailed t-test,  $p = 0.02$ ). There was no significant difference in pH; 4.27 at Bryn Du and 4.18  
160 at the Afon Ddu. Despite the significant difference in arylsulphatase activity, there was no  
161 significant difference in pore water sulphate concentrations between the two sites: mean  
162 concentrations for the period March-November 2011 (monthly sampling,  $n = 4$  per site) were  
163  $2.2 \text{ mg L}^{-1}$  at the Afon Ddu, and  $1.0 \text{ mg L}^{-1}$  at Bryn Du (with respective standard errors of 0.8  
164  $\text{mg L}^{-1}$  and  $0.5 \text{ mg L}^{-1}$ ). The only ion for which a significant difference was found was  
165 phosphate; concentrations at Bryn Du were often below the detection limit of the analyser  
166 (Table 1). There was no significant difference in the water content of soil samples (91.0%,  
167  $\text{SE} = 0.6\%$  at Bryn Du, 90.7%,  $\text{SE} = 0.8\%$  at the Afon Ddu).

168

#### 169 *3.2. Effect of ditch blocking on enzyme activity and phenolic compounds*

170 At the Afon Ddu experimental site 4-9 months after ditch-blocking, there was no  
171 significant difference between treatments for the activity of  $\beta$ -glucosidase, xylosidase or  
172 chitinase. There was a significant difference for arylsulphatase; activity was higher in the  
173 control ditches compared to the reprofiled ditches (Figure 3). Sulphate concentrations were



174 lowest for reprofiled ditches ( $1.8 \text{ mg L}^{-1}$  compared to  $2.2 \text{ mg L}^{-1}$  for open ditches and  $2.5 \text{ mg}$   
175  $\text{L}^{-1}$  for dammed ditches) but this difference was not significant.

176 There was no significant treatment effect on phenol oxidase activity (Figure 4).

177 There was no significant difference in ditch water pH between treatments; mean values for  
178 the length of the study were 4.21 (open), 4.34 (dam) and 4.20 (reprofiled). The depth to the  
179 water table was greatest for open ditches, with a mean of 14.8 cm (SE = 1.1 cm, min = 1.2  
180 cm, max = 46.5 cm) for the study period. Mean depth to the water table was 10.7 cm (SE =  
181 0.8 cm, min = 1.8 cm, max = 28.7 cm) for dammed ditches, and 9.9 cm (SE = 0.7 cm, min = -  
182 1.9 cm, max = 23.9 cm) for reprofiled ditches ( $n = 80$  for each treatment). The difference in  
183 water tables between open and blocked ditches was significant ( $p < 0.01$ ). Mean soil water  
184 content of samples was 90.7% (open, SE = 0.4%), 89.2% (dam, SE = 0.7%) and 88.1%  
185 (reprofiled, SE = 0.6%). Repeated-measures ANOVA showed no significant difference in  
186 mean water content. There was no significant difference between treatments for phenolic or  
187 DOC concentrations (Figures 5 and 6).

188

### 189 *3.3. Enzymatic controls on biogeochemistry*

190 A significant negative relationship was found between  $\beta$ -glucosidase activity and  
191 DOC concentration in both ditch and pore waters (Figure 7). No direct relationship was  
192 found between either phenol oxidase activity and DOC ( $r^2 = 0.02$ ) or phenol oxidase and  
193 phenolics ( $r^2 = 0.09$ ) for ditch water, but there was a significant negative relationship between  
194 the phenolic to DOC ratio and phenol oxidase activity (Figure 8). For pore water this was not  
195 the case; there was no correlation between phenolic to DOC ratio and phenol oxidase activity  
196 ( $r^2 = 0.05$ ), and the strongest relationship (highest  $r^2$  value) was between phenol oxidase  
197 activity and DOC concentration (Figure 8). There was a weak positive correlation between  
198 arylsulphatase activity and sulphate concentration in ditch water (Figure 9).

199

## 200 **4. Discussion**

### 201 *4.1. Effects of long term drainage*

202           Results from a comparison between an undrained site and a drained site support  
203 hypothesis 1; that drainage leads to lower phenolic concentrations, and enhanced activities of  
204 phenol oxidase and hydrolases. This is in agreement with Freeman *et al.* (2001a), who  
205 showed that increased oxygen availability following drainage stimulates phenol oxidase  
206 activity, which in turn degrades phenolics and removes the inhibition on hydrolase enzymes.  
207 The enhancement of hydrolase activity was partly controlled by increased  $\beta$ -glucosidase  
208 activity, a response which has been observed before (Fenner *et al.*, 2005). Additionally,  
209 long-term drainage leads to greater water table fluctuations (Holden *et al.*, 2011) which can  
210 exacerbate the effects of seasonal drought, leading to an associated increase in oxygen  
211 availability of a magnitude to override pH controls and consequently stimulate phenol  
212 oxidase activity. As an aside, it should be noted that phenolics were measured in pore water  
213 at the undrained site and ditch water at the drained site; this will somewhat confound the  
214 results, as pore water and surface water would have some natural differences. However, this  
215 does not impinge on the enzyme data where methods were identical at both sites.

216           It is important to acknowledge that the observed differences in biogeochemistry may  
217 not have been due to drainage, as this was a limited comparison of two sites (i.e. with no data  
218 from before the Afon Ddu catchment was drained), with pseudoreplication (i.e. sampling  
219 over time) rather than true replication. The sites are close together and share the same  
220 climate and similar peat characteristics, and the only difference in pore-water ion  
221 concentration was observed for phosphate. Nevertheless, it could be that some other factor is  
222 responsible for the differences in enzyme activity.

223

#### 224 4.2. Effect of ditch blocking

225           Although ditch blocking appeared to lower the activity of each of the hydrolase  
226 enzymes studied, arylsulphatase was the only enzyme to show a statistically significant  
227 difference. As such we are unable to find support for hypothesis 2: that ditch blocking would  
228 suppress phenol oxidase activity, leading to a subsequent increase in phenolics and lowered  
229 hydrolase activities. Fenner & Freeman (2011) noted that upon rewetting after drought,  
230 phenol oxidase activity did not immediately decline, and remained high (for a period of  
231 months to years) as a legacy from the previous aerobic conditions. It should be noted that  
232 there was no significant difference in soil moisture between the blocked and open ditches,  
233 despite the fact that the depth to the water table was significantly greater around open ditches.  
234 It could be that a lack difference in soil moisture is due to the fact that water tables were  
235 relatively high for all treatments, therefore making soil moisture insensitive to ditch blocking.  
236 Additionally, Holden *et al.* (2011) suggest that ditch blocking only partially restores the  
237 hydrological functioning of blanket bog, and other evidence suggests that it could be several  
238 years before the rewetting suppresses enzyme activity (Fenner & Freeman, 2011). It might  
239 be expected that enzyme activity would increase in the reprofiled ditches due to the  
240 disturbance that this method involves; large volumes of peat are removed from the adjacent  
241 borrow pits to infill the ditch, which might theoretically allow some oxygen infiltration.  
242 However, the enzyme response was identical for the dammed ditches and the reprofiled  
243 ditches, suggesting this was not the case. As such, it may be that the ditch blocking was on  
244 wet and dense peat, and therefore very little air entered or became trapped in the peat.

245           The suppression of arylsulphatase activity in the reprofiled ditches could have  
246 repercussions on CH<sub>4</sub> fluxes. Raising the water table will alter the redox conditions and  
247 stimulate the methanogenic community, thus increasing CH<sub>4</sub> emissions (Komulainen *et al.*,  
248 1998, Urbanová *et al.*, 2011). Coupled to this, arylsulphatase releases sulphate which is

249 implicated in reduced CH<sub>4</sub> emissions when the water table falls. The suppression of  
250 arylsulphatase following ditch blocking could result in a reduced rate of sulphate production  
251 which would then contribute to the enhanced CH<sub>4</sub> fluxes (Freeman *et al.*, 1997). A weak but  
252 significant, positive relationship was found between arylsulphatase activity and sulphate  
253 concentrations in ditch water, but no significant difference in sulphate concentration was  
254 detected between treatments.

255         We observed no change in ditch water DOC concentrations immediately after ditch  
256 blocking, and this is similar to studies of blanket bogs that have noted small changes in DOC  
257 following ditch blocking (i.e. differences of approximately 1 mg L<sup>-1</sup>, e.g. Gibson *et al.*, 2009,  
258 Ramchunder *et al.*, 2012) or even small increases (e.g. Wilson *et al.*, 2011b). The lack of  
259 change in DOC concentration can be explained partly by the overall lack of response in  
260 enzyme activities. Considering that other ditch blocking studies have speculated that the  
261 action of enzymes could be involved in any restoration-induced changes in DOC dynamics  
262 (e.g. Wallage *et al.*, 2006, Worrall *et al.*, 2007), it is interesting to note that there has  
263 apparently been only one other study that investigated the response of enzymes to ditch  
264 blocking. Bonnett *et al.* (2008) compared hydrolase activities around a natural gully and  
265 around a ditch that had been blocked twelve years previously. They noted no difference in  
266 hydrolase activities in surface peat samples, but some differences at depth; for instance, β-  
267 glucosidase activity was lower around the blocked ditch at both 25 cm and 45 cm. Some  
268 studies have suggested that DOC concentrations are lowered following restoration; Wallage  
269 *et al.* noted substantially lower pore water DOC (60-70% compared to a drained site)  
270 concentrations at a blanket bog where ditch blocking had occurred 6 years previously. This  
271 could be indicative of suppressed enzyme activities in the longer term following blocking.  
272 However, another study at the same site found similar fluxes and concentrations of DOC in  
273 ditches (Armstrong *et al.*, 2010), thus adding further complexity to the issue.

274 It should be noted that the early post-restoration measurements of DOC concentration  
275 and water table that are reported here are part of a long-term experiment. It may well be that  
276 the short-term response of these variables is different to that of any long-term response.

277

#### 278 4.3. Enzymatic controls on biogeochemistry

279 For both pore water and ditch water a weak negative relationship was observed  
280 between  $\beta$ -glucosidase activity and DOC concentration. Freeman *et al.* (1997) found the  
281 same relationship for a peatland in mid Wales, and concluded that DOC represented a  
282 substrate for  $\beta$ -glucosidase, with the metabolic products then being microbially degraded  
283 under anaerobic conditions.

284 There have been conflicting reports of the effect of phenol oxidase on phenolic  
285 concentrations. Freeman *et al.* (2001a) originally showed that increased phenol oxidase  
286 activity led to decreased phenolic concentrations, a result replicated by Fenner *et al.* (2005).  
287 However, Toberman *et al.* (2008) found a positive relationship between phenol oxidase  
288 activity and phenolics, and speculated that it could be possible for phenol oxidase to partially  
289 degrade complex phenolic compounds, thus releasing smaller, soluble phenolics. We found  
290 no relationship between phenol oxidase and phenolics. However, phenolics are a component  
291 of DOC, and (because DOC concentrations vary according to season and weather events)  
292 phenolic concentrations will also fluctuate. As such, by taking the phenolic to DOC ratio (as  
293 in Peacock *et al.*, 2013) then a significant negative relationship was observed with phenol  
294 oxidase, for ditch water. This observation suggests that phenol oxidase did not absolutely  
295 lower phenolic concentrations, but that it lowered phenolic concentrations relative to total  
296 DOC concentration. For pore water this relationship was not found; instead there was a  
297 significant negative relationship between phenol oxidase activity and DOC concentration. It

298 has been suggested previously that the phenolic to DOC ratio is an important factor in  
299 enzymatic degradation (Freeman *et al.*, 1990).

300         These results suggest that the action of enzymes on DOC/phenolics is complicated,  
301 occasionally contradictory, and sometimes unrelated. Indeed, Kane *et al.* (2014) emphasise  
302 the complexity of these interactions, and point out that positive feedbacks can exist between  
303 the release of labile DOC and enzyme activities. Although the relationships reported here  
304 between enzyme activities and DOC/phenolics are only weak, this is perhaps to be expected.  
305 In a natural system there will be multiple drivers that interact in a complex way to control  
306 fluvial carbon losses, with enzymes playing only a small part in the overall system.

307         It is useful to consider that drainage in this context can be used as an analogue for a  
308 prolonged drought event. Climate change in Europe is likely to result in more frequent and  
309 prolonged droughts (Alcamo *et al.*, 2007). Our findings thus agree with others (e.g. Freeman  
310 *et al.*, 2001a, Fenner & Freeman, 2011) in suggesting that future climate change may  
311 stimulate the activities of phenol oxidase,  $\beta$ -glucosidase and arylsulphatase. These changes  
312 could result in enhanced losses of gaseous and fluvial carbon from peatlands, although the  
313 increased activity of arylsulphatase in the drained site might be expected to suppress CH<sub>4</sub>  
314 fluxes (Freeman *et al.*, 2007). As a proxy for a recovery from severe drought, our data show  
315 that the activity of carbon-cycling enzymes remain high as a legacy of the previous aerobic  
316 conditions. The only significant change was a reduction in arylsulphatase activity in  
317 reprofiled ditches, which might therefore contribute to the enhanced CH<sub>4</sub> fluxes that are  
318 sometimes seen following ditch blocking (e.g. Green *et al.*, 2014, Cooper *et al.*, 2014).

319

#### 320 *4.4. Conclusions*

321         Our results suggest that drainage increased enzyme activity, specifically phenol  
322 oxidase,  $\beta$ -glucosidase and arylsulphatase. Enhanced activities of these enzymes could result

323 in increased losses of greenhouse gases (Freeman *et al.*, 2001a) and DOC (Freeman *et al.*,  
324 2001b). Following ditch blocking there was no evidence that enzyme activities were  
325 suppressed, apart from lowered arylsulphatase activities in reprofiled ditches. The absence  
326 of an effect on enzyme activities may have been due to a legacy of enhanced enzyme activity  
327 that was stimulated through drainage, combined with the absence of any post-blocking  
328 change in soil moisture. Furthermore, any changes may have been mediated by the weather  
329 during the monitoring period.

330         It is clear that long term monitoring is necessary to elucidate exactly when peatland  
331 restoration will begin to influence the activity of extracellular enzymes, as changes can create  
332 both positive and negative feedbacks to ecosystem processes (Sinsabaugh, 2010). Finally,  
333 the fact that arylsulphatase activity responded to both drainage and ditch blocking lends some  
334 evidence to suggest that it may be more sensitive to environmental change than other  
335 hydrolases.

336

### 337 **Acknowledgements**

338 The authors thank Trystan Edwards, Helen Buckingham and Andrew Roberts of the National  
339 Trust for their involvement in the project. DOC and water table data were collected as part of  
340 the UK's Defra (Department for Environment, Food and Rural Affairs) SP1202 project. This  
341 is a long-term project, and the analysis of early post-restoration data in this paper should not  
342 be taken to imply ditch blocking effects on DOC and water tables will be the same over two  
343 to four years as they are over the first year. Dr Richard Smart was involved with field data  
344 collection on SP1202. This work was written up through the assistance of a Knowledge  
345 Economy Skills Scholarship (KESS) PhD awarded to Mike Peacock. We thank two  
346 anonymous reviewers for their constructive comments.

347

348 **References**

349 Alcamo, J., Moreno, J.M., Nováky, B., Bindi, M., Corobov, R., Devoy, R.J.N.,  
350 Giannakopoulos, C., Martin, E., Olesen, J.E., Shvidenko, A. 2007: Europe. *Climate Change*  
351 *2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the*  
352 *Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, M.L. Parry,  
353 O.F. Canziani, J.P. Palutikof, P.J. van der Linden and C.E. Hanson, Eds., Cambridge  
354 University Press, Cambridge, UK, 541-580.

355

356 Armstrong, A. 2000. DITCH: a model to simulate field conditions in response to ditch levels  
357 managed for environmental aims. *Agriculture, Ecosystems and Environment*, **77**, 179-192.

358

359 Armstrong, A., Holden, J., Kay, P., Francis, B., Foulger, M., Gledhill, S., McDonald, A.T.,  
360 Walker, A., 2010. The impact of peatland drain-blocking on dissolved organic carbon loss  
361 and discolouration of water; results from a national survey. *Journal of Hydrology*, **381**, 112-  
362 120.

363

364 Billett, M.F., Charman, D.J., Clark, J.M., Evans, C.D., Evans, M.G., Ostle, N.J., Worrall, F.,  
365 Burden, A., Dinsmore, K.J., Jones, T., McNamara, N.P., Parry, L., Rowson, J.G., Rose, R.  
366 2010. Carbon balance of UK peatlands: current state of knowledge and future research  
367 challenges. *Climate Research*, **45**, 13-29.

368

369 Bonnett, S.A.F., Leah, R., Maltby, E., O'Connor, M. 2008. Short-term effectiveness of drain-  
370 blocking in suppressing enzymic peat decomposition and DOC export. Report for AONB  
371 Partnership's Peatscapes project. University of Liverpool.

372



373 Box, J.D., 1983. Investigation of the Folin-Ciocalteu phenol reagent for the determination of  
374 polyphenolic substances in natural waters. *Water Research*, **17**, 511-525.

375

376 Bussell, J., Jones, D.L., Healey, J.R., Pullin, A. 2010. How do draining and re-wetting affect  
377 carbon stores and greenhouse gas fluxes in peatland soils? CEE Review 08-012 (SR49).  
378 Collaboration for Environmental Evidence. [www.environmentalevidence.org/SR49.html](http://www.environmentalevidence.org/SR49.html).

379

380 Chow, A.T., Kanji, K.K., and Gao, K.K.T., 2003. Production of dissolved organic carbon  
381 (DOC) and trihalomethane (THM) precursor from peat soils. *Water Research*, **37**, 4475-4485.

382

383 Cooper, M.D.A., Evans, C.D., Zielinski, P., Levy, P.E., Gray, A., Peacock, M., Norris, D.,  
384 Fenner, N., Freeman, C. 2014. Infilled ditches are hotspots of landscape methane flux  
385 following peatland re-wetting. *Ecosystems*, DOI 10.1007/s10021-014-9791-3 .

386

387 Dunn, D., Jones, T.G., Girard, A., Freeman, C. 2013. Methodologies for extracellular enzyme  
388 assays from wetland soils. *Wetlands*, **34**, 9-17.

389

390 Fenner, N., Freeman, C., Reynolds, B., 2005. Hydrological effects on the diversity of  
391 phenolic degrading bacteria in a peatland: implications for carbon cycling. *Soil Biology and*  
392 *Biochemistry*, **37**, 1277-1287.

393

394 Fenner, N., and Freeman, C., 2011. Drought-induced carbon loss in peatlands. *Nature*  
395 *Geoscience*, **4**, 895-900.

396

397 Freeman, C., Lock, M.A., Marxsen, J., Jones, S.E. 1990. Inhibitory effects of high molecular

398 weight dissolved organic matter upon metabolic processes in biofilms from contrasting rivers  
399 and streams. *Freshwater Biology*, **24**, 159-166.

400

401 Freeman, C., Lock, M.A., and Reynolds, B., 1993. Fluxes of CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O from a  
402 Welsh peatland followed simulation of water table draw-down: potential feedback to climate  
403 change. *Biogeochemistry*, **19**, 51-60.

404

405 Freeman, C., Liska, G., Ostle, N.J., Jones, S.E., and Lock, M.A., 1995. The use of  
406 fluorogenic substrates for measuring enzyme activity in peatlands. *Plant and Soil*, **175**, 147-  
407 152.

408

409 Freeman, C., Liska, G., Ostle, N.J., Lock, M.A., Hughes, S., Reynolds, B., and Hudson, J.,  
410 1997. Enzymes and biogeochemical cycling in wetlands during a simulated drought.  
411 *Biogeochemistry*, **39**, 177-187.

412

413 Freeman, C., Ostle, N., and Kang, H., 2001a. An enzymic 'latch' on a global carbon store.  
414 *Nature*, **409**, 149.

415

416 Freeman, C., Evans, C.D., Monteith, D.T., Reynolds, B., Fenner, N., 2001b. Export of  
417 organic carbon from peat soils. *Nature*, **412**, 785-786.

418

419 Gibson, H.S., Worrall, F., Burt, T.P., Adamson, J.K. 2009. DOC budgets of drained peat  
420 catchments: implications for DOC production in peat soils. *Hydrological Processes*, **23**,  
421 1901-1911.

422

423 Glatzel, S., Kalbitz, K., Dalva, M., Moore, T., 2003. Dissolved organic matter properties and  
424 their relationship to carbon dioxide efflux from restored peat bogs. *Geoderma*, **113**, 397-411.  
425

426 Green, S.M., Baird, A.J., Boardman, C.P., Gauci, V. 2014. A mesocosm study of the effect of  
427 restoration on methane (CH<sub>4</sub>) emissions from blanket peat. *Wetlands Ecology and*  
428 *Management*, DOI 10.1007/s11273-014-9349-3.  
429

430 Holden, J., Wallage, Z.E., Lane, S.N., McDonald, A.T., 2011. Water table dynamics in  
431 undisturbed, drained and restored blanket peat. 2011. *Journal of Hydrology*, **402**, 103-114.  
432

433 Hope, D., Billett, M.F., Cresser, M.S. 1994. A review of the export of carbon in river water:  
434 fluxes and processes. *Environmental Pollution*, **84**, 301-324.  
435

436 Kane, E.S., Mazzoleni, L.R., Kratz, C.J., Hribljan, J.A., Johnson, C.P., Pypker, T.G.,  
437 Chimner, R. 2014. Peat porewater dissolved organic carbon concentration and lability  
438 increase with warming: a field temperature manipulation experiment in a poor fen.  
439 *Biogeochemistry*, DOI 10.1007/s10533-014-9955-4.  
440

441 Karlsson, J., Byström, P., Ask, J., Persson, L., Jansson, M. 2009. Light limitation of nutrient-  
442 poor lake ecosystems. *Nature*, **460**, 506-509.  
443

444 Komulainen, V-M., Nykänen, H., Martikainen, P.J., Laine, J. 1998. Short-term effect of  
445 restoration on vegetation change and methane emissions from peatlands drained for forestry  
446 in southern Finland. *Canadian Journal of Forest Research*, **28**, 402-411.  
447

448 Martikainen, P.J., Nykänen, H., Alm, J., and Silvola, J., 1995. Changes in fluxes of carbon  
449 dioxide, methane and nitrous oxide due to forest drainage of mire sites of different trophy.  
450 1995. *Plant and Soil*, **168-169**, 571-577.

451

452 McDonald, A.T., Mitchell, G.N., Naden, P.S., Martin, D.S.J., 1991. Discoloured Water  
453 Investigations. Final Report to Yorkshire Water plc. 432 pp.

454

455 Peacock, M., Burden, A., Cooper, M., Dunn, C., Evans, C.D, Fenner, N., Freeman, C.,  
456 Gough, R., Hughes, D., Hughes, S., Jones, T., Robinson, I., West, M., Zieliński, P., 2013.  
457 Quantifying dissolved organic carbon concentrations in upland catchments using phenolic  
458 proxy measurements. *Journal of Hydrology*, **477**, 251-260.

459

460 Pind, A., Freeman, C., and Lock, M.A., 1994. Enzymic degradation of phenolic materials in  
461 peatlands – measurement of phenol oxidase activity. *Plant and Soil*, **159**, 227-231.

462

463 Price, J.S., Heathwaite, A.L., Baird, A.J. 2003. Hydrological processes in abandoned and  
464 restored peatlands: an overview of management approaches. *Wetlands Ecology and*  
465 *Management*, **11**, 65-83.

466

467 Ramchunder, S.J., Brown, L.E., Holden, J. 2012. Catchment-scale peatland restoration  
468 benefits stream ecosystem biodiversity. *Journal of Applied Ecology*, **49**, 182-191.

469

470 Sinsabaugh, R.L., 2010. Phenol oxidase, peroxidase and organic matter dynamics of soil. *Soil*  
471 *Biology and Biochemistry*, **42**, 391-404.

472

473 Stewart, A.J.A., Lance, A.N., 1991. Effects of moor-draining on the hydrology and  
474 vegetation of northern Pennine blanket bog. *Journal of Applied Ecology*, **28**, 1105-1117.  
475

476 Tahvanainen, T., Haraguchi, A. 2013. Effect of pH on phenol oxidase activity on decaying  
477 *Sphagnum* mosses. *European Journal of Soil Biology*, **54**, 41-47.  
478

479 Toberman, H., Evans, C.D., Freeman, C., Fenner, N., White, M., Emmett, B.A., Artz, R.R.E.,  
480 2008. Summer drought effects upon soil and litter extracellular phenol oxidase activity and  
481 soluble carbon release in an upland *Calluna* heathland. *Soil Biology and Biochemistry*, **40**,  
482 1519-1532.  
483

484 Wallage, Z.E., Holden, J., McDonald, A.T., 2006. Drain blocking: an effective treatment for  
485 reducing dissolved organic carbon loss and water discolouration in a drained peatland.  
486 *Science of the Total Environment*, **367**, 811-821.  
487

488 Wetzel, R.G., 1992. Gradient-dominated ecosystems: sources and regulatory functions of  
489 dissolved organic matter in freshwater ecosystems. *Hydrobiologia*, **229**, 181-198.  
490

491 Wilson, L., Wilson, J., Holden, J., Johnstone, I., Armstrong, A., Morris, M., 2011a. Recovery  
492 of water tables in Welsh blanket bog after drain blocking: discharge rates, time scales and the  
493 influence of local conditions. *Journal of Hydrology*, **391**, 377-386.  
494

495 Wilson, L., Wilson, J., Holden, J., Johnstone, I., Armstrong, A., Morris, M., 2011b. Ditch  
496 blocking, water chemistry and organic carbon flux: evidence that blanket bog restoration  
497 reduces erosion and fluvial carbon loss. *Science of the Total Environment*, **409**, 2010-2018.

498

499 Worrall, F., Armstrong, A., and Holden, J., 2007. Short-term impact of peat drain-blocking  
500 on water colour, dissolved organic carbon concentration, and water table depth. *Journal of*  
501 *Hydrology*, **337**, 315-325.


502

503 Table 1. Pore water ion concentrations and standard errors ( $\text{mg L}^{-1}$ ) for the undrained Bryn Du site and  
504 piezometers associated with open ditches at the Afon Du. Values are means from monthly sampling for March-  
505 July 2011 ( $n = 20$ ), except for chloride, phosphate and sulphate where extra data were available; these ions were  
506 measured monthly March-November 2011 ( $n = 32$ ). For each site and month  $n = 4$ . The only significant  
507 difference between sites was found for phosphate (Mann-Whitney U test,  $p = 0.001$ ).

	Bryn Du	Afon Du
Sodium	4.00 $\pm 0.21$	4.24 $\pm 0.28$
Ammonium	0.01 $\pm 0.00$	0.01 $\pm 0.01$
Potassium	0.10 $\pm 0.04$	0.29 $\pm 0.20$
Magnesium	0.54 $\pm 0.09$	0.59 $\pm 0.07$
Calcium	0.34 $\pm 0.08$	0.60 $\pm 0.1$
Chloride	5.26 $\pm 0.47$	5.29 $\pm 0.27$
Bromide	0.01 $\pm 0.00$	0.02 $\pm 0.00$
Nitrate	0.01 $\pm 0.00$	0.02 $\pm 0.02$
Phosphate	0.00 $\pm 0.00$	0.25 $\pm 0.05$
Sulphate	1.02 $\pm 0.46$	2.22 $\pm 0.83$

508


509



510


511 Figure 1. Mean hydrolase activities ( $\text{nmol g}^{-1} \text{min}^{-1} \text{MUF released}$ ) for drained and undrained sites. Error bars  
 512 show standard error of the mean. Data are mean from two sampling dates,  $n = 8$  for each treatment, except  
 513 chitinase which is from one sampling date ( $n = 4$ ). There were significant differences (\*) between sites for  $\beta$ -  
 514 glucosidase (one-tailed t-test,  $p = 0.01$ ) and arylsulphatase (one-tailed t-test,  $p = 0.02$ ).

515




516

517 Figure 2. Mean phenol oxidase activity ( $\text{nmol dicq g}^{-1} \text{min}^{-1}$ ) ( $n = 8$ ) and total mean hydrolase activity ( $\text{nmol g}^{-1}$   
 518  $\text{min}^{-1} \text{MUF released}$ ) (i.e. sum of mean  $\beta$ -glucosidase, arylsulphatase, xylosidase and chitinase activity,  $n = 28$ )  
 519 for drained and undrained sites. Error bars show standard error of the mean. The difference is significant (\*)  
 520 for phenol oxidase (one-tailed t-test,  $p = 0.01$ ) and hydrolases (one-tailed t-test,  $p = 0.01$ ).



522


523 Figure 3. Mean hydrolase activities (nmol g<sup>-1</sup> min<sup>-1</sup> MUF released) for open control ditches, dammed ditches  
 524 and reprofiled ditches. Errors bars show standard error of the mean. Data are mean of five (approximately  
 525 monthly) sampling dates. *n* = 20 for each treatment. The only significant difference (\*) was for arylsulphatase  
 526 (repeated -measures ANOVA with Tukey HSD, *p* < 0.05).



527

528 Figure 4. Mean phenol oxidase activity (nmol dicq g<sup>-1</sup> min<sup>-1</sup>) for open control ditches, dammed ditches and  
 529 reprofiled ditches. Errors bars show standard error of the mean. Data are mean of five (approximately monthly)  
 530 sampling dates. *n* = 20 for each treatment.






531

532

533

534

Figure 5. Mean phenolic concentrations ( $\text{mg L}^{-1}$ ) for open control ditches, dammed ditches and reprofiled ditches. Errors bars show standard error of the mean. Data are mean of five (approximately monthly) sampling dates.  $n = 20$  for each treatment.




535

536

537


538

Figure 6. Mean DOC concentrations ( $\text{mg L}^{-1}$ ) for open control ditches, dammed ditches and reprofiled ditches. Errors bars show standard error of the mean. Data are mean of five (approximately monthly) sampling dates.  $n = 20$  for each treatment.




539

540 Figure 7. Relationship between  $\beta$ -glucosidase activity and DOC concentration in ditch and pore waters. Data  
 541 are from five sampling trips between June and October 2011. For ditch water  $n = 60$ ,  $r^2 = 0.20$ ,  $p < 0.05$ ,  $y = -$   
 542  $1.05x + 39.36$ . For pore water  $n = 29$ ,  $r^2 = 0.35$ ,  $p < 0.01$ ,  $y = -2.74x + 68.43$ .



543

544 Figure 8. Relationship between phenol oxidase activity and the ratio of phenolic compounds to DOC in ditch  
 545 waters, and the relationship between phenol oxidase activity and DOC concentration in pore waters. Data are  
 546 from five sampling trips between June and October 2011. For ditch water  $n = 56$ ,  $r^2 = 0.19$ ,  $p < 0.01$ ,  $y = -$   
 547  $0.0009x + 0.1939$ . For pore water  $n = 27$ ,  $r^2 = 0.17$ ,  $p < 0.05$ ,  $y = -1.21x + 62.64$ .



548

549 Figure 9. Relationship between arylsulphatase activity ( $\text{nmol g}^{-1} \text{min}^{-1}$  MUF released) and sulphate  
550 concentration in ditch waters. Data are from five sampling trips between June and October 2011.  $n = 56$ ,  $r^2 =$   
551  $0.19$ ,  $p < 0.01$ ,  $y = 0.064 x + 0.563$ .  
552