



Structural and Functional Changes in the Coupling of Fascial Tissue, Skeletal Muscle, and Nerves During Aging

Alberto Zullo^{1,2}, Johannes Fleckenstein³, Robert Schleip^{4,5}, Kerstin Hoppe⁶, Scott Wearing^{4,7} and Werner Klingler^{4,7,8,9*}

¹Department of Sciences and Technologies, University of Sannio, Benevento, Italy, ²CEINGE Advanced Biotechnologies, Naples, Italy, ³Department of Sports Medicine, Institute of Sports Sciences, Goethe-University Frankfurt, Frankfurt, Germany, ⁴Department of Sport and Health Sciences, Technical University Munich, Munich, Germany, ⁵Department of Sports Medicine and Health Promotion, Friedrich-Schiller University Jena, Jena, Germany, ⁶Department of Anaesthesiology, Würzburg University, Würzburg, Germany, ⁷Faculty of Health School, Queensland University of Technology, Brisbane, QLD, Australia, ⁸Fascia Research Group, Department of Experimental Anaesthesiology, Ulm University, Ulm, Germany, ⁹Department of Anaesthesiology, SRH Hospital Sigmaringen, Sigmaringen, Germany

OPEN ACCESS

Edited by:

Can A. Yucesoy,
Boğaziçi University, Turkey

Reviewed by:

Filiz Ates,
Mayo Clinic, United States
Laszlo Csernoch,
University of Debrecen,
Hungary

*Correspondence:

Werner Klingler
werner.klingler@uni-ulm.de

Specialty section:

This article was submitted to
Striated Muscle Physiology,
a section of the journal
Frontiers in Physiology

Received: 05 February 2020

Accepted: 11 May 2020

Published: 24 June 2020

Citation:

Zullo A, Fleckenstein J, Schleip R, Hoppe K, Wearing S and Klingler W (2020) Structural and Functional Changes in the Coupling of Fascial Tissue, Skeletal Muscle, and Nerves During Aging. *Front. Physiol.* 11:592. doi: 10.3389/fphys.2020.00592

Aging is a one-way process associated with profound structural and functional changes in the organism. Indeed, the neuromuscular system undergoes a wide remodeling, which involves muscles, fascia, and the central and peripheral nervous systems. As a result, intrinsic features of tissues, as well as their functional and structural coupling, are affected and a decline in overall physical performance occurs. Evidence from the scientific literature demonstrates that senescence is associated with increased stiffness and reduced elasticity of fascia, as well as loss of skeletal muscle mass, strength, and regenerative potential. The interaction between muscular and fascial structures is also weakened. As for the nervous system, aging leads to motor cortex atrophy, reduced motor cortical excitability, and plasticity, thus leading to accumulation of denervated muscle fibers. As a result, the magnitude of force generated by the neuromuscular apparatus, its transmission along the myofascial chain, joint mobility, and movement coordination are impaired. In this review, we summarize the evidence about the deleterious effect of aging on skeletal muscle, fascial tissue, and the nervous system. In particular, we address the structural and functional changes occurring within and between these tissues and discuss the effect of inflammation in aging. From the clinical perspective, this article outlines promising approaches for analyzing the composition and the viscoelastic properties of skeletal muscle, such as ultrasonography and elastography, which could be applied for a better understanding of musculoskeletal modifications occurring with aging. Moreover, we describe the use of tissue manipulation techniques, such as massage, traction, mobilization as well as acupuncture, dry needling, and nerve block, to enhance fascial repair.

Keywords: aging, connective tissue, fascia, skeletal muscle, nerve

INTRODUCTION

In animal species, aging is associated with substantial modifications to the molecular determinants of cells, which alter their morphology, activity, and functional properties. These events affect the characteristics of tissues and organs, thus resulting in overall decayed performance of the organism (Narici et al., 2008; Zullo et al., 2017). Indeed, during aging, human skeletal muscle, connective tissue, and the nervous system undergo consistent modifications, which can impair the ability to perform daily activities and, as a consequence, their health-related quality of life (Netuveli et al., 2006; Reid and Fielding, 2012; Diehr et al., 2013). To promote healthy aging and reduce the burden of aging, it would be important to define the time-dependent alterations impairing the neuromuscular system. Also, sensitive, specific, and non-invasive methods to investigate the structural and morphological changes occurring in these tissues during aging are needed, and therapeutic interventions should be implemented. In this review, we outline the structural and functional changes occurring within skeletal muscle, fascial tissue, and nervous system during aging. Moreover, we summarize the evidence on the age-dependent impairment of the coupling between these tissues and the effect of inflammation in aging. Finally, we describe some approaches for the study and the treatment of age-associated modifications in the neuromuscular system, such as imaging and ultrasound-based methods, as well as tissue manipulation techniques.

SKELETAL MUSCLE AND AGING

Sarcopenia, a gradual loss in muscle mass occurring in humans during aging, is detectable in the third decade of life and progressively increases with age (Janssen et al., 2000; Narici and Maffulli, 2010). It has been estimated that, during aging, there is a 30–50% reduction in the number and 10–40% decrease in the size of skeletal muscle fibers, which is associated with a decay of muscle performance (Lexell et al., 1988; Lexell, 1995; Janssen et al., 2002; Doherty, 2003). In particular, a reduction in skeletal muscle mass has been estimated to be 0.37 and 0.47% per year in women and men, respectively (Mitchell et al., 2012). Clinical studies in old adults have shown that muscle strength declines at a greater rate than muscle mass (Frontera et al., 2000; Goodpaster et al., 2006; Delmonico et al., 2009). Therefore, factors other than muscle mass must also contribute to the decay of muscle performance in aging.

In humans, three isoforms (type I, IIa, and IIx) of the myosin heavy chain (MHC) are present, with different combinations and proportions in individual muscles (Schiaffino and Reggiani, 1994, 1996). Each isoform has specific molecular features, which in turn, determine muscle characteristics (Trappe, 2009). There is evidence that the composition of the contractile apparatus of muscle, i.e., the isoforms of the MHC, varies with aging (Trappe, 2009). In the vastus lateralis of old adults, the percentage of muscle fibers expressing both MHC types I (MHC-I) and IIa (MHC-IIa) and MHC-IIa and IIx (MHC-IIx) is higher, and the percentage of muscle fibers expressing only MHC-I and MHC-IIa

is lower than those of young adults (Klitgaard et al., 1990). Moreover, in the sternocleidomastoid muscle (SCM) of old, compared to young, adults, the proportion of slow-twitch fibers is higher, the share of hybrid MHC-IIa/MHC-IIx fibers is reduced, the proportion of fast-twitch fibers with MHC-IIa and IIx is lower, and the proportion of fibers with the isoform MHC-IIx is reduced (Meznaric et al., 2018). This leads to a shift toward a slower phenotype of muscle fibers and suggests that aging induces the replacement of fast-twitch with slow-twitch motor units (Meznaric et al., 2018). Interestingly, the age-dependent decrease in the number and size of muscle fibers involves mainly type II, but not type I muscle fibers (Larsson, 1978; Lexell et al., 1988; Lexell, 1995; Martel et al., 2006; Snijders et al., 2009).

As for fiber structure, it has been demonstrated that pennation angles in the vastus lateralis muscle diminish with advancing age in women (Kubo et al., 2003). Such architectural modification results in a reduced number of parallel fibers and negatively impacts on muscle force generation (Thom et al., 2007). Moreover, besides changes in structural proteins, alteration of contractile and metabolic proteins, as well as calcium-handling proteins of skeletal muscle cells also take place in the elderly (Godard et al., 2002; Thompson, 2002). The impairment of excitation-contraction coupling, the increase of intermuscular adipose tissue, the alteration of intramuscular adipose tissue, and the amount of extracellular water relative to muscle volume with aging could also promote the decline in muscle performance (Rice et al., 1989; Yamada et al., 2010; Marcus et al., 2012; Beavers et al., 2013; Hausman et al., 2014; Tieland et al., 2018).

As for muscle function, data from studies in animal models and humans demonstrated that the strength and the shortening velocity of muscle fibers decrease with advancing age (Lauretani et al., 1985; Frontera et al., 2000; Thompson, 2002; D'Antona et al., 2003; Miljkovic et al., 2015; Del Campo et al., 2018). A study in a large cohort of adults demonstrated a significant reduction in isometric, concentric, and eccentric peak torque in the extensor muscles of the knee in old, compared to young, adults (Lindle et al., 1997). Also, a 16.6–40.9% reduction in muscular strength has been estimated by comparing young adults (<40 years) with old one (>40 years) (Keller and Engelhardt, 2014). A research employing magnetic resonance imaging (MRI) has demonstrated that maximum voluntary contraction, when normalized to the anatomical cross-sectional area of muscle, is lower in the elderly than healthy young adults (Morse et al., 2005). Although several pieces of evidence have been accumulated on the functional decline of skeletal muscle during aging, this issue is still disputed (Lexell et al., 1988; Trappe et al., 2003; Short et al., 2005a; Canepari et al., 2010). Indeed, some studies report that the power of human muscle fibers, normalized for cell size, does not change with age (Trappe et al., 2003; Malisoux et al., 2007; Raue et al., 2009).

The wide remodeling of skeletal muscle occurring with age also influences its mechanical properties. Kocur and colleagues demonstrated that stiffness and tone of the upper trapezius (UT) and SCM increase with age, and elasticity decreases (Kocur et al., 2017). On the contrary, the viscoelastic features of these muscles during movement are not significantly affected (Kocur et al., 2017). The analysis of muscle tone and

mechanical properties of rectus femoris and biceps brachii muscles in old and young adults revealed that aging is associated with an increase of muscle tone, and stiffness, and a decrease in elasticity (Agyapong-Badu et al., 2016). Also, the sternocleidomastoid and upper trapezius muscles undergo a decrease in elasticity and an increase in stiffness in women in a seated position (Kocur et al., 2019). These parameters varied about 1.5% per year, and, among all the possible etiologic factors analyzed (BMI, age, and head posture), only aging was found to be a major correlate (Kocur et al., 2019). Neuromuscular performance is also impaired with aging, as demonstrated in a study that evaluated both the time needed to perform specific muscular activities and muscular functional features (Haus et al., 2007b). The authors showed that the time to climb stairs, rise from a chair, and walk a set distance significantly increased and the quadriceps muscle volume, strength, and power significantly decreased in aged adults (Haus et al., 2007b). These alterations did not match changes in intramuscular endomysial collagen nor in enzymatically-mediated collagen cross-linking, which remained constant, but matched a 200% increase in the advanced glycation end-product (AGE), pentosidine (Haus et al., 2007b). The accumulation of AGEs in the intramuscular connective tissue, probably due to age-dependent decrease in collagen turnover, is associated with the formation of non-enzymatically-mediated collagen cross-linking (Verzijl et al., 2000; Haus et al., 2007b). This observation led the authors to hypothesize that the accumulation of AGEs in the intramuscular connective tissue of old adults might be responsible for the increase in muscle tissue stiffness and decrease in passive viscoelastic properties, thus impairing muscle function.

Skeletal muscle is highly responsive to activity, i.e., it remodels and adapts its metabolism and structure to meet the body's needs (Thompson, 2002; Potthoff et al., 2007). Aging affects the architecture and the function of the skeletal muscle, but there are conflicting reports about the level of muscular adaptive capacity in the elderly (Rogers and Evans, 1993; Kim and Thompson, 2013; Gheller et al., 2016). However, based on some pieces of evidence, aged skeletal muscle still maintains a certain degree of plasticity; therefore, it can undergo structural and functional changes in response to physiological stimuli (Joanisse et al., 2017). In particular, exercise training has been demonstrated to stimulate skeletal muscle fiber hypertrophy and increase mitochondrial oxidative capacity in aged humans and animal models, resulting in increased muscle mass and strength (Frontera et al., 1988; Joseph et al., 2016; White et al., 2016; Joanisse et al., 2017; Kim et al., 2017). At the same time, inactivity induces a decrease in skeletal muscle mass, and strength independently of age, but the recovery after immobility is impaired in aged humans and animal models (Zhang et al., 2018a,b; Oikawa et al., 2019).

At the molecular level, reactive oxygen species (ROS) and PGC-1 α signaling, as well as the autophagic machinery, have been suggested as important determinants contributing to metabolic adaption and remodeling of skeletal muscle after physical activity (Ferraro et al., 2014). Skeletal muscle fibers during their activity produce ROS, mainly due to the activity of mitochondria, NADPH oxidase, xanthine oxidase, and

phospholipase A2 (Szentesi et al., 2019). In turn, ROS activate several signaling pathways, which are pivotal for different cellular processes, such as proliferation, protein synthesis and degradation, ROS detoxification, and muscle fiber contraction (McDonagh, 2016). ROS, at physiological levels, have a crucial role in the activation of adaptive responses to exercise in skeletal muscle cells (Ji, 2015). Indeed, mitochondrial biogenesis, antioxidant activity, inflammation, protein turnover, apoptosis, and autophagy are upregulated. The principal molecular elements of these signaling cascades are NF- κ B, MAPK, and PGC-1 α . When the levels of ROS greatly exceed the cellular antioxidant capacity, deleterious oxidative modifications occur to the structures and the molecules of the skeletal muscle fibers, thus impairing their function. Aging is associated with increased levels of oxidative stress in the skeletal muscle tissue, which is the accumulation of reactive oxygen and nitrogen species. As a result, the skeletal muscle undergoes structural and functional modifications, which reduce muscle mass, strength, and function (Ji, 2015). Indeed, satellite cells become dysfunctional, the rate of protein breakdown prevails over that of protein synthesis, muscle fibers activate apoptotic processes, intracellular calcium homeostasis is altered, and excitation-contraction coupling is impaired (Szentesi et al., 2019). Studies in humans and mice showed that mitochondria of aged skeletal muscles have impaired mitochondrial NADH redox potential and reduced oxidative capacity (Conley et al., 2000; Clafin et al., 2015; Nelson et al., 2019). As a consequence, aged muscles accumulate oxidized proteins, and mitochondrial dysfunction, which, in turn, lead to increased ROS production and oxidative damage in the cells (Short et al., 2005b; Cerullo et al., 2012; Bratic and Larsson, 2013; Gomes et al., 2017). Besides, other cellular antioxidant enzymes, such as catalase, glutathione transferase, and superoxide dismutase, are also significantly reduced in skeletal muscle fibers of old adults, and exercise-activated redox signaling pathways are weakened (Cerullo et al., 2012; Ji, 2015). A proteomic study in mice demonstrated that aging is correlated with reduced levels of redox-sensitive proteins involved in the energy metabolism of skeletal muscle fibers, thus affecting the cellular responses to oxidants (McDonagh et al., 2014). These data were also supported by a recent study in humans, which showed that sarcopenia is associated with altered expression of genes regulating mitochondrial energy production in skeletal muscle (Migliavacca et al., 2019). Indeed, the PGC-1 α /ERR α signaling, oxidative phosphorylation, and proteostasis are downregulated. As a result, mitochondria are dysfunctional and their number is reduced, thus further exposing skeletal muscle fibers to oxidative damage (Migliavacca et al., 2019).

Experimental studies indicate a negative effect of aging also on skeletal muscle plasticity, which is the potential of this tissue to modify its structural and functional features in response to environmental changes (Verdijk et al., 2007; Jee and Kim, 2017; Suetta, 2017). The regrowth response to physical inactivity-induced atrophy is also hindered in aged muscles (Pisot et al., 2016).

As for the regenerative potential of skeletal muscle during aging, it has been reported that satellite cells, the quiescent adult stem cells of skeletal muscle, gradually lose their potential to regenerate skeletal muscle with advancing age

(Yin et al., 2013; Sousa-Victor et al., 2014; Joannis et al., 2016). Indeed, experimental evidence from studies in both humans and animal models suggests that aging is associated with a reduction in satellite cell self-renewal and myogenic competences, and possibly a decrease in their number, thus resulting in an impaired regeneration of skeletal muscle tissue (Corbu et al., 2010; Sousa-Victor et al., 2014; Brack and Muñoz-Cánoves, 2016; Joannis et al., 2016). At the molecular level, aging drives a dysregulation of important molecular signaling pathways, such as FGF2/Sprouty1, Notch, p38 MAPK, JAK-STAT3, and p16(INK4a) in satellite cells, and impairs the microenvironment feeding and regulating the muscle stem cell niche (Sousa-Victor et al., 2014; Parker, 2015; Snijders et al., 2015; Brack and Muñoz-Cánoves, 2016; Joannis and Parise, 2016; Stearns-Reider et al., 2017; Etienne et al., 2020; Levi et al., 2020).

To counteract the impairment of skeletal muscle with aging, three main approaches have been pursued: physical activity, antioxidant dietary supplementation, and regenerative medicine therapies. Some pieces of evidence showed that regular exercise can reduce the detrimental effects of aging on skeletal muscle. In particular, it has been reported, both in humans and animal models, that regular physical activity can attenuate the age-dependent decrease in the number of mitochondria and proteins involved in the excitation-contraction coupling, thus delaying the impairment in muscle function (Zampieri et al., 2015; Csernoch et al., 2017). Physical inactivity also damages the structure and function of skeletal muscle fibers. Moreover, in old adults, besides a weakened ROS scavenging system, dietary intake of antioxidants is often reduced (Damiano et al., 2019). In this context, data from studies in humans and animal models indicate that muscular activity and/or dietary supplementation with antioxidant compounds, such as L-ascorbic acid, tocopherols, carotenoids, flavonoids, or polyphenols, exert some beneficial effects on age-related skeletal muscle decline (Cesari et al., 2004; Pierno et al., 2014; Anton et al., 2015; Cartee et al., 2016; Jee and Kim 2017; Muhammad and Allam, 2018; Damiano et al., 2019). However, it should be taken into account that the effects of physical exercise and/or dietary supplementation with antioxidants on skeletal muscle aging are still debated due to discordant results (Marcell, 2003; Ji, 2015; Cartee et al., 2016; Flack et al., 2016; White et al., 2016; Distefano and Goodpaster, 2018; Damiano et al., 2019; Szentesi et al., 2019).

In the last decades, thanks to the progress of regenerative medicine, new approaches for improving skeletal muscle regeneration in humans have been developed. Indeed, cell therapies based on the delivery of myogenic stem cells have been largely used also in preclinical studies. Several sources of myogenic cells have been employed, so far: satellite cells, muscle-derived stem cells, myoblasts, mesoangioblasts, hTERT-immortalized muscle precursor cells, pericytes, CD133⁺ cells, hematopoietic stem cells, mesenchymal stem cells, perivascular stem cells, interstitial cells, pluripotent stem cells isolated from the dental pulp, embryonic stem cells, and induced pluripotent stem cells (iPSCs) (Hall et al., 2017; Del Carmen Ortuño-Costela et al., 2019; Marg et al., 2019; Mueller and Bloch, 2019). As a result, the delivery of these myogenic cells led to some improvements in muscle function and regeneration in mammalian animal

models (Rao et al., 2018; Mueller and Bloch, 2019). To enhance the effectiveness of these regenerative therapies, bioactive molecules, such as TGF- β , IGF-I, fibrin, keratine, as well as ECM-based bioscaffolds, have been combined with myogenic cells (Brown et al., 2012; Dziki et al., 2016; Fuoco et al., 2016; Jiao et al., 2018; Pollot et al., 2018; Mueller and Bloch, 2019). However, we are still far from effective treatments to use in clinical practice for skeletal muscle regeneration due to the difficulties associated with cell preparation, the limited engraftment of myogenic cells, the incomplete differentiation of myogenic cells *in situ*, the partial integration with host cells, and tumorigenic risk of immortalized cells. In the context of pharmacological intervention, many different drugs have been tested for improving muscle regeneration. Indeed, anti-inflammatory drugs have been used for blocking the anti-regenerative effect of chronic inflammation, hormones, and growth factors for stimulating cell growth; urocortin II for stimulating muscle fiber growth pathways; and myostatin for activating satellite cells. Indeed, urocortin II and myostatin showed an interesting potential for improving muscle hypertrophy and hindering the age-dependent loss of muscle tissue in animal models (Hinkle et al., 2003; Rodriguez et al., 2014; Cohen et al., 2015; Naranjo et al., 2017; Saul et al., 2019).

CONNECTIVE TISSUE AND AGING

Besides its influence on skeletal muscle, aging also results in modification of cells and extracellular matrix of myofascia and tendons (Barros et al., 2002; Haus et al., 2007a; Trappe, 2009; Kragstrup et al., 2011). Muscular fascia is composed of many different molecules, such as structural proteins (collagens, laminins, fibronectin, vitronectin, tenascin, and elastin), growth factors (TGFs and IGFs) glycosaminoglycans, proteoglycans, metalloproteinases, cytokines, and water (Eyre et al., 1984; Reiser et al., 1992; McCormick, 1999). Fascia, due to its structure and composition, has elastic, viscoelastic and plastic properties that strongly influence the biomechanical features of the locomotory apparatus, as demonstrated both in humans and animal models (Kovanen et al., 1988; Goldspink et al., 1994; Kjaer, 2004; Avila Gonzalez et al., 2018; Blottner et al., 2019; Schleip et al., 2019). Muscular connective tissue changes with advancing age, as reported also in studies in animal models. In particular, its thickness and the amount of collagen cross-linking increase, and its elasticity decreases (Alnaqeeb et al., 1984; Gosselin et al., 1998; Ducomps et al., 2003; Wilke et al., 2019; Etienne et al., 2020). Additionally, the composition of the muscular connective tissue varies; as an example, the amount of collagen type IV rises, and that of type VI reduces (Etienne et al., 2020; Levi et al., 2020). As a result, the extracellular matrix becomes more rigid and muscles increase their stiffness, thus resulting in impaired muscle function (Alnaqeeb et al., 1984; Kovanen et al., 1984; Gosselin et al., 1998; Willems et al., 2001; Ducomps et al., 2003; Etienne et al., 2020).

Studies in humans and animal models showed that with advancing age the basal lamina becomes thicker and destructured, and the content of collagen type IV, laminin, and the antimyogenic cytokine, osteopontin, increases in skeletal muscle, thus hindering

its regenerative potential (Kovanen et al., 1988; Grzelkowska-Kowalczyk, 2016). Moreover, a decrease in the number of fibroblasts and stem cells in tendon during aging has been also reported in humans and animal models (Squier and Magnes, 1983; Nakagawa et al., 1994; Zhou et al., 2010; Ruzzini et al., 2013).

A recent study by Wilke and collaborators demonstrated that aging is associated with variation in the thickness of the fascia. Indeed, the age-related modifications are specific for different body sites. Fascial thickness of the lower limb decreases with age (−12.3–25.8%), whereas fascia of the low back region increases (+40.0–76.7%) (Wilke et al., 2019). These changes in connective tissue have been suggested to reduce joint flexibility (Wilke et al., 2019). Moreover, the amount of intramuscular connective tissue in the gastrocnemius muscle of elderly adults is reportedly higher than that of young adults (Csapo et al., 2014). Although connective tissue content may change with aging, the amount of cross-linking of intramuscular collagen remains stable, according to the results of a study in the vastus lateralis of young and old adults (Haus et al., 2007a). Aging results also in a reduction in water content within tendon, as reported in the Achilles tendon of old rabbits (Ippolito et al., 1980), an increase in cross-links between tendon fibrils and a decrease in collagen content and density (Vogel, 1978; Haut et al., 1992; Coupe et al., 2009; Svensson et al., 2015). Reducing sugars can link to amino acids of collagen fibers and generate AGEs, which accumulate throughout life and result in increased tissue stiffness and strength (Bank et al., 1999; DeGroot et al., 2001; Suzuki et al., 2008).

In addition, reduced elasticity and increased stiffness of aged musculoskeletal system can also be caused by degeneration of connective tissue, which leads to reduced joint mobility in the elderly (Kocur et al., 2019). At the molecular level, it has been reported that Wnt signaling, the expression of metalloproteinase (MMP) and tissue inhibitor of metalloproteinase genes, which regulate ECM degradation and remodeling, change with advancing age (Phillip et al., 2015; Birch, 2018; de Sousa Neto et al., 2018). Barros and colleagues demonstrated that elastic fibers of the cervical interspinous ligaments undergo fragmentation and degeneration, and oxytalan fibers, which confer resistance to the tissue, degrade during aging (Barros et al., 2002). As a consequence, ligaments are more susceptible to ruptures following mechanical stress (Barros et al., 2002).

Despite age-related modifications of connective tissue, the effect of these changes on the mechanical properties of tendons, such as strength, stiffness, and elasticity, is still debated, due to conflicting reports (Svensson et al., 2015). Interestingly, a recent pilot study demonstrated that acute resistance exercise differentially affected young and old adults in the context of metalloproteinase gene expression (Wessner et al., 2019). These results support the evidence of stimulus-dependent ECM remodeling in the elderly.

COUPLING BETWEEN FASCIA, SKELETAL MUSCLE AND AGING

Muscles and fascia cooperate for the correct functioning of the locomotory apparatus. Their intimate relationship makes the

performance of movement strictly dependent on the status of both of them. As an example, appropriate preparation of fascial structures by a warm-up and stretching protocols is essential for optimal results and minimal risk of injury in physical exercises (Yahia et al., 1993; Mattieni et al., 2009; Wang et al., 2009). The structural and molecular changes of skeletal muscle and fascia with advancing age influence the transmission of force in the locomotory system (Wilke et al., 2018). Intramuscular fascia connects different muscle fibers and muscle bundles within the muscle to form the force-generating structure connected to the bone (Turrina et al., 2013). Indeed, studies in rats showed that the force generated by muscle fiber contraction is transmitted both longitudinally and laterally, through the intramuscular fascia, to the surrounding muscle fibers, till the tendon and the bones (Huijing and Jaspers, 2005). Moreover, molecular features, structure, and orientation of fascial fibers determine how force is transmitted and transferred through connective tissue to other surrounding elements (Stecco et al., 2013; Wilke et al., 2018). In humans, recent data showed that the length of muscle fascicles is affected by the coupling between muscle and fascia (Pamuk et al., 2016; Karakuzu et al., 2017). Indeed, heterogeneous fascicle strains have been detected in the medial gastrocnemius muscle following submaximal plantar flexion activity or knee extension. This result has been explained by epimuscular force transmission (Pamuk et al., 2016; Karakuzu et al., 2017). Intermuscular mechanical interactions also have important implications in patients affected by cerebral palsy (CP), who suffer from motor disability. It has been suggested that the co-activation of antagonistic muscles of the knee, which causes the limited joint motion in CP patients, is due to altered force transmission through myofascial structures (Kaya et al., 2019, 2020). This makes fascia a key structural and functional element of the contractile apparatus of muscle. As a consequence, the mechanical features of the entire muscle cannot be merely described as the sum of the isolated fascia and muscle fibers.

The fluid component of fascia assists in the hydrostat function of the tissue and dissipates energy, whereas the elastic component, made by fibrous protein structures, can store and release elastic energy (Huijing et al., 2003; Özkaya et al., 2017). Besides the capacity of fascial components to generate force by myofibroblast contraction, fascia can also modulate its composition and structure in response to biomechanical stimulation, thus allowing its adaptation over time to meet the body's needs (Blottner et al., 2019). Several studies have demonstrated that muscles located in anatomically separate body regions can exchange tensional stress through a tight link and cooperation with fascial structures, thus contributing to the accomplishment of locomotory tasks and movement proprioception (Vleeming et al., 2014; Blottner et al., 2019). In the anatomical compartment comprised between leg and trunk, this complex architecture has been described as a myofascial force transmission chain (Myers, 2013; Wilke et al., 2016).

Data collected so far suggest that myofascial force transmission is extremely relevant during muscle lengthening, such as in eccentric contractions and stretching activity (Yucesoy et al., 2005; Huijing, 2009; Yucesoy, 2010; Wilke et al., 2018). Indeed, the presence of intermuscular mechanical interactions in the lower

leg muscles under passive joint motion has been clearly demonstrated by several ultrasound-based studies (Bojsen-Møller et al., 2010; Tian et al., 2012; Le Sant et al., 2017; Ateş et al., 2018).

Some pieces of evidence indicate that the close interaction between muscular and fascial structures and the physical continuity of connective tissue along the myofascial chain weakens with age, thus reducing the magnitude of mechanical force transmission (Wilke et al., 2019). In this context, it has been shown that alteration in myofascial force transmission can influence the effects of nonlocal exercises, as the self-myofascial release of the plantar fascia on hamstring extensibility (Wilke et al., 2019). Fascial tissue can densify and develop fibrosis with age, thus reducing muscular force production and joint range of motion (Pavan et al., 2014; Zhang and Gao, 2014). Moreover, decreased physical mobility occurring in the elderly could be partially explained by increased stiffness and reduced elasticity of the extracellular matrix due to dehydration and increased collagen content (Sölch, 2015). Indeed, during aging, the connective tissue of patellar tendon increases fiber cross-links and reduces collagen content, and Achilles tendon and plantar fascia diminish their connecting fibers (Snow et al., 1995; Coupepe et al., 2009).

CONNECTIVE TISSUE, INFLAMMATION, AND AGING

A key feature of aging tissue is, also, the so-called inflammaging, which describes a low-grade chronic systemic inflammation in the absence of overt infection. This “sterile” inflammation is a highly significant risk factor for morbidity and mortality in elderly people. Chronic inflammation influences tissue functions *via* several mechanisms: persistent production of reactive molecules by infiltrating leucocytes might damage structural elements of tissues; production of cytokines might modulate inflammatory responses and alter phenotypes of nearby cells (Franceschi and Campisi, 2014). Most of the inflammatory responses take place in the extracellular matrix, which can interact with immune cells and change their functions, thereby influencing tissue regeneration. Although early inflammation after tissue damage is important for remodeling and adaptations, decreased inflammation seems to be associated with improved tissue regeneration and gains of muscle strength (Zügel et al., 2018).

MOTOR SYSTEM AND AGING

As for the effect of aging on the motors system, data showed that atrophy of the motor cortex, modifications of neurotransmission, reduced motor cortical excitability, and plasticity occur in the nervous system. These changes also result in impaired muscle performance (Tieland et al., 2018). Alterations in nervous system structure and function contribute to a decline in skeletal muscle efficiency with age, *via* a reduction in motor coordination and muscle strength.

Loss of neural processes impairs the control and activity of skeletal muscle fibers. Indeed, the occurrence of age-related reduction in descending drive and an increase in the threshold of excitability of the corticospinal tract has been reported by studies in humans (Rossini et al., 1992; Clark and Taylor, 2011; Unhjem et al., 2015). As demonstrated in mouse and rat models, aging is also associated with a decline in reinnervation of muscle fibers, accumulation of denervated fibers, and muscle atrophy. This phenomenon is probably caused by a reduced neurotrophin-mediated axonal sprouting following denervation (Aare et al., 2016). Moreover, aging leads to motor unit remodeling, which leads to an alteration in the fiber type composition, i.e., the replacement of type II with type I muscle fibers (Thompson, 2002). Reduction in the number of motor units is associated with the decay of muscle size and function in aged mice (Sheth et al., 2018). Aging also results in impaired movement coordination, due to a decreased control of motor output, as demonstrated in the first dorsal interosseous muscle of elderly subjects during submaximal contractions (Galganski et al., 1993). Old adults, have 40% reduction in the total number of motor units, 50% enlargement of remaining motor units (low- and moderate-threshold motor units), and an increase in fiber density (Piasecki et al., 2016). Also, differences in inter-joint coordination during squat jumps have been found between young and old adults (Argaud et al., 2019). A study investigating neuromuscular coordination demonstrated a loss and reorganization of muscular patterns in old adults (Vernooij et al., 2016).

Neuromuscular junctions (NMJs) are affected by detrimental modifications in their activity and conformation during aging (Li et al., 2018). Studies in animal models showed that the structural integrity of NMJs is affected by aging. The area of the motor nerve terminal and that of post-synaptic folds is reduced and both nerve terminals and the post-synaptic cluster of acetylcholine receptors (AChRs) undergo fragmentation. Moreover, in aged rodent it has been reported an increase in the branching of nerve terminals, and extra-junctional AChRs, and a reduction in myofibre nerve supply (Kreko-Pierce and Eaton, 2018). Interestingly, the degree of age-dependent degeneration of NMJs shows a certain variability between different muscles and even within the same muscle or the motor unit (Taetzsch and Valdez, 2018). The cause of this NMJs degeneration is still a matter of study, but some pieces of evidence indicate that impaired autophagic pathway and agrin signaling might cause NMJ decline (Li et al., 2018). Data from animal models showed that aged NMJs have a reduced number of mitochondria and accumulate morphological abnormal mitochondria and oxidative damage in the presynaptic plaque (Gonzalez-Freire et al., 2014). Also, it has been reported that aging is associated with altered expression of neurotrophin genes in the neuromuscular system. In particular, neurotrophin-3, neurotrophin-4, and brain-derived neurotrophic factors are downregulated and glial cell-derived neurotrophic factor is upregulated in aged rodent muscles (Kreko-Pierce and Eaton, 2018). The effect of aging on neuromuscular junction transmissions is still debated, due to conflicting results

(Piasecki et al., 2016; Willadt et al., 2016). The results collected so far demonstrate that age-dependent changes in the nervous system could promote an impairment in skeletal muscle function (Piasecki et al., 2016). However, in the perspective of healthy aging, several studies in mice demonstrated that the deleterious effects of aging on the structure and function of NMJs can regress due to physical exercise (Fahim, 1997; Valdez et al., 2010; Nishimune et al., 2012; Cheng et al., 2013).

IMAGING AND ULTRASOUND METHODS IN THE ELDERLY

From the clinical perspective, it is crucial to identify all the alterations that occur in the myofascial tissue with aging. Technological advancements allowed the development of high-resolution instruments for the morphological characterization of myofascia tissue and the assessment of factors that influence the “contractile quality” of skeletal muscle. Conventional imaging methods, such as MRI, the current “gold standard” for the analysis of muscle size, computed tomography (CT), and dual-energy X-ray absorptiometry (DXA), provide precise measures of muscle mass but are suboptimal in reflecting the composition of myofascial tissue. For instance, Hounsfield units in CT images (Engel et al., 2018) or MRI signal intensity can be used to evaluate the composition of the skeletal muscle (Carrier et al., 2016). However, it is difficult to obtain these values for many community-dwelling older people because of the need for special equipment and the relatively high measurement burden (time, cost, and radiation exposure) and the lack of standardization of measurements (Lee et al., 2019). Recently, ultrasound attenuation and echo intensity have emerged as potentially useful indicators of muscle composition. Ultrasonography is non-invasive, easily accessible, and relatively inexpensive compared with other imaging systems such as CT. Several groups of researchers have used ultrasound echo intensity as an index of skeletal muscle composition; whereby, increased amounts of intramuscular fat and fibrosis that occur with pathological aging result in increasing echo-reflection and a brighter ultrasound image (Reimers et al., 1993; Pillen et al., 2009; Watanabe et al., 2018). Although such measures have also been related to muscle strength and functional performance in elderly populations in some studies (Fukumoto et al., 2012; Watanabe et al., 2013; Lopez et al., 2017), they provide only indirect measures of the contractile quality of skeletal muscle *in vivo*. Several alternative ultrasound-based methods have recently been developed to quantify the contractile quality of muscle during dynamic contraction. Elastographic methods are increasingly used to assess the viscoelastic properties of muscle in aging but, as with echo-based measures, their widespread implementation in clinical and research settings is currently limited by low reproducibility (Zaidman et al., 2012; Alfuraih et al., 2018). Although other emerging ultrasound-based approaches including speckle-tracking (Frich et al., 2019), tissue Doppler imaging (Eranki et al., 2013), and axial transmission techniques

(Wearing et al., 2016) arguably provide more direct measures of the contractile quality of the muscle-tendon unit *in vivo*, these techniques also suffer from limitations, including susceptibility to decorrelation (speckle tracking), user intervention (feature-based tracking), and dependence on insonation angle (tissue Doppler) (Sikdar et al., 2014). Nonetheless, ultrasound-based measures appear to be promising approaches for quantifying contractile properties of myofascial tissues and have the potential to enhance our understanding of musculoskeletal function with aging and pathology.

MANIPULATION TECHNIQUES AND FASCIA REPAIR

Over the course of a lifetime, fascia can be injured due to excessive or prolonged loading, traumatic events, strenuous physical activity, and surgical procedures. As a consequence of damage, repair mechanisms are activated to restore the original structural and functional features of the tissue (Zullo et al., 2017). Impairment of this process can cause a reduction in the performance of the locomotory apparatus and musculoskeletal disorders (Zuegel et al., 2018). Therefore, strategies improving myofascial regeneration are pivotal. A broad range of tissue manipulation techniques has been proposed to enhance fascial repair. A study in the treatment of tension-type headache suggests a combination of soft tissue techniques and neural mobilization to be most promising in relieving myofascial-induced pain and dysfunction (Ferragut-Garcias et al., 2017). The authors highlight the importance of the treatment stimulus to mechanically stimulate nerve and fascia. Indeed, a study performed on people suffering from delayed-onset muscle soreness showed that the sensitivity of high-threshold mechanosensitive receptors is a predictor of pain and motor impairment (Fleckenstein et al., 2017). Thus, approaches modulating the activity of these receptors may be helpful for functional recovery. This is in line with recent clinical data showing that hands-on based conservative treatments can be effective in relieving pain in injured athletes (Fleckenstein and Banzer, 2019). These treatments include nerve block, injection, ultrasound and laser therapies, manipulation, mobilization, massage, and traction, as well as acupuncture and dry needling. However, it is not clear whether aging influences the effect of these treatments. There is evidence that multimodal rehabilitation, including classic massage, transcutaneous electrical nerve stimulation, and ultrasound therapy improves pain and function in older women (aged >60) suffering from back pain (Cichon et al., 2019). A systematic review found limited evidence of pain-reducing effects of physical therapy (three studies, two of them applying soft tissue treatments) among older adults with dementia (Coronado et al., 2019). In summary, we can only hypothesize that the fascial tissue of elderly people is susceptible to soft tissue stimuli, but its effect has to be determined.

CONCLUSIONS

Aging is associated with metabolic, structural, and functional modifications of cells, tissues, and organs, which lead to a gradual decline in psycho-physical performance. In particular, the locomotory apparatus loses its effectiveness, due to the molecular and cellular changes occurring in the myofascia, the skeletal muscle tissue, the nervous system, and their structural and functional coupling. Genetics, epigenetics, environment, diseases, lifestyle, nutrition, and injuries also have a prominent role in tissue remodeling occurring with aging (Jee and Kim, 2017). Thanks to recent scientific progress, many phenomena and mechanisms associated with aging have been defined, but still much remains to be investigated. From the perspective of healthy aging, it is crucial to identify and hinder all the age-dependent modifications through specific strategies targeting etiologic factors, and also psycho-social issues. Indeed, ultrasound-based techniques can provide a detailed morphological characterization of skeletal muscle and connective tissue, thus allowing a specific analysis of the detrimental changes occurring in the myofascia with aging. Moreover, tissue

manipulation techniques might contribute to improving myofascial regeneration in the elderly. Physical activity has been also suggested as an effective strategy for counteracting the deleterious consequences of aging, given that skeletal muscle plasticity might be only partially lost in elderly individuals (Distefano and Goodpaster, 2018). However, this issue is still debated, due to contradictory results. Nevertheless, the constant progress in technology and biomedical research holds great promise for fighting the burden of aging by targeted therapeutic interventions.

AUTHOR CONTRIBUTIONS

All authors contributed substantially to the design and concept of the review. WK, AZ, RS, and KH drafted the manuscript. SW and JF critically revised the manuscript. All authors approved the content of the final version of the manuscript. All authors agree to be accountable for all aspects for the work and ensure that all questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

REFERENCES

- Aare, S., Spendiff, S., Vuda, M., Elkrief, D., Perez, A., Wu, Q., et al. (2016). Failed reinnervation in aging skeletal muscle. *Skelet. Muscle* 6:29. doi: 10.1186/s13395-016-0101-y
- Agyapong-Badu, S., Warner, M., Samuel, D., and Stokes, M. (2016). Measurement of ageing effects on muscle tone and mechanical properties of rectus femoris and biceps brachii in healthy males and females using a novel hand-held myometric device. *Arch. Gerontol. Geriatr.* 62, 59–67. doi:10.1016/j.archger.2015.09.011
- Alfuraih, A. M., O'Connor, P., Hensor, E., Tan, A. L., Emery, P., and Wakefield, R. J. (2018). The effect of unit, depth, and probe load on the reliability of muscle shear wave elastography: variables affecting reliability of SWE. *J. Clin. Ultrasound* 46, 108–115. doi: 10.1002/jcu.22534
- Alnaqeeb, M. A., Al Zaid, N. S., and Goldspink, G. (1984). Connective tissue changes and physical properties of developing and ageing skeletal muscle. *J. Anat.* 139, 677–689.
- Anton, S. D., Woods, A. J., Ashizawa, T., Barb, D., Buford, T. W., Carter, C. S., et al. (2015). Successful aging: advancing the science of physical independence in older adults. *Ageing Res. Rev.* 24, 304–327. doi: 10.1016/j.arr.2015.09.005
- Argaud, S., Pairot de Fontenay, B., Blache, Y., and Monteil, K. (2019). Age-related differences of interjoint coordination in elderly during squat jumping. *PLoS One* 14:e0221716. doi: 10.1371/journal.pone.0221716
- Ateş, F., Andrade, R. J., Freitas, S. R., Hug, F., Lacourpaille, L., Gross, R., et al. (2018). Passive stiffness of monoarticular lower leg muscles is influenced by knee joint angle. *Eur. J. Appl. Physiol.* 118, 585–593. doi: 10.1007/s00421-018-3798-y
- Avila Gonzalez, C. A., Driscoll, M., Schleip, R., Wearing, S., Jacobson, E., Findley, T., et al. (2018). Frontiers in fascia research. *J. Bodyw. Mov. Ther.* 22, 873–880. doi: 10.1016/j.jbmt.2018.09.077
- Bank, R. A., TeKoppele, J. M., Oostingh, G., Hazleman, B. L., and Riley, G. P. (1999). Lysylhydroxylation and non-reducible cross-linking of human supraspinatus tendon collagen: changes with age and in chronic rotator cuff tendinitis. *Ann. Rheum. Dis.* 58, 35–41.
- Barros, E. M., Rodrigues, C. J., Rodrigues, N. R., Oliveira, R. P., Barros, T. E., and Rodrigues, A. J. Jr. (2002). Aging of the elastic and collagen fibers in the human cervical interspinous ligaments. *Spine J.* 2, 57–62. doi: 10.1016/S1529-9430(01)00167-X
- Beavers, K. M., Beavers, D. P., Houston, D. K., Harris, T. B., Hue, T. F., Koster, A., et al. (2013). Associations between body composition and gait-speed decline: results from the health, aging, and body composition study. *Am. J. Clin. Nutr.* 97, 552–560. doi: 10.3945/ajcn.112.047860
- Birch, H. L. (2018). Extracellular matrix and ageing. *Subcell. Biochem.* 90, 169–190. doi: 10.1007/978-981-13-2835-0_7
- Blottner, D., Huang, Y., Trautmann, G., and Sun, L. (2019). The fascia: Continuum linking bone and myofascial bag for global and local body movement control on Earth and in Space. A scoping review. *REACH.* 14–15. doi: 10.1016/j.reach.2019.100030
- Bojsen-Møller, J., Schwartz, S., Kalliokoski, K. K., Finni, T., and Magnusson, S. P. (2010). Intermuscular force transmission between human plantarflexor muscles in vivo. *J. Appl. Physiol.* 109, 1608–1618. doi: 10.1152/jappphysiol.01381.2009
- Brack, A. S., and Muñoz-Cánoves, P. (2016). The ins and outs of muscle stem cell aging. *Skelet. Muscle* 6:1. doi: 10.1186/s13395-016-0072-z
- Bratic, A., and Larsson, N. G. (2013). The role of mitochondria in aging. *J. Clin. Invest.* 123, 951–957. doi: 10.1172/JCI64125
- Brown, B. N., Ratner, B. D., Goodman, S. B., Amar, S., and Badylak, S. F. (2012). Macrophage polarization: an opportunity for improved outcomes in biomaterials and regenerative medicine. *Biomaterials* 33, 3792–3802. doi: 10.1016/j.biomaterials.2012.02.034
- Canepari, M., Pellegrino, M. A., D'Antona, G., and Bottinelli, R. (2010). Single muscle fiber properties in aging and disuse. *Scand. J. Med. Sci. Sports* 20, 10–19. doi: 10.1111/j.1600-0838.2009.00965.x
- Carlier, P. G., Marty, B., Scheidegger, O., Loureiro de Sousa, P., Baudin, P. Y., Snezhko, E., et al. (2016). Skeletal muscle quantitative nuclear magnetic resonance imaging and spectroscopy as an outcome measure for clinical trials. *J. Neuromuscul. Dis.* 3, 1–28. doi: 10.3233/JND-160145
- Cartee, G. D., Hepple, R. T., Bamman, M. M., and Zierath, J. R. (2016). Exercise promotes healthy aging of skeletal muscle. *Cell Metab.* 23, 1034–1047. doi: 10.1016/j.cmet.2016.05.007
- Cerullo, F., Gambassi, G., and Cesari, M. (2012). Rationale for antioxidant supplementation in sarcopenia. *J. Aging Res.* 2012:316943. doi: 10.1155/2012/316943
- Cesari, M., Pahor, M., Bartali, B., Cherubini, A., Penninx, B. W., Williams, G. R., et al. (2004). Antioxidants and physical performance in elderly persons: the invecchiare in chianti (InCHIANTI) study. *Am. J. Clin. Nutr.* 79, 289–294. doi: 10.1093/ajcn/79.2.289
- Cheng, A., Morsch, M., Murata, Y., Ghazanfari, N., Reddel, S. W., and Phillips, W. D. (2013). Sequence of age-associated changes to the mouse neuromuscular junction and the protective effects of voluntary exercise. *PLoS One* 8:e67970. doi: 10.1371/journal.pone.0067970
- Cichon, D., Ignasiak, Z., Fugiel, J., Kochan, K., and Ignasiak, T. (2019). Efficacy of physiotherapy in reducing Back pain and improve joint mobility in older women. *Ortop. Traumatol. Rehabil.* 21, 45–55. doi: 10.5604/01.3001.0013.1115

- Clafin, D. R., Jackson, M. J., and Brooks, S. V. (2015). Age affects the contraction-induced mitochondrial redox response in skeletal muscle. *Front. Physiol.* 6:21. doi: 10.3389/fphys.2015.00021
- Clark, B. C., and Taylor, J. L. (2011). Age-related changes in motor cortical properties and voluntary activation of skeletal muscle. *Curr. Aging Sci.* 4, 192–199. doi: 10.2174/1874609811104030192
- Cohen, S., Nathan, J. A., and Goldberg, A. L. (2015). Muscle wasting in disease: molecular mechanisms and promising therapies. *Nat. Rev. Drug Discov.* 14, 58–74. doi: 10.1038/nrd4467
- Conley, K. E., Jubrias, S. A., and Esselman, P. C. (2000). Oxidative capacity and ageing in human muscle. *J. Physiol.* 526, 203–210. doi: 10.1111/j.1469-7793.2000.t01-1-00203.x
- Corbu, A., Scaramozza, A., Badiali-DeGiorgi, L., Tarantino, L., Papa, V., Rinaldi, R., et al. (2010). Satellite cell characterization from aging human muscle. *Neurol. Res.* 32, 63–72. doi: 10.1179/174313209X385725
- Coronado, R. A., Albers, H. E., Allen, J. L., Clarke, R. G., Estrada, V. A., Simon, C. B., et al. (2019). Pain-reducing effects of physical therapist-delivered interventions: a systematic review of randomized trials among older adults with dementia. *J. Geriatr. Phys. Ther.* doi: 10.1519/JPT.0000000000000235 [Epub ahead of print].
- Coupe, C., Hansen, P., Kongsgaard, M., Kovanen, V., Suetta, C., Aagaard, P., et al. (2009). Mechanical properties and collagen cross-linking of the patellar tendon in old and young men. *J. Appl. Physiol.* 107, 880–886. doi: 10.1152/jappphysiol.00291.2009
- Csapo, R., Malis, V., Sinha, U., Du, J., and Sinha, S. (2014). Age-associated differences in triceps surae muscle composition and strength - an MRI-based cross-sectional comparison of contractile, adipose and connective tissue. *BMC Musculoskelet. Disord.* 15:209. doi: 10.1186/1471-2474-15-209
- Csernoch, L., Fodor, J., Al-Gaadi, D., Czirájk, T., Oláh, T., Dienes, B., et al. (2017). Modified calcium homeostasis in aged mouse skeletal muscle. *Biophys. J.* 112:99a. doi: 10.1016/j.bpj.2016.11.573
- D'Antona, G., Pellegrino, M. A., Adami, R., Rossi, R., Carlizzi, C. N., Canepari, M., et al. (2003). The effect of ageing and immobilization on structure and function of human skeletal muscle fibres. *J. Physiol.* 552, 499–511. doi: 10.1113/jphysiol.2003.046276
- Damiano, S., Muscariello, E., La Rosa, G., Di Maro, M., Mondola, P., and Santillo, M. (2019). Dual role of reactive oxygen species in muscle function: can antioxidant dietary supplements counteract age-related sarcopenia? *Int. J. Mol. Sci.* 20:E3815. doi: 10.3390/ijms20153815
- de Sousa Neto, I. V., Durigan, J. L. Q., Guzzoni, V., Tibana, R. A., Prestes, J., de Araujo, H. S. S., et al. (2018). Effects of resistance training on matrix metalloproteinase activity in skeletal muscles and blood circulation during aging. *Front. Physiol.* 9:190. doi: 10.3389/fphys.2018.00190
- DeGroot, J., Verzijl, N., Jacobs, K. M., Budde, M., Bank, R. A., Bijlsma, J. W., et al. (2001). Accumulation of advanced glycation endproducts reduces chondrocyte-mediated extracellular matrix turnover in human articular cartilage. *Osteoarthritis. Cartil.* 9, 720–726. doi: 10.1053/joca.2001.0469
- Del Campo, A., Contreras-Hernández, I., Castro-Sepúlveda, M., Campos, C. A., Figueroa, R., Tevy, M. F., et al. (2018). Muscle function decline and mitochondria changes in middle age precede sarcopenia in mice. *Aging* 10, 34–55. doi: 10.18632/aging.101358
- Del Carmen Ortuño-Costela, M., García-López, M., Cerrada, V., and Gallardo, M. E. (2019). iPSCs: a powerful tool for skeletal muscle tissue engineering. *J. Cell. Mol. Med.* 23, 3784–3794. doi: 10.1111/jcmm.14292
- Delmonico, M., Harris, T. B., Visser, M., Park, S. W., Conroy, M. B., Velasquez-Mieyer, P., et al. (2009). Longitudinal study of muscle strength, quality, and adipose tissue infiltration. *Am. J. Clin. Nutr.* 90, 1579–1585. doi: 10.3945/ajcn.2009.28047
- Diehr, P. H., Thielke, S. M., Newman, A. B., Hirsch, C., and Tracy, R. (2013). Decline in health for older adults: five-year change in 13 key measures of standardized health. *J. Gerontol. A. Biol. Sci. Med. Sci.* 68, 1059–1067. doi: 10.1093/gerona/glt038
- Distefano, G., and Goodpaster, B. H. (2018). Effects of exercise and aging on skeletal muscle. *Cold Spring Harb. Perspect. Med.* 8:a029785. doi: 10.1101/cshperspect.a029785
- Doherty, T. J. (2003). Invited review: aging and sarcopenia. *J. Appl. Physiol.* 95, 1717–1727. doi: 10.1152/jappphysiol.00347.2003
- Ducomps, C., Mauriege, P., Darche, B., Combes, S., Lebas, F., and Doutreloux, J. P. (2003). Effects of jump training on passive mechanical stress and stiffness in rabbit skeletal muscle: role of collagen. *Acta Physiol. Scand.* 178, 215–224. doi: 10.1046/j.1365-201X.2003.01109.x
- Dziki, J., Badylak, S., Yabroudi, M., Sicari, B., Ambrosio, F., Stearns, K., et al. (2016). An acellular biologic scaffold treatment for volumetric muscle loss: results of a 13-patient cohort study. *NPJ Regen. Med.* 1:16008. doi: 10.1038/npjregenmed.2016.8
- Engel, K., Musey, O., Wang, L., and Laredod, J.-D. (2018). Quantitative analysis of skeletal muscle by computed tomography imaging—state of the art. *J. Orthop. Translat.* 15, 91–103. doi: 10.1016/j.jot.2018.10.004
- Eranki, A., Cortes, N., Ferenček, Z. G., and Sikdar, S. (2013). A novel application of musculoskeletal ultrasound imaging. *J. Vis. Exp.* 79:50595. doi: 10.3791/50595
- Etienne, J., Liu, C., Skinner, C. M., Conboy, M. J., and Conboy, I. M. (2020). Skeletal muscle as an experimental model of choice to study tissue aging and rejuvenation. *Skelet. Muscle* 10:4. doi: 10.1186/s13395-020-0222-1
- Eyre, D. R., Paz, M. A., and Gallop, P. M. (1984). Cross-linking in collagen and elastin. *Annu. Rev. Biochem.* 53, 717–748. doi: 10.1146/annurev.bi.53.070184.003441
- Fahim, M. A. (1997). Endurance exercise modulates neuromuscular junction of C57BL/6Nnia aging mice. *J. Appl. Physiol.* 83, 59–66. doi: 10.1152/jappl.1997.83.1.59
- Ferragut-Garcias, A., Plaza-Manzano, G., Rodriguez-Blanco, C., Velasco-Roldan, O., Pecos-Martin, D., Oliva-Pascual-Vaca, J., et al. (2017). Effectiveness of a treatment involving soft tissue techniques and/or neural mobilization techniques in the management of tension-type headache: a randomized controlled trial. *Arch. Phys. Med. Rehabil.* 98, 211–9.e2. doi: 10.1016/j.apmr.2016.08.466
- Ferraro, E., Giammarioli, A. M., Chiandotto, S., Spoletini, I., and Rosano, G. (2014). Exercise-induced skeletal muscle remodeling and metabolic adaptation: redox signaling and role of autophagy. *Antioxid. Redox Signal.* 21, 154–176. doi:10.1089/ars.2013.5773
- Flack, K. D., Davy, B. M., DeBerardinis, M., Boutagy, N. E., McMillan, R. P., Hulver, M. W., et al. (2016). Resistance exercise training and in vitro skeletal muscle oxidative capacity in older adults. *Phys. Rep.* 4:e12849. doi: 10.14814/phy2.12849
- Fleckenstein, J., and Banzer, W. (2019). A review of hands-on based conservative treatments on pain in recreational and elite athletes. *Sci. Sports* 34, E77–E100. doi: 10.1016/j.scispo.2018.08.004
- Fleckenstein, J., Simon, P., König, M., Vogt, L., and Banzer, W. (2017). The pain threshold of high-threshold mechanosensitive receptors subsequent to maximal eccentric exercise is a potential marker in the prediction of DOMS associated impairment. *PLoS One* 12:e0185463. doi: 10.1371/journal.pone.0185463
- Franceschi, C., and Campisi, J. (2014). Chronic inflammation (inflammaging) and its potential contribution to age-associated diseases. *J. Gerontol. A. Biol. Sci. Med. Sci.* 63(Suppl. 1), 54–59. doi: 10.1093/gerona/glu057
- Frich, L. H., Lambertsen, K. L., Hjarbaek, J., Dahl, J. S., and Holsgaard-Larsen, A. (2019). Musculoskeletal application and validation of speckle-tracking ultrasonography. *BMC Musculoskelet. Disord.* 20:192. doi: 10.1186/s12891-019-2562-8
- Frontera, W. R., Huges, V. A., Fielding, R. A., Fiatarone, M. A., Evans, W. J., and Roubenoff, R. (2000). Aging of skeletal muscle: a 12-yr longitudinal study. *J. Appl. Physiol.* 88, 1321–1326. doi: 10.1152/jappl.2000.88.4.1321
- Frontera, W. R., Meredith, C. N., O'Reilly, K. P., Knuttgen, H. G., and Evans, W. J. (1988). Strength conditioning in older men: skeletal muscle hypertrophy and improved function. *J. Appl. Physiol.* 64, 1038–1044.
- Fukumoto, Y., Ikezoe, T., Yamada, Y., Tsukagoshi, R., Nakamura, M., Mori, N., et al. (2012). Skeletal muscle quality assessed from echo intensity is associated with muscle strength of middle-aged and elderly persons. *Eur. J. Appl. Physiol.* 112, 1519–1525. doi: 10.1007/s00421-011-2099-5
- Fuoco, C., Petrilli, L. L., Cannata, S., and Gargioli, C. (2016). Matrix scaffolding for stem cell guidance toward skeletal muscle tissue engineering. *J. Orthop. Surg. Res.* 11:86. doi: 10.1186/s13018-016-0421-y
- Galganski, M. E., Fuglevand, A. J., and Enoka, R. M. (1993). Reduced control of motor output in a human hand muscle of elderly subjects during submaximal contractions. *J. Neurophysiol.* 69, 2108–2115. doi: 10.1152/jn.1993.69.6.2108
- Gheller, B. J., Riddle, E. S., Lem, M. R., and Thalacker-Mercer, A. E. (2016). Understanding age-related changes in skeletal muscle metabolism: differences between females and males. *Annu. Rev. Nutr.* 36, 129–156. doi: 10.1146/annurev-nutr.071715-050901
- Godard, M. P., Gallagher, P. M., Raue, U., and Trappe, S. W. (2002). Alterations in single muscle fiber calcium sensitivity with resistance training in older women. *Pflugers Arch.* 444, 419–425. doi: 10.1007/s00424-002-0821-1

- Goldspink, G., Fernandes, K., Williams, P. E., and Wells, D. J. (1994). Age-related changes in collagen gene expression in the muscles of mdx dystrophic and normal mice. *Neuromuscul. Disord.* 4, 183–191. doi: 10.1016/0960-8966(94)90019-1
- Gomes, M. J., Martinez, P. F., Pagan, L. U., Damatto, R. L., Cezar, M. D. M., Lima, A. R. R., et al. (2017). Skeletal muscle aging: influence of oxidative stress and physical exercise. *Oncotarget* 8, 20428–20440. doi: 10.18632/oncotarget.14670
- Gonzalez-Freire, M., de Cabo, R., Studenski, S. A., and Ferrucci, L. (2014). The neuromuscular junction: aging at the crossroad between nerves and muscle. *Front. Aging Neurosci.* 6:208. doi: 10.3389/fnagi.2014.00208
- Goodpaster, B. H., Park, S. W., Harris, T. B., Kritchevsky, S. B., Nevitt, M., Schwartz, A. V., et al. (2006). The loss of skeletal muscle strength, mass, and quality in older adults: the health, aging and body composition study. *Gerontol. A. Biol. Sci. Med. Sci.* 61, 1059–1064. doi: 10.1093/gerona/61.10.1059
- Gosselin, L. E., Adams, C., Cotter, T. A., McCormick, R. J., and Thomas, D. P. (1998). Effect of exercise training on passive stiffness in locomotor skeletal muscle: role of extracellular matrix. *J. Appl. Physiol.* 85, 1011–1016. doi: 10.1152/jap.1998.85.3.1011
- Grzelkowska-Kowalczyk, K. (2016). “The importance of extracellular matrix in skeletal muscle development and function” in *Composition and function of the extracellular matrix in the human body*. ed. F. Travasico (London, UK: IntechOpen).
- Hall, M. N., Hall, J. K., Cadwallader, A. B., Pawlikowski, B. T., Doles, J. D., Elston, T. L., et al. (2017). Transplantation of skeletal muscle stem cells. *Methods Mol. Biol.* 1556, 237–244. doi: 10.1007/978-1-4939-6771-1_12
- Haus, J. M., Carrithers, J. A., Miller, B. I., and Trappe, T. A. (2007a). Intramuscular collagen cross-linking in aging men and women. *FASEB J.* 21, LB118–LB118.
- Haus, J. M., Carrithers, J. A., Trappe, S. W., and Trappe, T. A. (2007b). Collagen, cross-linking, and advanced glycation end products in aging human skeletal muscle. *J. Appl. Physiol.* 103, 2068–2076. doi: 10.1152/jap.2007.103.4.2068
- Hausman, G. J., Basu, U., Du, M., Fernyhough-Culver, M., and Dodson, M. V. (2014). Intermuscular and intramuscular adipose tissues: bad vs. good adipose tissues. *Adipocytes* 3, 242–255. doi: 10.4161/adip.28546
- Haut, R. C., Lancaster, R. L., and DeCamp, C. E. (1992). Mechanical properties of the canine patellar tendon: some correlations with age and the content of collagen. *J. Biomech.* 25, 163–173. doi: 10.1016/0021-9290(92)90273-4
- Hinkle, R. T., Donnelly, E., Cody, D. B., Bauer, M. B., and Isfort, R. J. (2003). Urocortin II treatment reduces skeletal muscle mass and function loss during atrophy and increases nonatrophying skeletal muscle mass and function. *Endocrinology* 144, 4939–4946. doi: 10.1210/en.2003-0271
- Huijing, P. A. (2009). Epimuscular myofascial force transmission: a historical review and implications for new research. International Society of Biomechanics Muybridge Award Lecture, Taipei, 2007. *J. Biomech.* 42, 9–21. doi: 10.1016/j.jbiomech.2008.09.027
- Huijing, P. A., and Jaspers, R. T. (2005). Adaptation of muscle size and myofascial force transmission: a review and some new experimental results. *Scand. J. Med. Sci. Sports* 15, 349–380. doi: 10.1111/j.1600-0838.2005.00457.x
- Huijing, P. A., Maas, H., and Baan, G. C. (2003). Compartmental fasciotomy and isolating a muscle from neighboring muscles interfere with myofascial force transmission within the rat anterior crural compartment. *J. Morphol.* 256, 306–321. doi: 10.1002/jmor.10097
- Ippolito, E., Natali, P. G., Postacchini, F., Accinni, L., and De Martino, C. (1980). Morphological, immunochemical, and biochemical study of rabbit achilles tendon at various ages. *J. Bone Joint Surg. Am.* 62, 583–598. doi: 10.2106/00004623-198062040-00014
- Janssen, I., Heymsfield, S. B., and Ross, R. (2002). Low relative skeletal muscle mass (sarcopenia) in older persons is associated with functional impairment and physical disability. *J. Am. Geriatr. Soc.* 50, 889–896. doi: 10.1046/j.1532-5415.2002.50216.x
- Janssen, I., Heymsfield, S. B., Wang, Z. M., and Ross, R. (2000). Skeletal muscle mass and distribution in 468 men and women aged 18–88 yr. *J. Appl. Physiol.* 89, 81–88. doi: 10.1152/jap.2000.89.1.81
- Jee, H., and Kim, J. H. (2017). A mini-overview of single muscle fibre mechanics: the effects of age, inactivity and exercise in animals and humans. *Swiss Med. Wkly.* 147:w14488. doi: 10.4414/SMW.2017.14488
- Ji, L. L. (2015). Redox signaling in skeletal muscle: role of aging and exercise. *Adv. Physiol. Educ.* 39, 352–359. doi: 10.1152/advan.00106.2014
- Jiao, A., Moerk, C. T., Penland, N., Perla, M., Kim, J., Smith, A. S. T., et al. (2018). Regulation of skeletal myotube formation and alignment by nanotopographically controlled cell-seeded extracellular matrix. *J. Biomed. Mater. Res. A* 106, 1543–1551. doi: 10.1002/jbm.a.36351
- Joanisse, S., Nederveen, J. P., Baker, J. M., Sniijders, T., Iacono, C., and Parise, G. (2016). Exercise conditioning in old mice improves skeletal muscle regeneration. *FASEB J.* 30, 3256–3268. doi: 10.1096/fj.201600143RR
- Joanisse, S., Nederveen, J. P., Sniijders, T., McKay, B. R., and Parise, G. (2017). Skeletal muscle regeneration, repair and remodelling in aging: the importance of muscle stem cells and vascularization. *Gerontology* 63, 91–100. doi: 10.1159/000450922
- Joanisse, S., and Parise, G. (2016). Cytokine mediated control of muscle stem cell function. *Adv. Exp. Med. Biol.* 900, 27–44. doi: 10.1007/978-3-319-27511-6_2
- Joseph, A. M., Adhithetty, P. J., and Leeuwenburgh, C. (2016). Beneficial effects of exercise on age-related mitochondrial dysfunction and oxidative stress in skeletal muscle. *J. Physiol.* 594, 5105–5123. doi: 10.1113/JP270659
- Karakuzu, A., Pamuk, U., Ozturk, C., Acar, B., and Yucesoy, C. A. (2017). Magnetic resonance and diffusion tensor imaging analyses indicate heterogeneous strains along human medial gastrocnemius fascicles caused by submaximal plantar-flexion activity. *J. Biomech.* 57, 69–78. doi: 10.1016/j.jbiomech.2017.03.028
- Kaya, C. S., Bilgili, F., Akalan, N. E., Temelli, Y., Ateş, F., and Yucesoy, C. A. (2019). Intraoperative experiments combined with gait analyses indicate that active state rather than passive dominates the spastic gracilis muscle’s joint movement limiting effect in cerebral palsy. *Clin. Biomech.* 68, 151–157. doi: 10.1016/j.clinbiomech.2019.06.005
- Kaya, C. S., Bilgili, F., Akalan, N. E., and Yucesoy, C. A. (2020). Intraoperative testing of passive and active state mechanics of spastic semitendinosus in conditions involving intermuscular mechanical interactions and gait relevant joint positions. *J. Biomech.* 103:109755. doi: 10.1016/j.jbiomech.2020.109755
- Keller, K., and Engelhardt, M. (2014). Strength and muscle mass loss with aging process. Age and strength loss. *Muscles Ligaments. Tendons J.* 3, 346–350. doi: 10.32098/mltj.04.2013.17
- Kim, J. H., and Thompson, L. V. (2013). Inactivity, age, and exercise: single-muscle fiber power generation. *J. Appl. Physiol.* 114, 90–98. doi: 10.1152/jap.2012.114.1.90
- Kim, Y., Triolo, M., and Hood, D. A. (2017). Impact of aging and exercise on mitochondrial quality control in skeletal muscle. *Oxidative Med. Cell. Longev.* 2017:3165396. doi: 10.1155/2017/3165396
- Kjaer, M. (2004). Role of extracellular matrix in adaptation of tendon and skeletal muscle to mechanical loading. *Physiol. Rev.* 84, 649–698. doi: 10.1152/physrev.00031.2003
- Klitgaard, H., Zhou, M., Schiaffino, S., Betto, R., Salviati, G., and Saltin, B. (1990). Ageing alters the myosin heavy chain composition of single fibres from human skeletal muscle. *Acta Physiol. Scand.* 140, 55–62. doi: 10.1111/j.1748-1716.1990.tb08975.x
- Kocur, P., Grzeskowiak, M., Wiernicka, M., Goliwas, M., Lewandowski, J., and Łochyński, D. (2017). Effects of aging on mechanical properties of sternocleidomastoid and trapezius muscles during transition from lying to sitting position—a cross-sectional study. *Arch. Gerontol. Geriatr.* 70, 14–18. doi: 10.1016/j.archger.2016.12.005
- Kocur, P., Tomczak, M., Wiernicka, M., Goliwas, M., Lewandowski, J., and Łochyński, D. (2019). Relationship between age, BMI, head posture and superficial neck muscle stiffness and elasticity in adult women. *Sci. Rep.* 9:8515. doi: 10.1038/s41598-019-44837-5
- Kovanen, V., Suominen, H., and Heikkinen, E. (1984). Mechanical properties of fast and slow skeletal muscle with special reference to collagen and endurance training. *J. Biomech.* 17, 725–735. doi: 10.1016/0021-9290(84)90103-9
- Kovanen, V., Suominen, H., Risteli, J., and Risteli, L. (1988). Type IV collagen and laminin in slow and fast skeletal muscle in rats—effects of age and life-time endurance training. *Coll. Relat. Res.* 8, 145–153. doi: 10.1016/S0174-173X(88)80026-8
- Kragstrup, T. W., Kjaer, M., and Mackey, A. L. (2011). Structural, biochemical, cellular and functional changes in skeletal muscle extracellular matrix with aging. *Scand. J. Med. Sci. Sports* 21, 749–757. doi: 10.1111/j.1600-0838.2011.01377.x
- Kreko-Pierce, T., and Eaton, B. A. (2018). Rejuvenation of the aged neuromuscular junction by exercise. *Cell Stress* 2, 25–33. doi: 10.15698/cst2018.02.123

- Kubo, K., Kanehisa, H., Azuma, K., Ishizu, M., Kuno, S. Y., Okada, M., et al. (2003). Muscle architectural characteristics in women aged 20–79 years. *Med. Sci. Sports Exerc.* 35, 39–44. doi: 10.1097/00005768-200301000-00007
- Larsson, L. (1978). Morphological and functional characteristics of the ageing skeletal muscle in man. A cross-sectional study. *Acta Physiol. Scand. Suppl.* 457, 1–36.
- Lauretani, F., Russo, C. R., Bandinelli, S., Bartali, B., Cavazzini, C., Di Iorio, A., et al. (1985). Age-associated changes in skeletal muscles and their effect on mobility: an operational diagnosis of sarcopenia. *J. Appl. Physiol.* 95, 1851–1860. doi: 10.1152/jappphysiol.00246.2003
- Le Sant, G., Nordez, A., Andrade, R., Hug, F., Freitas, S., and Gross, R. (2017). Stiffness mapping of lower leg muscles during passive dorsiflexion. *J. Anat.* 230, 639–650. doi: 10.1111/joa.12589
- Lee, K., Shin, Y., Huh, J., Sung, Y. S., Lee, I. S., Yoon, K. H., et al. (2019). Recent issues on body composition imaging for sarcopenia evaluation. *Korean J. Radiol.* 20, 205–217. doi: 10.3348/kjr.2018.0479
- Levi, N., Papismaiov, N., Solomonov, L., Sagi, I., and Krizhanovsky, V. (2020). The ECM path of senescence in aging: components and modifiers. *FEBS J.* doi: 10.1111/febs.15282
- Lexell, J. (1995). Human aging, muscle mass, and fiber type composition. *Gerontol. A. Biol. Sci. Med. Sci.* 50, 11–16. doi: 10.1093/gerona/50a.special_issue.11
- Lexell, J., Taylor, C. C., and Sjöström, M. (1988). What is the cause of the ageing atrophy? Total number, size and proportion of different fiber types studied in whole vastus lateralis muscle from 15- to 83-year-old men. *J. Neurol. Sci.* 84, 275–294. doi: 10.1016/0022-510x(88)90132-3
- Li, L., Xiong, W. C., and Mei, L. (2018). Neuromuscular junction formation, aging, and disorders. *Annu. Rev. Physiol.* 80, 159–188. doi: 10.1146/annurev-physiol-022516-034255
- Lindle, R. S., Metter, E. J., Lynch, N. A., Fleg, J. L., Fozard, J. L., Tobin, J., et al. (1997). Age and gender comparisons of muscle strength in 654 women and men aged 20–93 yr. *J. Appl. Physiol.* 83, 1581–1587. doi: 10.1152/jappphysiol.1997.83.5.1581
- Lopez, P., Wilhelm, E. N., Rech, A., Minozzo, F., Radaelli, R., and Pinto, R. S. (2017). Echo intensity independently predicts functionality in sedentary older men. *Muscle Nerve* 55, 9–15. doi: 10.1002/mus.25168
- Malisoux, L., Jamart, C., Delplace, K., Nielsen, H., Francaux, M., and Theisen, D. (2007). Effect of long-term muscle paralysis on human single fiber mechanics. *J. Appl. Physiol.* 102, 340–349. doi: 10.1152/jappphysiol.00609.2006
- Marcell, T. J. (2003). Sarcopenia: causes, consequences, and preventions. *J. Gerontol. A. Biol. Sci. Med. Sci.* 58, M911–M916. doi: 10.1093/gerona/58.10.M911
- Marcus, R. L., Brixner, D. I., Ghate, S., and Lastayo, P. (2012). Fat modulates the relationship between sarcopenia and physical function in nonobese older adults. *Curr. Gerontol. Geriatr. Res.* 2012:216185. doi: 10.1155/2012/216185
- Marg, A., Escobar, H., Karaiskos, N., Grunwald, S. A., Metzler, E., Kieshauer, J., et al. (2019). Human muscle-derived CLEC14A-positive cells regenerate muscle independent of PAX7. *Nat. Commun.* 10:5776. doi: 10.1038/s41467-019-13650-z
- Martel, G. F., Roth, S. M., Ivey, F. M., Lemmer, J. T., Tracy, B. L., Hurlbut, D. E., et al. (2006). Age and sex affect human muscle fibre adaptations to heavy-resistance strength training. *Exp. Physiol.* 91, 457–464. doi: 10.1113/expphysiol.2005.032771
- Matteini, P., Dei, L., Carretti, E., Volpi, N., Goti, A., and Pini, R. (2009). Structural behavior of highly concentrated hyaluronan. *Biomacromolecules* 10, 1516–1522. doi: 10.1021/bm900108z
- McCormick, R. J. (1999). Extracellular modifications to muscle collagen: implications for meat quality. *Poult. Sci.* 78, 785–791. doi: 10.1093/ps/78.5.785
- McDonagh, B. (2016). Editorial: redox regulation in skeletal muscle aging and exercise. *Front. Physiol.* 7:5. doi: 10.3389/fphys.2016.00005
- McDonagh, B., Sakellariou, G. K., Smith, N. T., Brownridge, P., and Jackson, M. J. (2014). Differential cysteine labeling and global label-free proteomics reveals an altered metabolic state in skeletal muscle aging. *J. Proteome Res.* 13, 5008–5021. doi: 10.1021/pr5006394
- Meznaric, M., Eržen, I., Karen, P., and Cvetko, E. (2018). Effect of ageing on the myosin heavy chain composition of the human sternocleidomastoid muscle. *Ann. Anat.* 216, 95–99. doi: 10.1016/j.aanat.2017.12.001
- Migliavacca, E., Tay, S. K. H., and Patel, H. P. (2019). Mitochondrial oxidative capacity and NAD⁺ biosynthesis are reduced in human sarcopenia across ethnicities. *Nat. Commun.* 10:5808. doi:10.1038/s41467-019-13694-1
- Miljkovic, N., Lim, J. Y., Miljkovic, I., and Frontera, W. R. (2015). Aging of skeletal muscle fibers. *Ann. Rehabil. Med.* 39, 155–162. doi: 10.5535/arm.2015.39.2.155
- Mitchell, W. K., Williams, J., Atherton, P., Larvin, M., Lund, J., and Narici, M. (2012). Sarcopenia, dynapenia, and the impact of advancing age on human skeletal muscle size and strength; a quantitative review. *Front. Physiol.* 3:260. doi: 10.3389/fphys.2012.00260
- Morse, C. I., Thom, J. M., Reeves, N. D., Birch, K. M., and Narici, M. V. (2005). In vivo physiological cross-sectional area and specific force are reduced in the gastrocnemius of elderly men. *J. Appl. Physiol.* 99, 1050–1055. doi: 10.1152/jappphysiol.01186.2004
- Mueller, A. L., and Bloch, R. J. (2019). Skeletal muscle cell transplantation: models and methods. *J. Muscle Res. Cell Motil.* doi: 10.1007/s10974-019-09550-w
- Muhammad, M. H., and Allam, M. M. (2018). Resveratrol and/or exercise training counteract aging-associated decline of physical endurance in aged mice; targeting mitochondrial biogenesis and function. *J. Physiol. Sci.* 68, 681–688. doi: 10.1007/s12576-017-0582-4
- Myers, T. W. (2013). *Anatomy trains*. Churchill Livingstone, Elsevier USA.
- Nakagawa, Y., Majima, T., and Nagashima, K. (1994). Effect of ageing on ultrastructure of slow and fast skeletal muscle tendon in rabbit Achilles tendons. *Acta Physiol. Scand.* 152, 307–313. doi: 10.1111/j.1748-1716.1994.tb09810.x
- Naranjo, J. D., Dziki, J. L., and Badylak, S. F. (2017). Regenerative medicine approaches for age-related muscle loss and sarcopenia: a mini-review. *Gerontology* 63, 580–589. doi: 10.1159/000479278
- Narici, M. V., and Maffulli, N. (2010). Sarcopenia: characteristics, mechanisms and functional significance. *Br. Med. Bull.* 95, 139–159. doi: 10.1093/bmb/ldq008
- Narici, M. V., Maffulli, N., and Maganaris, C. N. (2008). Ageing of human muscles and tendons. *Disabil. Rehabil.* 30, 1548–1554. doi: 10.1080/09638280701831058
- Nelson, G., Ngo, S., Pleasants, T., Raymond, F., Sayer, A. A., Ming Sim, C., et al. (2019). Mitochondrial oxidative capacity and NAD⁺ biosynthesis are reduced in human sarcopenia across ethnicities. *Nat. Commun.* 10:5808. doi: 10.1038/s41467-019-13694-1
- Netuveli, G., Wiggins, R. D., Hildon, Z., Montgomery, S. M., and Blane, D. (2006). Quality of life at older ages: evidence from the English longitudinal study of aging (wave 1). *J. Epidemiol. Community Health* 60, 357–363. doi: 10.1136/jech.2005.040071
- Nishimune, H., Numata, T., Chen, J., Aoki, Y., Wang, Y., Starr, M. P., et al. (2012). Active zone protein bassoon co-localizes with presynaptic calcium channel, modifies channel function, and recovers from aging related loss by exercise. *PLoS One* 7:e38029. doi: 10.1371/journal.pone.0038029
- Oikawa, S. Y., Holloway, T. M., and Phillips, S. M. (2019). The impact of step reduction on muscle health in aging: protein and exercise as countermeasures. *Front. Nutr.* 6:75. doi: 10.3389/fnut.2019.00075
- Özkaya, L. D., Goldsheyder, N. D., and Nordin, M. (2017). *Fundamentals of biomechanics, equilibrium, motion and deformation*. Edinburgh: Springer.
- Pamuk, U., Karakuzu, A., Ozturk, C., Acar, B., and Yucesoy, C. A. (2016). Combined magnetic resonance and diffusion tensor imaging analyses provide a powerful tool for in vivo assessment of deformation along human muscle fibers. *J. Mech. Behav. Biomed. Mater.* 63, 207–219. doi: 10.1016/j.jmbbm.2016.06.031
- Parker, M. H. (2015). The altered fate of aging satellite cells is determined by signaling and epigenetic changes. *Front. Genet.* 6:59. doi:10.3389/fgene.2015.00059
- Pavan, P. G., Stecco, A., Stern, R., and Stecco, C. (2014). Painful connections: densification versus fibrosis of fascia. *Curr. Pain Headache Rep.* 18:441. doi: 10.1007/s11916-014-0441-4
- Phillip, J. M., Aifuwa, I., Walston, J., and Wirtz, D. (2015). The Mechanobiology of aging. *Annu. Rev. Biomed. Eng.* 17, 113–141. doi: 10.1146/annurev-bioeng-071114-040829
- Piasecki, M., Ireland, A., Jones, D. A., and McPhee, J. S. (2016). Age-dependent motor unit remodelling in human limb muscles. *Biogerontology* 17, 485–496. doi: 10.1007/s10522-015-9627-3
- Pierno, S., Tricarico, D., Liantonio, A., Mele, A., Digennaro, C., Rolland, J. F., et al. (2014). An olive oil-derived antioxidant mixture ameliorates the age-related decline of skeletal muscle function. *Age* 36, 73–88. doi: 10.1007/s11357-013-9544-9

- Pillen, S., Tak, R. O., Zwartz, M. J., Lammens, M. M., Verrijp, K. N., Arts, I. M., et al. (2009). Skeletal muscle ultrasound: correlation between fibrous tissue and echo intensity. *Ultrasound Med. Biol.* 35, 443–446. doi: 10.1016/j.ultrasmedbio.2008.09.016
- Pisot, R., Marusic, U., Biolo, G., Mazzucco, S., Lazzar, S., Grassi, B., et al. (2016). Greater loss in muscle mass and function but smaller metabolic alterations in older compared with younger men following 2 wk of bed rest and recovery. *J. Appl. Physiol.* 120, 922–929. doi: 10.1152/jappphysiol.00858.2015
- Pollo, B. E., Rathbone, C. R., Wenke, J. C., and Guda, T. (2018). Natural polymeric hydrogel evaluation for skeletal muscle tissue engineering. *J. Biomed. Mater. Res. B. Appl. Biomater.* 106, 672–679. doi: 10.1002/jbmb.33859
- Potthoff, M. J., Olson, E. N., and Bassel-Duby, R. (2007). Skeletal muscle remodeling. *Curr. Opin. Rheumatol.* 19, 542–549. doi: 10.1097/BOR.0b013e3282efb761
- Rao, L., Qian, Y., Khodabukus, A., Ribar, T., and Bursac, N. (2018). Engineering human pluripotent stem cells into a functional skeletal muscle tissue. *Nat. Commun.* 9:126. doi: 10.1038/s41467-017-02636-4
- Raue, U., Slivka, D., Minchev, K., and Trappe, S. (2009). Improvements in whole muscle and myocellular function are limited with high-intensity resistance training in octogenarian women. *J. Appl. Physiol.* 106, 1611–1617. doi: 10.1152/jappphysiol.91587.2008
- Reid, K. F., and Fielding, R. A. (2012). Skeletal muscle power: a critical determinant of physical functioning in older adults. *Exerc. Sport Sci. Rev.* 40, 4–12. doi: 10.1097/JES.0b013e31823b5f13
- Reimers, K., Reimers, C. D., Wagner, S., Paetzke, I., and Pongratz, D. E. (1993). Skeletal muscle sonography: a correlative study of echogenicity and morphology. *J. Ultrasound Med.* 12, 73–77. doi: 10.7863/jum.1993.12.2.73
- Reiser, K., McCormick, R. J., and Rucker, R. B. (1992). Enzymatic and nonenzymatic cross-linking of collagen and elastin. *FASEB J.* 6, 2439–2449. doi: 10.1096/fasebj.6.7.1348714
- Rice, C. L., Cunningham, D. A., Paterson, D. H., and Lefcoe, M. S. (1989). Arm and leg composition determined by computed tomography in young and elderly men. *Clin. Physiol.* 9, 207–220. doi: 10.1111/j.1475-097X.1989.tb00973.x
- Rodriguez, J., Vernus, B., Chel, I., Cassar-Malek, I., Gabillard, J. C., Hadj Sassi, A., et al. (2014). Myostatin and the skeletal muscle atrophy and hypertrophy signaling pathways. *Cell. Mol. Life Sci.* 71, 4361–4371. doi: 10.1007/s00018-014-1689-x
- Rogers, M. A., and Evans, W. J. (1993). Changes in skeletal muscle with aging: effects of exercise training. *Exerc. Sport Sci. Rev.* 21, 65–102.
- Rossini, P. M., Desiato, M. T., and Caramia, M. D. (1992). Age-related changes of motor evoked potentials in healthy humans: non-invasive evaluation of central and peripheral motor tracts excitability and conductivity. *Brain Res.* 593, 14–19. doi: 10.1016/0006-8993(92)91256-E
- Ruzzini, L., Abbruzzese, F., Rainer, A., Longo, U. G., Trombetta, M., Maffulli, N., et al. (2013). Characterization of age-related changes of tendon stem cells from adult human tendons. *Knee Surg. Sports Traumatol. Arthrosc.* 22, 2856–2866. doi: 10.1007/s00167-013-2457-4
- Saul, D., Geisberg, L. K., Gehle, T., Hoffmann, D. B., Tezval, M., Sehmsch, S., et al. (2019). Changes in musculoskeletal system and metabolism in osteoporotic rats treated with urocortin. *Front. Endocrinol.* 10:400. doi: 10.3389/fendo.2019.00400
- Schiaffino, S., and Reggiani, C. (1994). Myosin isoforms in mammalian muscle. *J. Appl. Physiol.* 77, 493–501. doi: 10.1152/jappphysiol.1994.77.2.493
- Schiaffino, S., and Reggiani, C. (1996). Molecular diversity of myofibrillar proteins: gene regulation and functional significance. *Physiol. Rev.* 76, 371–423. doi: 10.1152/physrev.1996.76.2.371
- Schleip, R., Gabbiani, G., Wilke, J., Naylor, I., Hinz, B., Zorn, A., et al. (2019). Fascia is able to actively contract and may thereby influence musculoskeletal dynamics: a histochemical and mechanographic investigation. *Front. Physiol.* 10:336. doi: 10.3389/fphys.2019.00336
- Sheth, K. A., Iyer, C. C., Wier, C. G., Crum, A. E., Bratasz, A., Kolb, S. J., et al. (2018). Muscle strength and size are associated with motor unit connectivity in aged mice. *Neurobiol. Aging* 67, 128–136. doi: 10.1016/j.neurobiolaging.2018.03.016
- Short, K. R., Bigelow, M. L., Kahl, J., Singh, R., Coenen-Schimke, J., Raghavakaimal, S., et al. (2005b). Decline in skeletal muscle mitochondrial function with aging in humans. *Proc. Natl. Acad. Sci. U. S. A.* 102, 5618–5623. doi: 10.1073/pnas.0501559102
- Short, K. R., Vittone, J. L., Bigelow, M. L., Proctor, D. N., Coenen-Schimke, J. M., Rys, P., et al. (2005a). Changes in myosin heavy chain mRNA and protein expression in human skeletal muscle with age and endurance exercise training. *J. Appl. Physiol.* 99, 95–102. doi: 10.1152/jappphysiol.00129.2005
- Sikdar, S., Wei, Q., and Cortes, N. (2014). Dynamic ultrasound imaging applications to quantify musculoskeletal function. *Exerc. Sport Sci. Rev.* 42, 126–135. doi: 10.1249/JES.0000000000000015
- Snijders, T., Nederveen, J. P., McKay, B. R., Joannisse, S., Verdijk, L. B., van Loon, L. J., et al. (2015). Satellite cells in human skeletal muscle plasticity. *Front. Physiol.* 6:283. doi: 10.3389/fphys.2015.00283
- Snijders, T., Verdijk, L. B., and van Loon, L. J. (2009). The impact of sarcopenia and exercise training on skeletal muscle satellite cells. *Ageing Res. Rev.* 8, 328–338. doi: 10.1016/j.arr.2009.05.003
- Snow, S. W., Bohne, W. H., DiCarlo, E., and Chang, V. K. (1995). Anatomy of the Achilles tendon and plantar fascia in relation to the calcaneus in various age groups. *Foot Ankle Int.* 16, 418–421. doi: 10.1177/107110079501600707
- Sölch, D. (2015). Ageing and restricted mobility. Frailty from the perspective of myofascial structural models. *Z. Gerontol. Geriatr.* 48, 35–40. doi: 10.1007/s00391-013-0555-z
- Sousa-Victor, P., Gutarra, S., Garcia-Prat, L., Rodriguez-Ubreva, J., Ortet, L., Ruiz-Bonilla, V., et al. (2014). Geriatric muscle stem cells switch reversible quiescence into senescence. *Nature* 506, 316–321. doi: 10.1038/nature13013
- Squier, C. A., and Magnes, C. (1983). Spatial relationships between fibroblasts during the growth of rat-tail tendon. *Cell Tissue Res.* 234, 17–29. doi: 10.1007/BF00217399
- Stearns-Reider, K. M., D'Amore, A., Beezhold, K., Rothrauff, B., Cavalli, L., Wagner, W. R., et al. (2017). Aging of the skeletal muscle extracellular matrix drives a stem cell fibrogenic conversion. *Ageing Cell* 16, 518–528. doi: 10.1111/acel.12578
- Stecco, C., Corradin, M., Macchi, V., Morra, A., Porzionato, A., Biz, C., et al. (2013). Plantar fascia anatomy and its relationship with Achilles tendon and paratenon. *J. Anat.* 223, 665–676. doi: 10.1111/joa.12111
- Suetta, C. (2017). Plasticity and function of human skeletal muscle in relation to disuse and rehabilitation: influence of ageing and surgery. *Dan. Med. J.* 64:B5377.
- Suzuki, D., Takahashi, M., Abe, M., and Nagano, A. (2008). Biochemical study of collagen and its cross-links in the anterior cruciate ligament and the tissues used as a graft for reconstruction of the anterior cruciate ligament. *Connect. Tissue Res.* 49, 42–47. doi: 10.1080/03008200701820799
- Svensson, R. B., Couppé, C., and Magnusson, S. P. (2015). “Mechanical properties of the aging tendon” in *Mechanical properties of aging soft tissues, engineering materials and processes*. eds. B. Derby and R. Akhtar (Switzerland: Springer International Publishing). doi: 10.1007/978-3-319-03970-1_6
- Szentesi, P., Csernoch, L., Dux, L., and Keller-Pintér, A. (2019). Changes in redox signaling in the skeletal muscle with aging. *Oxidative Med. Cell. Longev.* 2019:4617801. doi: 10.1155/2019/4617801
- Taetzsch, T., and Valdez, G. (2018). NMJ maintenance and repair in aging. *Curr. Opin. Physiol.* 4, 57–64. doi: 10.1016/j.cophys.2018.05.007
- Thompson, L. V. (2002). Skeletal muscle adaptations with age, inactivity, and therapeutic exercise. *J. Orthop. Sports Phys. Ther.* 32, 44–57. doi: 10.2519/jospt.2002.32.2.44
- Thom, J. M., Morse, C. I., and Birch, K. M. (2007). Influence of muscle architecture on the torque and power-velocity characteristics of young and elderly men. *Eur. J. Appl. Physiol.* 100, 613–619. doi: 10.1007/s00421-007-0481-0
- Tian, M., Herbert, R. D., Hoang, P., Gandevia, S. C., and Bilston, L. E. (2012). Myofascial force transmission between the human soleus and gastrocnemius muscles during passive knee motion. *J. Appl. Physiol.* 113, 517–523. doi: 10.1152/jappphysiol.00111.2012
- Tieland, M., Trouwborst, I., and Clark, B. C. (2018). Skeletal muscle performance and ageing. *J. Cachexia Sarcopenia Muscle* 9, 3–19. doi:10.1002/jcsm.12238
- Trappe, S., Gallagher, P., Harber, M., Carrithers, J., Fluckey, J., and Trappe, T. (2003). Single muscle fibre contractile properties in young and old men and women. *J. Physiol.* 552, 47–58. doi: 10.1113/jphysiol.2003.044966
- Trappe, T. (2009). Influence of aging and long-term unloading on the structure and function of human skeletal muscle. *Appl. Physiol. Nutr. Metab.* 34, 459–464. doi: 10.1139/H09-041
- Turrina, A., Martinez-Gonzalez, M. A., and Stecco, C. (2013). The muscular force transmission system: role of the intramuscular connective tissue. *J. Bodyw. Mov. Ther.* 17, 95–102. doi: 10.1016/j.jbmt.2012.06.001
- Unhjem, R., Lundestad, R., Fimland, M. S., Mosti, M. P., and Wang, E. (2015). Strength training-induced responses in older adults: attenuation of descending neural drive with age. *Age* 37:9784. doi: 10.1007/s11357-015-9784-y

- Valdez, G., Tapia, J. C., Kang, H., Clemenson, G. D. Jr., Gage, F. H., Lichtman, J. W., et al. (2010). Attenuation of age-related changes in mouse neuromuscular synapses by caloric restriction and exercise. *Proc. Natl. Acad. Sci. U. S. A.* 107, 14863–14868. doi: 10.1073/pnas.1002220107
- Verdijk, L. B., Koopman, R., Schaart, G., Meijer, K., Savelberg, H. H., and van Loon, L. J. (2007). Satellite cell content is specifically reduced in type II skeletal muscle fibers in the elderly. *Am. J. Physiol. Endocrinol. Metab.* 292, E151–E157. doi: 10.1152/ajpendo.00278.2006
- Vernooij, C. A., Rao, G., Berton, E., Retornaz, F., and Temprado, J.-J. (2016). The effect of aging on muscular dynamics underlying movement patterns changes. *Front. Aging Neurosci.* 8:309. doi: 10.3389/fnagi.2016.00309
- Verzijl, N., DeGroot, J., Thorpe, S. R., Bank, R. A., Shaw, J. N., Lyons, T. J., et al. (2000). Effect of collagen turnover on the accumulation of advanced glycation end products. *J. Biol. Chem.* 275, 39027–39031. doi: 10.1074/jbc.M006700200
- Vleeming, A., Schuenke, M. D., Danneels, L., and Willard, F. H. (2014). The functional coupling of the deep abdominal and paraspinal muscles: the effects of simulated paraspinal muscle contraction on force transfer to the middle and posterior layer of the thoracolumbar fascia. *J. Anat.* 225, 447–462. doi:10.1111/joa.12227
- Vogel, H. G. (1978). Influence of maturation and age on mechanical and biochemical parameters of connective tissue of various organs in the rat. *Connect. Tissue Res.* 6, 161–166. doi: 10.3109/03008207809152626
- Wang, H. Q., Wei, Y. Y., Wu, Z. X., and Luo, Z. J. (2009). Impact of leg lengthening on viscoelastic properties of the deep fascia. *BMC Musculoskelet. Disord.* 10:105. doi: 10.1186/1471-2474-10-105
- Watanabe, Y., Ikenaga, M., Yoshimura, E., Yosuke Yamada, Y., and Kimura, M. (2018). Association between echo intensity and attenuation of skeletal muscle in young and older adults: a comparison between ultrasonography and computed tomography. *Clin. Interv. Aging* 13, 1871–1878. doi: 10.2147/CIA.S173372
- Watanabe, Y., Yamada, Y., Fukumoto, Y., Ishihara, T., Yokoyama, K., Yoshida, T., et al. (2013). Echo intensity obtained from ultrasonography images reflecting muscle strength in elderly men. *Clin. Interv. Aging* 8, 993–998. doi: 10.2147/CIA.S47263
- Wearing, S. C., Hooper, S. L., Smeathers, J. E., Pourcelot, P., Crevier-Denoix, N., and Brauner, T. (2016). Tendinopathy alters ultrasound transmission in the patellar tendon during squatting. *Scand. J. Med. Sci. Sports* 26, 1415–1422. doi: 10.1111/sms.12602
- Wessner, B., Liebensteiner, M., Nachbauer, M., and Csapo, R. (2019). Age-specific response of skeletal muscle extracellular matrix to acute resistance exercise: a pilot study. *Eur. J. Sport Sci.* 19, 354–364. doi: 10.1080/17461391.2018.1526974
- White, Z., Terrill, J., White, R. B., McMahon, C., Sheard, P., Grounds, M. D., et al. (2016). Voluntary resistance wheel exercise from mid-life prevents sarcopenia and increases markers of mitochondrial function and autophagy in muscles of old male and female C57BL/6J mice. *Skelet. Muscle* 6:45. doi: 10.1186/s13395-016-0117-3
- Wilke, J., Kalo, K., Niederer, D., Vogt, L., and Banzer, W. (2019). Gathering hints for myofascial force transmission under in vivo conditions: are remote exercise effects age dependent? *J. Sport Rehabil.* 28, 758–763. doi: 10.1123/jsr.2018-0184
- Wilke, J., Krause, F., Vogt, L., and Banzer, W. (2016). What is evidence-based about Myofascial chains: a systematic review. *Arch. Phys. Med. Rehabil.* 97, 454–461. doi: 10.1016/j.apmr.2015.07.023
- Wilke, J., Schleip, R., Yucesoy, C. A., and Banzer, W. (2018). Not merely a protective packing organ? A review of fascia and its force transmission capacity. *J. Appl. Physiol.* 124, 234–244. doi: 10.1152/jappphysiol.00565.2017
- Willadt, S., Nash, M., and Slater, C. R. (2016). Age-related fragmentation of the motor endplate is not associated with impaired neuromuscular transmission in the mouse diaphragm. *Sci. Rep.* 6:24849. doi: 10.1038/srep24849
- Willems, M. E., Miller, G. R., and Stauber, W. T. (2001). Force deficits after stretches of activated rat muscle-tendon complex with reduced collagen cross-linking. *Eur. J. Appl. Physiol.* 85, 405–411. doi: 10.1007/s004210100480
- Yahia, L. H., Pigeon, P., and DesRosiers, E. A. (1993). Viscoelastic properties of the human lumbodorsal fascia. *J. Biomed. Eng.* 15, 425–429. doi: 10.1016/0141-5425(93)90081-9
- Yamada, Y., Schoeller, D. A., Nakamura, E., Morimoto, T., Kimura, M., and Oda, S. (2010). Extracellular water may mask actual muscle atrophy during aging. *Gerontol. A Biol. Sci. Med. Sci.* 65, 510–516. doi: 10.1093/gerona/glq001
- Yin, H., Price, F., and Rudnicki, M. A. (2013). Satellite cells and the muscle stem cell niche. *Physiol. Rev.* 93, 23–67. doi: 10.1152/physrev.00043.2011
- Yucesoy, C. A. (2010). Epimuscular myofascial force transmission implies novel principles for muscular mechanics. *Exerc. Sport Sci. Rev.* 38, 128–134. doi: 10.1097/JES.0b013e3181e372ef
- Yucesoy, C. A., Baan, G. C., Koopman, B. H., Grootenboer, H. J., and Huijting, P. A. (2005). Pre-strained epimuscular connections cause muscular myofascial force transmission to affect properties of synergistic EHL and EDL muscles of the rat. *J. Biomech. Eng.* 127, 819–828. doi: 10.1115/1.1992523
- Zaidman, C. M., Holland, M. R., and Hughes, M. S. (2012). Quantitative ultrasound of skeletal muscle: reliable measurements of calibrated muscle backscatter from different ultrasound systems. *Ultrasound Med. Biol.* 38, 1618–1625. doi: 10.1016/j.ultrasmedbio.2012.04.020
- Zampieri, S., Pietrangelo, L., Loeffler, S., Fruhmann, H., Vogelauer, M., Burggraf, S., et al. (2015). Lifelong physical exercise delays age-associated skeletal muscle decline. *J. Gerontol. A Biol. Sci. Med. Sci.* 70, 163–173. doi: 10.1093/gerona/glu006
- Zhang, C., and Gao, Y. (2014). Effects of aging on the lateral transmission of force in rat skeletal muscle. *J. Biomech.* 47, 944–948. doi: 10.1016/j.jbiomech.2014.01.026
- Zhang, X., Trevino, M. B., Wang, M., Gardell, S. J., Ayala, J. E., Han, X., et al. (2018a). Impaired mitochondrial energetics characterize poor early recovery of muscle mass following hind limb unloading in old mice. *J. Gerontol. A Biol. Sci. Med. Sci.* 73, 1313–1322. doi: 10.1093/gerona/gly051
- Zhang, S. F., Zhang, Y., Li, B., and Chen, N. (2018b). Physical inactivity induces the atrophy of skeletal muscle of rats through activating AMPK/FoxO3 signal pathway. *Eur. Rev. Med. Pharmacol. Sci.* 22, 199–209. doi: 10.26355/eurrev_201801_14118
- Zhou, Z., Akinbiyi, T., Xu, L., Ramcharan, M., Leong, D. J., Ros, S. J., et al. (2010). Tendon-derived stem/progenitor cell aging: defective self-renewal and altered fate. *Aging Cell* 9, 911–915. doi: 10.1111/j.1474-9726.2010.00598.x
- Zügel, M., Maganaris, C. N., Wilke, J., Jurkat-Rott, K., Klingler, W., Wearing, S. C., et al. (2018). Fascial tissue research in sports medicine: from molecules to tissue adaptation, injury and diagnostics: consensus statement. *Br. J. Sports Med.* 52:1497. doi: 10.1136/bjsports-2018-099308
- Zullo, A., Mancini, F. P., Schleip, R., Wearing, S., Yahia, L., and Klingler, W. (2017). The interplay between fascia, skeletal muscle, nerves, adipose tissue, inflammation and mechanical stress in musculo-fascial regeneration. *J. Gerontol. Geriatr.* 65, 271–283.

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Zullo, Fleckenstein, Schleip, Hoppe, Wearing and Klingler. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.