



Queensland University of Technology
Brisbane Australia

This is the author's version of a work that was submitted/accepted for publication in the following source:

[Harkin, Damien G.](#), [Foy, Leanne](#), [Bray, Laura J.](#), Sutherland, Allison J., Li, Fiona J., & Cronin, Brendan G.

(2015)

Can mesenchymal stromal cells differentiate into corneal cells? A systematic review of published data.

Stem Cells, 33(3), pp. 785-791.

This file was downloaded from: <http://eprints.qut.edu.au/78134/>

© Copyright 2014 AlphaMed Press

Notice: *Changes introduced as a result of publishing processes such as copy-editing and formatting may not be reflected in this document. For a definitive version of this work, please refer to the published source:*

<http://doi.org/10.1002/stem.1895>

Concise Review:

Can Mesenchymal Stromal Cells Differentiate into Corneal Cells?

A systematic review of published data.

Damien G Harkin,^{a,b,c} Leanne Foyn,^{a,c} Laura J Bray,^{a,b,c,d} Allison J Sutherland,^c Fiona J Li,^c
and Brendan G Cronin^c

^aSchool of Biomedical Sciences, Queensland University of Technology, Brisbane, Australia.

^bInstitute of Health & Biomedical Innovation, Queensland University of Technology.

^cQueensland Eye Institute, South Brisbane, Australia.

^dLeibniz Institute for Polymer Research Dresden, Max Bergmann Center of Biomaterials, Dresden, Germany.

Author contributions:

Damien Harkin: Conception and design, data analysis and interpretation, manuscript writing, final approval of manuscript.

Leanne Foyn: Administrative support, collection and assembly of data, data analysis and interpretation.

Laura Bray: Data analysis and interpretation, manuscript writing, final approval of manuscript.

Allison Sutherland: Data analysis and interpretation, manuscript writing, final approval of manuscript.

Fiona Li: Data analysis and interpretation.

Brendan Cronin: Data analysis and interpretation, manuscript writing, final approval of manuscript.

Corneal differentiation of MSC

Corresponding author:

Damien G Harkin, BSc PhD

School of Biomedical Sciences, Queensland University of Technology, 2 George Street,

Brisbane, Queensland, 4000, Australia.

Telephone: +61 7 3239 5023

Fax: +61 7 3138 6030

E-mail: d.harkin@qut.edu.au

Grant support:

National Health and Medical Research Council of Australia Project Grant 1049050.

Key words:

Mesenchymal stromal cells, cornea, transdifferentiation, systematic review, stem cell therapy.

Abstract

The majority of stem cell therapies for corneal repair are based upon the use of progenitor cells isolated from corneal tissue, but a growing body of literature suggests a role for mesenchymal stromal cells (MSC) isolated from non-corneal tissues. While the mechanism of MSC action seems likely to involve their immuno-modulatory properties, claims have emerged of MSC transdifferentiation into corneal cells. Substantial differences in methodology and experimental outcomes, however, have prompted us to perform a systematic review of the published data. Key questions used in our analysis included; the choice of markers used to assess corneal cell phenotype, the techniques employed to detect these markers, adequate reporting of controls, and tracking of MSC when studied *in vivo*. Our search of the literature revealed 28 papers published since 2006, with half appearing since 2012. MSC cultures established from bone marrow and adipose tissue have been best studied (22 papers). Critically, only 11 studies employed appropriate markers of corneal cell phenotype, along with necessary controls. Ten out of these 11 papers, however, contained positive evidence of corneal cell marker expression by MSC. The clearest evidence is observed with respect to expression of markers for corneal stromal cells by MSC. In comparison, the evidence for MSC conversion into either corneal epithelial cells or corneal endothelial cells is often inconsistent or inconclusive. Our analysis clarifies this emerging body of literature and provides guidance for future studies of MSC differentiation within the cornea as well as other tissues.

Introduction

The cornea has been extensively studied as a tissue for stem cell therapies. To date, the majority of this research has focused on corneal epithelial progenitor cells located at the peripheral edge, or so-called limbus, where the cornea adjoins the sclera [1, 2]. As such, cultivated epithelial autografts have become widely used as a standard treatment for repairing the ocular surface [3]. Exploring deeper, a number of groups have more recently identified corneal/limbal stromal cells with stem cell properties [4-10] and similar studies are also being pursued for the innermost cellular layer, the corneal endothelium [11]. Nevertheless, the limited availability and sensitive location of corneal tissue present significant challenges for autologous corneal stem cell therapies, particularly in cases of bilateral disease.

Given the limited availability of a patient's own corneal stem cells, a number of non-corneal tissues have been investigated as potential sources of epithelial progenitor cells for repairing the ocular surface including the oral mucosa [12]. More recently, however, several groups have evaluated the potential of mesenchymal stromal cell (MSC) cultures derived from tissues of non-corneal origin [13-16]. While much of this research has centered on exploiting the immuno-regulatory properties of MSC to encourage corneal healing, claims have emerged that mesenchymal cells of non-corneal origin have the capacity to transdifferentiate into corneal cells [17-24]. Such a conclusion, if confirmed, would not only have important implications for the treatment of corneal diseases, but would have a significant impact on our understanding of general MSC biology. Upon initial engagement with this literature, however, we have noted substantial differences in experimental design and reported outcomes that hamper a clear interpretation of the data. The goal of this concise review, therefore, is to systematically evaluate this recent body of literature for evidence of non-corneal MSC differentiation into corneal cells.

Literature search, terms of reference and method of analysis

Our study design is based upon published guidelines for the conduct of systematic reviews [25]. The initial ‘identification stage’ consisted of searching the PubMed database for studies where the terms ‘mesenchymal’ and ‘cornea’ had been used. This search retrieved 296 studies published since 1950 (as of July 25th 2014). We subsequently screened these publications for studies where cultures of non-transformed mesenchymal stromal cells, established from tissues of non-corneal origin, had been examined either *in vitro* or *in vivo* for their ability to transdifferentiate into corneal cells. Reports of efficacy alone, while interesting, were nonetheless excluded from our subsequent analysis. All literature pertaining to these terms of reference were included in this systematic review, irrespective of the primary language in which the article was published. The evidence presented in each study was evaluated using four standard questions.

- (1) Have appropriate markers been used to determine transformation to a corneal phenotype?
- (2) By what methods has the expression of these corneal markers been evaluated?
- (3) Have appropriate controls been reported to validate these results? For example, positive results obtained by immunostaining should be validated through demonstration of a negative control. Likewise, a negative result should be validated through demonstration of a positive control.
- (4) In the case of *in vivo* studies, has the provenance of observed “corneal cells” been traced back to the MSC of non-corneal origin by using some form of marker?

Definition of corneal cell phenotype

Our definition of corneal cell phenotype is based upon the following considerations. To begin, the transcription factor paired box 6 (Pax-6) is widely regarded as the canonical marker of eye

Corneal differentiation of MSC

tissue development and is retained to varying degrees by mature corneal cells. A variety of molecules including the transcription factor p63 [26] and cytokeratin 19 [27] have been used as markers for corneal epithelial progenitor cells, but neither protein is specific to the cornea. Corneal epithelium is therefore defined by expression of the tissue-specific differentiation markers cytokeratin 3 (K3) and cytokeratin 12 (K12) [28]. On a technical note, while K3 expression in corneal epithelial cells can be reliably studied using the monoclonal antibody AE5, care must be taken when applying this antibody to detect K3 in other cell types, since it is known to cross-react with cytokeratin 2p/76 (K2p/76). Corneal stromal cells (keratocytes) are generally defined by expression of CD34, keratocan, lumican, and aldehyde dehydrogenase (ALDH), but during wound healing are known to differentiate into wound repair fibroblasts (CD34⁻/CD90⁺) and myofibroblasts (expressing alpha-smooth muscle actin or α -sma) [29]. Nevertheless, since both CD90 and α -sma are expressed by MSC cultures [30], they have been excluded as valid markers of corneal differentiation. While specific markers for the corneal endothelium are emerging [31], the expression of N-cadherin, zonula occludens-1 (ZO-1) and sodium/potassium ATPase are more generally employed to identify these cells [32]. Of the three markers used, the presence of sodium/potassium ATPase is perhaps most important given the role of this protein in maintaining the pump function of corneal endothelial cells required for corneal transparency.

Overview of published literature

Our literature search identified 28 papers published between January 2006 and June 2014, with half of these having been reported since January 2012 [13, 14, 17-20, 22-24, 33-51]. The essential details for each study are summarized in the first 5 columns of Table 1. The majority of studies have been performed using MSC derived from either bone marrow (13 studies) [13, 14, 17, 22-24, 38-40, 43, 47, 48, 50] or adipose tissue (9 studies) [19, 33, 35, 42, 44-46, 49,

Corneal differentiation of MSC

51], with the remainder using MSC extracted from either umbilical cord tissues (4 studies) [20, 34, 36, 41] or dental pulp (2 studies) [18, 37]. Most studies have utilized cultures of MSC established from human tissues (19 studies) [13, 18-20, 23, 24, 33-39, 41, 42, 45, 46, 49, 51], with the balance having been established from rabbits (5 studies) [14, 17, 43, 44, 48], rats (3 studies) [40, 47, 50] and mice (1 study) [22].

A large proportion of studies (19 studies) [13, 14, 17, 18, 20, 22, 24, 33-35, 37, 38, 40, 42-44, 47-49] have involved *in vivo* experiments, with 11 studies involving administration of human MSC into rabbits (6 studies) [18, 33, 35, 37, 38, 42], rats (3 studies) [13, 24, 49] or mice (2 studies) [20, 34]. Routes of administration include topical application (10 studies) [13, 17, 18, 24, 37, 38, 40, 47-49], typically following alkali burn (9 studies) [13, 14, 17, 18, 24, 38, 40, 42, 49], with or without carrier/adjunct materials including amniotic membrane (7 studies) [13, 18, 24, 37, 38, 47, 48] and fibrin (1 study) [17]. Other methods employed include direct injection into the cornea (4 studies) [20, 22, 33, 34] or adjacent conjunctiva (1 study) [42], stromal implants consisting of MSC cultured within synthetic scaffolds (2 studies) [35, 44], and intravenous injection (1 study) [14]. Significantly, the fate of administered MSC was traced using some form of temporary (DiI or BrdU) or permanent marker (green fluorescent protein, human nuclear antigen or sex-linked DNA marker) in only 15 out of 19 studies conducted *in vivo* [13, 14, 17, 18, 20, 22, 33-35, 37, 38, 43, 44, 47, 49].

Reports of MSC displaying evidence of corneal phenotype *in vitro* (15 studies) [17-19, 23, 24, 36, 39-41, 45-48, 50, 51] have utilized a variety of induction methods including co-culture in the presence of corneal cells (2 studies) [40, 51], treatment with ocular cell conditioned media (3 studies) [17, 41, 46], cultivation in either specialized epithelial cell growth media or keratocyte growth media (4 studies) [19, 23, 24, 42], and corneal organ culture (2 studies) [39, 41]. Only two studies have utilized specialized enrichment techniques such as magnetic assisted cell sorting (MACS for stage-specific embryonic antigen-4 [23]) or

Corneal differentiation of MSC

flow cytometry (side population cells [19]) prior to cultivation (under epithelial or keratocyte growth conditions respectively). Surprisingly, 3 studies [18, 45, 48] have claimed evidence of corneal phenotype markers being expressed when MSC were apparently maintained in their standard growth medium.

In terms of target tissue, 17 studies [13, 17, 18, 23, 24, 36-40, 42, 45-50] have presented data relevant to epithelial differentiation, 9 studies [14, 19, 20, 22, 33-35, 44, 51] have examined differentiation into corneal stromal cells (keratocytes), and 2 studies [41, 43] have examined the potential of MSC to produce corneal endothelium.

The last five columns of Table 1 present the results from our analysis of published data using the standard set of four questions. For convenience, the highlights from this analysis are discussed below according to MSC tissue of origin and are summarized in Table 2.

Corneal differentiation of bone marrow-derived MSC (BM-MS)

Of the 13 studies involving BM-MS, only 7 papers contain data that according to our analysis has been validated through use of appropriate markers and controls [13, 17, 22-24, 38, 50]. Six of these papers have addressed conversion of MSC into corneal epithelium with the remaining paper exploring keratocyte differentiation *in vivo*.

While one study found negative evidence of human BM-MS differentiation into corneal epithelium when applied to the ocular surface of rats (validated by positive control) [13], the combined data from an additional 5 studies involving cells from rabbits [17], rats [50] or human subjects [23, 24, 38] provides partial evidence of K3 and/or K12 expression under either *in vitro* or *in vivo* conditions. Nevertheless, the level of K3 or K12 expression observed in these papers is consistently less than that for corneal epithelium and often limited to a subset of cells. At the very least, therefore, it appears that BM-MS have some ability to

Corneal differentiation of MSC

produce low levels of cornea-specific keratins, but it remains unclear as to whether this level of expression represents true conversion to a functional corneal epithelial cell phenotype.

The single paper containing evidence of BM-MSC differentiation into keratocytes *in vivo* is convincing given that the cells were implanted into *keratocan* $-/-$ null mutant mice [22]. Thus, the subsequent observation of keratocan expression in conjunction with DiI labeled cells is justifiably explained by conversion of BM-MSC into keratocytes. Nevertheless, it is unclear as to what percentage of administered MSC adopted a keratocyte phenotype.

Only one study has examined the potential of BM-MSC to transdifferentiate into corneal endothelial cells [43]. In this study (published in Chinese language), the authors report improvements in corneal clarity and thickness when autologous BM-MSC cultured on gelatin membranes are adhered to the posterior surface of corneal buttons implanted into rabbits. While these results are encouraging in terms of efficacy, the phenotype of implanted cells was only examined using a combination of morphological techniques (live confocal imaging and scanning electron microscopy).

Corneal differentiation of MSC-derived from adipose tissue (A-MSC)

Reports of A-MSC conversion into corneal cells are, according to our criteria, also often lacking appropriate markers of cell phenotype and necessary controls (5 out of 9 studies). Moreover, the results for a further 3 studies are clouded by either inconsistent or unclear data.

The case for A-MSC differentiation into corneal epithelial cells is particularly weak. Only one *in vitro* study contains validated data in support of this hypothesis [45] and even in this paper there are inconsistencies with respect to the measurement of corneal phenotype markers by immunocytochemistry, Western blotting and RT-PCR. In short, while “moderate” levels of K3 were reported using immunocytochemistry and Western blotting, the mRNA transcripts required for producing this protein were not detected by RT-PCR. Nevertheless,

Corneal differentiation of MSC

weak expression was reported by RT-PCR for K12 and this was confirmed by sequencing. These results were apparently obtained for A-MSC grown in their standard expansion medium. Further studies are therefore required to clarify whether A-MSC have potential to produce corneal epithelium and ideally this research should be extended to *in vivo* models as well.

One study has reported elevated levels of both ALDH and keratocan by flow cytometry when A-MSC are co-cultured in the presence of corneal stromal cells, but no change in transcript numbers for either protein was detected by RT-qPCR [51]. Likewise, while a faint band for human keratocan was detected by Western blot after injecting human A-MSC into rabbit corneas, background fluorescence hampers a clear interpretation of the images reported to show DiI-labeled cells [33]. Nevertheless, a separate study utilizing the side population fraction of human A-MSC isolated by flow cytometry, provides good evidence of keratocan and ALDH expression using a combination of techniques [19]. It therefore seems quite plausible that A-MSC could be used as a source of keratocytes, but this case could also be strengthened by more data including studies *in vivo*.

Corneal differentiation of MSC-derived from umbilical cord (UC-MSC)

Research into stem cells derived from umbilical cord is a complex topic as mesenchymal cells with progenitor cell properties have been isolated from the cord blood as well as the surrounding primitive connective tissue. In addition, epithelial progenitor cells have been isolated from umbilical cords. All three sources of progenitor cells have been examined as tools for corneal reconstruction. For the purpose of this systematic review, however, we have focused on the evidence arising from studies using the mesenchymal cell populations obtained from umbilical cord tissues.

Corneal differentiation of MSC

Out of the 4 studies identified by our literature search, only one report contains evidence that has been validated through use of appropriate markers and controls [20]. In this study, human UC-MSC loaded with DiI were injected into dysfunctional corneas of *lum -/-* and *keratocan -/-* null mutant mice. The labeled cells subsequently became integrated within the host tissue and adopted a morphology similar to that expected for keratocytes. Evidence of transdifferentiation was provided by co-localization of the DiI tracker dye with observed patterns of immunohistochemistry for keratocan, lumican and CD34, with lumican and keratocan expression being confirmed by Western blotting. A subsequent study by this research group using a mouse model of lysosomal enzyme dysfunction has produced similar results, but specific markers of corneal phenotype were not used on this occasion [34].

One additional paper requires mention since it is one of the few studies to have evaluated the potential of MSC derived from any non-corneal tissue to adopt a corneal endothelial cell phenotype [41]. In this study, a sub-culture of UC-MSC was examined for evidence of ZO-1 and N-cadherin expression prior to and following treatment with lens epithelium conditioned medium. GFP-labeled cells were also examined for these same markers following two weeks cultivation upon the wounded posterior surface of donor human corneas. While both ZO-1 and N-cadherin were detected in un-treated cultures (validated by controls) these proteins reportedly became more distributed to cell boundaries (both in standard and organ cultures) in the presence of conditioned medium. Moreover, the results from a microarray analysis of 250 genes indicated a shift towards a phenotype closer to that of corneal endothelial cells when treated with conditioned medium. Interestingly, the integration of labeled UC-MSC with the organ cultured endothelial cells was best encouraged by attachment to the surface of damaged cells rather than by contact with exposed areas of Descemet's membrane. These findings support the theory of 'licensing' whereby MSC are activated or primed by local signals such as those produced by damaged and necrotic cells.

Corneal differentiation of MSC

While these results are somewhat encouraging, the presence of ZO-1 and N-cadherin in untreated cultures raises questions about the degree of transdifferentiation. When taken together with the functional data observed using BM-MSC [43], however, these findings suggest that MSC might at very least provide a partial surrogate for corneal endothelial cells in the event that transdifferentiation does not occur.

Corneal differentiation of MSC-derived from dental pulp (DP-MSC)

While only two studies by one group have explored the potential of DP-MSC as a source of corneal tissue [18, 37], both papers contain validated evidence of cornea-specific keratin expression. In the first study [18], while only traces of K3/K12 were detected in cultures of human DP-MSC by immunostaining (with transcripts for K12 detected by RT-PCR), co-expression of K3 and human nuclear antigen was detected by immunostaining following application to the wounded ocular surface of rabbits. These *in vivo* findings were essentially confirmed a year later in the group's second study [37].

Conclusions and recommendations

While there is evidence that MSC derived from tissues of non-corneal origin have some ability to produce proteins associated with corneal phenotype, only 10 out of the 28 papers that we analysed presented validated data in support of this hypothesis (summarized in Table 2). The strongest evidence exists with regard to expression of markers associated with corneal stromal cells (keratocytes), largely owing to the use of null mutant animal models and multiple sources of MSC having been tested with positive results. Although a greater number of studies have observed expression of markers associated with corneal epithelium, there are often significant gaps in the evidence provided, with many studies relying solely on immunohistochemistry data and several reports indicating that only a subpopulation of MSC

may be involved. Despite some interesting preliminary data, convincing evidence of MSC conversion into corneal endothelial cells has yet to be published.

On weight of evidence, we must therefore conclude that there is indeed value in pursuing the use of MSC from tissues of non-corneal origin as a potential source of corneal cells, and especially in the case of stromal tissue reconstruction. Nevertheless, the evidence for MSC differentiation into either corneal epithelial cells or corneal endothelial cells is relatively less clear. With regard to future studies, a number of key recommendations can be made.

1. First and foremost, specific markers of corneal phenotype should be used and ideally the expression of Pax-6, in conjunction with its recently identified regulator Wnt7A [52]), should also be examined as the canonical marker of ocular tissue development. In doing so, these studies will be consistent with the best evidence-based strategies being used for studying directed differentiation of induced pluripotent stem (iPS) cells.
2. Given the technical limitations of some antibodies used for detection of corneal cell phenotype (e.g. AE5 antibody to K3/K2p/76), the expression of corneal specific markers should always be confirmed at the transcriptional level. Moreover, reporting of experimental controls should be mandatory.

Failure to comply with either of these first two recommendations, risks production of data that would be unlikely to pass quality control measures required under good manufacturing practice (GMP) for clinical applications.

3. In the case of studies being conducted *in vivo*, it is essential that the provenance of observed “corneal cells” be traced back to the MSC administered to the animal.
4. Finally, while the majority of positive evidence has been obtained for MSC cultures established from bone marrow, this trend no doubt arises from the wider availability of

this resource and thus should not be taken to indicate an optimal MSC type for corneal reconstruction. Indeed, it could well be argued that MSC isolated from craniofacial tissues provide a more appropriate source, given their shared embryonic origin with corneal stromal cells and corneal endothelial cells (cranial neural crest).

In drawing our conclusions, we must stress that the findings of this systematic review in no-way detract from the broader potential use of MSC as a therapeutic agent for corneal repair through their proven abilities to modulate immune responses. Indeed, in the course of reviewing the 28 papers we noted several claims of improvements in corneal structure and/or function following administration of MSC [18, 20, 34, 42, 47, 49]. Moreover, there have been case reports of clinical efficacy outside the scope of our systematic review [53]. The question of whether or not transdifferentiation is necessary for a therapeutic effect is therefore perhaps academic, but we trust that our analysis provides some much needed clarity with respect to the mechanism of MSC action within the cornea.

Acknowledgements

This work was supported by a Project Grant (APP1049050) awarded from the National Health and Medical Research Council of Australia.

References

1. Pellegrini G, Traverso CE, Franzi AT et al. Long-term restoration of damaged corneal surfaces with autologous cultivated corneal epithelium. *Lancet* 1997;349:990-993.
2. Schwab IR. Cultured corneal epithelia for ocular surface disease. *Trans Am Ophthalmol Soc* 1999;97:891-986.

3. Angunawela RI, Mehta JS, Daniels JT. Ex-vivo ocular surface stem cell therapies: current techniques, applications, hurdles and future directions. *Expert Rev Mol Med* 2013;15:e4.
4. Choong PF, Mok PL, Cheong SK et al. Mesenchymal stromal cell-like characteristics of corneal keratocytes. *Cytherapy* 2007;9:252-258.
5. Polisetty N, Fatima A, Madhira SL et al. Mesenchymal cells from limbal stroma of human eye. *Mol Vis* 2008;14:431-442.
6. Lu JM, Zhou ZY, Zhang XR et al. A preliminary study of mesenchymal stem cell-like cells derived from murine corneal stroma. *Graefes Arch Clin Exp Ophthalmol* 2010;248:1279-1285.
7. Branch MJ, Hashmani K, Dhillon P et al. Mesenchymal stem cells in the human corneal limbal stroma. *Invest Ophthalmol Vis Sci* 2012;53:5109-5116.
8. Garfias Y, Nieves-Hernandez J, Garcia-Mejia M et al. Stem cells isolated from the human stromal limbus possess immunosuppressant properties. *Mol Vis* 2012;18:2087-2095.
9. Li GG, Zhu YT, Xie HT et al. Mesenchymal stem cells derived from human limbal niche cells. *Invest Ophthalmol Vis Sci* 2012;53:5686-5697.
10. Bray LJ, Heazlewood CF, Munster DJ et al. Immunosuppressive properties of mesenchymal stromal cell cultures derived from the limbus of human and rabbit corneas. *Cytherapy* 2014;16:64-73.
11. Walshe J, Harkin DG. Serial explant culture provides novel insights into the potential location and phenotype of corneal endothelial progenitor cells. *Experimental Eye Research* 2014;127:9-13.

12. Inatomi T, Nakamura T, Kojoyo M et al. Ocular surface reconstruction with combination of cultivated autologous oral mucosal epithelial transplantation and penetrating keratoplasty. *Am J Ophthalmol* 2006;142:757-764.
13. Ma Y, Xu Y, Xiao Z et al. Reconstruction of chemically burned rat corneal surface by bone marrow-derived human mesenchymal stem cells. *Stem Cells* 2006;24:315-321.
14. Ye J, Yao K, Kim JC. Mesenchymal stem cell transplantation in a rabbit corneal alkali burn model: engraftment and involvement in wound healing. *Eye (Lond)* 2006;20:482-490.
15. Oh JY, Kim MK, Shin MS et al. The anti-inflammatory and anti-angiogenic role of mesenchymal stem cells in corneal wound healing following chemical injury. *Stem Cells* 2008;26:1047-1055.
16. Ye J, Lee SY, Kook KH et al. Bone marrow-derived progenitor cells promote corneal wound healing following alkali injury. *Graefes Arch Clin Exp Ophthalmol* 2008;246:217-222.
17. Gu S, Xing C, Han J et al. Differentiation of rabbit bone marrow mesenchymal stem cells into corneal epithelial cells in vivo and ex vivo. *Mol Vis* 2009;15:99-107.
18. Monteiro BG, Serafim RC, Melo GB et al. Human immature dental pulp stem cells share key characteristic features with limbal stem cells. *Cell Prolif* 2009;42:587-594.
19. Du Y, Roh DS, Funderburgh ML et al. Adipose-derived stem cells differentiate to keratocytes in vitro. *Mol Vis* 2010;16:2680-2689.
20. Liu H, Zhang J, Liu CY et al. Cell therapy of congenital corneal diseases with umbilical mesenchymal stem cells: lumican null mice. *PLoS One* 2010;5:e10707.
21. Zhang X, Sun H, Li X et al. Utilization of human limbal mesenchymal cells as feeder layers for human limbal stem cells cultured on amniotic membrane. *J Tissue Eng Regen Med* 2010;4:38-44.

22. Liu H, Zhang J, Liu CY et al. Bone marrow mesenchymal stem cells can differentiate and assume corneal keratocyte phenotype. *J Cell Mol Med* 2012;16:1114-1124.
23. Katikireddy KR, Dana R, Jurkunas UV. Differentiation potential of limbal fibroblasts and bone marrow mesenchymal stem cells to corneal epithelial cells. *Stem Cells* 2014;32:717-729.
24. Rohaina CM, Then KY, Ng AM et al. Reconstruction of limbal stem cell deficient corneal surface with induced human bone marrow mesenchymal stem cells on amniotic membrane. *Transl Res* 2014;163:200-210.
25. Moher D, Liberati A, Tetzlaff J et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6:e1000097.
26. Pellegrini G, Dellambra E, Golisano O et al. p63 identifies keratinocyte stem cells. *Proc Natl Acad Sci U S A* 2001;98:3156-3161.
27. Barnard Z, Apel AJ, Harkin DG. Phenotypic analyses of limbal epithelial cell cultures derived from donor corneoscleral rims. *Clin Experiment Ophthalmol* 2001;29:138-142.
28. Schermer A, Galvin S, Sun TT. Differentiation-related expression of a major 64K corneal keratin in vivo and in culture suggests limbal location of corneal epithelial stem cells. *J Cell Biol* 1986;103:49-62.
29. Fini ME. Keratocyte and fibroblast phenotypes in the repairing cornea. *Prog Retin Eye Res* 1999;18:529-551.
30. Bray LJ, Heazlewood CF, Atkinson K et al. Evaluation of methods for cultivating limbal mesenchymal stromal cells. *Cytherapy* 2012;14:936-947.
31. Cheong YK, Ngoh ZX, Peh GS et al. Identification of cell surface markers glypican-4 and CD200 that differentiate human corneal endothelium from stromal fibroblasts. *Invest Ophthalmol Vis Sci* 2013;54:4538-4547.

32. Watanabe R, Hayashi R, Kimura Y et al. A novel gelatin hydrogel carrier sheet for corneal endothelial transplantation. *Tissue Eng Part A* 2011;17:2213-2219.
33. Arnalich-Montiel F, Pastor S, Blazquez-Martinez A et al. Adipose-derived stem cells are a source for cell therapy of the corneal stroma. *Stem Cells* 2008;26:570-579.
34. Coulson-Thomas VJ, Caterson B, Kao WW. Transplantation of human umbilical mesenchymal stem cells cures the corneal defects of mucopolysaccharidosis VII mice. *Stem Cells* 2013;31:2116-2126.
35. Espandar L, Bunnell B, Wang GY et al. Adipose-derived stem cells on hyaluronic acid-derived scaffold: a new horizon in bioengineered cornea. *Arch Ophthalmol* 2012;130:202-208.
36. Garzon I, Martin-Piedra MA, Alfonso-Rodriguez C et al. Generation of a biomimetic human artificial cornea model using Wharton's Jelly mesenchymal stem cells. *Invest Ophthalmol Vis Sci* 2014;55:4073-4083.
37. Gomes JA, Geraldine Monteiro B, Melo GB et al. Corneal reconstruction with tissue-engineered cell sheets composed of human immature dental pulp stem cells. *Invest Ophthalmol Vis Sci* 2010;51:1408-1414.
38. Guo T, Wang W, Zhang J et al. [Experimental study on repairing damage of corneal surface by mesenchymal stem cells transplantation]. *Zhonghua Yan Ke Za Zhi* 2006;42:246-250.
39. Hou GH, Ye N, Wu J et al. [Preliminary study on human bone marrow mesenchymal stem cells differentiation into epithelial-like cells]. *Zhonghua Yan Ke Za Zhi* 2010;46:719-724.
40. Jiang TS, Cai L, Ji WY et al. Reconstruction of the corneal epithelium with induced marrow mesenchymal stem cells in rats. *Mol Vis* 2010;16:1304-1316.

41. Joyce NC, Harris DL, Markov V et al. Potential of human umbilical cord blood mesenchymal stem cells to heal damaged corneal endothelium. *Mol Vis* 2012;18:547-564.
42. Lin HF, Lai YC, Tai CF et al. Effects of cultured human adipose-derived stem cells transplantation on rabbit cornea regeneration after alkaline chemical burn. *Kaohsiung J Med Sci* 2013;29:14-18.
43. Liu XW, Zhao JL. [Transplantation of autologous bone marrow mesenchymal stem cells for the treatment of corneal endothelium damages in rabbits]. *Zhonghua Yan Ke Za Zhi* 2007;43:540-545.
44. Ma XY, Bao HJ, Cui L et al. The graft of autologous adipose-derived stem cells in the corneal stroma after mechanical damage. *PLoS One* 2013;8:e76103.
45. Martinez-Conesa EM, Espel E, Reina M et al. Characterization of ocular surface epithelial and progenitor cell markers in human adipose stromal cells derived from lipoaspirates. *Invest Ophthalmol Vis Sci* 2012;53:513-520.
46. Nieto-Miguel T, Galindo S, Reinoso R et al. In vitro simulation of corneal epithelium microenvironment induces a corneal epithelial-like cell phenotype from human adipose tissue mesenchymal stem cells. *Curr Eye Res* 2013;38:933-944.
47. Pinarli FA, Okten G, Beden U et al. Keratinocyte growth factor-2 and autologous serum potentiate the regenerative effect of mesenchymal stem cells in cornea damage in rats. *Int J Ophthalmol* 2014;7:211-219.
48. Reinshagen H, Auw-Haedrich C, Sorg RV et al. Corneal surface reconstruction using adult mesenchymal stem cells in experimental limbal stem cell deficiency in rabbits. *Acta Ophthalmol* 2011;89:741-748.

Corneal differentiation of MSC

49. Zeppieri M, Salvetat ML, Beltrami AP et al. Human adipose-derived stem cells for the treatment of chemically burned rat cornea: preliminary results. *Curr Eye Res* 2013;38:451-463.
50. Zhang J, Huang C, Feng Y et al. Comparison of beneficial factors for corneal wound-healing of rat mesenchymal stem cells and corneal limbal stem cells on the xenogeneic acellular corneal matrix in vitro. *Mol Vis* 2012;18:161-173.
51. Zhang S, Espandar L, Imhof KM et al. Differentiation of Human Adipose-derived Stem Cells along the Keratocyte Lineage. *J Clin Exp Ophthalmol* 2013;4:11435.
52. Ouyang H, Xue Y, Lin Y et al. WNT7A and PAX6 define corneal epithelium homeostasis and pathogenesis. *Nature* 2014;511:358-361.
53. Agorogiannis GI, Alexaki VI, Castana O et al. Topical application of autologous adipose-derived mesenchymal stem cells (MSCs) for persistent sterile corneal epithelial defect. *Graefes Arch Clin Exp Ophthalmol* 2012;250:455-457.

Table 1. Results of systematic review for prior studies of non-corneal MSC conversion into corneal cells.

Tissue of MSC origin	Study	Species of MSC origin	Study design	Host species	Correct markers used	Adequate controls used	Origin of cells tracked	Our conclusions	Result
Bone marrow	[38]	Human	<i>In vivo</i>	Rabbit	Yes	Yes	Yes (Human antigen)	Transplanted cells formed epithelial-like structure on ocular surface. Cells were immunoreactive for K12 and human nuclear antigen.	√Ep
	[14]	Rabbit	<i>In vivo</i>	Rabbit	No (α -sma)	-	Yes (DiI)	Evidence of MSC survival and conversion into myofibroblasts.	Inc. S
	[13]	Human	<i>In vivo</i>	Rat	Yes	Yes	Yes (Human antigen)	Transplanted cells formed epithelial-like structure on ocular surface, but staining for cytokeratin was negative.	XEp
	[43]	Rabbit	<i>In vivo</i>	Rabbit <i>Auto</i>	No (Morph.)	No	Yes (BrdU)	Some evidence of ability to substitute for corneal endothelial cell function, but poor	?En

								evidence of transdifferentiation.	
[17]	Rabbit	<i>In vivo</i>	Rabbit	Yes	No	Yes	Evidence of MSC survival <i>in vivo</i> .	√Ep	
		<i>In vitro</i>	-	Yes	Yes	(BrdU) -	10% of cells express K3 <i>in vitro</i> .		
[40]	Rat	<i>In vivo</i>	Rat	Yes	Yes	No	Poor evidence without tracking of cells.	Inc. Ep	
		<i>In vitro</i>		Yes	No	-	Poor evidence without control data.		
[39]	Human	<i>Organ culture</i>	Pig	Yes	Yes	No	Cultures grown on denuded pig corneas displayed positive staining for K12 but origin of cells is unclear.	Inc. Ep	
[48]	Rabbit	<i>In vivo</i>	Rabbit	Yes	No	No	Poor evidence without control/tracking.	Inc. Ep	
		<i>In vitro</i>	-	Yes	No	-	Cultured MSC reported to express K3/K12, but data not shown.		
[50]	Rat	<i>In vitro</i>	-	Yes	Yes	-	K3 expression (IHC, no controls) supported by RT-PCR and WB.	√Ep	
[22]	Mouse	<i>In vivo</i>	Mouse <i>Kera -/-</i>	Yes	Yes	Yes (DiI)	Evidence of conversion into keratocytes.	√S	

	[24]	Human	<i>In vivo</i>	Rat	Yes	Yes	No	Poor evidence without tracking.	
			<i>In vitro</i>	-	Yes	Yes	-	Evidence of K3 <i>in vitro</i> .	√Ep
	[23]	Human	<i>In vitro</i>	-	Yes	Yes	-	K3 expression confirmed by ICC and RT-qPCR, but less than for control tissue.	√Ep
	[47]	Rat	<i>In vivo</i>	Rat	No	No	Yes (male DNA)	Traces of male DNA detected <i>in vivo</i> , but K19 not specific for corneal cells.	Inc. Ep
		<i>Male</i>		<i>Female</i>	(K19)				
			<i>In vitro</i>	-	No	No	-		
					(K19)				
Adipose tissue	[33]	Human	<i>In vivo</i>	Rabbit	Yes	Yes	Yes (DiI)	IHC data Inconclusive due to high background. Faint band for human keratocan detected by RT-PCR.	?S
	[19]	Human	<i>In vitro</i>	-	Yes	Yes	-	Keratocan expression confirmed by WB, RT-PCR and RT-qPCR.	√S
	[45]	Human	<i>In vitro</i>	-	Yes	Yes	-	Inconsistencies are reported with respect to K3 expression when measured by ICC, WB, RT-PCR and RT-qPCR.	?Ep

	[35]	Human	<i>In vivo</i>	Rabbit	Yes	No	Yes (DiI)	Poor evidence without control data.	Inc. S
	[42]	Human	<i>In vivo</i>	Rabbit	No (H&E)	-	No	Poor evidence without appropriate markers and tracking of cells.	Inc. Ep
	[46]	Human	<i>In vitro</i>	-	Yes	Yes	-	Inconclusive owing to expression of K3/K12 throughout all cultures tested including limbal fibroblasts.	Inc. Ep
	[44]	Rabbit	<i>In vivo</i>	Rabbit <i>Auto</i>	Yes	No	Yes (GFP)	Co-localization of GFP with keratocan and ALDH difficult to interpret without adequate controls.	Inc. S
	[49]	Human	<i>In vivo</i>	Rat	No (H&E)	-	Yes (CFSE)	Traces of label detected, but data inconclusive without exploring expression of cell phenotype.	Inc. Ep
	[51]	Human	<i>In vitro</i>	-	Yes	Yes	-	Inconsistencies are reported with respect to measurement of ALDH and keratocan by RT-qPCR and flow cytometry.	?S

Umbilical cord	[20]	Human	<i>In vivo</i>	Mice <i>Lum -/-</i> <i>Kera -/-</i>	Yes	Yes	Yes (DiI)	Good evidence provided of cells differentiating into functioning keratocytes.	√S
	[34]	Human	<i>In vivo</i>	Mice	No (F-actin)	No	Yes (DiI)	Evidence of cell survival and appropriate morphology. No markers of phenotype assessed.	Inc. S
	[36]	Human	<i>In vitro</i>	-	Yes	No	-	Poor evidence without control data.	Inc. Ep
Umbilical cord blood	[41]	Human	<i>Organ culture</i>	Human	Yes	Yes	Yes (GFP)	A shift observed towards a phenotype similar to corneal endothelial cells.	?En
Dental pulp	[18]	Human	<i>In vivo</i>	Rabbit	Yes	Yes	Yes (Human antigen)	Co-expression of K3 and human antigen	√Ep
			<i>In vitro</i>		Yes	Yes	-	Traces of K3/K12 in a few cells.	
	[37]	Human	<i>In vivo</i>	Rabbit	Yes	Yes	Yes (Human antigen)	Co-expression of K3 and human antigen	√Ep

							antigen)		
--	--	--	--	--	--	--	----------	--	--

Abbreviations. *Host species:* *Kera* *-/-* = keratocan null mutant mice, *Auto* = autologous transplant, *Lum* *-/-* = lumican null mutant mice. *Markers of cell phenotype:* ALDH = aldehyde dehydrogenase, α -sma = alpha smooth muscle actin, F-actin = filamentous actin, H&E = hematoxylin and eosin staining, K3 = cytokeratin 3, K12 = cytokeratin 12, K19 = cytokeratin 19, Morph. = morphology assessed. *Cell tracker reagents:* CFSE = Carboxyfluorescein succinimidyl ester, BrdU = bromo deoxyuridine, DiI = a fluorescent lipophilic dialkylcarbocyanine dye, GFP = green fluorescent protein. *Conclusions comments:* MSC = mesenchymal stromal cells of non-corneal origin, IHC = immunohistochemistry, RT-qPCR = quantitative (real time) reverse transcriptase polymerase chain reaction, RT-PCR = reverse transcriptase polymerase chain reaction, WB = Western blotting, ICC = immunocytochemistry. *Result comments:* Inc. = Inconclusive evidence, XEp = negative evidence for epithelial differentiation, $\sqrt{\text{Ep}}$ = positive evidence for epithelial differentiation, $\sqrt{\text{S}}$ = positive evidence for stromal cell (keratocyte) differentiation, ?S = partial evidence of stromal differentiation, ?Ep = partial evidence of epithelial differentiation, ?En = partial evidence of differentiation into corneal endothelial cells.

Table 2. Summary of validated evidence for MSC conversion to corneal phenotype.

	BM-MS	A-MS	UC-MS	DP-MS
Corneal epithelium	X √√√√√*	?		√√
Corneal stromal cells	√	√??	√	
Corneal endothelium	?		?	

Source of MSC: BM-MS = derived from bone marrow, A-MS = derived from adipose tissue, U-MS = derived from umbilical cord, DP-MS = derived from dental pulp.

Scoring: X = study reporting validated negative evidence, √ = study reporting validated positive evidence, ? = partial evidence of conversion.

*Note: In the majority of these studies, the levels of expression reported for corneal specific keratins (K3/K12) are substantially lower than that seen in corneal epithelial cells (positive control) and immunoreactivity is often limited to a small subset of cells.