



**Queensland University of Technology**  
Brisbane Australia

This is the author's version of a work that was submitted/accepted for publication in the following source:

Chan, Raymond J., McCarthy, Alexandra L., Devenish, Jackie, Sullivan, Karen A., & Chan, Alexandre  
(2015)  
Systematic review of pharmacologic and non-pharmacologic interventions to manage cognitive alterations after chemotherapy for breast cancer.  
*European Journal of Cancer*.  
(In Press)

This file was downloaded from: <http://eprints.qut.edu.au/79677/>

**© Copyright 2015 The Authors**

NOTICE: this is the author's version of a work that was accepted for publication in *European Journal of Cancer*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *European Journal of Cancer*, [in press] DOI: 10.1016/j.ejca.2014.12.017

**Notice:** *Changes introduced as a result of publishing processes such as copy-editing and formatting may not be reflected in this document. For a definitive version of this work, please refer to the published source:*

<http://dx.doi.org/10.1016/j.ejca.2014.12.017>

**Title:** Systematic review of pharmacologic and non-pharmacologic interventions to manage cognitive alterations after chemotherapy for breast cancer

**Running title:** Cognitive enhancement interventions after breast cancer chemotherapy

Raymond J. Chan  
NHMRC Research Fellow, Institute of Health and Biomedical Innovation,  
Queensland University of Technology  
Deputy Director of Research, Cancer Care Services, Royal Brisbane and  
Women's Hospital  
Herston, Queensland, Australia 4029

Alexandra L. McCarthy  
Chair of Cancer Nursing  
Princess Alexandra Hospital and the Institute of Health and Biomedical  
Innovation, Queensland University of Technology  
Victoria Park Rd  
Kelvin Grove, Queensland, Australia 4059

Jackie Devenish  
Liaison Librarian  
The University of Queensland Library Herston Health Sciences Library  
Royal Brisbane and Women's Hospital  
Herston, Queensland, Australia 4029

Karen A. Sullivan  
Professor, School of Psychology and Counseling  
Queensland University of Technology  
Victoria Park Rd  
Kelvin Grove, Queensland, Australia 4059

Alexandre Chan  
Associate Professor and Assistant Head  
Department of Pharmacy  
National University of Singapore  
Specialist Pharmacist (Oncology Pharmacy), Department of Pharmacy  
National Cancer Centre, Singapore 117543

Disclaimer: The authors have no conflicts of interest that are directly relevant to the content of this study.

Word count (abstract): 249

Word count (text): 3862

References: 56

Tables: 4

Figure : 1

**Keywords:** Breast cancer, Chemotherapy, Cognitive function, Cognitive training, Physical activity, Psychostimulants

## **Abstract (250 words)**

**Purpose:** Cognitive alterations are reported in breast cancer patients receiving chemotherapy. This has adverse effects on patients' quality of life and function. This systematic review investigates the effectiveness of pharmacologic and non-pharmacologic interventions to manage cognitive alterations associated with breast cancer treatment.

**Methods:** Medline via EBSCOhost, CINAHL and Cochrane CENTRAL were searched for the period January 1999 to May 2014 for prospective randomized controlled trials related to the management of chemotherapy-associated cognitive alterations. Included studies investigated the management of chemotherapy-associated cognitive alterations and used subjective or objective measures in patients with breast cancer during or after chemotherapy. Two authors independently extracted data and assessed the risk of bias.

**Results:** Thirteen studies involving 1138 participants were included. Overall, the risk of bias for the 13 studies were either high (n=11) or unclear (n=2). Pharmacologic interventions included psychostimulants (n=4), epoetin alfa (n=1), and Ginkgo biloba (n=1). Non-pharmacologic interventions were cognitive training (n=5) and physical activity (n=2). Pharmacologic agents were ineffective except for self-reported cognitive function in an epoetin alfa study. Cognitive training interventions demonstrated benefits in self-reported cognitive function, memory, verbal function and language and orientation/attention. Physical activity interventions were effective in improving executive function and self-reported concentration.

**Conclusion:** Current evidence does not favor the pharmacologic management of cognitive alterations associated with breast cancer treatment. Cognitive training and physical activity

interventions appear promising, but additional studies are required to establish their efficacy. Further research is needed to overcome methodological shortfalls such as heterogeneity in participant characteristics and non-standardized neuropsychological outcome measures.

## **INTRODUCTION**

Alterations in cognitive function are often observed in patients receiving chemotherapy, particularly those treated for breast cancer.[1] These changes can comprise poor word or name recall, difficulty staying focused, diminished ability to learn new things and a decreased ability to multitask.[2] Other alterations in executive function, information processing speed, language, motor function and spatial skills are documented. Depending on the nature of the malignancy and the treatment regimen, the time of onset, severity and duration of these changes are highly variable,[3] as are its affective, functional and psychosocial outcomes.[4]

Depending on the type of cancer investigated, estimates of the prevalence of cancer treatment-related alterations in cognitive function range from 16% to 75% during treatment,[5] although they can endure beyond treatment. Supported by findings from neuropsychological tests, reports indicate that individuals can experience longer-term cognitive changes for as long as 21 years after chemotherapy for breast cancer.[6] In addition, imaging research has reported a correlation between deficits in cognitive function and white matter changes in the brain.[7]

A number of systematic reviews and meta-analyses have investigated the prevalence of cognitive alteration and its association with treatment in cancer patients.[2, 8, 9] One systematic review[10] and one non-systematic narrative review, which discussed unpublished and ongoing studies,[11] focused on interventions to enhance cognitive function. Both reviews, however, are limited in that they included non-randomized controlled trials. Furthermore, Hines et al. limited their studies to cognitive behavioral therapy (CBT), which does not encompass the full range of interventions available.[10] In summary, a high quality,

comprehensive systematic review of interventions for managing chemotherapy-associated cognitive alterations is lacking.

This clinical problem has significant adverse effects on the post-treatment quality of life and function of patients with cancer; hence, interventions to prevent or manage it are warranted. Over the next decade, the number of individuals living with a cancer diagnosis is projected to increase by 31%, with a high proportion being patients with breast cancer.[12] Treatment-associated adverse effects in this growing population have significant public health implications if they are not well managed. In this paper, we systematically review the effectiveness of pharmacologic and non-pharmacologic interventions to manage alterations of cognitive function associated with breast cancer treatment.

## **METHOD**

This systematic review adhered to the PRISMA statement[13] for reporting systematic reviews.

### **Search strategy**

A medical librarian (JD) searched Medline via EBSCOhost, CINAHL and Cochrane CENTRAL for studies published between January 1999 and May 2014. The key search terms were chemotherapy, antineoplastic agents, chemoradiotherapy, cancer, neoplasms, randomized controlled trial, cognitive impairment, cognitive dysfunction, cognitive disorder, cognitive loss, cognitive deficit, and memory disorder. The search was limited to prospective randomized controlled trials (RCTs) published in English that investigated the management of chemotherapy-associated cognitive alterations (as primary or secondary outcomes).

Further manual searches of the reference lists of the relevant studies and reviews were undertaken by authors AC, RC and AM.

### **Study selection**

Three authors (RC, AM and AC) pre-screened all of the search results (titles and abstracts) and after consensus was reached for possible inclusion, the full text of all selected papers was assessed. Studies were included if they were prospective RCTs; reported pharmacologic or non-pharmacologic interventions for cognitive alterations in breast cancer patients during or after chemotherapy or multimodal therapy including chemotherapy; and used subjective or objective measures of cognitive function. Investigations of patients with secondary brain metastases and studies with less than 50% of breast cancer patients in the sample or with patients receiving radiation monotherapy were excluded. Unpublished reports, letters to the editor, retrospective chart reviews, and non-RCTs were also excluded.

### **Data extraction and rating of articles for risk of bias**

Two authors (RC, AM or AC) independently extracted the data using a pre-designed, piloted form. Studies were independently rated according to the Cochrane Collaboration's risk of bias (ROB) criteria for a high, low or unclear ROB with respect to random sequence generation, allocation concealment, blinding, incomplete outcome data, selective outcome reporting and other sources of bias.[14] An 'unclear' ROB was assigned to a study if the risk was unclear in one or more domains, with no domain rated as a high risk. A 'high' ROB was assigned to a study if the risk was high in one or more domains. A 'low' ROB was assigned to a study rated low risk in all domains.[15] Differences in ratings were settled by discussion or by a third person if consensus was not achieved by the two primary reviewers.



Studies that compared an intervention to usual care, placebo or another intervention and that presented adequate data for the calculation of effect size were evaluated. To determine the effects of the interventions in any of the included studies, effect sizes (mean difference [MD] or relative risk with a 95% confidence interval [CI]) were calculated using Review Manager 5.[14] We classified outcome assessments of <3 months, 3-6 months and  $\geq 6$  months as short-, medium- and long-term time points, respectively. If more than one measurement was reported within the defined period, the latest assessment was extracted. When published articles presented insufficient data to calculate the effect sizes, the authors were contacted for the required information. Although some studies reported statistical analyses for within-group changes from baseline, between-group differences were analyzed to determine the effects of the interventions (positive, negative or inconclusive). Data elicited from screening measures (e.g., Mini Mental Status Exam or High Sensitivity Cognitive Screen) were not extracted or analyzed. Objective outcome data were classified into the seven pre-defined domains of cognitive function recommended by Lezak et al.[16] and Hodgson et al.[17]

The interventions and outcome measurements reported in these studies were heterogeneous. Therefore, meta-analysis was not undertaken.

## **RESULTS**

Screening of 555 citations identified a total of 29 potentially relevant papers, the full texts of which were retrieved. Thirteen of the 29 studies were excluded as the majority of the included participants did not have breast cancer;[18-30] two studies did not include cognitive function measurement;[31, 32] and one study was not an RCT.[33] Thirteen studies met the inclusion criteria for quantitative and qualitative analyses. A flowchart detailing the identification of studies is provided in Figure 1.

Thirteen studies with a total of 1,138 participants were included in this review.[34-46] Eleven were undertaken in North America, one in Japan and the other in France. Six studies evaluated pharmacologic interventions (psychostimulants, n=4; erythropoietin stimulating agent, n=1, Gingko biloba, n=1). Seven studies investigated non-pharmacologic interventions, five of which involved cognitive training through forms of cognitive behavioral (n=4) or mindfulness therapy (n=1) and two of which explored physical activity. The characteristics of the included studies, the age of participants, treatments received, time since chemotherapy, sample sizes, assessment outcomes and time points for assessments are listed in Table 1.

Ten studies provided specific information on how the random sequence was generated. One provided sufficient information on allocation concealment.[38] The blinding of participants, personnel and outcome assessments was achieved in the six pharmacologic placebo trials but was not possible in the seven non-pharmacologic interventions. The risk of incomplete data outcome reporting bias was detected in six trials,[34-37, 44, 45] which did not provide reasons for participant dropout, or did not undertake intention-to-treat analyses. Three studies[35, 37, 38] had selective outcome reporting bias, as they did not report the data on all outcomes measured. Overall, the ROB for the 13 studies was either high[34-40, 43-46] (n=11) or unclear[41, 42] (n=2) (Table 2).

## **Pharmacologic interventions**

### *Psychostimulants*

Psychostimulants including d-methylphenidate (d-MPH) (n=2), methylphenidate (n=1) and modafinil (n=1) were evaluated. Two studies[41, 42] evaluated the ability of d-MPH to enhance cognitive function. In Lower et al.'s study,[41] patients were begun on 5 mg d-MPH twice daily with doses titrated weekly to a maximum of 50 mg/day over 8 weeks. In the other d-MPH study, conducted by Mar Fan et al.,[42] patients who demonstrated compliance through a placebo run-in phase were randomized to d-MPH 5 mg twice daily or to matched placebo. Doses were titrated to a maximum of 10 mg twice daily until the end of the final cycle of chemotherapy. In another cross-over study,[37] breast cancer patients were randomized to methylphenidate 18 mg/day for 2 weeks followed by placebo for 2 weeks or vice versa. Modafinil was trialed in a study involving two phases.[40] In the first phase, all patients received modafinil 100 mg once daily for 3 days and 200 mg once daily during an open-label period of 4 weeks. In the subsequent phase, patients who achieved a positive response in terms of attention and memory in the first phase were randomized to an additional 4 weeks of modafinil 200 mg/day or placebo. In the assessment of short-term and medium-term cognitive measures between the psychostimulants and controls, there was no statistically significant difference in cognitive measures in any of the studies (Table 3).

#### *Epoetin alfa (EPO)*

O'Shaughnessy et al.[45] evaluated whether epoetin alfa (EPO) could enhance cognitive and execution function in patients with breast cancer receiving adjuvant chemotherapy. Patients were randomized to receive 40,000 U of EPO subcutaneously weekly or a comparable volume of placebo during adjuvant or neoadjuvant chemotherapy over a maximum of 12 weeks. EPO doses were titrated according to hemoglobin levels. An improvement in self-

perceived cognitive function (EXIT-25) was noted in patients receiving EPO compared to a placebo (MD=-1.60 [95% CI, -2.81 to -0.39]).

### *Ginkgo biloba*

Barton et al.[34] investigated whether Ginkgo biloba could prevent cognitive impairment in breast cancer patients receiving adjuvant chemotherapy. Patients were randomized to receive Ginkgo biloba 60 mg twice daily or a placebo. The intervention commenced at the second cycle of chemotherapy and continued until 1 month after the completion of chemotherapy. The authors concluded that there were no significant differences in either subjective and objective cognitive measures between the two groups.

### *Toxicities of pharmacologic interventions*

Five of the six studies[34, 37, 41, 42, 45] evaluating pharmacologic interventions reported adverse events associated with the interventions and the placebo. The adverse events were generally mild, with few Common Terminology Criteria for Adverse Events (CTCAE) Grade 3 or 4 toxicities reported. In the two studies investigating d-MPH versus a placebo,[41, 42] higher incidences of dry mouth, nausea, dizziness, insomnia, anxiety and nervousness were reported among patients receiving d-MPH compared to the placebo. Similar findings were found in the methylphenidate study.[37] In the EPO study, O'Shaughnessy et al.[45] reported a fatal cerebrovascular accident in one patient in the EPO group. Ginkgo biloba was generally well tolerated compared to the placebo, with the exception of nausea, which was worse in the placebo group.[34]

### **Non-pharmacologic interventions**

### *Cognitive training*

In their three-arm study, Von Ah et al.[46] delivered memory training in one arm and speed processing training in the other. Memory training entailed the teaching of strategies to remember word lists, sequences and text material and learning how to apply the principles of meaningfulness, organization, visualization and association to these activities. Strategies focused on multiple mnemonic techniques. The intervention comprised 10 sessions, the first five comprising strategy instruction and practice, and the last five comprising practical exercises. This study reported a significant improvement in memory using objective neuropsychological testing compared to the control group, measured using composite scores for both immediate memory recall (MD=0.31, [95% CI 0.04 to 0.58]) and long term delayed memory (MD=0.46, [95% CI 0.12 to 0.80]). When self-perceived cognition for this intervention was measured with the Functional Assessment of Cancer Therapy-Cognitive Function (FACT-Cog), the results demonstrated a statistically significant improvement in the intervention group compared to the control group (MD=9.85 [95% CI, 1.67 to 18.03]). (Table 4) The other arm of the study involved speed processing training, which aimed to systematically reduce stimulus duration during a series of progressively more difficult computerized information processing tasks. Exercises comprised time-order judgment, discrimination, spatial match, forward span, instruction following and narrative memory tasks. Although the speed processing training did not target specifically at memory improvement, the training yielded positive improvement in memory, measured using composite scores for both immediate memory recall (MD=0.43, [95% CI 0.16 to 0.70]) and long term delayed memory (MD=0.47, [95% CI 0.13 to 0.81]) .

Kesler et al.[39] targeted executive function, which in their study encompassed working memory, cognitive flexibility, multitasking, planning and attention. The intervention comprised an online computerized training program undertaken in the participants' homes. It spanned 48 sessions of 20-30 minutes each over 6 weeks, with each session comprising combinations of 13 different exercises to enhance core executive function. Each participant logged in four times per week to complete five separate exercises. Exercises involved computerized visual stimuli that required a motor response such as a mouse click plus immediate feedback and reinforcement. The exercises were adaptive to individual activity, with the level of difficulty and complexity increasing according to a pre-determined algorithm. Compared to controls, the intervention participants demonstrated statistically significant improvements in executive function as measured by the Delis-Kaplan measure of verbal function and language skills (MD=2.00 [95% CI, 0.78 to 3.22]) and the Symbol Search measure of orientation and attention (MD=2.00 [95% CI, 0.16 to 3.84]).

Dolbeault et al.[36] delivered a CBT-based stress management intervention in which cognition, a secondary objective, was measured with the European Organisation for Research and Treatment of Cancer (EORTC)-Cognitive Functioning subscale. The results were not statistically significant for enhancements in cognitive function. In another study, rather than traditional repetitive techniques of mental exercise in CBT that aim to repair damaged neuro-circuitry to recover memory function, Ferguson et al.[38] taught strategies for cognitive processing and new behavior that compensated for chronic memory dysfunction. This intervention entailed the participants monitoring their cognitive failures and learning new processes to succeed in daily activities in which memory was required. The participants undertook twice weekly face-to-face sessions of 30-50 minutes each, with reinforcing phone

contacts between each visit. Differences in all outcome measures between intervention and control were not significant.

Milbury et al.[44] delivered a Tibetan sound meditation intervention, based on the premise that the focused concentration of such meditation, coupled with awareness, stress reduction and relaxation techniques would improve objective cognitive performance. Each participant undertook 60-minute meditation classes twice weekly for 6 weeks. Compared to controls, the intervention did not result in significant differences in objectively or subjectively measured cognitive function.

#### *Physical activity*

One physical activity intervention[43] comprised speed feedback therapy with a bicycle ergometer connected to a computer. Participants pedaled the bike to match the target arbitrarily displayed on the computer screen, which appeared as a pathway. The participants were instructed to pedal while visually tracking the path, and they undertook one pedaling session per week for 4 weeks. The exercise load was pre-set, with the participants pedaling for 5 minutes each session. Compared to controls, the intervention participants had improved executive function and motor function as measured by the Frontal Assessment Battery (MD=-2.50 [95% CI, -4.56 to -0.44]).

Culos-Reed et al.[35] delivered a program of modified hatha yoga, which focused on relaxation and awareness of breathing, body sensations and thoughts, to enhance post-treatment quality of life. Participants progressively built flexibility, strength and balance while maintaining awareness and relaxation. A reduction of cognitive disorganization (as

measured by the Profile of Mood State [POMS] Concentration subscale) was demonstrated in the intervention group compared to the control immediately on conclusion of the program (MD=-2.50 [95% CI, -4.56 to -0.44]).

## **DISCUSSION**

Current evidence does not favor the pharmacologic management of cognitive alteration associated with breast cancer treatment. The inherent variability of the psychology-derived cognitive training interventions makes it difficult to determine their role in practice. Some forms of cognitive training, particularly those that focus on quality of life enhancements, hold potential. For example, one study demonstrated a clinically important (i.e., subjectively reported) and statistically significant benefit in cognition-related quality of life.[46] Physical activity interventions also appear promising; however, methodological challenges in these studies preclude any concrete recommendations for practice.

Psychostimulants effectively manage cognitive issues related to attention deficit hyperactivity disorder and neurodegenerative diseases. The studies included in this review hypothesized that these agents are as effective in treating chemotherapy-associated cognitive alterations. These drugs include methylphenidate and d-MPH, which are sympathomimetic amines that modulate neurotransmitters in the brain. They are short-acting and were prescribed for a limited time during chemotherapy in these studies; therefore, the long-term benefits were not assessed. The long-term benefits of psychostimulants have not been established,[47] which suggests limited clinical benefits for individuals previously treated for breast cancer.[37, 41, 42] Similar to the sympathomimetics, modafinil improves wakefulness by acting on specific pathways in the brain that regulate sleep-wake patterns, without increasing the risk of the



extrapyramidal side effects that are commonly observed with sympathomimetics. Although patients receiving this treatment achieved a level of improvement in the open-label phase of the study,[40] this review did not detect any subsequent benefit in the randomized phase. In summary, the role of these agents is limited. In addition to conventional medications, herbal supplements such as Ginkgo biloba were also investigated as potential cognitive enhancers.[34] The literature indicates that Ginkgo biloba may improve cognitive function in patients with mild or moderate Alzheimer's disease or dementia.[48, 49] However, no benefits were observed in the study by Barton et al.[34] The authors proposed that the mechanisms underpinning chemotherapy-induced cognitive changes are different from those associated with dementia.[34]

In terms of non-pharmacologic interventions, cognitive training is useful in a range of conditions such as traumatic brain injury, which, like chemotherapy-associated dysfunction, demonstrate more subtle cognitive impairment.[39] Physical activity and cognitive training techniques involve repeated skills and awareness practice, adaptive difficulty levels and an engaging and rewarding environment. It is possible that these aspects of the interventions might not necessarily target cognitive function. However, they could yield positive benefits in cognitive organization due to overall enhancement of self-reported quality of life.[38] Given that quality of life was a primary or secondary endpoint in six of the seven non-pharmacologic studies in this review, and that improvements in the participants' quality of life were integral to many of these interventions, this assumption is worthy of empirical investigation.

A number of other interventions not included in this review also warrant further exploration. For example, the effectiveness of cholinesterase inhibitors (such as donepezil) and antioxidants (such as vitamin E) were investigated in the prevention of cognitive decline in patients with small cell lung cancer.[50] Unfortunately, poor patient accrual led to the early closure of the study. The results of trials of granulocyte macrophage-colony stimulating factor,[50] memantine[19] and medical qigong[29] are also promising, and further evaluations are required.

A number of methodological limitations featured in the included studies. First, there was at least one ROB in all of the studies. Second, the treatment characteristics of the participants were variable (e.g., they were at different stages of the disease, or received different treatment regimens). Third, the studies did not explain whether the participants were primed for cognitive impairment, with the entry criteria of many studies stipulating self-reported cognitive function. Fourth, the participants could not be blinded to the intervention in the non-pharmacologic studies. Fifth, many interventions required an intense commitment and repeat visits from participants, yet their sustainability over time is hard to determine, particularly where losses to follow-up were not documented.[34-37, 44, 45] Eleven studies did not evaluate the sustainability of effects beyond 3 months, by which time most of the interventions had ceased. Sixth, the majority of studies involved less than 50 participants per arm, although we recognize that many were pilot and feasibility investigations, which are integral components of high-quality research programs. We also recognize that there may be a potential risk of publication bias with studies reporting negative results remain unpublished. Seventh, some of the included studies did not include cognitive function as a primary endpoint. Trials are often not powered to detect differences in secondary outcomes. However,

we included these studies due to the potential for meta-analysis. Finally, the majority of the studies were undertaken in North America. Given that a patient's symptom experience is often culturally specific,[51] the generalizability of these results to other sociocultural contexts is uncertain.

The problems reflected in the range of methodologies and different cognitive outcomes reported in these studies could be addressed through the harmonization of intervention studies. The International Cognition and Cancer Task Force[52] provide some useful guidance in this respect. They recommend for observational studies that pre-treatment cognitive function is assessed, that intervention and control groups are standardized in terms of regimen and type of cancer, and that neuropsychological outcome measures are harmonized. These principles are equally germane to intervention studies.

This review suggests that in any intervention study in this field, the patient cohort requires careful consideration in terms of the stage of cancer and time since diagnosis. Studies could incorporate a screen for expectancy effects prior to randomization that are controlled for during data analysis. Expectancies and stereotypes including those associated with diagnoses are known to influence cognitive profiles[53]. Screening should also assess premorbid cognitive function if possible.[54] Subjective and objective measurements appear to be equally important in detecting effects. Self-reported measures detect outcomes that are clinically significant to patients, whereas objective neuropsychological tests remain the gold standard.[52] Utilization of validated tools would also be essential for future interventional studies.

Future studies would benefit from the addition of an attention control arm to address the bias inherent in the inability to blind in non-pharmacologic studies.[55, 56] The potential uptake of the intervention should also be carefully considered. Aside from feasibility studies to determine this, interventions need to be accessible and easy for patients to undertake. Technology-enhanced interventions have promise, particularly multimodal programs that combine physical activity and cognitive training.

In summary, the burden associated with this commonly reported problem in the breast cancer community is significant. The science to address this problem, however, is imprecise. Well-designed clinical studies are clearly warranted to enhance the quality of life and function of this growing population.

## References

1. Ahles TA, Root JC, Ryan EL. Cancer- and cancer treatment-associated cognitive change: an update on the state of the science. *J Clin Oncol* 2012; 30: 3675-3686.
2. Jansen CE, Miaskowski C, Dodd M et al. A metaanalysis of studies of the effects of cancer chemotherapy on various domains of cognitive function. *Cancer* 2005; 104: 2222-2233.
3. Wefel JS, Saleeba AK, Buzdar AU, Meyers CA. Acute and late onset cognitive dysfunction associated with chemotherapy in women with breast cancer. *Cancer* 2010; 116: 3348-3356.
4. Jacobs SR, Jacobsen PB, Booth-Jones M et al. Evaluation of the Functional Assessment of Cancer Therapy Cognitive Scale with hematopoietic stem cell transplant patients. *Journal of Pain and Symptom Management* 2007; 33: 13-23.
5. Cheung YT, Chui WK, Chan A. Neuro-cognitive impairment in breast cancer patients: pharmacological considerations. *Crit Rev Oncol Hematol* 2012; 83: 99-111.
6. Koppelmans V, Breteler MM, Boogerd W et al. Neuropsychological performance in survivors of breast cancer more than 20 years after adjuvant chemotherapy. *Journal of Clinical Oncology* 2012; 30: 1080-1086.
7. Deprez S, Amant F, Smeets A et al. Longitudinal assessment of chemotherapy-induced structural changes in cerebral white matter and its correlation with impaired cognitive functioning. *J Clin Oncol* 2012; 30: 274-281.
8. Jim HS, Phillips KM, Chait S et al. Meta-analysis of cognitive functioning in breast cancer survivors previously treated with standard-dose chemotherapy. *J Clin Oncol* 2012; 30: 3578-3587.
9. McGinty HL, Phillips KM, Jim HS et al. Cognitive functioning in men receiving androgen deprivation therapy for prostate cancer: a systematic review and meta-analysis. *In Support Care Cancer*. 2014.
10. Hines S, Ramis MA, Pike S, Chang AM. The Effectiveness of Psychosocial Interventions for Cognitive Dysfunction in Cancer Patients Who Have Received Chemotherapy: A Systematic Review. *Worldviews Evid Based Nurs* 2014.
11. Gehring K, Roukema JA, Sitskoorn MM. Review of recent studies on interventions for cognitive deficits in patients with cancer. *Expert Rev Anticancer Ther* 2012; 12: 255-269.
12. de Moor JS, Mariotto AB, Parry C et al. Cancer survivors in the United States: prevalence across the survivorship trajectory and implications for care. *Cancer Epidemiol Biomarkers Prev* 2013; 22: 561-570.
13. Liberati A, Altman DG, Tetzlaff J et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ* 2009; 339: b2700.
14. Higgins J, Green S. *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 In. The Cochrane Collaboration 2011.
15. Garcia MK, McQuade J, Haddad R et al. Systematic review of acupuncture in cancer care: a synthesis of the evidence. *J Clin Oncol* 2013; 31: 952-960.
16. Lezek M, Howieson D, Bigler E, Tranel D. *Neuropsychological Assessment*. USA: Oxford University Publisher 2012.
17. Hodgson KD, Hutchinson AD, Wilson CJ, Nettelbeck T. A meta-analysis of the effects of chemotherapy on cognition in patients with cancer. *Cancer Treat Rev* 2013; 39: 297-304.


18. Boele F, Douw L, de Groot M et al. The effect of modafinil on fatigue, cognitive functioning, and mood in primary brain tumor patients: a multicenter randomized controlled trial. *Neuro-Oncology* 2013; 15: 1420-1428.
19. Brown PD, Pugh S, Laack NN et al. Memantine for the prevention of cognitive dysfunction in patients receiving whole-brain radiotherapy: a randomized, double-blind, placebo-controlled trial. *Neuro-Oncology* 2013; 15: 1429-1437.
20. Capuron L, Gummnick JF, Musselman DL et al. Neurobehavioral effects of interferon-alpha in cancer patients: phenomenology and paroxetine responsiveness of symptom dimensions. *Neuropsychopharmacology* 2002; 26: 643-652.
21. Cherrier MM, Anderson K, David D et al. A randomized trial of cognitive rehabilitation in cancer survivors. *Life Sciences* 2013; 93: 617-622.
22. Gehring K, Patwardhan SY, Collins R et al. A randomized trial on the efficacy of methylphenidate and modafinil for improving cognitive functioning and symptoms in patients with a primary brain tumor. *Journal Of Neuro-Oncology* 2012; 107: 165-174.
23. Goedendorp MM, Knoop H, Gielissen MFM et al. The effects of cognitive behavioral therapy for postcancer fatigue on perceived cognitive disabilities and neuropsychological test performance. *Journal of Pain & Symptom Management* 2014; 47: 35-44.
24. Green HJ, Pakenham KI, Headley BC et al. Altered cognitive function in men treated for prostate cancer with luteinizing hormone-releasing hormone analogues and cyproterone acetate: a randomized controlled trial. *BJU international* 2002; 90: 427-432.
25. Jarden M, Nelausen K, Hovgaard D et al. The effect of a multimodal intervention on treatment-related symptoms in patients undergoing hematopoietic stem cell transplantation: a randomized controlled trial. *Journal of Pain & Symptom Management* 2009; 38: 174-190.
26. Jim H, Boyd T, Booth-Jones M et al. Granulocyte Macrophage Colony Stimulating Factor Treatment is Associated with Improved Cognition in Cancer Patients. *Brain Disorders & Therapy* 2012; 1: 1-10.
27. Lundorff LE, Jønsson BH, Sjøgren P. Modafinil for attentional and psychomotor dysfunction in advanced cancer: a double-blind, randomised, cross-over trial. *Palliative Medicine* 2009; 23: 731-738.
28. McDougall GJ, Becker H, Acee TW et al. Symptom management of affective and cognitive disturbance with a group of cancer survivors. *Arch Psychiatr Nurs* 2011; 25: 24-35.
29. Oh B, Butow P, Mullan B et al. Effect of medical Qigong on cognitive function, quality of life, and a biomarker of inflammation in cancer patients: a randomized controlled trial. *Supportive Care In Cancer: Official Journal Of The Multinational Association Of Supportive Care In Cancer* 2012; 20: 1235-1242.
30. Poppelreuter M, Weis J, Bartsch HH. Effects of specific neuropsychological training programs for breast cancer patients after adjuvant chemotherapy. *J Psychosoc Oncol* 2009; 27: 274-296.
31. Doorenbos A, Given B, Given C et al. Reducing symptom limitations: a cognitive behavioral intervention randomized trial. *Psychooncology* 2005; 14: 574-584.
32. Given CW, Sikorskii A, Tamkus D et al. Managing symptoms among patients with breast cancer during chemotherapy: results of a two-arm behavioral trial. *J Clin Oncol* 2008; 26: 5855-5862.
33. Schuurs A, Green H. A feasibility study of group cognitive rehabilitation for cancer survivors: enhancing cognitive function and quality of life. *Psycho-Oncology* 2013; 22: 1043-1049.

34. Barton DL, Burger K, Novotny PJ et al. The use of Ginkgo biloba for the prevention of chemotherapy-related cognitive dysfunction in women receiving adjuvant treatment for breast cancer, N00C9. *Support Care Cancer* 2013; 21: 1185-1192.
35. Culos-Reed SN, Carlson LE, Daroux LM, Hatley-Aldous S. A pilot study of yoga for breast cancer survivors: physical and psychological benefits. *Psycho-oncology* 2006; 15: 891-897.
36. Dolbeault S, Cayrou S, BrÈdard A et al. The effectiveness of a psycho-educational group after early-stage breast cancer treatment: results of a randomized French study. *Psycho-oncology* 2009; 18: 647-656.
37. Escalante CP, Meyers C, Reuben JM et al. A Randomized, Double-blind, 2-Period, Placebo-Controlled Crossover Trial of a Sustained-Release Methylphenidate in the Treatment of Fatigue in Cancer Patients. *Cancer Journal* 2014; 20: 8-14.
38. Ferguson RJ, McDonald BC, Rocque MA et al. Development of CBT for chemotherapy-related cognitive change: results of a waitlist control trial. *Psycho-Oncology* 2012; 21: 176-186.
39. Kesler S, Hadi Hosseini SM, Heckler C et al. Cognitive training for improving executive function in chemotherapy-treated breast cancer survivors. *Clinical Breast Cancer* 2013; 13: 299-306.
40. Kohli S, Fisher SG, Tra Y et al. The effect of modafinil on cognitive function in breast cancer survivors. *Cancer* 2012; 115: 2605-2616.
41. Lower EE, Fleishman S, Cooper A et al. Efficacy of dexamethylphenidate for the treatment of fatigue after cancer chemotherapy: a randomized clinical trial. *J Pain Symptom Manage* 2009; 38: 650-662.
42. Mar Fan HG, Clemons M, Xu W et al. A randomised, placebo-controlled, double-blind trial of the effects of d-methylphenidate on fatigue and cognitive dysfunction in women undergoing adjuvant chemotherapy for breast cancer. *Supportive care in cancer* 2012; 16: 577-583.
43. Miki E, Kataoka T, Okamura H. Feasibility and efficacy of speed-feedback therapy with a bicycle ergometer on cognitive function in elderly cancer patients in Japan. *Psycho-Oncology* 2014.
44. Milbury K, Chaoul A, Biegler K et al. Tibetan sound meditation for cognitive dysfunction: Results of a randomized controlled pilot trial. *Psycho-oncology* 2013; 22: 2354-2363.
45. O'Shaughnessy JA, Vukelja SJ, Holmes FA et al. Feasibility of quantifying the effects of epoetin alfa therapy on cognitive function in women with breast cancer undergoing adjuvant or neoadjuvant chemotherapy. *Clinical Breast Cancer* 2005; 5: 439-446.
46. Von Ah D, Carpenter JS, Saykin A et al. Advanced cognitive training for breast cancer survivors: a randomized controlled trial. *Breast Cancer Research And Treatment* 2012; 135: 799-809.
47. Hazell P. The challenges to demonstrating long-term effects of psychostimulant treatment for attention-deficit/hyperactivity disorder. *Curr Opin Psychiatry* 2011; 24: 286-290.
48. Janssen IM, Sturtz S, Skipka G et al. Ginkgo biloba in Alzheimer's disease: a systematic review. *Wien Med Wochenschr* 2010; 160: 539-546.
49. Weinmann S, Roll S, Schwarzbach C et al. Effects of Ginkgo biloba in dementia: systematic review and meta-analysis. *BMC Geriatr* 2010; 10: 14.

50. Jatoi A, Kahanic SP, Frytak S et al. Donepezil and vitamin E for preventing cognitive dysfunction in small cell lung cancer patients: preliminary results and suggestions for future study designs. *Supportive care in cancer* 2005; 13: 66-69.
51. Cheung YT, Shwe M, Tan YP et al. Cognitive changes in multiethnic Asian breast cancer patients: a focus group study. *Ann Oncol* 2012; 23: 2547-2552.
52. Wefel JS, Vardy J, Ahles TA, Schagen SB. International Cognition and Cancer Task Force recommendations to harmonise studies of cognitive function in patients with cancer. *Lancet Oncology* 2011; 12: 703-708.
53. Suhr JA, Wei C. Response expectancies and their potential influence in neuropsychological evaluation. In Arnett PA (ed) *Secondary influences on neuropsychological test performance: Research findings and practical applications*. New York: Oxford University Press 2013.
54. Axelrod BN, Vanderploeg RD, Schinka JA. Comparing methods for estimating premorbid intellectual functioning. *Arch Clin Neuropsychol* 1999; 14: 341-346.
55. Boutron I, Guttet L, Estellat C et al. Reporting methods of blinding in randomized trials assessing nonpharmacological treatments. *PLoS Med* 2007; 4: e61.
56. Freedland KE, Mohr DC, Davidson KW, Schwartz JE. Usual and unusual care: existing practice control groups in randomized controlled trials of behavioral interventions. *Psychosom Med* 2011; 73: 323-335.



Figure 1. PRISMA Flow Diagram



**Table 1. Characteristics of Included Studies (n=13)**

Study	Participants	Prior treatments	Time since chemotherapy	Comparisons	Domains examined	Assessed time point
Barton (2013)[34]  USA	Breast CA: Both arms: 100%  Age: I: ≥50 years: 50% C: ≥50 years: 50%	I: AC: 33%; AC and taxane: 52% Others: 15% Tam planned: 51%  C: AC: 36%; AC and taxane: 52% Others: 12% Tam planned: 54%	During chemo	Ginkgo biloba 60 mg vs placebo twice daily (started at the second cycle of chemo and ceased at 1 month after completion of chemo)  I: n=107 C: n=103	Orientation and attention, self-reported cognitive function	Short-term, medium-term, long-term
Culos-Reed (2006)[35]  Canada	Breast CA: Both arms: 100%  Age: Both arms: 51.2 (10.3)	Both arms: Chemo: percentage not stated	>3 months post chemo	7-week yoga program vs control  I: n=10 C: n=10	Self-reported cognitive function	Short-term
Dolbeault (2009)[36]  France	Breast CA: Both arms: 100%  Age: I: 54.5 (9.3) C: 51.6 (9.6)	I: Chemo: 45.1% C: Chemo: 61.4%	Not stated	A CBT-based psycho-educational group intervention (8 weekly 2-hour sessions) vs control (usual care)  I: n=102 C: n=101	Self-reported cognitive function	Short-term
Escalante (2014)[37]  USA	Breast CA: Both arms: 100%  Age: Both arms: Median 57 (Range: 32-79)	Both groups: Chemo: 100%	Either undergoing or completed treatment in the previous 12 months	Methylphenidate (18 mg daily) vs placebo for 14 days  n=42 (cross-over design)	Orientation and attention	Short-term
Ferguson (2012)[38]  USA	Breast CA: Both arms: 100%  Age: I: 51.2 (7.3) C: 49.4 (5.1)	I: Received AC/FAC C: Received AC/FAC	>18 months post chemo	8-week Memory and Attention Adaptive Training (MAAT) vs waitlist control  I: n=19 C: n=21	Orientation and attention, Executive function and motor function, memory, self-reported cognitive function	Short-term
Kesler (2013)[39]	Breast CA:	I: 100% chemo	Mean (SD): 6.0 (3)	Online computerized training program (5	Orientation and attention,	Short-term

USA	Both arms: 100% Age: I: 55.0 (7) C: 56.0 (6)	70% RT 60% HT  C: 100% chemo 63% RT 63% HT	months	exercises 4 times weekly for 12 weeks) vs usual care  I: n=21 C: n=21	executing functioning and motor function, verbal function and language skills, memory, self-reported cognitive function	
Kohli (2012)[40]  USA	Breast CA: Both arms: 100%  Age: I: 54.0 (10.3) C: 56.35 (11.4)	I: 100% chemo, 82% RT C: 100% chemo, 85% RT	>30 days post chemo	Modafinil 200 mg daily vs placebo for 4 weeks  I: n=34 C: n=34	Orientation and attention, memory	Short-term
Lower (2009)[41]  USA	Breast CA: I: 78% C: 73%  Age: I: 52.5 (10.2) C: 53.2 (8.4)	I: 100% chemo C: 100% chemo	Mean (SD): 115.3 (106.5) weeks	D-methylphenidate vs placebo for 8 weeks; dose modifications were allowed; max 50 mg/day  I: n=76 C: n=78	Orientation and attention	Short-term
Mar Fan (2008)[42]  Canada	Breast CA: Both arms: 100%  Age: I: Median=50, Range= 36-72 C: Median=51, Range= 37-74	Both arms: 100% Chemo  Both arms: AC: 100% Cy: 96.5% 5FU: 33.3% Taxane: 31.6%	I: Median (range): 84 (23-141) days post chemo  C: Median (range): 85 (26-131) days post chemo	D-methylphenidate (titration: 5 to 10 mg twice daily) vs placebo until final cycle of chemo  I: n=28 C: n=29	Memory	Short-term
Miki (2014)[43]  Japan	Breast CA: I: 55.3% C: 55.0%  Age: I: 72.97 (4.57) C: 75.45 (6.57)	I: 81.6% Surgery 23.7% Chemo 68.4% HT 68.4% RT  C: 72.5% Surgery 27.5% Chemo 80.0% HT 4.00% RT	Not stated	4-week Speed-feedback therapy with a bicycle ergometer vs usual care  I: n=38 C: n=40	Executing function and motor function	Short-term
Milbury	Breast CA:	I: 100% chemo	6-60 months post	Tibetan sound meditation (2x weekly sessions	Orientation and attention,	Short-term

(2014)[44] USA	Both arms: 100% Age: I: 53.0 (6.6) C: 54.1 (8.6)	73.9% RT 87% Surgery C: 100% chemo 79.2% RT 100% Surgery	chemo	for 6 weeks) vs waitlist control I: n=18 C: n=24	Memory, verbal functions and language skills, self-reported cognitive function	
O'Shaughnessy (2005)[45] USA	Breast CA: Both arms: 100% Age: I: 53.3 (9.7) C: 54.3 (12)	Both arms: 100% chemo Both arms: Doxorubicin/epirubicin:100% Cy: 96.8% 5-FU: 11.7% Taxane: 24.5%	Undergoing chemo	40,000 U epoetin alfa subcutaneous weekly vs placebo (started on D1 of cycle 1 of 4 cycles of chemo, and continued for a maximum of 12 weeks). I: n=47 C: n=47	Executive function and motor function, self-reported cognitive function	Short-term, long-term
Von Ah (2012)[46] USA	Breast CA: All three arms: 100% Age: I1: 55.19 (7.58) I2: 56.93 (7.83) C: 57.21 (9.8)	All three arms: 100% chemo	1 year post chemo	Memory training (Advanced Cognitive Training for Independent and Vital Elderly (ACTIVE) program (10 sessions for 6-8 weeks) vs speed of processing training (Posit Science®) (10 sessions for 6-8 weeks) vs waitlist group I1: n=29 I2: n=30 C: n=29	Memory, self-reported cognitive function	Short-term

Abbreviations: 5-FU: 5-Fluorouracil; AC: Anthracycline and cyclophosphamide; C: Control; CBT: Cognitive behavioral therapy; Cy: Cyclophosphamide; Chemo: Chemotherapy; FAC: 5-Fluorouracil, anthracycline and cyclophosphamide; HT: Hormonal therapy, I: Intervention, RT: Radiation therapy. Tam: Tamoxifen. Measurement time points: short term: less than 3 months; medium term: 3-6 months; long term: beyond 6 months.

**Table 2. Risk of bias (ROB) table for included studies (n=13)**

Study	Sequence generation	Allocation concealment	Blinding of participants and personnel	Blinding of outcome assessment	Incomplete outcome data	Selective outcome reporting	Other sources of bias	Level of risk
Barton (2013)[34]	+	0	+	+	-	+	+	H
Culos-Reed (2006)[35]	0	0	0	-	-	-	+	H
Dolbeault (2009)[36]	+	0	-	-	-	+	+	H
Escalante (2014)[37]	0	0	+	+	0	-	+	H
Ferguson (2012)[38]	+	+	+	+	+	-	+	H
Kesler (2013)[39]	+	0	-	0	+	+	+	H
Kohli (2012)[40]	+	0	+	+	+	+	-	H
Lower (2009)[41]	+	0	+	+	+	+	+	U
Mar Fan (2008)[42]	0	0	+	+	+	+	+	U
Miki (2014)[43]	+	0	-	+	+	+	+	H
Milbury (2014)[44]	+	0	-	0	0	+	+	H
O'Shaughnessy (2005)[45]	+	0	+	+	-	+	+	H
Von Ah (2012)[46]	+	0	-	-	+	+	+	H

0 represents an unclear ROB, - represents a high ROB, and + represents a low ROB. Abbreviations: H, high ROB (-) for one or more domains; L, low ROB (+) for all domains; U, unclear ROB for one or more domains.

**Table 3. Pharmacologic interventions and outcomes (n=6)**

Categories	Intervention	Tool	Domains examined	Assessment time	Effect size	Conclusion	
Psychostimulants	Dexamethylphenidate[41]	Modified Swanson, Nelson and Pelham Attention Deficit/Hyperactivity Scale (SNAP)	Orientation and attention	Short-term	MD=0.30 (95% CI, -2.19 to 2.79)	Negative	
	Dexamethylphenidate[42]	Hopkins Verbal Learning Test-Revised (HVLTR)	Memory	Medium-term	RR=1.20 (95% CI, 0.72-2.00)	Negative	
	Methylphenidate[37]	Digit Span	Orientation and attention	Short-term	Data not reported	Inconclusive	
	Modafinil[40]		Cognitive Drug Research Computerized Assessment System <sup>#</sup> (Power of Attention)	Orientation and attention	Short-term	MD=-27.64 (95% CI, -89.66 to 34.38)	Negative
			Cognitive Drug Research Computerized Assessment System <sup>#</sup> (Continuity of Attention)	Orientation and attention	Short-term	MD=0.65 (95% CI: -0.65 to 1.95)	Negative
			Cognitive Drug Research Computerized Assessment System <sup>#</sup> (Episodic Secondary Memory)	Memory	Short-term	MD=-4.52 (95% CI, -29.84 to 20.80)	Negative
			Cognitive Drug Research Computerized Assessment System <sup>#</sup> (Working Memory)	Orientation and attention	Short-term	MD=0.12 (95% CI, -0.06 to 0.30)	Negative
		Cognitive Drug Research Computerized Assessment System <sup>#</sup> (Speed of Memory)	Orientation and attention	Short-term	MD=-103.46 (95% CI, -567.03 to 360.10)	Negative	
Erythropoietic stimulating agent	Epoetin alfa[45]	Executive Clock Drawing Task 1 (CLOX1)	Executive function and motor function	Short-term	MD=0.10 (95% CI, -0.93 to 1.13)	Negative	
		Executive Clock Drawing Task 1 (CLOX1)	Executive function and motor function	Long-term	MD=-0.80 (95% CI, -1.83 to 0.23)	Negative	
		The Executive Interview (EXIT-25)	Self-reported cognitive function	Short-term	MD=-1.60 (95% CI, -2.81 to -0.39)*	Positive	
		The Executive Interview (EXIT-25)	Self-reported cognitive function	Long-term	MD=-0.10 (95% CI, -1.35 to 1.15)	Negative	
Complementary alternative	Gingko biloba[34]	Trail Making Test-A	Orientation and attention	Short-term, medium-term, long-	Data not extractable	Inconclusive	

medicine		Trail Making Test-B	Orientation and attention	term Short-term, medium-term, long-term	Data not extractable	Inconclusive
----------	--	---------------------	---------------------------	--	----------------------	--------------

\* $p < 0.05$ ; #These subtest names do not appear in the original paper—they are derivative measures of a factor analysis. Positive represents ‘favors intervention’ and Negative represents ‘does not favor intervention’. Measurement time points: short-term: less than 3 months; medium-term: 3-6 months; long-term: beyond 6 months.

**Table 4. Non-pharmacologic interventions and outcomes (n=7)**

Categories	Intervention	Tool	Domains examined	Assessment time point	Effect size	Conclusion
Cognitive training	Computerized cognitive training[39]	Wisconsin Card Sorting Test	Executive function and motor function	Short-term	MD=3.00 (95% CI, -1.49 to 7.49)	Negative
		Delis-Kaplan (letter fluency)	Verbal function and language skills	Short-term	MD=2.00 (95% CI, 0.78 to 3.22)*	Positive
		Digit Span	Orientation and attention	Short-term	MD=0.90 (95% CI, -3.17 to 1.17)	Negative
		Symbol Search	Orientation and attention	Short-term	MD=2.00 (95% CI, 0.16 to 3.84)*	Positive
		Hopkins Verbal Learning Test-Revised (HVLTR)	Memory	Short-term	MD=1.00 (95% CI, -5.95 to 3.95)	Negative
		BRIEF GEC	Self-reported cognitive function	Short-term	MD=-2.00 (95% CI, -10.37 to 6.37)	Negative
		BRIEF GEC (plan and organize)	Self-reported cognitive function	Short-term	MD=-5.00 (95% CI -13.30 to 3.30)	Negative
		BRIEF GEC (task monitor)	Self-reported cognitive function	Short-term	MD=-4.00 (95% CI, -10.75 to 2.75)	Negative
	Cognitive-behavioral therapy[38]	Trail Making Test-B	Orientation and attention	Short-term	MD=3.22 (95% CI, -7.9 to 14.34)	Negative
		Color Word Trail (D-KEFS subset)	Executive function and motor function	Short-term	MD=-0.73 (95% CI, -6.44 to 4.98)	Negative
		Color Word Switching Trail (D-KEFS subset)	Executive function and motor function	Short-term	MD=0.53 (95% CI, -6.81 to 7.87)	Negative
		Digit Symbol-Coding (subtest of Wechsler Adult Intelligence Scale)	Orientation and attention	Short-term	MD=1.09 (95% CI, -0.75 to 2.93)	Negative
		California Verbal Learning Test (CVLT) II	Memory	Short-term	MD=4.26 (95% CI, -2.32 to 10.84)	Negative
		MASQ scores	Self-reported cognitive function	Short-term	MD=4.02 (95% CI, -8.83 to 16.87)	Negative
	Memory training[46]	FACT-Cog	Self-reported cognitive function	Short-term	MD=9.85 (95% CI, 1.67 to 18.03)*	Positive
Composite score: Rey Auditory Verbal Learning Test (AVLT) (sum recall, short delay and recognition		Memory	Short-term	MD=0.31 (95% CI 0.04 to 0.58)*	Positive	



		score and Rivermead Behavioral Paragraph Recall Test (immediate recall score)				
		Composite score: Rey Auditory Verbal Learning Test (AVLT) (long term delay score) and Rivermead Behavioral Paragraph Recall Test (long term delay score)	Memory	Short-term	MD=0.46 (95% CI, 0.12 to 0.80)*	Positive
	Speed of processing training[46]	FACT-Cog	Self-reported cognitive function	Short-term	MD=6.66 (95% CI, -1.43 to 14.75)	Negative
		Composite score: Rey Auditory Verbal Learning Test (AVLT) (sum recall, short delay and recognition score and Rivermead Behavioral Paragraph Recall Test (immediate recall score)	Memory	Short-term	MD=0.43 (95% CI, 0.16 to 0.70)*	Positive
		Composite score: Rey Auditory Verbal Learning Test (AVLT) (long term delay score) and Rivermead Behavioral Paragraph Recall Test (long term delay score)	Memory	Short-term	MD=0.47 (95% CI, 0.13 to 0.81)*	Positive
	Psycho-education[36]	EORTC-CF	Self-reported cognitive function	Short-term	MD=-0.16 (95% CI, -0.38 to 0.06)	Negative
	Tibetan sound[44]	Rey Auditory Verbal Learning Test (AVLT)	Memory	Short-term	Data not extractable	Inconclusive
		Digit Span	Orientation and attention	Short-term	MD=0.43 (95% CI, -0.17 to 1.03)	Negative
		Digit Symbol-Coding (subtest of Wechsler Adult Intelligence Scale)	Orientation and attention	Short-term	MD=0.46 (95% CI, -0.14 to 1.06)	Negative
		Controlled Oral Word Association Test	Verbal function and language skills	Short-term	MD=0.00 (95% CI, -0.66 to 0.66)	Negative
		FACT-Cog (Impairment)	Self-reported cognitive function	Short-term	MD=3.70 (95% CI, -8.20 to 15.60)	Negative
		FACT-Cog (Ability)	Self-reported cognitive function	Short-term	MD=1.40 (95% CI, -3.06 to 5.86)	Negative
		FACT-Cog (Other Comment)	Self-reported cognitive function	Short-term	MD=0.40 (95% CI, -1.57 to 2.37)	Negative
		FACT-Cog (Impact)	Self-reported cognitive function	Short-term	MD=0.10 (95% CI, -4.28 to	Negative

			function		4.48)	
Exercise	Speed-feedback therapy with a bicycle ergometer[43]	Frontal Assessment Battery	Executive function and motor function	Short-term	MD=1.66 (95% CI, 0.84 to 2.48)*	Positive
	Yoga[35]	Symptoms of Stress Inventory (SOSI) Cognition Subscale	Self-reported cognitive function	Short-term	MD=-1.67 (95% CI, -3.66 to 0.32)	Negative
		Profile of Moods Scale (POMS) Concentration Subscale	Self-reported cognitive function	Short-term	MD=-2.50 (95% CI, -4.56 to -0.44)*	Positive

\* $p < 0.05$ ; Positive represents 'favors intervention' and Negative represents 'does not favor intervention'. Abbreviations: BRIEF GEC: Behavior Rating Inventory of Executive Function Global Executive Composite; EORTC-CF: European Organisation for Research and Treatment of Cancer-Cognitive functioning subscale; FACT-Cog: Functional Assessment of Cancer Therapy-Cognitive Function; GEC: Global Executive Composite; MASQ: Multiple Abilities Self-Report Questionnaire. Measurement time point: short-term: less than 3 months.