



Detection and Attribution Model Intercomparison Project (DAMIP)

Nathan P. Gillett¹, Hideo Shiogama², Bernd Funke³, Gabriele Hegerl⁴, Reto Knutti⁵, Katja Matthes⁶, Benjamin D. Santer⁷, Daithi Stone⁸, Claudia Tebaldi⁹

5 ¹Canadian Centre for Climate Modelling and Analysis, Environment and Climate Change Canada, University of Victoria, Victoria, V8W 2Y2, Canada

²National Institute for Environmental Studies, Tsukuba, 305-8506, Japan

³Instituto de Astrofísica de Andalucía, CSIC, Glorieta de la Astronomía s/n, 18008 Granada, Spain

⁴School of GeoSciences, University of Edinburgh, James Hutton Rd, Edinburgh, EH9 3FE, UK

10 ⁵Institute for Atmospheric and Climate Science, ETH Zürich, Universitätstrasse 16, 8092 Zürich, Switzerland

⁶GEOMAR Helmholtz Centre for Ocean Research Kiel, and Christian-Albrechts-Universität zu Kiel, 24105 Kiel, Germany

⁷Lawrence Livermore National Laboratory, Livermore, California 94550, USA

⁸Lawrence Berkeley National Laboratory, Berkeley, California, 94720, USA

⁹Climate and Global Dynamics Laboratory, National Center for Atmospheric Research, Boulder, CO, 80305, USA

15

Correspondence to: N. P. Gillett (nathan.gillett@canada.ca)

Abstract. Detection and attribution (D&A) simulations were important components of CMIP5 and underpinned the climate change detection and attribution assessments of the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. The primary goals of the Detection and Attribution Model Intercomparison Project (DAMIP) are to facilitate improved estimation of the contributions of anthropogenic and natural forcing changes to observed global warming as well as to observed global and regional changes in other climate variables; to contribute to the estimation of how historical emissions have altered and are altering contemporary climate risk; and to facilitate improved observationally-constrained projections of future climate change. D&A studies typically require unforced control simulations and historical simulations including all major anthropogenic and natural forcings. Such simulations will be carried out as part of the DECK and the CMIP6 historical simulation. In addition D&A studies require simulations covering the historical period driven by individual forcings or subsets of forcings only: such simulations are proposed here. Key novel features of the experimental design

20
25



presented here include: new historical simulations of aerosols-only, stratospheric-ozone-only, CO₂-only, solar-only and volcanic-only forcing, facilitating an improved estimation of the climate response to individual forcing; future single forcing experiments, allowing observationally-constrained projections of future climate change; and an experimental design which allows models with and without coupled atmospheric chemistry to be compared on an equal footing.

5

1 Introduction

Research into the detection and attribution (D&A) of climate change is concerned with diagnosing the existence of forced changes in the observed climate record and assessing the roles of various possible contributors to those observed changes (Hegerl et al. 2010). This research is key for our understanding of anthropogenic climate change, as evidenced by a dedicated chapter in every assessment report of the Intergovernmental Panel of Climate Change (IPCC) since the first report published in 1990. Over this time the resources available to this area of research have developed considerably. Together with longer and improved observations of the climate system, D&A research builds upon the analysis of climate model simulations of how the climate should have responded in the presence and absence of various factors which are expected to have affected the climate. D&A analysis specifically compares these retrospective predictions against the available observational record, and thus serves as a comprehensive evaluation of our understanding of how the climate system responds to anthropogenic interference. Confidence in our ability to project future climate change hinges strongly on D&A conclusions.

This paper describes a new coordinated international project to conduct D&A simulations with the next generation of climate models, to be conducted as part of the Sixth Coupled Model Intercomparison Project (CMIP6, Eyring et al. 2016). The most basic sets of simulations required for D&A analysis first became available from a large number of climate models through the Third Coupled Model Intercomparison Project (CMIP3, Meehl et al. 2007). At that time, individual forcing simulations allowing attribution investigations were available from over half a dozen models (Hegerl et al. 2007), but few of these were publicly available. It was not until the Fifth Coupled Model Intercomparison Project (CMIP5, Taylor et al. 2012) that the full suite of simulations needed for an assessment of the role of anthropogenic interference, as well as of greenhouse gas emissions specifically, in observed climate trends was conducted using multiple climate models under a common experimental design and, moreover, made publicly available en masse.

D&A studies using CMIP5 simulations underpinned several key high-level findings of the IPCC Fifth Assessment Report (AR5), including for example the assessment that it is extremely likely that human influence has been the dominant cause of the observed warming since the mid-20th century, and a figure showing estimated contributions of greenhouse gases, aerosols and natural forcings to observed temperature trends included as one of fourteen summary figures in the summary of the IPCC AR5 Synthesis Report (IPCC, 2014). D&A studies have also underpinned attribution assessments across a range of



variables and regions (Bindoff et al., 2013), and have also been used to constrain near-term projected warming (Stott and Jones, 2012; Gillett et al., 2012; Stott et al., 2013; Shiogama et al., 2016), as well as climate system parameters such as Transient Climate Response (TCR) and the Transient Climate Response to Emissions (TCRE) (Allen et al., 2009; Gillett et al., 2013). Hence D&A results remain of key interest and relevance both scientifically and to policymakers. The only
5 simulations targeted uniquely towards D&A included as part of the CMIP5 experimental design were historical simulations with natural forcing changes only, and historical simulations with greenhouse gas changes only, which were used together with historical simulations and pre-industrial control simulations to support these D&A analyses. But research carried out since the design of CMIP5 has highlighted several key research questions which may be addressed by the inclusion of additional simulations in CMIP6.

10 The separate quantification of greenhouse gas and aerosol contributions to observed global warming is important both for understanding past climate change, and, since aerosol forcing is projected to decline while greenhouse gas forcing increases in the future, for constraining projections of future warming. While some earlier studies were able to clearly separate greenhouse gas and aerosol influence on observed temperature changes using individual models (Stott et al., 2006), more recent studies using newer models and a longer period of observations have identified substantial uncertainties in the
15 separate estimation of greenhouse gas and aerosol contributions (Jones et al., 2013; Gillett et al., 2013; Ribes et al., 2015). These larger uncertainties stem at least in part from uncertainties and inter-model differences in the simulated spatio-temporal pattern of response to aerosols (e.g., Ribes et al. 2015; Boucher et al. 2013), which may have been exacerbated by the large amount of sampling variability in estimates of the aerosol response derived from a difference between the historical simulations and historical simulations with greenhouse gases only and natural forcings only (e.g., Ribes et al., 2015).
20 Simulations of the response to historical changes in aerosols alone will allow the calculation of the response to aerosol forcing with less contamination from internal variability, and without conflating the effects of aerosols with those of ozone and land use changes. Moreover, comparisons of these simulations with proposed Radiative Forcing Model Intercomparison Project (RFMIP) simulations will allow the separation of uncertainty in the simulated aerosol response into a component associated with differences in the simulated distribution of aerosols and a component associated with differences in the
25 simulated climate response to a given aerosol distribution. These advances should allow a more robust quantification of the response to aerosol forcings and its uncertainties.

A further key contribution of D&A studies to the findings of the IPCC AR5 was through the use of observationally-constrained climate projections to inform the assessed range of near-term warming (Stott and Jones, 2012; Gillett et al., 2012; Stott et al., 2013; see also Shiogama, et al. 2016). This is achieved by scaling the projected responses to greenhouse
30 gases and aerosols by their respective regression coefficients derived from a regression analysis over the historical period (Allen et al., 2000; Stott and Kettleborough, 2002; Kettleborough et al., 2007). Such analyses rely on both individual forcing simulations covering the historical period, which were included as part of CMIP5, and individual-forcing simulations of the



future, which were not. The latter are required for constraints on projections for scenarios of future emissions in which the relative importance of greenhouse gas to the other the other anthropogenic forcings changes from that in the historical experiments. Such studies carried out to date have relied on simulations provided by a small number of climate models (Bindoff et al. 2013; Shiogama et al. 2016): the inclusion of future simulations with individual forcings in CMIP6 will allow
5 the derivation of more robust observationally-constrained projections based on a broader range of models, as well as facilitating studies of the aerosol contribution to projected future climate change, which is an area of increasing scientific interest (Shiogama et al., 2010a; Shiogama et al., 2010b; Gillett and von Salzen, 2013; Gagné et al., 2015; Rotstayn, 2013; Myhre, et al. 2015).

The experiments proposed as part of DAMIP (Table 1 and Figure 1) will facilitate a number of D&A analyses of
10 anthropogenic and natural forcing influences on historical climate changes. These analyses are expected to address historical changes in temperature, the hydrological cycle, the atmospheric circulation, ocean properties, cryospheric variables, extreme indices and other variables, from global to regional scales (Bindoff et al., 2013). The extension of DAMIP experiments from 2005 in CMIP5 to 2020 with updated climate forcings will support better understanding of the early 21st-century slowdown of climate warming (Meehl et al., 2011; Watanabe et al., 2014; Huber and Knutti, 2014; Schmidt et al., 2014) and improve
15 the signal-to-noise ratio for D&A of changes in high-noise variables such as precipitation (Zhang et al., 2007). These DAMIP experiments are particularly relevant to two of the major CMIP6 science questions. Analysis of individual forcing simulations, and attribution studies of observed changes will help us to understand how the Earth system responds to forcing. And their use to derive observationally-constrained projections will improve our assessments of future climate change. The DAMIP experiments are also very relevant to the WCRP Grand Challenge on extremes: DAMIP simulations will support
20 attribution studies of changes in temperature, hydrological and other extremes and will improve understanding of the drivers of observed changes in extremes, as well as improving our assessment of the present-day probabilities of extremes. This effort will be further facilitated by using output from DAMIP simulations as input to the C20C+ Detection and Attribution Project (Stone and Pall, 2016.) and other efforts using atmosphere-land-only models to sample extreme weather. Attribution studies based on DAMIP output of hydrological changes and cryospheric changes will also address WCRP Grand
25 Challenges on water availability and cryospheric change.

We do not include an analysis plan in this proposal, because the field of detection and attribution is well-established, and past experience indicates that individual groups are able to self-organise and to carry out attribution studies on variables and regions of interest. Numerous attribution studies were carried out using the CMIP5 attribution simulations (Bindoff et al., 2013), and it is expected that this number will only increase for DAMIP. D&A activities are coordinated internationally in
30 part by the International Detection and Attribution Group (Barnett et al., 2005), and the long-standing interest of the IPCC in attribution will likely also prompt analysis. Therefore our scope here is restricted to explaining and justifying the planned DAMIP simulations.



2 Experimental design

There are two possible frameworks for designing climate model experiments for D&A analysis: the "only" approach, in which simulations are driven with changes only in the forcing of interest, while all other forcings are held at pre-industrial values; and the "all-but" approach, in which simulations are driven with changes in all forcings except the forcing of interest.

5 The appropriateness of each approach depends on the scientific question being asked, for instance whether the intention is to understand how the climate system responds to a given factor, or to diagnose the relative contributions of the various forcings to observed climate change. DAMIP will follow a combination of the "only" and "all-but" approaches. For instance, the total contribution of anthropogenic forcing to observed climate change can be diagnosed by taking the difference of the histALL and histNAT experiments, an "all-but" design. However, the response to greenhouse gas forcing
10 should be determined from the histGHG experiment, an "only" design. A major consideration is that the proposed design allows direct comparison against the attribution-relevant simulations of CMIP5. The final design reflects a goal for DAMIP to be relevant for a number of scientific questions, whilst limiting the demand of the experiments to a feasible computational expense.

A common practice has been to assume linear additivity, i.e. that the climate response to the combined forcing is equal to
15 the sum of the responses to the individual forcings (e.g., Rogelj et al. 2012, Knutti and Hegerl 2008,). This assumption appears to hold for certain forcings, magnitudes and variables (e.g., Meehl et al. 2004, Shiogama et al. 2013), but may not hold for others (Schaller et al. 2013, Schaller et al. 2014, Marvel et al. 2015b).

2.1 Tier 1 Experiments

20 DAMIP simulations build on the piControl simulation which forms part of the DECK, and the CMIP6 Historical Simulation (Eyring et al. 2016) on which all DAMIP historical simulations are based. All simulations used in DAMIP are CO₂-concentration driven for ESMs. We request at least 3 ensemble members with different initial conditions for each historical individual forcing experiment, and recommend that modeling groups which cannot afford to do this for all requested runs start by carrying out at least 3-member ensembles of the Tier 1 simulations.

25 In order to maximise the signal-to-noise ratio, and to facilitate the comparison with the most recent climate observations, it is often desirable to include data from the most recent years in D&A analyses. Given that the CMIP6 historical simulation finishes in 2014, we therefore request that modelling groups extend all DAMIP historical simulations to 2020 using the SSP2-4.5 forcings. A similar approach was applied in CMIP5 with D&A simulations extended from 2006 to 2012 using RCP 4.5 forcings. While this approach has the disadvantage that forcings in the final years of the simulations are estimated rather
30 than directly observed, and in the case of CMIP5 there was discussion about whether differences in forcing over the post-2005 period could contribute to differences in simulated and observed temperature trends over the slowdown period (Santer et al., 2013; Huber and Knutti, 2014; Schmidt et al., 2014) in practice, barring a major volcanic eruption such forcings are



unlikely to diverge strongly from reality (see Box 9.2 of Flato et al. 2013). Moreover, as long as the use of projected forcings in the post-2014 period is made clear, as it is here, investigators can make their own informed decision of whether or not to include the post-2014 period in their analyses. SSP2-4.5 was chosen because of its intermediate level of greenhouse gas emissions and its future aerosol and land-use changes which are more representative of a broader range of SSP-based integrated assessment models projections..

Four experiments are requested under Tier One, each of which should cover the 1850-2020 period and comprise at least three simulations with different initial conditions.

CMIP6 historical (1850-2014) & SSP2-4.5 of ScenarioMIP (2015-2020) (histALL): This includes the CMIP6 historical simulations (Eyring et al. 2016), the extension of that simulation through 2020 under the SSP2-4.5 scenario, and the generation of at least two additional (thus three total) members with different initial conditions. For brevity, we call this “CMIP6 historical + SSP2-4.5” experiment as histALL in this paper. Modelling groups should provide the output data under the labels of the CMIP6 historical runs (1850-2014) and the SSP2-4.5 runs of ScenarioMIP (2015-2020).

histNAT: These historical natural-only simulations resemble the histALL simulations but instead are forced with only solar and volcanic forcings from the historical simulations, similarly to the CMIP5 historicalNat experiment. Together with the historical and piControl simulations, such simulations will allow the attribution of observed changes to anthropogenic and natural influences.

histGHG: These historical greenhouse-gas only simulations resemble the histALL simulations but instead are forced by *well-mixed* greenhouse gas changes only, similarly to the CMIP5 historicalGHG experiment. histALL, histNAT and histGHG will allow the attribution of observed climate change to natural, greenhouse gas and other anthropogenic components. Models with interactive chemistry schemes should either turn off the chemistry or use a preindustrial climatology of stratospheric and tropospheric ozone in their radiation schemes. This will ensure that ozone is fixed in all these simulations, and simulated responses in models with and without coupled chemistry are comparable. By comparison, in CMIP5 some models included changes in ozone in their greenhouse-gas only simulations while others did not (Gillett et al., 2013; Jones et al., 2013; Bindoff et al., 2013), making it harder to compare responses between models.

histAER: These historical aerosol-only simulations resemble the histALL simulations but instead are forced by changes in aerosol forcing only. As discussed in the introduction, such simulations should allow the response to aerosols to be better constrained, and physically understood, and may also allow the response to greenhouse gases to be better constrained (e.g., Ribes et al., 2015). Two experimental designs are proposed for anthropogenic aerosol-only runs depending on whether the model concerned includes a complete representation of atmospheric chemistry. For models in which greenhouse gas concentrations do not influence aerosol concentrations, and aerosol concentrations do not influence ozone concentrations, we request historical simulations forced by anthropogenic aerosol concentrations only or aerosol and aerosol precursor



emissions only as in the historical simulation (sulfate, black carbon, organic carbon, ammonia, NO_x and VOCs). For models with interactive atmospheric chemistry in which aerosol and greenhouse-gas concentrations interact, we recommend an alternative experimental design. Changes in well-mixed-GHGs, aerosol precursors and ozone precursors should be prescribed as in the historical simulations. However, in the radiation scheme, the concentrations of well-mixed-GHGs and the ozone climatology from the piControl runs should be used. This procedure will allow the simulation of aerosol burdens and the associated climate influence consistent with the historical simulations, hence allowing output from historical simulations and aerosol-only simulations to be meaningfully compared.

2.2 Tier 2 Experiments

DAMIP Tier 2 Experiments include experiments to enable observationally-constrained projections (Stott and Kettleborough, 2002; Stott and Jones, 2012; Gillett, et al. 2012; Shiogama et al., 2016) and experiments to facilitate the attribution of observed changes to stratospheric ozone changes (Gillett et al., 2013; Lott et al., 2013). Before performing the future simulations of DAMIP, modelling groups are asked to complete at least one SSP2-4.5 simulation of ScenarioMIP up to 2100.

ssp245GHG: This comprises an extension of at least one histGHG simulation to 2100 using the SSP2-4.5 greenhouse gas concentrations. ssp245GHG, ssp245NAT and SSP2-4.5 will allow the simulated future responses to greenhouse gases, natural-forcing and other forcing to be separated and constrained separately based on regression coefficients derived from observations, hence allowing observationally-constrained projections to be derived. As in histGHG, models with interactive chemistry schemes should either run with the chemistry scheme turned off or use a preindustrial climatology of ozone in the radiation scheme.

histSOZ: These simulations resemble the histALL simulations but are forced by changes in stratospheric ozone concentrations only. They will allow an improved characterization of the response to stratospheric ozone changes, which have played an important role in driving circulation changes in the Southern Hemisphere and temperature changes in the stratosphere, as well as facilitating attribution studies of the response to stratospheric ozone change (e.g., Gillett et al., 2013). Such experiments were not included as standard in CMIP5, although a small number of modelling groups carried them out. In models with coupled chemistry, the chemistry scheme should be turned off, and the simulated ensemble mean monthly mean 3D stratospheric ozone concentrations from the historical simulations should be prescribed, since previous studies have indicated that the 3-D structure of ozone trends is an important driver of the tropospheric response (e.g. Waugh et al., 2009, Crook et al., 2008). Tropospheric ozone should be fixed at 3D long-term monthly mean piControl values, with grid cells having an ozone concentration below 100 ppbv in the piControl climatology for a given month classed as tropospheric. In models without coupled chemistry the same stratospheric ozone prescribed in the historical simulation should be prescribed.



ssp245SOZ: These simulations are extensions of the histSOZ simulations to 2100 following the ozone concentrations specified for the SSP2-4.5 scenario. Stratospheric ozone is projected to recover following the successful implementation of the Montreal Protocol and its amendments (WMO, 2014). These simulations will facilitate a robust multi-model assessment of the climate effects of this recovery on Southern Hemisphere climate and stratospheric temperature.

5

2.3 Tier 3 Experiments

DAMIP Tier 3 experiments consist of solar, volcanic and CO₂ individual forcing experiments, an extension of the aerosol-only and natural-only simulation to 2100 and perturbed forcing experiments.

10 **histSOL:** These simulations resemble the histNAT simulations except that histSOL simulations are driven by solar forcing only. The importance of solar forcing in particular for regional climate variability is becoming increasingly evident (Gray et al., 2010; Seppälä et al., 2014). Because of its prominent 11-year cycle, solar variability could offer a degree of predictability for regional climate variability. Foreseeable fluctuations in solar output could help reduce the uncertainty of future regional climate predictions on decadal time scales. However there are still large uncertainties in the atmospheric solar signal and its transfer mechanisms including changes in total and spectral solar irradiance as well as in solar-driven energetic particles. Recent work in addition suggests a lagged response in the North Atlantic European region due to atmosphere-ocean coupling (Gray et al., 2013, Scaife et al., 2013) as well as a synchronization of decadal NAO variability through the solar cycle (Thiéblemont et al., 2015). Recent modeling efforts have made progress in defining the pre-requisites to simulate solar influence on regional climate more realistically but the lessons learned from CMIP5 show that a more systematic analysis of climate models within CMIP6 is required to better understand the differences in model responses to solar forcing (Mitchell et al., 2015; Misios et al., 2015; Hood et al., 2015). In particular the role of solar induced ozone changes and the need to prescribe spectrally resolved solar irradiance variations and therefore the need for a suitable resolution in the model's shortwave radiation scheme is becoming increasingly evident. The proposed histSOL experiment will facilitate the unambiguous characterization of each model's solar signal and particularly separate clearly solar and volcanic effects. Hence it will allow for a more systematic analysis of the differences in model responses.

15
20
25
histVLC: The histVLC simulations resemble the histNAT simulations except that the histVLC simulations are driven by volcanic forcing only. The histVOL experiments will allow the characterisation of and attribution to volcanic influences. Careful validation of a model's volcanic response may inform its use for geoengineering simulations, such as those in GeoMIP (Kravitz et al., 2013). In addition, it will be possible to test linear additivity of the responses to these natural forcings (Gillett et al., 2004; Shiogama et al., 2013) by comparing the sum of the histVLC and histSOL responses with the histNAT response.

histCO2: Historical simulations driven by observed changes in CO₂ concentration only as in histALL. One approach to observationally-constraining the ratio of warming to cumulative CO₂ emissions, a policy-relevant climate metric known as



the Transient Climate Response to Emissions (TCRE), requires an estimate of historical CO₂-attributable warming, but detection and attribution analyses typically only provide an estimate of warming attributable to changes in all well-mixed greenhouse gases (Gillett et al., 2013a). Together with histGHG simulations these simulations would allow the ratio of CO₂-attributable to GHG-attributable warming to be estimated, and hence more robust estimates of TCRE to be obtained. Further, observationally-constrained estimates of the Transient Climate Response (TCR) typically assume a perfect correlation between TCR and warming in histGHG across models, but in fact there is considerable spread in this ratio (Gillett et al., 2013a). These simulations will allow the reasons for this spread to be investigated, and hence for the better characterisation of uncertainties in TCR.

ssp245AER: This involves an extension of at least one of the histAER simulations to 2100 following SSP2-4.5 aerosol concentrations/emissions. Together with SSP2-4.5 and ssp245NAT, this will allow observationally-constrained projections of future climate to be derived based on separating past and future climate change into components associated with natural forcings, aerosols, and other anthropogenic forcings (greenhouse gases including ozone and land-use change). Such an approach may give more accurate estimates of uncertainties than the more usual approach in which climate change is separated into natural, greenhouse gas, and other anthropogenic (aerosols, ozone and land-use change) components (e.g. Ribes et al., 2015). These simulations will also allow the more robust characterisation of the response to future aerosol changes, without conflating these changes with the responses to ozone and land-use changes.

ssp245NAT: This involves an extension of at least one of the histNAT simulations to 2100 following SSP2-4.5 solar and volcanic forcing. The future solar forcing data recommended for CMIP6 has a downward trend (Matthes et al., 2016). ssp245NAT will be useful to investigate effects of this declining solar forcing on future climate change projections. Together with SSP2-4.5 and ssp245GHG, ssp245NAT will allow observationally-constrained projections of future climate to be derived based on separating past and future climate change into components associated with natural forcing, well-mixed greenhouse gases and other anthropogenic forcing factors.

histALLestAER2 and histALLestNAT2: The final two sets of simulations, histALLestAER2 and histALLestNAT2, are identical to the histALL historical simulation (including the extension to 2020 with SSP2-4.5), except that they contain an alternative estimate of the aerosol forcing and natural forcings, respectively. Standard attribution analyses sample over internal variability, and in some cases sample over model uncertainty and observational uncertainty (e.g., Bindoff et al., 2013). However, if these analyses use a multi-model ensemble in which all models use the same set of forcings, then the contribution of forcing uncertainty to the full uncertainty in the results is neglected. Hence we propose simulations with different estimates of historical forcings to explore this source of uncertainty. We focus here on the uncertainties in aerosol emissions/concentrations and natural forcings since these sources of forcing uncertainty are expected to be the most important for global climate. Investigators could for example carry out attribution analyses using the histALLestAER2, histGHG and histNAT simulations to address the contribution of aerosol forcing uncertainty to global attribution results, and similarly use the histALLestNAT2, histGHG and histAER simulations to address the contribution of natural forcing uncertainty to global attribution results. These simulations could also be used to examine the role of forcing uncertainties in



simulating climate trends over particular periods, such as global warming over the slowdown period. The exact method for sampling uncertainty in the tropospheric aerosol, volcanic, and solar forcings is being developed in cooperation with the groups developing forcing data sets for CMIP6.

5 3. Synergies with other MIPs

Synergies between the CMIP6 endorsed MIPs are important to address the science questions of the WCRP grand challenges and CMIP6. Table 2 shows the examples of synergies between DAMIP, the other MIPs and research activities.

The Decadal Climate Prediction Project (DCPP) and DAMIP together propose the enlargement of ensemble size of histALL and SSP2-4.5 to investigate the importance of internal variability in the past and the near future, providing reduced total computing costs if both MIPs are pursued.

The Global Monsoons Modeling Intercomparison Project (GMMIP) has proposed the pacemaker 20th century historical runs (Kosaka and Xie, 2013) to understand the influences of the Interdecadal Pacific Oscillation (IPO) and the Atlantic Multidecadal Oscillation (AMO) on the global monsoons. The combination between GMMIP and DAMIP facilitate assessments of the relative contributions of external forcing factors and internal variability to historical climate change.

Closely collaborating with DAMIP, the Effective Radiative Forcing subproject of RFMIP (RFMIP-ERF) has proposed experiments to estimate effective radiative forcing for all forcings, well-mixed greenhouse gas forcing, natural forcing and anthropogenic aerosol forcing. Combining radiative forcing estimated from RFMIP-ERF and transient climate responses from DAMIP, we can investigate transient climate sensitivities (or forcing efficacies) (Hansen et al., 2005; Yoshimori and Broccoli, 2008; Shindell, 2014) for those forcing agents, which are essential parameters for the observational constraints of TCR and effective climate sensitivity (Shindell, 2014; Kummer and Dessler, 2014; Gregory et al. 2015; Marvel et al., 2015).

LUMIP proposes historical ALL forcing experiments without land-use land-cover changes. Combinations of this and the DAMIP experiments allow the separation of climate responses to the land-use land-cover changes and the other forcing agents.

While VolMIP includes simulations of individual eruptions it does not include simulations of the transient response to historical eruptions, and its focus is on 19th century eruptions. histVLC of DAMIP allows better validation of long-term transient effects against observations,

4. New variables requested by DAMIP

The specific questions addressed by the histSOL experiment, in particular the attribution of differences in the model responses to the solar forcing and its link to different transfer mechanisms, require additional model output related to the radiation scheme and calculated or prescribed ozone chemistry. This includes zonal mean short- and longwave heating rates, as well as ozone fields (prescribed or interactively calculated). Further, a reduced set of new chemistry variables has been proposed for models with interactive chemistry schemes, including O₂ and O₃ photolysis rates, as well as odd oxygen total



loss and production rates. These new variables will be also of interest to investigate and understand the volcanic response in the DAMIP simulations.

5. Summary

5 DAMIP will coordinate the climate model simulations needed to more robustly attribute global and regional climate change to anthropogenic and natural causes, to derive observationally-constrained projections of future climate change, and to improve understanding of the mechanisms by which particular forcings affect climate. The Tier One simulations differ from those included in CMIP5 only by the inclusion of a set of simulations with aerosol changes only, which will help constrain the climate response to aerosol forcing, and also by an experimental design that ensures that results from models with and
10 without coupled chemistry are comparable. Tiers Two and Three include individual forcing simulations covering the future through the end of the century, which are needed to observationally constrain the future response to greenhouse gases, aerosols, stratospheric ozone and natural forcing based on observed historical changes. These tiers also include additional simulations of the response to historical variations in stratospheric ozone, CO₂, volcanoes and solar forcing individually, and perturbed forcing simulations to allow the contribution of forcing uncertainty to uncertainty in attribution results to be
15 assessed for the first time.

Website of DAMIP: Updated details on the project and its progress will be available at <http://damip.lbl.gov>.

Data Availability: The model output from the DAMIP simulations described in this paper will be distributed through the
20 Earth System Grid Federation (ESGF) with digital object identifiers (DOIs) assigned. As in CMIP5, the model output will be freely accessible through data portals after registration. In order to document DAMIP's scientific impact and enable ongoing support of DAMIP, users are obligated to acknowledge CMIP6, DAMIP, the participating modelling groups, and the ESGF centres (see details on the CMIP Panel website at <http://www.wcrp-climate.org/index.php/wgcm-cmip/about-cmip>).

25

References

- Allen, M. R., Stott, P. A., Mitchell, J. F. B., Schnur, R., and Delworth, T. L.: Quantifying the uncertainty in forecasts of anthropogenic climate change, *Nature*, 407, 617-620, 10.1038/35036559, 2000.
- Allen, M. R., Frame, D. J., Huntingford, C., Jones, C. D., Lowe, J. A., Meinshausen, M., and Meinshausen, N.: Warming
30 caused by cumulative carbon emissions towards the trillionth tonne, *Nature*, 458, 1163-1166, 10.1038/nature08019, 2009.
- Barnett, T., Zwiers, F., Hegerl, G., Allen, M., Crowley, T., Gillett, N., Hasselmann, K., Jones, P., Santer, B., Schnur, R., Scott, P., Taylor, K., Tett, S., and Int Ad Hoc Detention, A.: Detecting and attributing external influences on the climate system: A review of recent advances, *Journal of Climate*, 18, 1291-1314, 2005.



- Bindoff, N.L., P.A. Stott, K.M. AchutaRao, M.R. Allen, N. Gillett, D. Gutzler, K. Hansingo, G. Hegerl, Y. Hu, S. Jain, I.I. Mokhov, J. Overland, J. Perlwitz, R. Sebbari and X. Zhang, 2013: Detection and Attribution of Climate Change: from Global to Regional. In: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Boucher, O., D. Randall, P. Artaxo, C. Bretherton, G. Feingold, P. Forster, V.-M. Kerminen, Y. Kondo, H. Liao, U. Lohmann, P. Rasch, S.K. Satheesh, S. Sherwood, B. Stevens and X.Y. Zhang, 2013: Clouds and Aerosols. In: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Crook, J.A., Gillett, N.P. and Keeley, S.P.: Sensitivity of Southern Hemisphere climate to zonal asymmetry in ozone. *Geophysical Research Letters*, 35(7), L07806, 2008.
- Davin, E. L., de Noblet-Ducoudré N., Friedlingstein, P.: Impact of land cover change on surface climate: Relevance of the radiative forcing concept, *Geophys. Res. Lett.*, 34, L13702, doi:10.1029/2007GL029678, 2007.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organisation, *Geosci. Model Dev. Discuss.*, 8, 10539-10583, doi:10.5194/gmdd-8-10539-2015, 2015.
- Flato, G., J. Marotzke, B. Abiodun, P. Braconnot, S.C. Chou, W. Collins, P. Cox, F. Driouech, S. Emori, V. Eyring, C. Forest, P. Gleckler, E. Guilyardi, C. Jakob, V. Kattsov, C. Reason and M. Rummukainen, 2013: Evaluation of Climate Models. In: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Gagné, M. È., Gillett, N. P., and Fyfe, J. C.: Impact of aerosol emission controls on future Arctic sea ice cover, *Geophysical Research Letters*, 42, 8481-8488, 10.1002/2015gl065504, 2015.
- Gillett, N. P., Wehner, M. F., Tett, S. F. B., and Weaver, A. J.: Testing the linearity of the response to combined greenhouse gas and sulfate aerosol forcing, *Geophysical Research Letters*, 31, 10.1029/2004gl020111, 2004.
- Gillett, N. P., Arora, V. K., Flato, G. M., Scinocca, J. F., and von Salzen, K.: Improved constraints on 21st-century warming derived using 160 years of temperature observations, *Geophysical Research Letters*, 39, 10.1029/2011gl050226, 2012.
- Gillett, N. P., Arora, V. K., Matthews, D., and Allen, M. R.: Constraining the Ratio of Global Warming to Cumulative CO₂ Emissions Using CMIP5 Simulations, *Journal of Climate*, 26, 6844-6858, 10.1175/jcli-d-12-00476.1, 2013a.



- Gillett, N. P., Fyfe, J. C., and Parker, D. E.: Attribution of observed sea level pressure trends to greenhouse gas, aerosol, and ozone changes, *Geophysical Research Letters*, 40, 2302-2306, 10.1002/grl.50500, 2013b.
- Gillett, N. P., and Von Salzen, K.: The role of reduced aerosol precursor emissions in driving near-term warming, *Environmental Research Letters*, 8, 10.1088/1748-9326/8/3/034008, 2013.
- 5 Gray, L.J., J. Beer, M. Geller, J.D. Haigh, M. Lockwood, K. Matthes, U. Cubasch, D. Fleitmann, G. Harrison, L. Hood, J. Luterbacher, G. A. Meehl, D. Shindell, B. van Geel, and W. White: Solar Influences on Climate, *Rev. Geophys.*, 48, RG4001, doi:10.1029/2009RG000282, 2010.
- Gray, L.J. et al., A lagged response to the 11 year solar cycle in observed winter Atlantic/European weather patterns, *J. Geophys. Res.*, 118, doi:10.1002/2013JD020062, 2013. Gregory, J. M., Andrews, T., and Good, P.: The inconstancy of the transient climate response parameter under increasing CO₂, *Philosophical Transactions of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 373, 10.1098/rsta.2014.0417, 2015.
- 10 Hansen, J., Sato, M., Ruedy, R., Nazarenko, L., Lacis, A., Schmidt, G. A., Russell, G., Aleinov, I., Bauer, M., Bauer, S., Bell, N., Cairns, B., Canuto, V., Chandler, M., Cheng, Y., Del Genio, A., Faluvegi, G., Fleming, E., Friend, A., Hall, T., Jackman, C., Kelley, M., Kiang, N., Koch, D., Lean, J., Lerner, J., Lo, K., Menon, S., Miller, R., Minnis, P., Novakov, T., Oinas, V., Perlwitz, J., Rind, D., Romanou, A., Shindell, D., Stone, P., Sun, S., Tausnev, N., Thresher, D., Wielicki, B., Wong, T., Yao, M., and Zhang, S.: Efficacy of climate forcings, *J. Geophys. Res.-Atmos.*, 110, 10.1029/2005jd005776, 2005.
- Hood, L., S. Misios, D. Mitchel, L.J. Gray, K. Tourpali, K. Matthes, H. Schmidt, G. Chiodo, R. Thiéblemont, E. Rozanov, D. Shindell, A. Krivolutsky, Solar Signals in CMIP-5 Simulations: The Ozone Response, *Q. J. Roy. Met. Soc.*, DOI 10.1002/qj.2553, 2015.
- 20 Huber, M., and Knutti, R.: Natural variability, radiative forcing and climate response in the recent hiatus reconciled, *Nat. Geosci.*, 7, 651-656, 10.1038/ngeo2228, 2014.
- IPCC, 2014: Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, R.K. Pachauri and L.A. Meyer (eds.)].
- 25 IPCC, Geneva, Switzerland, 151 pp.
- Jones, G. S., Stott, P. A., and Christidis, N.: Attribution of observed historical near-surface temperature variations to anthropogenic and natural causes using CMIP5 simulations, *J. Geophys. Res.-Atmos.*, 118, 4001-4024, 10.1002/jgrd.50239, 2013.
- Kettleborough, J. A., Booth, B. B. B., Stott, P. A., and Allen, M. R.: Estimates of uncertainty in predictions of global mean surface temperature, *Journal of Climate*, 20, 843-855, 10.1175/jcli4012.1, 2007.
- 30 Kosaka, Y., and Xie, S.-P.: Recent global-warming hiatus tied to equatorial Pacific surface cooling. *Nature* 501, 403-407, 2013
- Kravitz, B., Caldeira, K., Boucher, O., Robock, A., Rasch, P. J., Alterskjær, K., Karam, D. B., Cole, J. N. S., Curry, C. L., Haywood, J. M., Irvine, P. J., Ji, D., Jones, A., Kristjánsson, J. E., Lunt, D. J., Moore, J. C., Niemeier, U., Schmidt, H.,



- Schulz, M., Singh, B., Tilmes, S., Watanabe, S., Yang, S., and Yoon, J.-H.: Climate model response from the Geoengineering Model Intercomparison Project (GeoMIP), *Journal of Geophysical Research: Atmospheres*, 118, 8320–8332, 10.1002/jgrd.50646, 2013.
- Kummer, J. R., and Dessler, A. E.: The impact of forcing efficacy on the equilibrium climate sensitivity, *Geophysical Research Letters*, 41, 3565–3568, 10.1002/2014gl060046, 2014.
- Knutti, R., Hegerl, G. C.: The equilibrium sensitivity of the Earth’s temperature to radiation changes, *Nature Geosci.* 1, 735–743, 2008.
- Knutti, R., Rugenstein, M.A.A.: Feedbacks, climate sensitivity and the limits of linear models, *Philosophical Transactions of the Royal Society*, 373, 20150146, doi: 10.1098/rsta.2015.0146, 2015.
- Lott, F. C., Stott, P. A., Mitchell, D. M., Christidis, N., Gillett, N. P., Haimberger, L., Perlwitz, J., and Thorne, P. W.: Models versus radiosondes in the free atmosphere: A new detection and attribution analysis of temperature, *Journal of Geophysical Research: Atmospheres*, 118, 2609–2619, 10.1002/jgrd.50255, 2013.
- Marvel, K., Schmidt, G. A., Miller, R. L., and Nazarenko, L. S.: Implications for climate sensitivity from the response to individual forcings, *Nature Clim. Change*, 10.1038/nclimate2888, 2015a.
- Marvel, K., Schmidt, G.A., Shindell, D., Bonfils, C., LeGrande, A.N., Nazarenko, L., Tsigaridis, K.: Do responses to different anthropogenic forcings add linearly in climate models? *Environ. Res. Lett.*, 10, no. 10, 104010, doi:10.1088/1748-9326/10/10/104010, 2015b.
- Matthes K., B. Funke, M. E. Anderson, L. Barnard, J. Beer, P. Charbonneau, M. A. Clilverd, T. Dudok de Wit, M. Haberreiter, A. Hendry, C. H. Jackman, M. Kretzschmar, T. Kruschke, M. Kunze, U. Langematz, A. Maycock, S. Misios, C. J. Rodger, A. A. Scaife, A. Seppälä, M. Shangguan, M. Sinnhuber, K. Tourpali, I. Usoskin, M. von der Kamp, P. T. Verronen, S. Versick: Solar Forcing for CMIP6, *Geosci. Model Dev.*, to be submitted
- Meehl, G. A., Arblaster, J. M., Fasullo, J. T., Hu, A., and Trenberth, K. E.: Model-based evidence of deep-ocean heat uptake during surface-temperature hiatus periods, *Nature Climate Change*, 1, 360–364, 10.1038/nclimate1229, 2011.
- Meehl, G. A., Washington, W.M., Ammann, C.M., Arblaster, J.M., Wigley, T. M. L., Tebaldi, C. Combinations of natural and anthropogenic forcings in Twentieth-Century Climate. *J. Climate*, 17, 3721–3727, 2004.
- Mitchell, D., S. Misios, L.J. Gray, K. Tourpali, K. Matthes, L. Hood, H. Schmidt, G. Chiodo, R. Thiéblemont, E. Rozanov, D. Shindell, A. Krivolutsky: Solar Signals in CMIP-5 Simulations: The Stratospheric Pathway, *Q. J. Roy. Met. Soc.*, doi:10.1002/qj.2530, 2015.
- Myhre, G., Boucher, O., Breon, F.-M., Forster, P., and Shindell, D.: Declining uncertainty in transient climate response as CO₂ forcing dominates future climate change, *Nat. Geosci.*, 8, 181–185, 10.1038/ngeo2371, 2015.
- Ribes, A., Gillett, N. P., and Zwiers, F. W.: Designing Detection and Attribution Simulations for CMIP6 to Optimize the Estimation of Greenhouse Gas–Induced Warming, *Journal of Climate*, 28, 3435–3438, 10.1175/jcli-d-14-00691.1, 2015.
- Rogelj, J., Meinshausen, M., Knutti, R.: Global warming under old and new scenarios using IPCC climate sensitivity range estimates, *Nature Climate Change*, 2, 248–253, 2012.



- Rotstayn, L. D., Collier, M. A., Chrastansky, A., Jeffrey, S. J., and Luo, J. J.: Projected effects of declining aerosols in RCP4.5: unmasking global warming?, *Atmospheric Chemistry and Physics*, 13, 10883-10905, 10.5194/acp-13-10883-2013, 2013.
- Santer, B. D., Painter, J. F., Bonfils, C., Mears, C. A., Solomon, S., Wigley, T. M. L., Gleckler, P. J., Schmidt, G. A.,
5 Doutriaux, C., Gillett, N. P., Taylor, K. E., Thorne, P. W., and Wentz, F. J.: Human and natural influences on the changing thermal structure of the atmosphere, *Proceedings of the National Academy of Sciences of the United States of America*, 110, 17235-17240, 10.1073/pnas.1305332110, 2013.
- Scaife, A. A., Ineson, S., Knight, J. R., Gray, L. J., Kodera, K., and Smith, D. M.: A mechanism for lagged North Atlantic climate response to solar variability, *Geophys. Res. Lett.*, 40, 1-6 doi:10.1002/grl.50099 195, 2013.
- 10 Schaller, N., Cermak, J., Wild, M., Knutti R.: The sensitivity of the energy budget and water cycle to CO₂ and solar forcing. *Earth System Dynamics*, 4, 253-266, 2013.
- Schaller, N., Sedlacek J., Knutti R.: The asymmetry of the climate system's response to solar forcing changes and its implications for geoengineering scenarios. *J. Geophys. Res.*, 119, 5171–5184, doi:10.1002/2013JD021258, 2014.
- Schmidt, G. A., Shindell, D. T., and Tsigaridis, K.: Reconciling warming trends, *Nat. Geosci.*, 7, 158-160,
15 10.1038/ngeo2105, 2014.
- Seppälä, A., K. Matthes, C. Randall, and I. Mironova: What is the solar influence on climate? – Overview of activities during CAWSES-II, special issue of *Progress in Earth and Planetary Science (PEPS)*, 1:24, doi:10.1186/s40645-014-0024-3, 2014.
- Sherwood, SC., Bony, S., Boucher, O., Bretherton, C., Forster, P.M., Gregory, J.M., Stevens, B.: Adjustments in the
20 Forcing-Feedback Framework for Understanding Climate Change. *Bull. Amer. Meteor. Soc.*, 96, 217–228, 2015.
- Shindell, D. T.: Inhomogeneous forcing and transient climate sensitivity, *Nature Climate Change*, 4, 274-277, 10.1038/nclimate2136, 2014.
- Shiogama, H., Emori, S., Takahashi, K., Nagashima, T., Ogura, T., Nozawa, T., and Takemura, T.: Emission scenario dependency of precipitation on global warming in the MIROC3.2 model, *Journal of Climate*, 23, 2404-2417, 2010a.
- 25 Shiogama, H., Hanasaki, N., Masutomi, Y., Nagashima, T., Ogura, T., Takahashi, K., Hijioaka, Y., Takemura, T., Nozawa, T., and Emori, S.: Emission scenario dependencies in climate change assessments of the hydrological cycle, *Climatic Change*, 99, 321-329, 10.1007/s10584-009-9765-1, 2010b.
- Shiogama, H., Stone, D. A., Nagashima, T., Nozawa, T., and Emori, S.: On the linear additivity of climate forcing-response relationships at global and continental scales, *International Journal of Climatology*, 33, 2542-2550, 2013.
- 30 Shiogama, H., Stone, D., Emori, S., Takahashi, K., Mori, S., Maeda, A., Ishizaki, Y., and Allen, M. R.: Predicting future uncertainty constraints on global warming projections, *Scientific Reports*, 6, 18903, 10.1038/srep18903, 2016.
- Stone, D. A., Pall., P.: A benchmark estimate of the effect of anthropogenic emissions on the ocean surface, submitted.



- Stott, P., Good, P., Jones, G., Gillett, N., and Hawkins, E.: The upper end of climate model temperature projections is inconsistent with past warming, *Environmental Research Letters*, 8, 10.1088/1748-9326/8/1/014024, 2013.
- Stott, P. A., and Kettleborough, J. A.: Origins and estimates of uncertainty in predictions of twenty-first century temperature rise, *Nature*, 416, 723-726, 10.1038/416723a, 2002.
- 5 Stott, P. A., Mitchell, J. F. B., Allen, M. R., Delworth, T. L., Gregory, J. M., Meehl, G. A., and Santer, B. D.: Observational constraints on past attributable warming and predictions of future global warming, *Journal of Climate*, 19, 3055-3069, 10.1175/jcli3802.1, 2006.
- Stott, P. A., and Jones, G. S.: Observed 21st century temperatures further constrain likely rates of future warming, *Atmos. Sci. Lett.*, 13, 151-156, 10.1002/asl.383, 2012.
- 10 Thiéblemont, R., K. Matthes, N. Omrani, K. Kodera, and F. Hansen: Solar forcing synchronizes decadal North Atlantic climate variability, *Nat. Comm.*, doi: 10.1038/ncomms9268, 2015.
- Watanabe, M., Shiogama, H., Tatebe, H., Hayashi, M., Ishii, M., and Kimoto, M.: Contribution of natural decadal variability to global warming acceleration and hiatus, *Nature Climate Change*, 2014.
- Waugh, D.W., Oman, L., Newman, P.A., Stolarski, R.S., Pawson, S., Nielsen, J.E. and Perlwitz, J.: Effect of zonal
15 asymmetries in stratospheric ozone on simulated Southern Hemisphere climate trends. *Geophysical Research Letters*, 36(18), L18701, 2009.
- World Meteorological Organization (WMO), Scientific Assessment of Ozone Depletion: 2014, World Meteorological Organization, Global Ozone Research and Monitoring Project—Report No. 55, 416 pp., Geneva, Switzerland, 2014.
- Yoshimori, M., and Broccoli, A. J.: Equilibrium response of an atmosphere-mixed layer ocean model to different radiative
20 forcing agents: Global and zonal mean response, *Journal of Climate*, 21, 4399-4423, 10.1175/2008jcli2172.1, 2008.
- Zhang, X., Zwiers, F. W., Hegerl, G. C., Lambert, F. H., Gillett, N. P., Solomon, S., Stott, P. A., and Nozawa, T.: Detection of human influence on twentieth-century precipitation trends, *Nature*, 448, 461-U464, 10.1038/nature06025, 2007.




Figure 1. Schematic of the relationships of the various experiments proposed under DAMIP and other MIPs. Solid arrows indicate the decomposition into separated forced responses.

5



Table 1. List of proposed DAMIP experiments

Name of exp.	Description	Tier	Start year	End year*	Min. Ens. size
CMIP6 historical simulation and SSP2-4.5	Enlarging ensemble size of the CMIP6 historical simulations (1850-2014) and the SSP2-4.5 simulations of ScenarioMIP (2015-2020). While these simulations are called histALL in this paper just for readability, please provide the output data separately as the CMIP6 historical simulation (-2014) and the SSP2-4.5 (2015-).	1	1850	2020	3
histNAT	Natural-only historical simulations	1	1850	2020	3
histGHG	Well-mixed greenhouse-gas-only historical simulations	1	1850	2020	3
histAER	Anthropogenic-Aerosol-only historical simulations	1	1850	2020	3
ssp245GHG	Extension of at least one histGHG simulation through the 21st century according to SSP2-4.5 greenhouse gas concentrations	2	2021	2100	1
histSOZ	Stratospheric-Ozone-only historical simulations	2	1850	2020	3
ssp245SOZ	Extension of at least one histSOZ simulation through the 21st century according to SSP2-4.5 stratospheric ozone concentrations	2	2021	2100	1
histSOL	Solar-only historical simulations	3	1850	2020	3
histVLC	Volcanic-only historical simulations	3	1850	2020	3
histCO2	CO ₂ -only historical simulations	3	1850	2020	3
ssp245AER	Extension of at least one histAER simulation through the 21st century according to SSP2-4.5 tropospheric aerosols concentrations/emissions	3	2021	2100	1
ssp245NAT	Extension of at least one histNAT simulation through the 21st century according to SSP2-4.5 solar and volcanic forcing	3	2021	2100	1
histALLestAER2	histALL with alternate estimates of anthropogenic aerosol emissions/concentrations	3	1850	2020	3
histALLestNAT2	histALL with alternate estimates of solar and volcanic forcing	3	1850	2020	3

*2015-2100 segments of the simulations are driven by the SSP2-4.5 emission scenario.

**Table 2.** Examples of synergies with DECK, other MIPs and research activities

MIP or Project	Simulations in MIP	Simulations in DAMIP	Area of synergy
DECK	piControl, 1petCO2	All	piControl is essential for estimating internal variability. Thus we recommend modelling groups to perform a 500-year or longer piControl run to allow robust estimates of internal variability. 1petCO2 is needed for observationally-constrained estimates of TCR and TCRE.
CMIP6 historical simulations	All	histALL, histNAT, histGHG, histAER, histSOZ, histVLC, histSOL, histCO2	All the historical experiments of DAMIP are based on the CMIP6 historical simulations and allow refined understanding of the contribution of individual forcing components to climate variations and change in the CMIP6 historical simulations..
ScenarioMIP	SSP2-4.5	histALL, histNAT, histGHG, histAER, ssp245GHG, ssp245AER, ssp245NAT	Allow observational-constraints of uncertainties in future projections.
AerChemMIP	HISTghg, HISTghgNtcf, WMFORCch4, WMFORCn2o	histALL, histGHG, histSOZ	HISTghg (AerChemMIP) is the same as histALL (DAMIP) but with 1850 aerosol and tropospheric ozone precursors. In HISTghgNtcf, only the aerosol precursors are kept at 1850, the ozone precursors follow the historical. WMFORCch4 and WMFORCn2o experiments should also include the chemical effects on strat+trop ozone and strat H2O.
DCPP	Historical+SSP2-4.5	histALL	DCPP proposes a 10-member ensemble of histALL up to 2030 also extended with SSP2-4.5.
GMMIP	AMIP20C, HIST-IPO, HIST-AMO	histALL, histNAT, histGHG, histAER	The combinations of DAMIP and GMMIP allow the assessments of the relative contributions of external forcing factors and the internal variability on the historical climate change.
LUMIP	LND_noLULCC_hist	histALL	"ALL minus land-use (LND_noLULCC_hist)" of LUMIP and histALL allow the separation of effects of land-use changes and the others.
RFMIP-ERF	Historical+SSP2-4.5, Natural, Aerosols, WMGHG	histALL, histNAT, histGHG, histAER, ssp245GHG, ssp245AER, ssp245NAT	Combining radiative forcing estimated from RFMIP-ERF and transient climate responses from DAMIP, we can investigate how feedbacks and adjustments vary with forcing factors.
RFMIP-Historical	Historical, Hist-Nat, Hist-Aer	histALL, histNAT, histAER	Combinations of DAMIP and RFMIP-Historical will allow us to separate uncertainties in climate response based on specified aerosol evolution from the overall uncertainties in climate response to specified aerosol precursor emissions.
GeoMIP & VolMIP	All	histVLC	The volcanic response of models can be validated against observations using histVLC, whereas GeoMIP experiments cannot. Thus histVLC experiments will provide useful context for interpreting simulated responses to stratospheric aerosol across models in the GeoMIP experiment. While VolMIP includes simulations of individual eruptions, it does not include simulations of the transient response to historical eruptions and its focus is on 19th century eruptions. histVLC facilitates validation of long-term transient effects against observations.