

# Neutrosophic Sets and Systems

---

Volume 33

Article 6

---

5-1-2020

## Systems of Neutrosophic Linear Equations

S. A. Edalatpanah

Follow this and additional works at: [https://digitalrepository.unm.edu/nss\\_journal](https://digitalrepository.unm.edu/nss_journal)

---

### Recommended Citation

Edalatpanah, S. A.. "Systems of Neutrosophic Linear Equations." *Neutrosophic Sets and Systems* 33, 1 (2020). [https://digitalrepository.unm.edu/nss\\_journal/vol33/iss1/6](https://digitalrepository.unm.edu/nss_journal/vol33/iss1/6)

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in Neutrosophic Sets and Systems by an authorized editor of UNM Digital Repository. For more information, please contact [amywinter@unm.edu](mailto:amywinter@unm.edu), [lsloane@salud.unm.edu](mailto:lsloane@salud.unm.edu), [sarahrk@unm.edu](mailto:sarahrk@unm.edu).



# Systems of Neutrosophic Linear Equations

S. A. Edalatpanah <sup>1,\*</sup>

<sup>1</sup> Department of Research Center, Ayandegan Institute of Higher Education, Tonekabon, Iran;

\* Correspondence: saedalatpanah@aihe.ac.ir; Tel.: (+98)11 5431 04 29

**Abstract:** In the present paper, for first time, a System of Neutrosophic Linear Equations (SNLE) is investigated based on the embedding approach. To this end, the  $(\alpha, \beta, \gamma)$ -cut is used for transformation of SNLE into a crisp linear system. Furthermore, the existence of a neutrosophic solution to  $n \times n$  linear system is proved in details and a computational procedure for solving the SNLE is designed. Finally, numerical experiments are presented to show the reliability and efficiency of the method.

**Keywords:** Neutrosophic set; Neutrosophic number; Neutrosophic linear equation; Neutrosophic linear system; Embedding method.

## 1. Introduction

A system of linear equations can be defined as:

$$Ax=b, \quad (1)$$

Various equations in the field of scientific modeling that describe the realistic issues like engineering problems and natural phenomena such as differential equations, computational fluid, circuit simulation, cryptography, quantum and structural mechanics, MRI reconstructions, vibroacoustics, linear and non-linear optimization, portfolios, economic modeling, astrophysics, Google page rank, image processing, nano-technology, natural language processing, deep learning, etc., must be solved mathematically. These issues can regularly be diminished to solving of linear systems. There are a huge amount of models to solve this problem, for more details, see [1-15] and the references therein.

Nevertheless, if the assessment of the coefficients of systems is uncertain and imprecise and just some ambiguous understanding regarding the real values of the parameters is accessible, it might be advantageous to characterize them with special numbers related to soft computing. Fuzzy set was introduced by Zadeh [16, 17], as a suitable instrument to express uncertainty in real life situation. After the introduction of fuzzy set, numerous scholars deliberate on this topic (information of some studies can be observed in [18-23]).

Numerous researchers also suggested several strategies to solve linear systems under fuzzy situation. Fuzzy linear systems emerged at least until 1980 [24]; however Friedman et al. [25] launched a particular model to solve a fuzzy linear system where, the matrix coefficient is crisp and the right-hand hand vector is a fuzzy number. Their model later modified by some other scholars; see [26-46].

However, when there is not clarity in information then the measure of non-membership is not the complement of the measure of membership. In these cases, individual measure of membership and non-membership are needed. Keeping this type of situation in consideration, intuitionistic fuzzy set (IFS) was established by Atanassov [47]. Nevertheless, in different branches of sciences and engineering, it was found that two mentioned components are not sufficient to represent some special types of information. In such cases, a component namely 'neutrality' is needed to represent the

information completely. Thus, to remove the limitation of IFS and to handle with more possible types of uncertainty in practical situation, Smarandache [48-51] initiated neutrosophic set (NS) as an extension of the classical and all types of fuzzy sets.

This concept divided into two category of the neutrosophic numbers (NNs) and the neutrosophic sets (NSs). The neutrosophic number (NN) introduce a concept of indeterminacy, denoted by  $A = m + nI$  ( $m, n \in R$ ), consists of its determinate part  $m$  and its indeterminate part  $nI$ . In the worst scenario,  $A$  can be unknown, *i.e.*,  $A = nI$ . However, when there is no indeterminacy related to  $A$ , in the best scenario, there is only its determinate part *i.e.*,  $A = m$  [50, 51]. But, the neutrosophic sets (NSs) represented by a truth-membership degree, an indeterminacy-membership degree, a falsity-membership degree and have some subclasses such as interval neutrosophic set [52-54], bipolar neutrosophic set [55-57], single-valued neutrosophic set [58-66], multi-valued neutrosophic set [67-98], and neutrosophic linguistic set [69-70] and applied to solve various problems; see [71-78]. It is worth mentioning that NSs and NNs are two different branches in neutrosophic theory and indicate different forms and concepts of information.

Like any other framework, system of linear equations has also been the topic of evolution. One of the important developments in this field related to situations that coefficients are defined under conditions of uncertainty and indeterminacy. In fact, one of the expectations of classic linear systems is their crispness of data. However, in circumstances where uncertainty and indeterminacy is an inevitable feature of a real life environment, the assumption of crispness of data seems questionable. Also, there is a lot of ambiguity, indeterminacy, and uncertainty in these problems. The system of linear equations under neutrosophic environment are more useful than crisp and other fuzzy linear systems because user in his/her formulation of the problem is not forced to make a delicate formulation. The use of system of Neutrosophic linear equations (SNLE) is recommended to avert unrealistic modeling. Though there are numerous methodologies to solve various issues under NSs and also some models presented to solve linear systems with NNs [79-80], but to the best of our knowledge, the SNLE has not been discussed sets until now. Therefore, the contributions of this study are as follow:

- (i) We present for first time, the system of Neutrosophic Linear Equations (SNLE) problem.
- (ii) Based on the  $(\alpha, \beta, \gamma)$ -cut, we design a strategy for solving SNLE with the single valued neutrosophic numbers (SVNNs).
- (iii) Some theorems about SNLE are investigated and the conditions of a strong neutrosophic solution to  $n \times n$  system of linear equations is proved in details.

This study prearranged as follows: some fundamental information, notions and operations on SVNNs are announced in Section 2. In Section 3, we introduce the SNLE and propose a general model to solve it. To show the efficiency and reliability of the method, numerical tests are provided in Section 4. Lastly, conclusions are offered in Section 5.

## 2. Some Basic Definitions and Arithmetic Operations

Here, we have deliberated some fundamental definitions regarding the neutrosophic sets and single-valued neutrosophic numbers.

**Definition 1** [48-49]. A neutrosophic set  $A$  in objects  $X$  is described by a truth-membership function  $T_A(x)$ , an indeterminacy-membership function  $I_A(x)$ , and a falsity-membership function  $F_A(x)$  where,  $T_A(x): X \rightarrow ]0^-, 1^+[$ ,  $I_A(x): X \rightarrow ]0^-, 1^+[$ , and  $F_A(x): X \rightarrow ]0^-, 1^+[$ , and

$$0^- \leq \sup T_A(x) + \sup I_A(x) + \sup F_A(x) \leq 3^+.$$

**Definition 2** [58]. When three membership functions of neutrosophic set A be singleton subsets in the real standard [0, 1], we have a single-valued neutrosophic set (SVNS) A that is denoted by

$$A = \{ (x, T_A(x), I_A(x), F_A(x)) \mid x \in X \}.$$

**Definition 3** [59]. A single valued triangular neutrosophic number (SVTrN-number) is denoted by  $A^{\mathbb{N}} = \langle (a, b, c), (\mu, \nu, \omega) \rangle$  whose its three membership functions are given as follows:

$$T_{A^{\mathbb{N}}}(x) = \begin{cases} \frac{(x-a)}{(b-a)}\mu & a \leq x < b, \\ \mu & x = b, \\ \frac{(c-x)}{(c-b)}\mu & b \leq x < c, \\ 0 & \text{otherwise.} \end{cases} \quad I_{A^{\mathbb{N}}}(x) = \begin{cases} \frac{(b-x)}{(b-a)}\nu & a \leq x < b, \\ \nu & x = b, \\ \frac{(x-c)}{(c-b)}\nu & b \leq x < c, \\ 1 & \text{otherwise.} \end{cases} \quad F_{A^{\mathbb{N}}}(x) = \begin{cases} \frac{(b-x)}{(b-a)}\omega & a \leq x < b, \\ \omega & x = b, \\ \frac{(x-c)}{(c-b)}\omega & b \leq x < c, \\ 1 & \text{otherwise.} \end{cases}$$

**Definition 3** [59]. Let  $A_1^{\mathbb{N}} = \langle (a_1, b_1, c_1), (\mu_1, \nu_1, \omega_1) \rangle$  and  $A_2^{\mathbb{N}} = \langle (a_2, b_2, c_2), (\mu_2, \nu_2, \omega_2) \rangle$  be two SVTrN-numbers. Then the arithmetic relations are defined as:

$$(i) A_1^{\mathbb{N}} \oplus A_2^{\mathbb{N}} = \langle (a_1 + a_2, b_1 + b_2, c_1 + c_2), (\mu_1 \wedge \mu_2, \nu_1 \vee \nu_2, \omega_1 \vee \omega_2) \rangle \tag{2}$$

$$(ii) \lambda A_1^{\mathbb{N}} = \begin{cases} \langle (\lambda a_1, \lambda b_1, \lambda c_1), (\mu_1, \nu_1, \omega_1) \rangle, & \text{if } \lambda > 0 \\ \langle (\lambda c_1, \lambda b_1, \lambda a_1), (\mu_1, \nu_1, \omega_1) \rangle, & \text{if } \lambda < 0 \end{cases} \tag{3}$$

**Definition 4** [59]. The  $(\alpha, \beta, \gamma)$ -cut Neutrosophic set F is denoted by  $F_{(\alpha, \beta, \gamma)}$ , where  $\alpha, \beta, \gamma \in [0, 1]$  and are fixed numbers such that  $\alpha + \beta + \gamma \leq 3$  is defined as by  $F_{(\alpha, \beta, \gamma)} = \{ \langle T_A(x), I_A(x), F_A(x) \rangle : x \in X, T_A(x) \geq \alpha, I_A(x) \leq \beta, F_A(x) \leq \gamma \}$ .

Also, If  $A^{\mathbb{N}} = \langle (a, b, c), (\mu, \nu, \omega) \rangle$  then  $(\alpha, \beta, \gamma)$ -cut is given by:

$$A^{\mathbb{N}}_{(\alpha, \beta, \gamma)} = \left\langle \begin{matrix} [(a + \alpha(b - a))\mu, (c - \alpha(c - b))\mu], \\ [(b - \beta(b - a))\nu, (b + \beta(c - b))\nu], \\ [(b - \gamma(b - a))\omega, (b + \gamma(c - b))\omega] \end{matrix} \right\rangle \tag{4}$$

### 3. System of Neutrosophic Linear Equations (SNLE)

Consider the  $n \times n$  linear system with the following equations:

$$\begin{cases} a_{11}x_1^{\mathbb{N}} + a_{12}x_2^{\mathbb{N}} + \dots + a_{1n}x_n^{\mathbb{N}} = b_1^{\mathbb{N}}, \\ a_{21}x_1^{\mathbb{N}} + a_{22}x_2^{\mathbb{N}} + \dots + a_{2n}x_n^{\mathbb{N}} = b_2^{\mathbb{N}}, \\ \cdot \\ \cdot \\ \cdot \\ a_{n1}x_1^{\mathbb{N}} + a_{n2}x_2^{\mathbb{N}} + \dots + a_{nn}x_n^{\mathbb{N}} = b_n^{\mathbb{N}}. \end{cases} \tag{5}$$

The matrix form of the Eq.(5) is as follows:

$$Ax^{\mathbb{N}} = b^{\mathbb{N}}, \tag{6}$$

where, the coefficient matrix  $A = (a_{ij})$  is a crisp  $n \times n$  matrix and  $b_i^{\mathbb{N}}, i = 1, 2, \dots, n$  is a neutrosophic number. The Eq.(6) is called a system of neutrosophic linear equations (SNLE).

Let the solution of the SNLE of Eq.(6) be  $x^{\mathbb{N}}$  and its  $(\alpha, \beta, \gamma)$ -cut be  $x_{(\alpha, \beta, \gamma)}^{\mathbb{N}} = ([\underline{x}^T(\alpha), \bar{x}^T(\alpha)], [\underline{x}^I(\beta), \bar{x}^I(\beta)], [\underline{x}^F(\gamma), \bar{x}^F(\gamma)])$ . If the  $(\alpha, \beta, \gamma)$ -cut of  $b^{\mathbb{N}}$  be  $x_{(\alpha, \beta, \gamma)}^{\mathbb{N}} = ([\underline{b}^T(\alpha), \bar{b}^T(\alpha)], [\underline{b}^I(\beta), \bar{b}^I(\beta)], [\underline{b}^F(\gamma), \bar{b}^F(\gamma)])$ , then The SNLE of (6) can be written as:

$$\begin{cases} \sum_{j=1}^n a_{ij} x_j^T(\alpha) = \sum_{j=1}^n a_{ij} x_j^T(\alpha) = \underline{b}_i^T(\alpha), \\ \sum_{j=1}^n a_{ij} x_j^T(\alpha) = \sum_{j=1}^n a_{ij} x_j^T(\alpha) = \bar{b}_i^T(\alpha), \\ \sum_{j=1}^n a_{ij} x_j^I(\beta) = \sum_{j=1}^n a_{ij} x_j^I(\beta) = \underline{b}_i^I(\beta), \\ \sum_{j=1}^n a_{ij} x_j^I(\beta) = \sum_{j=1}^n a_{ij} x_j^I(\beta) = \bar{b}_i^I(\beta), \\ \sum_{j=1}^n a_{ij} x_j^F(\gamma) = \sum_{j=1}^n a_{ij} x_j^F(\gamma) = \underline{b}_i^F(\gamma), \\ \sum_{j=1}^n a_{ij} x_j^F(\gamma) = \sum_{j=1}^n a_{ij} x_j^F(\gamma) = \bar{b}_i^F(\gamma). \end{cases} \tag{7}$$

If we define  $x_i^{\mathbb{N}} = (\underline{x}_1^T, \dots, \underline{x}_n^T, \bar{x}_1^T, \dots, \bar{x}_n^T, \underline{x}_1^I, \dots, \underline{x}_n^I, \bar{x}_1^I, \dots, \bar{x}_n^I, \underline{x}_1^F, \dots, \underline{x}_n^F, \bar{x}_1^F, \dots, \bar{x}_n^F)^T$  and  $b_i^{\mathbb{N}} = (\underline{b}_1^T, \dots, \underline{b}_n^T, \bar{b}_1^T, \dots, \bar{b}_n^T, \underline{b}_1^I, \dots, \underline{b}_n^I, \bar{b}_1^I, \dots, \bar{b}_n^I, \underline{b}_1^F, \dots, \underline{b}_n^F, \bar{b}_1^F, \dots, \bar{b}_n^F)^T$ , then following Friedman et al., (1998) we must solve an  $6n \times 6n$  crisp linear system as:

$$HX=B \tag{8}$$

Where,

$$H = \begin{bmatrix} D_{2n \times 2n} & [0]_{2n \times 2n} & [0]_{2n \times 2n} \\ [0]_{2n \times 2n} & D_{2n \times 2n} & [0]_{2n \times 2n} \\ [0]_{2n \times 2n} & [0]_{2n \times 2n} & D_{2n \times 2n} \end{bmatrix}, B = \begin{bmatrix} B^T \\ B^I \\ B^F \end{bmatrix}. \tag{9}$$

Also  $D = (d_{ij})$ , and obtain as follows:

$$\begin{cases} a_{ij} \geq 0 \rightarrow d_{ij} = a_{ij}, & d_{i+n,j+n} = a_{ij}, \\ a_{ij} < 0 \rightarrow d_{i,j+n} = -a_{ij}, & d_{i+n,j} = -a_{ij} \end{cases} \tag{10}$$

and any  $d_{ij}$  which is not determined by (10) is zero. Also:

$$D = \begin{bmatrix} D_1 & -D_2 \\ -D_2 & D_1 \end{bmatrix}, B^T = \begin{bmatrix} \underline{b}^T \\ \bar{b}^T \end{bmatrix}, B^I = \begin{bmatrix} \underline{b}^I \\ \bar{b}^I \end{bmatrix}, B^F = \begin{bmatrix} \underline{b}^F \\ \bar{b}^F \end{bmatrix}.$$

Where,  $D_1, D_2 \geq 0, D = D_1 - D_2$ .

Since H is a block diagonal matrix, to reduce the computational complexity, we need only to solve the following  $2n \times 2n$  crisp linear systems:

$$Dx^i = B^i, \quad i = T, I, F. \tag{11}$$

Worthy mentioning that the matrix  $D$  may be singular even if  $A$  is nonsingular; see the following example:

**Example 1.** The matrix  $A = \begin{bmatrix} 1 & -2 \\ 1 & 1 \end{bmatrix}$  of the SNLE is nonsingular, while  $D = \begin{bmatrix} 1 & 0 & 0 & -2 \\ 1 & 1 & 0 & 0 \\ 0 & -2 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{bmatrix}$  is

singular.

In other sense, a SNLE represented by a nonsingular matrix  $A$  may be have no solution or an infinite number of solutions. Next, following the Friedman et al., (1998), we study some theorems regarding the properties of  $D$ .

**Theorem 1.**  $D$  is nonsingular iff  $A = D_1 + D_2$  and  $D_1 - D_2$  are nonsingular.

**Theorem 2.** If  $D^{-1}$  exists it must have the same structure as  $D$ , i.e,

$$D^{-1} = \begin{pmatrix} E & F \\ F & E \end{pmatrix}$$

**Definition 5.** Let  $x_i^{\mathcal{N}} = (\underline{x}_1^T, \dots, \underline{x}_n^T, \bar{x}_1^T, \dots, \bar{x}_n^T, \underline{x}_1^I, \dots, \underline{x}_n^I, \bar{x}_1^I, \dots, \bar{x}_n^I, \underline{x}_1^F, \dots, \underline{x}_n^F, \bar{x}_1^F, \dots, \bar{x}_n^F)^T$

be the unique solution of Eq.(5). If  $\forall k \in \{1, 2, \dots, n\}$ :  $\underline{x}_k^T \leq \bar{x}_k^T, \underline{x}_k^I \leq \bar{x}_k^I$  and  $\underline{x}_k^F \leq \bar{x}_k^F$ , then the

solution  $x_i^{\mathcal{N}}$  is called a strong neutrosophic solution. Otherwise, it is a weak neutrosophic solution.

**Theorem 3.** Assume that  $D = \begin{pmatrix} D_1 & D_2 \\ D_2 & D_1 \end{pmatrix}$  be a nonsingular matrix. Then Eq.(5) has a strong solution

if and only if:

$$(D_1 - D_2)^{-1}(\underline{b}^i - \bar{b}^i) \leq 0, \quad i = T, I, F. \tag{12}$$

**Proof.** From the system (11) we obtain:

$$\begin{pmatrix} D_1 & D_2 \\ D_2 & D_1 \end{pmatrix} \begin{pmatrix} \underline{x}^i \\ \bar{x}^i \end{pmatrix} = \begin{pmatrix} \underline{b}^i \\ \bar{b}^i \end{pmatrix}, \quad i = T, I, F$$

Hence,

$$D_1 \underline{x}^i - D_2 \bar{x}^i = \underline{b}^i \tag{13}$$

$$-D_2 \underline{x}^i + D_1 \bar{x}^i = \bar{b}^i \tag{14}$$

From (13) and (14) we have:

$$\begin{cases} (D_1 + D_2)\underline{x}^i - (D_1 + D_2)\bar{x}^i = \underline{b}^i - \bar{b}^i, \\ (D_1 + D_2)(\underline{x} - \bar{x}) = \underline{b}^i - \bar{b}^i. \end{cases}$$

From Theorem 1,  $D_1 - D_2$  is nonsingular. So,

$$(\underline{x}^i - \bar{x}^i) = (D_1 - D_2)^{-1}(\underline{b}^i - \bar{b}^i) \tag{15}$$

By the Definition 5,  $\underline{x}^i - \bar{x}^i \leq 0$  if Eq. (5) has a strong solution. Henceforth (12) holds. Conversely,

if (12) holds, by Eq.(15), we have  $\underline{x}^i - \bar{x}^i \leq 0$ . □

From the theorems 1 and 3, we conclude this result:

**Theorem 4.** The SNLE has a strong solution if and only if the following conditions hold:

1. The matrices  $A = D_1 + D_2$  and  $D_1 - D_2$  are both nonsingular.
2.  $(D_1 - D_2)^{-1}(\underline{b}^i - \bar{b}^i) \leq 0$ .

#### 4. Numerical Example

Here, we provide an experiment to demonstrate the consequences gained in former sections.

**Example 2.** Consider the following SNLE:

$$\begin{cases} x_1^N - x_2^N = \langle (0,1,2); (0.9,0.4,0.2) \rangle, \\ x_1^N + 3x_2^N = \langle (4,5,7); (0.8,0.3,0.3) \rangle. \end{cases} \tag{16}$$

The extended  $4 \times 4$  matrix is

$$D = \begin{bmatrix} 1 & 0 & 0 & -1 \\ 1 & 3 & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 0 & 1 & 3 \end{bmatrix}.$$

Since the matrices  $A = D_1 + D_2$  and  $D_1 - D_2 = \begin{bmatrix} 1 & 1 \\ 1 & 3 \end{bmatrix}$  are both nonsingular, then by Theorem1,

it is easy to see that the matrix  $D$  is nonsingular. Therefore,  $D^{-1}$  exists and based on Theorem 2, it must have the same structure as  $D$ . If we obtain this inverse, we can see that the Theorem 2 is true:

$$D^{-1} = \begin{bmatrix} \frac{9}{8} & \frac{-1}{8} & \frac{-3}{8} & \frac{3}{8} \\ \frac{-3}{8} & \frac{3}{8} & \frac{1}{8} & \frac{-1}{8} \\ \frac{-3}{8} & \frac{3}{8} & \frac{9}{8} & \frac{-1}{8} \\ \frac{1}{8} & \frac{-1}{8} & \frac{-3}{8} & \frac{3}{8} \end{bmatrix}.$$

Now, we obtain the  $(\alpha, \beta, \gamma)$ -cut of the right hand side vector. By Definition 4, we get:

$$b_{1(\alpha, \beta, \gamma)}^N = \langle [0.9(\alpha), 0.9(2 - \alpha)], [0.4(1 - \beta), 0.4(1 + \beta)], [0.2(1 - \gamma), 0.2(1 + \gamma)] \rangle,$$

$$b_{2(\alpha, \beta, \gamma)}^N = \langle [0.8(4 + \alpha), 0.8(8 - 2\alpha)], [0.3(5 - \beta), 0.3(5 + 2\beta)], [0.3(5 - \gamma), 0.3(5 + 2\gamma)] \rangle.$$

Also,  $(D_1 + D_2)^{-1} = \begin{bmatrix} \frac{3}{2} & \frac{-1}{2} \\ \frac{-1}{2} & \frac{1}{2} \end{bmatrix}.$

So:

$$(D_1 + D_2)^{-1}(\underline{b}^T - \bar{b}^T) = \begin{bmatrix} \frac{3}{2} & \frac{-1}{2} \\ \frac{-1}{2} & \frac{1}{2} \end{bmatrix} \begin{bmatrix} \frac{9}{5}(\alpha - 1) \\ \frac{12}{5}(\alpha - 1) \end{bmatrix} = \begin{bmatrix} \frac{3}{2}(\alpha - 1) \\ \frac{3}{10}(\alpha - 1) \end{bmatrix} \leq \begin{bmatrix} 0 \\ 0 \end{bmatrix},$$

$$(D_1 + D_2)^{-1}(\underline{b}^I - \bar{b}^I) = \begin{bmatrix} \frac{3}{2} & \frac{-1}{2} \\ \frac{-1}{2} & \frac{1}{2} \end{bmatrix} \begin{bmatrix} -0.8\beta \\ -0.9\beta \end{bmatrix} = \begin{bmatrix} \frac{-3}{4}\beta \\ \frac{-1}{20}\beta \end{bmatrix} \leq \begin{bmatrix} 0 \\ 0 \end{bmatrix},$$

$$(D_1 + D_2)^{-1}(\underline{b}^F - \bar{b}^F) = \begin{bmatrix} \frac{3}{2} & \frac{-1}{2} \\ \frac{-1}{2} & \frac{1}{2} \end{bmatrix} \begin{bmatrix} -0.4\gamma \\ -0.9\gamma \end{bmatrix} = \begin{bmatrix} \frac{-3}{20}\gamma \\ \frac{-1}{4}\gamma \end{bmatrix} \leq \begin{bmatrix} 0 \\ 0 \end{bmatrix},$$



Therefore, by theorems 3 and 4, The SNLE (16) should has a strong solution. To obtain this solution, form Eq.(11) we have:

$$x^T = \begin{bmatrix} x_1^T \\ x_2^T \\ \bar{x}_1^T \\ \bar{x}_2^T \end{bmatrix} = D^{-1}B^T = \left[ \frac{1}{40}(26\alpha + 41), \frac{1}{40}(2\alpha + 29), \frac{1}{40}(-34\alpha + 101), \frac{1}{40}(-10\alpha - 41) \right]^t,$$

$$x^I = \begin{bmatrix} x_1^I \\ x_2^I \\ \bar{x}_1^I \\ \bar{x}_2^I \end{bmatrix} = D^{-1}B^I = \left[ \frac{-27}{80}(\beta - 2), \frac{1}{80}(\beta + 22), \frac{3}{80}(11\beta + 18), \frac{1}{80}(5\beta - 22) \right]^t,$$

$$x^F = \begin{bmatrix} x_1^F \\ x_2^F \\ \bar{x}_1^F \\ \bar{x}_2^F \end{bmatrix} = D^{-1}B^F = \left[ \frac{-3}{80}(\gamma - 14), \frac{-1}{80}(7\gamma + 26), \frac{3}{80}(3\gamma + 14), \frac{13}{80}(\gamma + 2) \right]^t.$$

$$x_{1(\alpha,\beta,\gamma)}^N = \langle \left[ \frac{1}{40}(26\alpha + 41), \frac{1}{40}(-34\alpha + 101) \right], \left[ \frac{-27}{80}(\beta - 2), \frac{3}{80}(11\beta + 18) \right], \left[ \frac{-3}{80}(\gamma - 14), \frac{3}{80}(3\gamma + 14) \right] \rangle,$$

$$x_{2(\alpha,\beta,\gamma)}^N = \langle \left[ \frac{1}{40}(2\alpha + 29), \frac{1}{40}(-10\alpha - 41) \right], \left[ \frac{1}{80}(\beta + 22), \frac{1}{80}(5\beta - 22) \right], \left[ \frac{-1}{80}(7\gamma + 26), \frac{13}{80}(\gamma + 2) \right] \rangle.$$

For different values of  $0 \leq \alpha, \beta, \gamma \leq 1$ , the graphical interpretation of the above results is shown in figures 1 and 2.




Figure 1. The value of  $x_1^N$ .




Figure 2. The value of  $x_2^N$ .

## 5. Conclusions

In this study, we present for first time, the system of Neutrosophic Linear Equations (SNLE) and establish a general model to solve it. Some theorems about SNLE are investigated and the conditions of a strong neutrosophic solution to  $n \times n$  system of linear equations is proved in details. Finally, from numerical and theoretical studies it can be concluded that the model is efficient and convenient.

**Funding:** This research received no external funding.

**Acknowledgments:** The author is most grateful to all anonymous referees for the very constructive criticism on a previous version of this work, which significantly improved the quality of the present paper.

**Conflicts of Interest:** The author declares no conflict of interest.

## References

- [1] Eisenstat, S. C., Elman, H. C., & Schultz, M. H. (1983). Variational iterative methods for nonsymmetric systems of linear equations. *SIAM Journal on Numerical Analysis*, 20(2), 345-357.
- [2] Barrett, R., Berry, M. W., Chan, T. F., Demmel, J., Donato, J., Dongarra, J., ... & Van der Vorst, H. (1994). *Templates for the solution of linear systems: building blocks for iterative methods* (Vol. 43). Siam.
- [3] Greenbaum, A. (1997). *Iterative methods for solving linear systems* (Vol. 17): Siam.
- [4] Saad, Y. (2003). *Iterative methods for sparse linear systems*: Siam.
- [5] Varga, R. S. (2009). *Matrix iterative analysis* (Vol. 27): Springer Science & Business Media.
- [6] Saberi Najafi, H., & Edalatpanah, S. A. (2011). Some Improvements In Preconditioned Modified Accelerated Overrelaxation (PMAOR) Method For Solving Linear Systems. *Journal of Information and Computing Science*, 6(1), 015-022 .

- [7] Najafi, H. S., & Edalatpanah, S. A. (2013). On the convergence regions of generalized accelerated overrelaxation method for linear complementarity problems. *Journal of Optimization Theory and Applications*, 156(3), 859-866.
- [8] Saberi Najafi, H., & Edalatpanah, S. A. (2014). A new modified SSOR iteration method for solving augmented linear systems. *International Journal of Computer Mathematics*, 91(3), 539-552 .
- [9] Najafi, H. S., Edalatpanah, S. A., & Gravvanis, G. A. (2014). An efficient method for computing the inverse of arrowhead matrices. *Applied mathematics letters*, 33, 1-5.
- [10] Najafi, H. S., & Edalatpanah, S. A. (2015). A new family of (I+ S)-type preconditioner with some applications. *Computational and Applied Mathematics*, 34(3), 917-931.
- [11] Saberi Najafi, H., & Edalatpanah, S. A. (2016). On the iterative methods for weighted linear least squares problem. *Engineering Computations*, 33(2), 622-639.
- [12] Saberi Najafi, H., Edalatpanah, S. A., & Refahisheikhani, A. H. (2018). An analytical method as a preconditioning modeling for systems of linear equations. *Computational and Applied Mathematics*, 37(2), 922-931.
- [13] Edalatpanah, S. A. (2018). On the modified methods for irreducible linear systems with L-matrices. *Bulletin of Computational Applied Mathematics*, 6(1).
- [14] Edalatpanah, S. A. (2020). On the preconditioned projective iterative methods for the linear complementarity problems. *RAIRO-Operations Research*, 54(2), 341-349.
- [15] Fallah, M., & Edalatpanah, S. A. (2020). On the Some New Preconditioned Generalized AOR Methods for Solving Weighted Linear Least Squares Problems. *IEEE Access*, 8, 33196-33201.
- [16] Zadeh, L. A. (1965). Fuzzy sets. *Information and control*, 8(3), 338-353 .
- [17] Zadeh, L. A. (1972). A fuzzy-set-theoretic interpretation of linguistic hedges .
- [18] Bellman, R. E., & Zadeh, L. A. (1970). Decision-making in a fuzzy environment. *Management science*, 17(4), B-141-B-164 .
- [19] Yager, R. R., & Filev, D. P. (1994). *Essentials of fuzzy modeling and control*. New York .
- [20] Hájek, P. (1998). *Metamathematics of fuzzy logic (Vol. 4)*: Springer Science & Business Media.
- [21] Cordon, O. (2001). *Genetic fuzzy systems: evolutionary tuning and learning of fuzzy knowledge bases (Vol. 19)*: World Scientific.
- [22] Zimmermann, H.-J. (2001). *Fuzzy set theory—and its applications*: Springer Science & Business Media.
- [23] Kaufmann, A., & Gupta, M. M. (1991). *Introduction to fuzzy arithmetic: theory and applications*: Arden Shakespeare.
- [24] Dubois, D., & Prade, H. (1980). Systems of linear fuzzy constraints. *Fuzzy sets and systems*, 3(1), 37-48 .
- [25] Friedman, M., Ming, M., & Kandel, A. (1998). Fuzzy linear systems. *Fuzzy sets and systems*, 96(2), 201-209 .
- [26] Allahviranloo, T. (2004). Numerical methods for fuzzy system of linear equations. *Applied mathematics and computation*, 155(2), 493-502.
- [27] Allahviranloo, T. (2005). Successive over relaxation iterative method for fuzzy system of linear equations. *Applied Mathematics and Computation*, 162(1), 189-196.
- [28] Allahviranloo, T., Ghanbari, M., Hosseinzadeh, A. A., Hagh, E., & Nuraei, R. (2011). A note on "Fuzzy linear systems". *Fuzzy sets and systems*, 177(1), 87-92.
- [29] Asady, B., Abbasbandy, S., & Alavi, M. (2005). Fuzzy general linear systems. *Applied Mathematics and Computation*, 169(1), 34-40.
- [30] Dehghan, M., & Hashemi, B. (2006). Iterative solution of fuzzy linear systems. *Applied mathematics and*

- computation, 175(1), 645-674.
- [31] Dehghan, M., Hashemi, B., & Ghatte, M. (2006). Computational methods for solving fully fuzzy linear systems. *Applied Mathematics and Computation*, 179(1), 328-343 .
- [32] Saberi Najafi, H., & Edalatpanah, S.A. (2012). The block AOR iterative methods for solving fuzzy linear systems. *The Journal of Mathematics and Computer Science*, 4(4), 527-535 .
- [33] Saberi Najafi, H., & Edalatpanah, S.A. (2012). Preconditioning strategy to solve fuzzy linear systems (FLS). *International Review of Fuzzy Mathematics*, 7(2), 65-80 .
- [34] Saberi Najafi, H., & Edalatpanah, S. A. (2014). H-Matrices in Fuzzy Linear Systems. *International Journal of Computational Mathematics*, 2014, Article ID 394135, 394136 pages .
- [35] Saberi Najafi, H., Edalatpanah, S. A., & Refahi Sheikhan, A. H. (2013). Application of homotopy perturbation method for fuzzy linear systems and comparison with Adomian's decomposition method. *Chinese Journal of Mathematics*, 2013.
- [36] Saberi Najafi, H., Edalatpanah, S. A. (2013). An improved model for iterative algorithms in fuzzy linear systems. *Computational Mathematics and Modeling*, 24(3), 443-451.
- [37] Salahshour, S., Ahmadian, A., Ismail, F., & Baleanu, D. (2016). A novel weak fuzzy solution for fuzzy linear system. *Entropy*, 18(3), 68.
- [38] Lodwick, W. A., & Dubois, D. (2015). Interval linear systems as a necessary step in fuzzy linear systems. *Fuzzy sets and systems*, 281, 227-251.
- [39] Behera, D., & Chakraverty, S. (2012). A new method for solving real and complex fuzzy systems of linear equations. *Computational Mathematics and Modeling*, 23(4), 507-518.
- [40] Chakraverty, S., & Behera, D. (2013). Fuzzy system of linear equations with crisp coefficients. *Journal of Intelligent & Fuzzy Systems*, 25(1), 201-207.
- [41] Wang, K., Chen, G., & Wei, Y. (2009). Perturbation analysis for a class of fuzzy linear systems. *Journal of Computational and Applied Mathematics*, 224(1), 54-65.
- [42] Rzeżuchowski, T., & Wa, J. (2008). Solutions of fuzzy equations based on Kaucher arithmetic and AE-solution sets. *Fuzzy Sets and Systems*, 159(16), 2116-2129.
- [43] Edalatpanah, S. A. (2017). Modified iterative methods for solving fully fuzzy linear systems. In *Fuzzy Systems: Concepts, Methodologies, Tools, and Applications* (pp. 55-73). IGI Global.
- [44] Jerković, V. M., Mihailović, B., & Malešević, B. (2017). A New Method for Solving Square Fuzzy Linear Systems. In *Advances in Fuzzy Logic and Technology 2017* (pp. 278-289). Springer, Cham.
- [45] Mihailović, B., Jerković, V. M., & Malešević, B. (2018). Solving fuzzy linear systems using a block representation of generalized inverses: The group inverse. *Fuzzy Sets and Systems*, 353, 66-85.
- [46] Akram, M., Muhammad, G., & Allahviranloo, T. (2019). Bipolar fuzzy linear system of equations. *Computational and Applied Mathematics*, 38(2), 69.
- [47] Atanassov, K. T. (1986). Intuitionistic fuzzy sets. *Fuzzy sets and Systems*, 20(1), 87-96.
- [48] Smarandache, F, A unifying field in logics. *Neutrosophy: Neutrosophic probability, set and logic*, American Research Press, Rehoboth 1999.
- [49] Smarandache, F, A unifying field in logics: neutrosophic logic. *Neutrosophy, neutrosophic set, neutrosophic probability and statistics*, third ed., Xiquan, Phoenix, 2003.
- [50] Smarandache, F. (2013). n-Valued Refined Neutrosophic Logic and Its Applications to Physics. *Prog Phys* 4,143-146.
- [51] Smarandache, F. (2015). Refined literal indeterminacy and the multiplication law of sub-indeterminacies. *Neutrosophic Sets Syst* 9,58-63.

- [52] Gallego Lupiáñez, F. (2009). Interval neutrosophic sets and topology. *Kybernetes*, 38(3/4), 621-624.
- [53] Tian, Z. P., Zhang, H. Y., Wang, J., Wang, J. Q., & Chen, X. H. (2016). Multi-criteria decision-making method based on a cross-entropy with interval neutrosophic sets. *International Journal of Systems Science*, 47(15), 3598-3608.
- [54] Garg, H. (2018). Non-linear programming method for multi-criteria decision making problems under interval neutrosophic set environment. *Applied Intelligence*, 48(8), 2199-2213.
- [55] Broumi, S., Smarandache, F., Talea, M., & Bakali, A. (2016). An introduction to bipolar single valued neutrosophic graph theory. In *Applied Mechanics and Materials* (Vol. 841, pp. 184-191). Trans Tech Publications.
- [56] Uluçay, V., Deli, I., & Şahin, M. (2018). Similarity measures of bipolar neutrosophic sets and their application to multiple criteria decision making. *Neural Computing and Applications*, 29(3), 739-748.
- [57] Abdel-Basset, M., Mohamed, M., Elhoseny, M., Chiclana, F., & Zaied, A. E. N. H. (2019). Cosine similarity measures of bipolar neutrosophic set for diagnosis of bipolar disorder diseases. *Artificial Intelligence in Medicine*, 101, 101735.
- [58] Ye, J. (2013). Multicriteria decision-making method using the correlation coefficient under single-valued neutrosophic environment. *International Journal of General Systems*, 42(4), 386-394.
- [59] Deli, I., & Şubaş, Y. (2017). A ranking method of single valued neutrosophic numbers and its applications to multi-attribute decision making problems. *International Journal of Machine Learning and Cybernetics*, 8(4), 1309-1322.
- [60] Edalatpanah, S. A. (2018). Neutrosophic perspective on DEA. *J. Appl. Res. Ind. Eng*, 5(4), 339-345.
- [61] Edalatpanah, S. A., & Smarandache, F. (2019). Data Envelopment Analysis for Simplified Neutrosophic Sets. *Neutrosophic Sets and Systems*, 29, 215-226.
- [62] Edalatpanah, S. A. (2019). A nonlinear approach for neutrosophic linear programming. *J. Appl. Res. Ind. Eng*, 6(4), 367-373.
- [63] Edalatpanah, S. A. (2020). Data envelopment analysis based on triangular neutrosophic numbers. *CAAI Transactions on Intelligence Technology*.2020, Article DOI: 10.1049/trit.2020.0016.
- [64] Edalatpanah, S. A. (2020). Neutrosophic structured element. *Expert Systems*.Article DOI: 10.1111/exsy.12542 .
- [65] Edalatpanah, S. A. (2020). A Direct Model for Triangular Neutrosophic Linear Programming. *International Journal of Neutrosophic Science*, 1(1), 19-28.
- [66] Yang, W., Cai, L., Edalatpanah, S. A., & Smarandache, F. (2020). Triangular Single Valued Neutrosophic Data Envelopment Analysis: Application to Hospital Performance Measurement. *Symmetry*, 12(4), 588.
- [67] Peng, J. J., Wang, J. Q., Wu, X. H., Wang, J., & Chen, X. H. (2015). Multi-valued neutrosophic sets and power aggregation operators with their applications in multi-criteria group decision-making problems. *International Journal of Computational Intelligence Systems*, 8(2), 345-363.
- [68] Ji, P., Zhang, H. Y., & Wang, J. Q. (2018). A projection-based TODIM method under multi-valued neutrosophic environments and its application in personnel selection. *Neural Computing and Applications*, 29(1), 221-234.
- [69] Tian, Z. P., Wang, J., Wang, J. Q., & Zhang, H. Y. (2017). Simplified neutrosophic linguistic multi-criteria group decision-making approach to green product development. *Group Decision and Negotiation*, 26(3), 597-627.
- [70] Wang, J. Q., Yang, Y., & Li, L. (2018). Multi-criteria decision-making method based on single-valued neutrosophic linguistic Maclaurin symmetric mean operators. *Neural Computing and Applications*, 30(5),

- 1529-1547.
- [71] Zhang, M., Zhang, L., & Cheng, H. D. (2010). A neutrosophic approach to image segmentation based on watershed method. *Signal Processing*, 90(5), 1510-1517.
- [72] Kumar, R., Edalatpanah, S.A., Jha, S., Broumi, S., Dey, A. (2018) Neutrosophic shortest path problem, *Neutrosophic Sets and Systems*, 23, 5-15.
- [73] Kumar, R., Edalatpanah, S. A., Jha, S., Broumi, S., Singh, R., & Dey, A. (2019). A Multi Objective Programming Approach to Solve Integer Valued Neutrosophic Shortest Path Problems. *Neutrosophic Sets and Systems*, 134.
- [74] Tey, D. J. Y., Gan, Y. F., Selvachandran, G., Quek, S. G., Smarandache, F., Abdel-Basset, M., & Long, H. V. (2019). A novel neutrosophic data analytic hierarchy process for multi-criteria decision making method: A case study in kuala lumpur stock exchange. *IEEE Access*, 7, 53687-53697.
- [75] Son, N. T. K., Dong, N. P., Abdel-Basset, M., Manogaran, G., & Long, H. V. (2020). On the stabilizability for a class of linear time-invariant systems under uncertainty. *Circuits, Systems, and Signal Processing*, 39(2), 919-960.
- [76] Abdel-Basset, M., Ali, M., & Atef, A. (2020). Uncertainty assessments of linear time-cost tradeoffs using neutrosophic set. *Computers & Industrial Engineering*, 141, 106286.
- [77] Abdel-Basset, M., & Mohamed, R. (2020). A novel plithogenic TOPSIS-CRITIC model for sustainable supply chain risk management. *Journal of Cleaner Production*, 247, 119586.
- [78] Abdel-Basset, M., Ali, M., & Atef, A. (2020). Resource levelling problem in construction projects under neutrosophic environment. *The Journal of Supercomputing*, 76(2), 964-988.
- [79] Ye, J. (2017). Neutrosophic linear equations and application in traffic flow problems. *Algorithms*, 10(4), 133.
- [80] Ye, J., & Cui, W. (2018). Neutrosophic state feedback design method for SISO neutrosophic linear systems. *Cognitive Systems Research*, 52, 1056-1065.

Received: Jul 7, 2019. Accepted: Apr 26, 2020