

Bacterial carbon sources in coastal sediments: a cross-system analysis based on stable isotope data of biomarkers

S. Bouillon^{1,*} and H. T. S. Boschker^{2,*}

¹Department of Analytical and Environmental Chemistry, Vrije Universiteit Brussel, Pleinlaan 2, B-1050 Brussels, Belgium

²Netherlands Institute of Ecology (NIOO-KNAW), Centre for Estuarine and Marine Ecology, PO Box 140, 4400 AC Yerseke, The Netherlands

*Both authors contributed equally to this work.

Received: 26 August 2005 – Published in Biogeosciences Discuss.: 14 October 2005

Revised: 18 January 2006 – Accepted: 6 March 2006 – Published: 3 May 2006

Abstract. Coastal ecosystems are typically highly productive, and the sediments in these systems receive organic matter from a variety of local and imported sources. To assess if general patterns are present in the origin of carbon sources for sedimentary bacteria and their relation to the origin of the sediment organic carbon pool, we compiled both literature and new data on $\delta^{13}\text{C}$ of bacterial biomarkers (the phospholipid derived fatty acids $i+a15:0$), along with $\delta^{13}\text{C}$ data on sediment organic carbon ($\delta^{13}\text{C}_{\text{TOC}}$) and macrophyte biomass from a variety of typical near-coastal systems. These systems included mangroves, salt marshes (both C3 and C4-dominated sites), seagrass beds, and macroalgae-based systems, as well as unvegetated sediments. First, our $\delta^{13}\text{C}_{i+a15:0}$ data showed large variability over the entire range of $\delta^{13}\text{C}_{\text{TOC}}$, indicating that in many settings, bacteria may depend on carbon derived from various origins. Secondly, systems where local macrophyte production is the major supplier of organic carbon for in situ decomposition are generally limited to organic carbon-rich, peaty sites ($\text{TOC} > 10 \text{ wt}\%$), which are likely to make up only a small part of the global area of vegetated coastal systems. These carbon-rich sediments also provided a field based estimate of isotopic fractionation between bacterial carbon sources and biomarkers ($-3.7 \pm 2.1\text{‰}$), which is similar to the expected value of about -3‰ associated with the biosynthesis of fatty acids. Thirdly, only in systems with low TOC (below $\sim 1 \text{ wt}\%$), we consistently found that bacteria were selectively utilizing an isotopically enriched carbon source, which may be root exudates but more likely is derived from microphytobenthos. In other systems with between ~ 1 and $10 \text{ wt}\%$ TOC, bacteria appear to show on average little selectivity and $\delta^{13}\text{C}_{i+a15:0}$ data generally follow the $\delta^{13}\text{C}_{\text{TOC}}$, even in sys-

tems where the TOC is a mixture of algal and macrophyte sources that generally are believed to have a very different degradability.

1 Introduction

The coastal zone is widely recognized as a biogeochemically active region, where organic carbon inputs from a variety of sources undergo intense biogeochemical processing. It forms a significant component in the global oceanic carbon budget despite a relatively small areal extent (e.g. Gattuso et al., 1998; Borges, 2005; Middelburg et al., 2004; Duarte et al., 2004). The coastal zone is suggested to be responsible for about 20% of the oceanic primary production and the vast majority of oceanic organic carbon burial (Gattuso et al., 1998; Duarte et al., 2004), and the most recent data compilation indicates that, although not well constrained, benthic mineralization in coastal sediments amounts to $620 \text{ Tmol C y}^{-1}$ or half of the total mineralization in marine sediments (Middelburg et al., 2004).

Various sources of organic matter enter the coastal zone, ranging from local primary production by phytoplankton or benthic microalgae, terrestrial inputs via river discharge to production by macrophyte systems such as seagrasses, macroalgal beds, mangroves and salt marshes. The sediment organic matter pool is therefore mostly derived from a mixture of source materials as a result of the intense mixing by currents. The identity and importance of the source materials that drive mineralization in sediments likely depends on a combination of their relative amounts and degradability. Carbon sources in coastal areas are characterized by a large variability in their composition and degradability, ranging from labile sources such as phytoplankton and benthic microalgae

Correspondence to: S. Bouillon
(steven.bouillon@vub.ac.be)

to less degradable sources such as macrophyte material and terrestrial C transported by rivers. Degradability can further be modified in time as less available fractions remain (Middelburg, 1989) or decreased by adsorption to clay minerals (Keil et al., 1994). Recent studies in estuaries have indicated that bacterial mineralization can be sustained both by aquatic primary production and by terrestrial C, and that it can have a large impact on the amount, composition, age, and lability of organic matter prior to its export into the coastal zone or ocean (e.g. Raymond and Bauer, 2001; McAllister et al., 2004; Boschker et al., 2005).

In coastal ecosystems, movement of carbon across ecosystem boundaries complicates budgeting studies, as mineralization may be partially sustained by non-local sources, necessitating the use of proxies to take these sources into account (e.g. Bouillon et al., 2004b). For a given system, it is not always straightforward to assess the origin of carbon driving benthic mineralization. The importance of non-local sources has been demonstrated both with stable isotope techniques (e.g. Boschker et al., 1999; Holmer et al., 2004) and based on mass balance considerations – i.e. when benthic mineralization rates considerably exceed rates of local primary production (e.g. Barrón et al., 2004).

Stable carbon isotope signatures ($\delta^{13}\text{C}$) of the various carbon inputs are often different, and despite some overlap between different sources, can be powerful tracers of carbon inputs in various ecosystem components (Fry and Sherr, 1984). Although bulk stable isotope measurements have been possible for several decades, the introduction of compound-specific $\delta^{13}\text{C}$ analyses as a tool to include microbial communities only started a decade ago (Freeman et al., 1990). PLFA (phospholipid derived fatty acids) in particular have become popular biomarkers for stable isotope studies since they are representative of live microbial biomass (fast degradation of phospholipids occurs after cell death), are suitable for gas chromatography – isotope ratio mass spectrometry (GC-IRMS) analysis after a derivatization procedure which introduces only one additional C atom, and because various PLFA can be linked to specific microbial groups (Boschker and Middelburg, 2002).

In this study, we have compiled 339 data on both bulk sedimentary biogeochemical parameters (total organic carbon content (%TOC) and the stable isotope composition of organic C, i.e. $\delta^{13}\text{C}_{\text{TOC}}$) and PLFA proxies for the isotope composition of sedimentary bacteria ($\delta^{13}\text{C}_{\text{i+a15:0}}$) from a variety of coastal ecosystems in order to identify general patterns in the sources of sedimentary carbon and their use by microbial communities across and within these coastal ecosystem types. To determine the origin of carbon supporting in situ bacterial populations, we selected iso- and anteiso-branched 15:0 (i+a15:0) PLFA because (i) these branched fatty acids have been well demonstrated to be of bacterial origin, (ii) they show no chromatographic interferences with other compounds under the analytical conditions used, (iii) they are ubiquitous in coastal marine sediments in concen-

trations suitable for $\delta^{13}\text{C}$ analysis, and (iv) fractionation data are available in the literature (Boschker et al., 1999). The i+a15:0 PLFA are found in a diverse range of important bacterial groups detected in sediments, such as the sulphate reducing bacteria of the δ -Proteobacteria, the CFB-bacteria and various groups in the Gram-positive bacteria (e.g. Kaneda, 1991; Bowman and McCuaig, 2003). In addition, several studies have shown good correlations in stable isotopic composition between i+a15:0 and other PLFA in bulk sediment (Boschker et al., 1999; Cifuentes and Salata, 2001; Bouillon et al., 2004a). The extensive dataset in this cross-system analysis is used to demonstrate general trends in carbon sources used by bacteria in near-coastal sediments.

2 Materials and methods

2.1 Data sources

Overall, our dataset contains 187 data from our own published work, 23 data from other literature sources, and 129 new observations (see supplemental file: <http://www.biogeosciences.net/3/175/bg-3-175-sp.pdf>). An overview of the various sampling locations and data sources is presented in Table 1. Only data where both $\delta^{13}\text{C}_{\text{TOC}}$ and $\delta^{13}\text{C}_{\text{i+a15:0}}$ were measured on sediment horizons in the 0 to 10 cm depth range were retained. Due to some missing data on other parameters (e.g. %TOC, $\delta^{13}\text{C}_{\text{plant}}$), the number of data points for each ecosystem in some of the graphs may differ slightly.

We gathered data from mangrove systems in the following locations: (i) a lagoonal mangrove system in southwest Sri Lanka (Pambala, see Bouillon et al., 2004a), (ii) estuarine mangrove sites in southeast India (Pichavaram and Chunnambar, see Bouillon et al., 2004a), (iii) various sites in an estuarine mangrove system with adjacent seagrass beds in southeast Kenya (Gazi Bay, see Bouillon et al., 2004b), and (iv) riverine mangrove forests along the Tana river (northeast Kenya) and estuarine mangrove sites in the Tana delta (northeast Kenya) collected in April 2004 (this study). For the latter sites, it is worth mentioning that the organic matter transported by Tana river contains a significant amount of C4-derived carbon, with river and mangrove creek particulate organic carbon having $\delta^{13}\text{C}$ values of $\sim -20\text{‰}$ (S. Bouillon, unpublished data).

Data from marshes dominated by C4 plants (*Spartina* spp) were obtained from a number of sites on both sides of the Atlantic Ocean. Literature data from the estuarine Waarde and the Kattedijke marshes (The Netherlands), the Great Marshes (MA, USA, see Boschker et al., 1999), and from the Gulf of Mexico (Cifuentes and Salata, 2001) were included together with a large number of data from unpublished work on various marshes in the Netherlands, France and the USA (Table 1). Sampling sites for C4 marshes were chosen to cover a range of sediment organic matter content from very organic-poor and recently colonized sites on sand beaches to

Table 1. Overview of study site characteristics and data sources.

Site	Latitude, Longitude	Number of samples per ecosystem type					Data source
		C4 marsh	C3 marsh	Mangrove	Seagrass	Unvegetated and macroalgae	
Schiermonnikoog (the Netherlands)	53°30' N 6°10' E		35			6	1
St Annaland (the Netherlands)	51°36' N 4°08' E	2			3	3	1, 2
Kattedijke (the Netherlands)	51°38' N 3°56' E	3				2	3
Waarde marsh (the Netherlands)	51.24° N 4°07' E	17	9			13	1, 3
Valkenisse (the Netherlands)	51°23' N 4°08' E	2				2	1
Ritthem (the Netherlands)	51°27' N 3°40' E	13				8	1
Mont St. Michel Bay (France)	48°36' N 1°48' E	3					1
Plum Island Sound (MA, USA)	42°45' N 70°50' W	13				10	1
Great Marshes (MA, USA)	41°43' N 70°21' W	3				2	3
Canary Creek (DE, USA)	38°47' N 75°09' W	2				2	1
North River (NC, USA)	34°45' N 76°35' W	2	2		2	1	4
North Inlet (SC, USA)	33°20' N 79°10' W	2				2	1
Chunnambar (India)	11°53' N 79°48' E			3			5
Pambala (Sri Lanka)	7°35' N 79°47' E			24			5
Pichavaram (India)	11°27' N 79°17' E			8			5
Gazi Bay (Kenya)	4°22' S 39°30' E			41	12		6
Tana delta (Kenya)	2°30' S, 40°30' E			15			1
Nyborg Fjord (Denmark)	55°17' N 10°49' E				6	6	2
Arcachon Bay (France)	44°40' N 1°10' E				9	9	2
Mallorca, various sites (Spain)	39°09' N 2°56' E				10	14	7
Laguna Madre (TX, USA)	26°09' N 97°12' W				8	4	8
Ban Pak Klok (Thailand)	8°03' N 98°25' E				6		1, 9

Data sources: 1: this study, 2: Boschker et al. (2000), 3: Boschker et al. (1999), 4: Cifuentes and Salata (2001), 5: Bouillon et al. (2004a), 6: Bouillon et al. (2004b), 7: Holmer et al. (2004), 8: Jones et al. (2003), 9: Holmer et al. (2001).

older, organic rich systems with silt or peat-rich sediments. Literature data from C3 marshes were very scarce (Waarde marsh, the Netherlands, see Boschker et al. (1999), and from the Gulf of Mexico, see Cifuentes and Salata, 2001), but a significant amount of new data have been gathered from the marsh on the island of Schiermonnikoog (Wadden Sea, the Netherlands) where an elevation and age gradient was sampled.

Finally, seagrass data were compiled from various temperate (Cifuentes and Salata, 2001; Boschker et al., 2000; Holmer et al., 2004) and tropical (Holmer et al., 2001; Jones et al., 2003; Bouillon et al., 2004b) systems. Both subtidal and intertidal seagrass beds from all climatic zones are represented in our data set. In addition to vegetated sediments, data from nearby unvegetated sites were also available for various marshes and seagrass beds. Many of the intertidal and subtidal unvegetated sediments were covered by benthic microalgae, mainly diatoms. Only a very limited number of data on macroalgae systems are available (Holmer et al., 2004), and in these cases the distinction between vegetated and unvegetated (i.e. mudflats and bare sub-tidal sediments) was not always straightforward. Consequently, no distinction was made, and the data from macroalgal systems were combined with those from the unvegetated sites.

It should be noted that for the data in Cifuentes and Salata (2001), we calculated some of the $\delta^{13}\text{C}_{i+a15:0}$ data as the average of tabulated $\delta^{13}\text{C}_{i15:0}$ and $\delta^{13}\text{C}_{a15:0}$ values. Since these show excellent correlation when sufficient chromatographic separation is achieved, any possible bias introduced by this procedure is likely irrelevant in the context of this meta-analysis. For all other sites, the more correct, concentration-based average $\delta^{13}\text{C}$ is reported for the bacterial markers i15:0 and a15:0. Finally, we also considered the possibility of compiling data on the $\delta^{13}\text{C}$ composition of the diatom marker 20:5 ω 3 to evaluate the coupling between microphytobenthos and bacteria. Although such data were available from a limited number of sites (see Boschker et al., 2000; Holmer et al., 2004), the low concentrations of this marker in other systems did not allow us to reliably use the resulting $\delta^{13}\text{C}$ data or to unambiguously ascribe the marker data to benthic diatoms, since it has also been reported to occur in other algae (Volkman et al., 1989) and heterotrophic micro-eukaryotes such as ciliates (Harvey et al., 1997). In addition, the 20:5 ω 3 PLFA is only an indicator for living biomass and not for algal detritus, which may be important in many of the studied systems.

Relationships between %TOC and $\delta^{13}\text{C}_{\text{TOC}}$ were tested with a Spearman Rank Correlation analysis, and a cumulative sum analysis (CUSUM) was performed on the %TOC- $\Delta\delta$




Fig. 1. Analytical scheme used for the stable isotope data collected in this study.

data (see discussion, Sect. 3.2). The latter approach is frequently used to detect shifts or deviations in processes. After ranking the data in order of increasing %TOC, a plot of the cumulative sequential differences between each data point and the overall average or median of the data is made. The slope of the CUSUM plot in a certain data range (positive, flat, or negative) then indicates that in this region, the average is, respectively, higher, similar, or lower than the overall average. This allowed us to define ranges of %TOC (between a marked change in the slope), after which a Dunn's multiple comparison test was performed on the $\Delta\delta$ data corresponding to from each %TOC interval.

2.2 Analytical techniques

For all new data in this study, sediment samples were collected with corers. The exact sediment horizons analyzed depends on the data set, but all samples presented here are from the 0 to 10 cm depth range. Samples for PLFA analysis were either directly transferred in the extraction solvents, or frozen, after which they were freeze-dried and stored frozen prior to extraction. Extraction and derivatisation of PLFA was performed using a modified Bligh and Dyer extraction, silica column partitioning, and mild alkaline transmethylation as described earlier (Boschker et al., 2004; Bouillon et al., 2004a). $\delta^{13}\text{C}$ of the resulting FAMES (fatty acid methyl esters) were determined on a ThermoFinnigan Delta type of GC-IRMS (gas chromatograph – isotope ratio mass spectrometer) in various configurations. All samples were run in splitless mode, using a HP-5 or BPX-70 column (30 or 60 m, 0.32 mm ID) with a He flow rate of 2 ml/min. $\delta^{13}\text{C}$ data of PLFA are corrected for the addition of the methyl group by simple mass balance and were calibrated by our own internal and external FAME standards. Reproducibility is estimated to be 0.6‰ or better. Elemental analyses (TOC, TN, as % of sediment dry weight) and bulk TOC $\delta^{13}\text{C}$ analyses were performed by elemental analyzer-IRMS. For a more elaborate description of sampling and analytical details, we refer to Boschker (2004) and Bouillon et al. (2004a).

3 Results and discussion

The two most important data types collect in our study are the stable carbon isotopic ratio of the sediment organic matter ($\delta^{13}\text{C}_{\text{TOC}}$) and of the bacterial PLFA ($\delta^{13}\text{C}_{\text{C}_{1+15:0}}$). The relationship between these two data types and the way we

analyze them is presented in Fig. 1. The TOC in sediments is generally a mixture of different source materials with different isotopic ratios, and the sediment isotopic ratio ($\delta^{13}\text{C}_{\text{TOC}}$) is therefore indicative for the carbon sources that contribute to the TOC pool. Bacteria will probably only utilize a fraction of the TOC as their substrate as the various source materials are characterized by differences in degradability or accessibility. It is generally accepted that isotopic ratios of bacteria or heterotrophic organisms in general reflects their substrate (with no discernable fractionation, e.g. Fry and Sherr, 1984; Hullar et al., 1996). However, we analyzed bacterial PLFA as representatives of the bacterial biomass, and there generally is an offset between the total biomass and PLFA (the latter being more ^{13}C -depleted) due to fractionation effects during fatty acid synthesis. This fractionation factor is not well constrained but appears fairly constant for diverse bacterial communities growing on complex substrates as found in sediments (Boschker et al., 1999). In order to relate the isotopic ratio of the bacterial PLFA to the substrates the bacteria were using, this offset must be known and relatively constant. In the discussion below, we will first discuss general trends in the isotopic composition of the TOC in our data set and then turn our attention to the bacterial biomarker data and their relationship with the TOC.

3.1 Sources of organic carbon in coastal sediments

Organic carbon in coastal sediments mostly consists of a mixture of different sources, including locally produced macrophyte material, microphytobenthos, and suspended organic matter, with a variable composition, imported from outside the ecosystem boundaries – through sedimentation in the water column (subtidal seagrass beds) or during tidal inundation (mangroves and salt marshes). A combination of sediment organic carbon concentrations and $\delta^{13}\text{C}$ of sediment TOC can be used to document the variation in the relative importance of local versus allochthonous carbon sources (Middelburg et al., 1997). For mangrove and salt marsh systems, a good overall relationship between $\delta^{13}\text{C}_{\text{TOC}}$ and sediment TOC levels was found (Fig. 2a; Spearman correlation coefficient = -0.46 , $p < 0.001$ for mangroves and C3 marshes; Spearman correlation coefficient = 0.38 , $p = 0.002$ for C4 marshes), with $\delta^{13}\text{C}_{\text{TOC}}$ values approaching those of the local macrophyte inputs in systems with high TOC of more than 10% (i.e. $\sim -26\text{‰}$ in mangroves and $\sim -14\text{‰}$ in *Spartina* marshes). In contrast, the $\delta^{13}\text{C}_{\text{TOC}}$ was much more variable in low TOC settings (up to between 5 and 10% TOC)




Fig. 2. Compilation of $\delta^{13}\text{C}$ signatures of (A) bulk sediments organic carbon and (B) $\delta^{13}\text{C}$ of bacterial PLFA as a function of sedimentary organic carbon content in different types of coastal ecosystems. Note the different scales on the Y-axes.

reflecting a variable contribution by benthic microalgae, phytoplankton and detrital inputs. Such allochthonous inputs can have a wide range of $\delta^{13}\text{C}$ signatures, and depending on the ecosystem considered may consist of terrestrial C (either from C3 or C4-dominated catchments), marine or estuarine phytodetritus, microphytobenthos and seagrass-derived C. Based on silt content data that were available for a subset of the sediments (not shown), this range of low TOC sediments also presents a transition from predominantly sandy to silty sediments.

The patterns observed in Fig. 2a confirm those reported earlier for *Spartina* marshes (Middelburg et al., 1997) and mangroves (Bouillon et al., 2003a, 2004a), whereby it was proposed that variations in sediment TOC and $\delta^{13}\text{C}_{\text{TOC}}$ can in general be adequately described as resulting from simple admixture of local macrophyte C and tidal inputs of suspended matter. Sediments where local macrophyte inputs dominate are typically peaty (i.e. high %TOC), with $\delta^{13}\text{C}_{\text{TOC}}$ close to those of the macrophyte vegetation, whereas the more mineral-rich sediments result largely from sedimentation and trapping of suspended material and its associated organic matter – hence, such sediments are characterized by a lower TOC content and highly variable $\delta^{13}\text{C}_{\text{TOC}}$, often deviating significantly from the $\delta^{13}\text{C}$ signature of the dominant vegetation. Moreover, for salt marshes where data from both vegetated systems and adjacent mudflats or unvegetated patches are available, unvegetated areas typically have a lower TOC content and show $\delta^{13}\text{C}$ signatures that deviate from those of the dominant vegetation, i.e. lower $\delta^{13}\text{C}$ for *Spartina* marshes and higher $\delta^{13}\text{C}$ for C3 marshes (Fig. 2a). The latter pattern indicates, as expected, that vegetated patches have an important but variable contribution from local macrophyte carbon and that unvegetated sites are more dominated by tidally imported C and, possibly, microphytobenthos.

For seagrass and unvegetated systems, such clear relationships between %TOC and $\delta^{13}\text{C}_{\text{TOC}}$ are not observed (Fig. 2a; Spearman correlation coefficient = -0.07 ; $p > 0.5$), but it should be kept in mind that (i) the overall range of TOC is much smaller than that in marshes and mangroves and (ii) the variability in macrophyte $\delta^{13}\text{C}$ values in seagrass systems is much larger, since the $\delta^{13}\text{C}$ of the seagrass biomass is in part determined by the $\delta^{13}\text{C}$ of the dissolved inorganic carbon pool (DIC) and the growth conditions, which can be highly variable in coastal settings (e.g. Hemminga and Mateo, 1996).

From the direct comparison of $\delta^{13}\text{C}_{\text{TOC}}$ with $\delta^{13}\text{C}_{\text{plant}}$ (Fig. 3a), it is clear that in most cases, local macrophyte production is not the dominant C input to the sediment TOC pool. If macrophyte material dominates the TOC pool then most of the data would plot close to the 1:1 line in Fig. 3a, i.e. sediment $\delta^{13}\text{C}_{\text{TOC}}$ would reflect the signature of the macrophyte vegetation. The deviation from this expected pattern for all systems considered (i.e. more positive $\delta^{13}\text{C}$ in C3 marshes and mangroves more negative $\delta^{13}\text{C}$ in C4 marshes, seagrasses and macroalgae) is consistent with extensive inputs from suspended organic C and/or microphytobenthos, and this pattern is similar to that recently reported based on a more comprehensive data compilation on seagrass sediment $\delta^{13}\text{C}$ data (Kennedy et al., 2004; Bouillon et al., 2004b). A second point worth noting in Fig. 2 is that settings in which local macrophyte inputs dominate the TOC pool (i.e. with high TOC content and $\delta^{13}\text{C}$ values close to those of the macrophyte vegetation) are overall quite scarce. Although it could be argued that the dataset here is too limited to generalize this conclusion, this pattern appears to be maintained if other datasets on TOC in coastal sediments are included (e.g. for mangrove systems; 65% of the data we have compiled show a TOC of less than 5%, and 82% of the data have less than 10% TOC, $n=650$). This implies that a substantial




Fig. 3. Plot of (A) $\delta^{13}\text{C}$ of bulk sediment TOC and (B) $\delta^{13}\text{C}$ of bacterial PLFA i+a15:0 versus $\delta^{13}\text{C}$ of the dominant macrophyte vegetation for all vegetated coastal ecosystems considered. Note that for C3 marshes, C4 marshes, and mangroves for which no direct measurements of plant $\delta^{13}\text{C}$ were available, an average value of the data from other sites was assigned (i.e. $-28.2 \pm 2.0\%$ ($n=471$) for mangroves, $-25.9 \pm 1.0\%$ ($n=44$) for C3 marshes, and $-13.2 \pm 0.4\%$ ($n=21$) for C4 marshes). The isoline in panel A is a 1:1 line, in panel B the isoline represents a shift of -3.7% to correct for fractionation between i+a15:0 and the substrate (see text for rationale). Symbols as in Fig. 2.

input of non-local organic carbon should be considered to be more rule than exception in coastal sediments (see also Duarte et al., 2004). The sediment organic matter in most systems therefore consists of a mixture of carbon sources potentially sustaining bacterial mineralization processes and growth.

3.2 Bacterial carbon sources in coastal sediments

Given the wide range of $\delta^{13}\text{C}_{\text{TOC}}$ and $\delta^{13}\text{C}_{\text{plant}}$ in coastal ecosystems, we used a compilation of concurrent $\delta^{13}\text{C}_{\text{TOC}}$ and $\delta^{13}\text{C}$ data of bacterial PLFA (i+a15:0, Fig. 2b) to determine the extent to which bacteria assimilate various available carbon sources. The coastal ecosystems covered here are typically very productive, and a substantial part of this production is by macrophytes (saltmarsh plants, mangroves and seagrasses). The compilation of data presented here, however, shows that this local macrophyte production is not the dominant carbon source in most systems and thereby generalizes and confirms some of our earlier case studies (e.g. Boschker et al., 1999; Bouillon et al., 2004a). When compared to the stable isotope signatures of the dominant vegetation (Fig. 3b), the $\delta^{13}\text{C}_{\text{i+a15:0}}$ data clearly demonstrate that in C3-dominated systems (mangroves and C3 marshes), bacteria typically consume carbon sources more enriched in ^{13}C than local macrophytes. Whereas in *Spartina* marshes, seagrass and macroalgae beds (where local macrophytes are characterized by heavy $\delta^{13}\text{C}$ signatures), the substrate used by bacteria is generally much more depleted in ^{13}C than the local macrophytes. The general trend is therefore that the bacterial PLFA show more average isotopic ratios, whereas the local macrophytes are found at the extremes of the range. Given the evidence mentioned above that tidally imported carbon sources form a significant and often isotopically dis-

tinct input in these systems (Fig. 2a), it is clear that imported C sources often are a major C source sustaining benthic mineralization.

Considering that a variety of C sources can be available in coastal sediments, and that such different sources may have a different lability or accessibility, the question arises if and to which extent bacteria make selective use of certain sources. In the simplest scenario wherein no selectivity would occur, we would expect to see an excellent relationship between $\delta^{13}\text{C}_{\text{TOC}}$ and $\delta^{13}\text{C}_{\text{i+a15:0}}$, with an offset caused by isotope fractionation during fatty acid biosynthesis (determined experimentally for i+a15:0 as $-5.6 \pm 1.8\%$ by Boschker et al., 1999). Although the entire dataset (Fig. 4) shows a reasonable positive relationship ($R^2=0.58$, slope close to unity) between $\delta^{13}\text{C}_{\text{TOC}}$ and $\delta^{13}\text{C}_{\text{i+a15:0}}$, the variability observed is much larger than would be expected given the analytical precision of both parameters (better than $\pm 0.2\%$ and $\pm 0.6\%$ respectively) and the variability in isotope fractionation from the experiments by Boschker et al. (1999) ($-5.6 \pm 1.8\%$) or estimated from our data set ($-3.7 \pm 2.1\%$, $n=29$, see further). For any given $\delta^{13}\text{C}_{\text{TOC}}$, the range in $\delta^{13}\text{C}_{\text{i+a15:0}}$ typically spans 10‰ or more, which indicates that bacteria in many cases do not assimilate carbon sources merely in proportion to their relative abundance in the sediment TOC pool. It is also apparent from Fig. 4 that the majority of the points are located above the expected line, which suggests that in a substantial number of sediments the bacteria preferentially utilize an isotopically enriched carbon source.

A further interesting pattern in bacterial selectivity can be discerned when plotting the difference between both bacterial PLFA and TOC (hereafter referred to as $\Delta\delta$, i.e. $\delta^{13}\text{C}_{\text{i+a15:0}} - \delta^{13}\text{C}_{\text{TOC}}$) as a function of the organic carbon content of the sediments considered (Fig. 5). As discussed above, if bacteria show no selectivity against the different




Fig. 4. $\delta^{13}\text{C}$ of bacterial PLFA (i+a15:0) versus $\delta^{13}\text{C}$ of the total sediment organic carbon pool (TOC), for various types of coastal ecosystems. The dotted line represents the expected $\delta^{13}\text{C}_{i+a15:0}$ when bulk TOC would be the main substrate (see text for details). Symbols as in Fig. 2.

carbon sources in the TOC, the $\Delta\delta$ values would be more or less constant and show a slightly negative offset due to isotopic fractionation in bacterial lipid synthesis. In order to evaluate possible trends in this dataset, we first performed a cumulative sum analysis based on the median value. This indicated that there were three distinct regions ($\% \text{TOC} < 0.8$, $0.8 < \% \text{TOC} < 2.3$, and $\% \text{TOC} > 2.3$), and $\Delta\delta$ values in these regions were found to differ significantly (Dunn's multiple comparison test, $p < 0.05$).

A first observation is that the $\Delta\delta$ data converge to a median value -3.8% (interquartile range: 1.5%) for the data where $\% \text{TOC}$ exceeds 2.3% (see Fig. 5b, where average $\Delta\delta$ values are plotted for binned data). If we consider only the data with a TOC content higher than 10% , and hence where macrophyte material is the dominant (only) C source available (Fig. 2a), the average $\Delta\delta$ values is $-3.7 \pm 2.1\%$ ($n=29$), and our dataset therefore provides an empirical verification of the fractionation between substrate and i+a15:0 under field conditions. This relationship is very robust as it holds for both sediments from C3-dominated mangroves and C4, *Spartina* marshes, which have very different isotopic ratios in the sediment TOC. This value is also within the range reported by Boschker et al. (1999) and the -3% value expected from fatty acid synthesis in general (Hayes, 2001) and confirms that possible variations in the degree of isotope fractionation with environmental conditions (e.g. under anoxic or oxic conditions (Teece et al., 1999)) are not likely to be a major complication in our interpretations. Moreover, this field verification suggests that the large range in lipid fractiona-




Fig. 5. (A) Plot of $\Delta\delta$ (defined as $\delta^{13}\text{C}_{i+a15:0} - \delta^{13}\text{C}_{\text{TOC}}$) as a function of the organic carbon content of coastal sediments. Note that the data for vegetated C3 marshes from one of the sites (Schiermonnikoog) fall outside the general pattern and are therefore plotted separately with small +. Symbols further as in Fig. 2. Panel (B) shows the binned averages ± 1 s.d., whereby each $\% \text{TOC}$ interval represents 20 datapoints (the Schiermonnikoog data were not included).

tions found in experiments with single (simple) substrates and/or specific bacterial strains (e.g. Pelz et al., 1997; Abraham et al., 1998; Teece et al., 1999) likely can be ruled out in natural, highly diverse communities where bacteria process more complex natural organic substrates. However, we do find a considerable variation around the mean $\Delta\delta$ values (SD of binned ranges vary between 1.8 and 3.3% Fig. 5b), which could be due to both a selective use of certain organic matter sources and to some variation in fractionation in lipid synthesis between samples.

Secondly, in all sediments with a TOC content above 0.8% , the difference between $\delta^{13}\text{C}$ for TOC and the bacterial biomarkers is found to be also fairly constant. A Dunn's multiple comparison test does indicate a slightly higher $\Delta\delta$ values in 0.8 to 2.3% TOC range compared to sediments with higher TOC, but this difference is small ($\sim 1\%$). This uniformity in $\Delta\delta$ values is remarkable because it suggests that in the intermediate range (roughly between 1 and 10% TOC) the bacteria on average also utilized the TOC as found in the sediment with little preference between the different source materials. However, our $\delta^{13}\text{C}_{\text{TOC}}$ data (Fig. 2a) suggest that

in this range the TOC consists of a mixture of source materials derived from various algal and macrophyte sources, which in general have a greatly different degradability. Material from algae is mostly much more available to bacteria and is degraded with a much higher rate than macrophyte derived materials (Schoenberg et al., 1990; Hee et al., 2001). This difference in degradability is thought to be the result of differences in biochemical composition between algal and macrophyte derived materials. These intrinsic differences in degradability appear, however, on average not to be expressed in the sediments that we studied, indicating that other mechanisms may determine the degradability of organic materials in these sediments. Most of the TOC found in sediments is sorbed to the mineral, clay phase of the sediment, and it has been shown that this greatly reduces the availability for bacterial degradation (Keil et al., 1994). A hypothesis to explain our results may be that this sorption determines the degradability of all source materials to a similar extent and that the availability of the organic matter is largely determined by the rate at which the sorbed substrates are released from the mineral phase.

Thirdly, below a TOC content of 0.8% there is a significant and large shift to more positive $\Delta\delta$ values (median $\Delta\delta$ is -0.2%). Under the assumption that isotope fractionation in lipid synthesis is similar as in the high TOC environments, this shift indicates that in these low TOC sediments, bacteria are preferentially utilizing an easily degradable, relatively ^{13}C -enriched carbon source with a limited abundance in the total TOC pool (Fig. 5). It is possible that this shift is partially because bacteria were preferentially using a specific component of the plant biomass that is isotopically enriched such as carbohydrates. However, the enrichment in carbohydrates ($0\text{--}2\%$ enriched compared to total biomass; Benner et al 1987) is much smaller compared to the shift of about 4% that we detected, and it is not clear why this would only become apparent in low % TOC sediments and not in the high TOC, peaty sediments. The data in this range of % TOC are mainly from C4 marshes and seagrass beds, which both have relatively enriched ratios in the local macrophyte material, and also from unvegetated sediments. The effect is also clearly seen in the $\delta^{13}\text{C}_{i+a15:0}$ data from C4, *Spartina* marshes in Fig. 2b, which show a substantial increase in low TOC sediments after reaching minimum values at approximately 5% TOC. At the vegetated sites, this may suggest that the enriched source material used by bacteria may be organic material, such as root exudates, released from seagrasses and C4 marsh plants (*Spartina* spp.), which are generally simple organic molecules that are readily available to bacteria. However, this explanation is not consistent with the pattern observed in the $\delta^{13}\text{C}_{i+a15:0}$ data presented in Fig. 3b. Especially for the unvegetated sediments, where a role for root exudates or other readily available macrophyte materials is unlikely, the data are also consistent with an important role for microphytobenthos-derived carbon as a carbon source for bacteria. Microphytobenthos in general has $\delta^{13}\text{C}$ signatures

generally more positive than those of the sediment TOC pool (typically between -20 and -13% , e.g. see France, 1995; Currin et al., 1995). The role of microphytobenthos as a carbon substrate for bacteria has been suggested previously in several studies (Boschker et al., 2000; Bouillon et al., 2004a; Cook et al., 2004), and this view is consistent with results from ^{13}C -labeling experiments where a very rapid transfer of microphytobenthos C to bacteria was found (Middelburg et al., 2000; our unpublished data from mangroves in Kenya).

It seems surprising that this selectivity is only expressed in the TOC poorest sediments where one would expect the microbial community to be more limited by organic carbon making preferential carbon degradation less likely. One possible explanation for this apparent paradox is that this selectivity is caused by an intrinsic higher degradability of a more labile carbon source, which is present in both low and high TOC sediments but in different relative abundances. Under such a scenario, the contrast in isotope signatures between the bulk TOC pool and that of the microbial community will per definition be more pronounced in settings where this labile C source makes up a more substantial part of the total TOC pool namely in low TOC sediments. Even if the difference in degradability between both sources is the same in both types of sediments, the much higher availability of less degradable C sources in higher TOC settings will dampen the resulting difference in $\delta^{13}\text{C}$ signatures between bulk TOC and microbial communities (i.e. $\Delta\delta$). For our data set, such a scenario would mean that the signal of a small pool of more easily degradable local materials, like root exudates or microphytobenthos, is more important for bacterial growth in low TOC sediments, whereas their signal in the bacteria is diluted in high TOC sediments with much higher concentrations of less degradable allochthonous sources.

The data from the C3 marshes in Schiermonnikoog (the Netherlands) do not follow the pattern as discussed above, as they show higher $\Delta\delta$ values compared the other data above a TOC of 1% (Fig. 5, although a cumulative sum analysis with or without the Schiermonnikoog data gave similar results). Since they are also distinct from the other C3 marsh data, we consider this to be a site-specific case for which we have no conclusive explanation. Unlike C4 marshes and seagrass beds, root exudation cannot be an explanation because the local C3 macrophyte material has a depleted signature. However, this marsh is situated next to the very extensive mud-flats of the Wadden Sea that are prone to wind induced erosion due to their long wind fetch. It has been shown that the seston in the tide water of the Wadden Sea contains high amounts of benthic diatoms eroded from the mud-flat surface, and benthic diatoms even dominate the plankton during high winds (de Jonge and van Beusekom, 1995). As sedimentation of suspended materials on salt marshes mainly occurs during high wind conditions, our data may indicate that the bacteria in the sediment of the Schiermonnikoog marsh thrive to a large extent on exceptionally high inputs of microphytobenthos material that was produced on the nearby mud-

flats. Another exception is the study by Cook et al. (2004) on a pristine intertidal mud-flat in Tasmania (data not shown in Fig. 5). Here the TOC was dominated by terrestrial C3 material, but the bacteria were probably mainly growing on material produced by benthic microalgae leading to $\Delta\delta$ ratios between 2 and 7‰.

4 Final remarks

4.1 Representativeness of the current data set

Despite the relatively large number of data compiled in this study (~340, from a variety of coastal systems), the question should be raised whether this dataset is sufficiently representative. For mangrove systems, we feel the data are likely to cover most types of settings, since they span the full range of %TOC and $\delta^{13}\text{C}_{\text{TOC}}$ encountered in the literature (e.g. see compilation in Bouillon et al., 2003a), but there are some less frequently encountered cases for which no $\delta^{13}\text{C}_{\text{C}_{i+a15:0}}$ are available. One such example is the situation described by Wooller et al. (2003), where high %TOC (29–36%) coincide with high $\delta^{13}\text{C}_{\text{TOC}}$ (–24.6 to –20.2‰) due to particularly large inputs of seagrass material. Similarly, considering the range of %TOC and $\delta^{13}\text{C}_{\text{TOC}}$ covered by our data (Fig. 2a), we can argue that *Spartina* marshes and seagrass systems are likely to be covered in a representative way. Data on C3 marshes are more scarce, however, and as discussed above, may be somewhat biased since the majority of data come from a single site with the possibility of an exceptionally high contribution of resuspended benthic diatoms in the tidal inputs.

Finally, we must stress that there is an almost complete lack of data on macroalgae-based systems, despite the fact that of all the vegetated coastal systems, they cover the largest surface area, and their integrated benthic mineralization rate ($247 \text{ Tmol C y}^{-1}$) is larger than that of the other systems combined ($208 \text{ Tmol C y}^{-1}$, Middelburg et al., 2004).

4.2 Distinction between local macrophyte production and terrestrial carbon

Terrestrial organic carbon sources transported to the coastal zone generally fall in two categories, C3 and C4-plant derived matter, each with a distinct and non-overlapping $\delta^{13}\text{C}$ range (typically ~–27 and –13‰, respectively). Since most of the data presented here come from regions where C3 vegetation dominates the catchment areas, we must keep in mind that terrestrial C and some local producers (C3 marshes and mangroves) are isotopically indistinguishable, and hence, that part of the C we ascribe to local macrophytes might in fact be terrestrial C (as part of the suspended matter pool deposited during inundation). One particular case, however, are the data from the Tana delta (northern Kenya), where a significant part of the catchment area is dominated by C4 grasslands, and where riverine suspended matter is comprised of

~50% C4-derived C (S. Bouillon, unpublished data). In this particular case, the contribution by C4-derived C is also reflected in the $\delta^{13}\text{C}_{\text{TOC}}$ data of the mangrove sediments (~–21‰) and in the $\delta^{13}\text{C}_{\text{C}_{i+a15:0}}$ data (~–25.5‰), which could indicate that this terrestrial carbon is a significant C source for sedimentary bacteria. It is not implausible that such a pattern is more widespread. Nevertheless, if the terrestrial C pool is derived from C3 vegetation, this would likely go undetected with the techniques used here except for seagrass systems and C4-marshes.

4.3 Conclusions and implications for carbon dynamics in the coastal zone

Our study clearly illustrates that mineralization in coastal sediments is often not fuelled by local macrophyte production and that in certain systems, bacteria may be selectively degrading more labile carbon sources such as microphytobenthos and carbon imported or settled from the water column. This may have implications for budgeting studies, since community respiration rates (where no source characterization is done) may overestimate the role of mineralization in the C budget of a particular ecosystems' production (e.g. see Bouillon et al., 2004b). Moreover, if such additional sources are preferentially mineralized, this implies that the fraction of carbon available for further export or long-term burial will differ in source, age, and composition to the total C pool available. Furthermore, these results may also have some implications for our understanding of N cycling in coastal systems, since the assimilated algae-derived material typically has a much higher N content than organic matter derived from vascular plants.

In summary, our cross-system analysis on bacterial carbon sources in near-coastal sediments demonstrates that:

1. $\delta^{13}\text{C}$ of bacterial PLFA show a large variability over the entire range of $\delta^{13}\text{C}_{\text{TOC}}$ data, indicating that in most settings, sedimentary bacteria may depend on C from various origins.
2. Systems where local macrophyte production is the major supplier of C for in situ decomposition are generally limited to organic carbon-rich sites (TOC>10%), which are likely to make up only a small part of the global areas of salt marsh and mangrove systems. In this respect, there appears to be a major difference in functioning between “open” and “closed” systems – the former with more pronounced exchange and subsidy of organic matter with adjacent systems.
3. In the majority of systems with ~1 to 10% TOC, bacterial PLFA $\delta^{13}\text{C}$ data indicate that non-macrophyte sources such as microphytobenthic production or imported carbon sources become important substrates but there is on average no apparent preferential use of the different source materials.

4. Only for sediments with less than ~1% TOC, bacteria clearly make preferential use of an isotopically heavy carbon source. These sediments were mostly from C4 *Spartina* marshes, seagrass beds and unvegetated sites suggesting that this heavy carbon source may be either root exudates from macrophytes or material produced by benthic diatoms. A similar effect was, however, also found for a C3-marsh where the macrophyte material is relatively depleted, suggesting that import or local production of microphytobenthos is a likely explanation.

Acknowledgements. Funding for this work was provided by the Fonds voor Wetenschappelijk Onderzoek (FWO-Vlaanderen, contract G.0118.02), EC-INCO project contract ERB IC18-CT98-0295, the NSF Microbial Observatories Project "Salt Marsh Microbes and Microbial Processes", the NIOZ-NIOO cooperation project and the NWO-VIDI grant of HTSB. SB is a postdoctoral fellow with the FWO-Vlaanderen. We are grateful to the following people for help during sample collection, fieldwork logistics, and analyses: I. De Mesel, A. Borges, J. Kairo, J. Bosire, A. Rao, F. Dahdouh-Guebas, V. Sharma, A. Wielemaker, J. de Brouwer, V. Créach, M. Steegstra, B. Schaub, S. Meger, H. van Wijnen, J. Bakker, M. Holmer, C. Hopkinson and J. Hobbie. B. Jones (Texas A&M University) kindly provided his raw data on sediment, macrophyte, and PLFA $\delta^{13}\text{C}$. M. Elskens has provided valuable advice on the statistical data treatment. A. Borges, F. Dehairs, D. Gillikin, P. Mangion, J. Middelburg, and N. Koedam provided useful comments on an earlier draft of this manuscript, and the manuscript benefited greatly from the constructive suggestions made by T. W. Lyons and two anonymous referees. This is publication 3778 Netherlands Institute of Ecology (NIOO-KNAW).

Edited by: T. W. Lyons

References

- Barrón, C., Marba, N., Terrados, J., Kennedy, H., and Duarte, C. M.: Community metabolism and carbon budget along a gradient of seagrass (*Cymodocea nodosa*) colonization, *Limnol. Oceanogr.*, 49, 1642–1651, 2004.
- Abraham, W. R., Hesse, C., and Pelz, O.: Ratios of carbon isotopes in microbial lipids as an indicator of substrate usage, *Appl. Environ. Microbiol.* 64, 4202–4209, 1998.
- Benner, R., Fogel, M. L., Sprague, E. K., and Hodson, R. E.: Depletion of ^{13}C in lignin and its implications for stable isotope studies, *Nature* 329, 708–710, 1987.
- Borges, A. V.: Do we have enough pieces of the jigsaw to integrate CO_2 fluxes in the coastal ocean?, *Estuaries*, 28, 3–27, 2005.
- Boschker, H. T. S., de Brouwer, J. F. C., and Cappenberg, T. E.: The contribution of macrophyte-derived organic matter to microbial biomass in salt-marsh sediments: stable carbon isotope analysis of microbial biomarkers, *Limnol. Oceanogr.*, 44, 309–319, 1999.
- Boschker, H. T. S., Wielemaker, A., Schaub, B. E. M., and Holmer, M.: Limited coupling of macrophyte production and bacterial carbon cycling in the sediments of *Zostera* spp. meadows, *Mar. Ecol. Prog. Ser.*, 203, 181–189, 2000.
- Boschker, H. T. S. and Middelburg, J. J.: Stable isotopes and biomarkers in microbial ecology, *FEMS Microbiol. Ecol.*, 40, 85–95, 2002.
- Boschker, H. T. S.: Linking microbial community structure and functioning: stable isotope (^{13}C) labeling in combination with PLFA analysis, in: *Molecular Microbial Ecology Manual II*, edited by: Kowalchuk, G. A., de Bruijn, F. J., Head, I. M., Akkermans, A. D., and van Elsas, J. D., Kluwer Academic Publishers, Dordrecht, The Netherlands, pp. 1673–1688, 2004.
- Boschker, H. T. S., Kromkamp, J., and Middelburg, J. J.: Biomarker and carbon isotopic constraints on bacterial and algal community structure and functioning in the highly polluted Scheldt estuary, *Limnol. Oceanogr.*, 50, 70–80, 2005.
- Bouillon, S., Dahdouh-Guebas, F., Rao, A. V. V. S., Koedam, N., and Dehairs, F.: Sources of organic carbon in mangrove sediments: variability and some possible implications for ecosystem functioning, *Hydrobiologia*, 495, 33–39, 2003a.
- Bouillon, S., Frankignoulle, M., Dehairs, F., Velimirov, B., Eiler, A., Etcheber, H., Abril, G., and Borges, A. V.: Inorganic and organic carbon biogeochemistry in the Gautami Godavari estuary (Andhra Pradesh, India) during pre-monsoon: the local impact of extensive mangrove forests, *Glob. Biogeochem. Cycles*, 17(4), 1114, doi:10.1029/2002GB002026, 2003b.
- Bouillon, S., Moens, T., Koedam, N., Dahdouh-Guebas, F., Baeyens, W., and Dehairs, F.: Variability in the origin of carbon substrates for bacterial communities in mangrove sediments, *FEMS Microbiol. Ecol.*, 49, 171–179, 2004a.
- Bouillon, S., Moens, T., and Dehairs, F.: Carbon sources sustaining benthic mineralization in mangrove and adjacent seagrass sediments (Gazi bay, Kenya), *Biogeosciences*, 1, 71–78, 2004b.
- Bowman, J. P. and McCuaig, R. D.: Biodiversity, community structural shifts, and biogeography of prokaryotes within Antarctic continental shelf sediment, *Appl. Environ. Microbiol.*, 69, 2463–2483, 2003.
- Cifuentes, L. A. and Salata, G. G.: Significance of carbon isotope discrimination between bulk carbon and extracted phospholipid fatty acids in selected terrestrial and marine environments, *Org. Geochem.*, 32, 613–621, 2001.
- Cook, P. L. M., Revill, A. R., Clementson, L. A., and Volkman, J. K.: Carbon and nitrogen cycling on intertidal mudflats of a temperate Australian estuary. III. Sources of organic matter, *Mar. Ecol. Prog. Ser.*, 280, 55–72, 2004.
- Currin, C. A., Newell, S. Y., and Paerl, H. W.: The role of standing dead spartina-alterniflora and benthic microalgae in salt-marsh food webs – considerations based on multiple stable-isotope analysis, *Mar. Ecol. Progr. Ser.*, 121, 99–116, 1995.
- Duarte, C. M., Middelburg, J. J., and Caraco, N.: Major role of marine vegetation on the oceanic carbon cycle, *Biogeosciences*, 2, 1–8, 2004.
- France, R. L.: C-13 Enrichment in benthic compared to planktonic algae – foodweb implications, *Mar. Ecol. Prog. Ser.*, 124, 307–312, 1995.
- Freeman, K. H., Hayes, J. M., Trendel, J. M., and Albrecht, P.: Evidence from carbon isotope measurements for diverse origins of sedimentary hydrocarbons, *Nature*, 343, 254–256, 1990.
- Fry, B. and Sherr, E. B.: $\delta^{13}\text{C}$ measurements as indicators of carbon flow in marine and freshwater ecosystems, *Contrib. Mar. Sci.*, 27, 13–47, 1984.

- Gattuso, J. P., Frankignoulle, M., and Wollast, R.: Carbon and carbonate metabolism in coastal aquatic ecosystems, *Annu. Rev. Ecol. Syst.*, 29, 405–434, 1998.
- Harvey, H. R., Ederington, M. C., and McManus, G. B.: Lipid composition of the marine ciliates *Pleuronema* sp. and *Fabrea salina*: shifts in response to diet, *J. Eukar. Microbiol.*, 44, 189–193, 1997.
- Hayes, J. M.: Fractionation of carbon and hydrogen isotopes in biosynthetic processes, in: *Organic geochemistry of contemporary and ancient sediments*, edited by: Valley, J. S. and Cole, D.R., Mineralogical Society of America, 225–277, 2001.
- Hee, C. A., Pease, T. K., Alperin, M. J., and Martens, C. S.: Dissolved organic carbon production and consumption in anoxic marine sediments: A pulsed-tracer experiment, *Limnol. Oceanogr.*, 46, 1908–1920, 2001.
- Hemminga, M. A. and Mateo, M. A.: Stable carbon isotopes in seagrasses: variability in ratios and use in ecological studies, *Mar. Ecol. Prog. Ser.*, 140, 285–298, 1996.
- Holmer, M., Andersen, F.O., Nielsen, S.L., and Boschker, H.T.S.: The importance of mineralization based on sulfate reduction for nutrient regeneration in tropical seagrass sediments, *Aquat. Bot.* 71, 1–17, 2001.
- Holmer, M., Duarte, C. M., Boschker, H. T. S., and Barrón, C.: Carbon cycling and bacterial carbon sources in pristine and impacted Mediterranean seagrass sediments, *Aquat. Microb. Ecol.*, 36, 227–237, 2004.
- Hullar, M. A. J., Fry, B., Peterson, B. J., and Wright, R. T.: Microbial utilization of estuarine dissolved organic carbon: A stable isotope tracer approach tested by mass balance, *Appl. Environ. Microbiol.*, 62, 2489–2493, 1996.
- Jones, W. B., Cifuentes, L. A., and Kaldy, J. E.: Stable carbon isotope evidence for coupling between sedimentary bacteria and seagrasses in a sub-tropical lagoon, *Mar. Ecol. Prog. Ser.*, 255, 15–25, 2003.
- Kaneda, T.: Iso-fatty and anteiso-fatty acids in bacteria - biosynthesis, function, and taxonomic significance. *Microbiol. Rev.* 55, 288–302, 1991.
- Keil, R. G., Monlucon, C. B., Prahl, G. F., and Hedges, J. I.: Sorptive preservation of labile organic matter in marine sediments, *Nature*, 370, 549–551, 1994.
- Kennedy, H., Gacia, E., Kennedy, D. P., Papadimitriou, S., and Duarte, C. M.: Organic carbon sources to SE Asian coastal sediments, *Estuar. Coast. Shelf. Sci.*, 60, 59–68, 2004.
- McAllister, S. L., Bauer, J. E., Cherrier, J. E., and Ducklow, H. W.: Assessing sources and ages of organic matter supporting river and estuarine bacterial production: a multiple-isotope ($\Delta^{14}\text{C}$, $\delta^{13}\text{C}$, and $\delta^{15}\text{N}$) approach, *Limnol. Oceanogr.*, 49, 1687–1702, 2004.
- Middelburg, J. J.: A simple rate model for organic-matter decomposition in marine-sediments, *Geochim. Cosmochim. Acta*, 53, 1577–1581, 1989.
- Middelburg, J. J., Barranguet, C., Boschker, H. T. S., Herman, P. M. J., Moens, T., and Heip, C. H. R.: The fate of intertidal microphytobenthos carbon: an in situ ^{13}C labeling study, *Limnol. Oceanogr.*, 45, 1224–1234, 2000.
- Middelburg, J. J., Duarte, C. M., and Gattuso, J. P.: Respiration in coastal benthic communities, in: *Respiration in aquatic ecosystems*, edited by: del Giorgio, P. A. and Williams, P. J. le B., Oxford University Press, 2004.
- Middelburg, J. J., Nieuwenhuize, J., Lubberts, R. K., and van de Plassche, O.: Organic carbon isotope systematics of coastal marshes, *Estuar. Coast. Shelf. Sci.*, 45, 681–687, 1997.
- Pelz, O., Hesse, C., Tesar, M., Coffin, R. B., and Abraham, W. R.: Development of methods to measure carbon isotope ratios of bacterial biomarkers in the environment, *Isotopes Environ. Health Stud.*, 33, 131–144, 1997.
- Raymond, P. A. and Bauer, J. E.: DOC cycling in a temperate estuary: a mass balance approach using natural ^{14}C and ^{13}C isotopes, *Limnol. Oceanogr.*, 46, 655–667, 2001.
- Schoenberg, S. A., Benner, R., Armstrong, A., Sobecky, P., and Hodson, R. E.: Effects of acid stress on aerobic decomposition of algal and aquatic macrophyte detritus – direct comparison in a radiocarbon assay, *Appl. Environ. Microbiol.*, 56, 237–244, 1990.
- Teece, M. A., Fogel, M. L., Dollhopf, M. E., and Nealson, K. H.: Isotopic fractionation associated with biosynthesis of fatty acids by a marine bacterium under oxic and anoxic conditions, *Org. Geochem.*, 30, 1571–1579, 1999.
- Volkman, J. K., Jeffrey, S. W., Nichols, P. D., Rogers, G. I., and Garland, C. D.: Fatty acids and lipid compositions of 10 species of microalgae used in mariculture, *J. Exp. Mar. Biol. Ecol.*, 128, 219–240, 1989.
- Wooller, M., Smallwood, B., Jacobson, M., and Fogel, M.: Carbon and nitrogen stable isotopic variation in *Laguncularia racemosa* from Florida and Belize: implications for trophic level studies, *Hydrobiologia*, 499, 13–23, 2003.

Electronic Supplement 1

Bouillon and Boschker

Bacterial carbon sources in coastal sediments: a review based on stable isotope data of biomarkers.

Site location	Ecosystem type	Depth range (cm)	% TOC	C/N (molar)	$\delta^{13}\text{C}_{\text{TOC}}$	$\delta^{13}\text{C}_{\text{macrophyte}}$	$\delta^{13}\text{C}_{\text{Frat15.0}}$	$\delta^{13}\text{C}_{\text{Frat15.0}} - \delta^{13}\text{C}_{\text{TOC}}$	Data source
Waarde marsh (the Netherlands)	C3 Marsh	0-8	0.8	13.2	-24.3	-24.8	-28.3	-4.1	Boschker et al (1999)
Waarde marsh (the Netherlands)	C3 Marsh	0-8	1.2	15.3	-25.5	-24.8	-27.7	-2.3	Boschker et al (1999)
Waarde marsh (the Netherlands)	C3 Marsh	0-8	0.9	12.4	-24.3	-24.8	-28.0	-3.8	Boschker et al (1999)
Waarde marsh (the Netherlands)	C3 Marsh	0-8	1.4	14.2	-24.4	-24.8	-23.4	1.0	Boschker et al (1999)
Waarde marsh (the Netherlands)	C3 Marsh	0-8	1.4	14.2	-24.4	-24.8	-26.5	-2.1	Boschker et al (1999)
Waarde marsh (the Netherlands)	C3 Marsh	0-8	1.4	14.2	-24.4	-24.8	-23.4	1.0	Boschker et al (1999)
Waarde marsh (the Netherlands)	C3 Marsh	0-8	1.8	14.2	-24.4	-24.8	-25.6	-1.2	Boschker et al (1999)
Waarde marsh (the Netherlands)	C3 Marsh	0-8	1.8	16.3	-24.5	-24.8	-27.9	-3.4	Boschker et al (1999)
Waarde marsh (the Netherlands)	C3 Marsh	0-8	1.3	11.7	-24.4	-24.8	-23.8	0.6	Boschker et al (1999)
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	0.3	15.6	-26.5	-27.5	-25.3	1.2	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	1.0	18.5	-25.5	-26.5	-24.9	0.6	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	0.9	13.5	-24.9	-28.5	-26.9	-2.0	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	0.7	15.6	-26.1	-26.3	-25.1	0.9	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	0.1	13.7	-26.9	-28.5	-24.4	2.5	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	0.9	16.0	-26.1	-26.6	-24.3	1.8	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	1.6	9.6	-25.5	-26.1	-24.0	1.5	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	1.2	16.3	-25.2	-26.1	-24.3	0.9	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	1.0	14.7	-25.4	-27.3	-23.5	1.9	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	1.0	11.6	-24.1	-26.4	-23.0	1.1	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	0.7	12.1	-22.5	-25.1	-22.3	0.1	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	1.4	13.5	-26.0	-26.7	-24.0	2.0	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	2.6	13.7	-26.0	-27.0	-24.6	1.4	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	7.8	14.7	-25.3	-26.5	-24.9	0.4	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	5.7	16.3	-26.4	-27.4	-25.5	0.9	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	4.0	12.0	-24.0	-25.5	-23.9	0.1	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	3.3	11.5	-23.2	-26.1	-22.6	0.6	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	1.8	13.7	-25.9	-25.9	-23.2	2.7	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	2.7	14.6	-26.0	-26.4	-22.8	3.1	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	8.7	14.2	-25.5	-25.6	-24.5	1.0	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	3.1	14.0	-25.6	-26.7	-23.9	1.7	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	4.4	12.8	-24.3	-24.9	-25.0	-0.7	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	5.9	13.5	-24.7	-25.6	-25.5	-0.8	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	6.6	13.1	-25.4	-24.9	-23.8	1.7	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	7.2	13.6	-25.0	-25.2	-23.3	1.7	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	6.7	13.9	-25.2	-25.7	-23.4	1.7	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	11.2	16.5	-25.1	-26.3	-23.2	1.9	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	3.1	11.9	-23.6	-25.5	-23.6	0.0	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	4.2	12.4	-23.6	-24.5	-22.8	0.7	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	2.3	13.3	-23.3	-26.3	-22.1	1.2	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	3.3	12.6	-23.5	-25.9	-22.6	0.9	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	5.6	14.0	-25.5	-26.1	-23.3	2.2	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	7.3	14.7	-25.1	-25.9	-23.9	1.2	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	3.5	13.4	-25.1	-24.8	-22.3	2.8	This study
Schiermonnikoog (the Netherlands)	C3 Marsh	0-10	4.6	14.8	-25.0	-25.9	-22.9	2.2	This study
North River (USA)	C3 Marsh				-25.3	-25.9	-33.9	-8.6	Cifuentes and Salata (2001)

Site location	Ecosystem type	Depth range (cm)	% TOC	C/N (molar)	$\delta^{13}\text{C}_{\text{TOC}}$	$\delta^{13}\text{C}_{\text{macrophyte}}$	$\delta^{13}\text{C}_{\text{P+I+S+O}}$	$\delta^{13}\text{C}_{\text{P+I+S+O}} - \delta^{13}\text{C}_{\text{TOC}}$	Data source
North River (USA)	C3 Marsh				-22.3	-25.9	-27.5	-5.2	Cifuentes and Salata (2001)
Schiermonnikoog (the Netherlands)	C3 marsh, unvegetated	0-10	0.1	5.8	-19.2		-17.6	1.7	This study
Schiermonnikoog (the Netherlands)	C3 marsh, unvegetated	0-10	0.1	7.1	-20.9		-17.9	3.0	This study
Schiermonnikoog (the Netherlands)	C3 marsh, unvegetated	0-10	0.1	6.4	-21.1		-18.3	2.8	This study
Schiermonnikoog (the Netherlands)	C3 marsh, unvegetated	0-10	0.2	7.5	-21.7		-16.9	4.8	This study
Schiermonnikoog (the Netherlands)	C3 marsh, unvegetated	0-10	0.1	9.1	-21.1		-17.0	4.1	This study
Schiermonnikoog (the Netherlands)	C3 marsh, unvegetated	0-10	0.1	6.2	-21.9		-17.4	4.5	This study
North River (USA)	C3 marsh, unvegetated				-20.5		-24.3	-3.8	Cifuentes and Salata (2001)
Waarde marsh (the Netherlands)	C4 marsh	0-8	2.0	14.0	-23.4	-12.7	-24.9	-1.5	Boschker et al. (1999)
Waarde marsh (the Netherlands)	C4 marsh	0-8	1.9	13.3	-22.7	-12.7	-26.1	-3.4	Boschker et al. (1999)
Waarde marsh (the Netherlands)	C4 marsh	0-8	2.2	15.4	-22.9	-12.7	-26.4	-3.5	Boschker et al. (1999)
Waarde marsh (the Netherlands)	C4 marsh	0-8	1.8	15.5	-22.2	-12.7	-24.4	-2.2	Boschker et al. (1999)
Waarde marsh (the Netherlands)	C4 marsh	0-8	1.8	15.5	-22.2	-12.7	-24.3	-2.1	Boschker et al. (1999)
Waarde marsh (the Netherlands)	C4 marsh	0-8	1.8	15.5	-22.2	-12.7	-24.0	-1.8	Boschker et al. (1999)
Waarde marsh (the Netherlands)	C4 marsh	0-8	1.8	15.5	-22.2	-12.7	-23.3	-1.1	Boschker et al. (1999)
Waarde marsh (the Netherlands)	C4 marsh	0-8	1.8	18.0	-20.0	-12.7	-23.6	-3.6	Boschker et al. (1999)
Waarde marsh (the Netherlands)	C4 marsh	0-8	2.0	13.9	-23.1	-12.7	-24.3	-1.2	Boschker et al. (1999)
Kattedijke (the Netherlands)	C4 marsh	0-8	0.4	14.0	-17.9	-12.8	-21.7	-3.8	Boschker et al. (1999)
Kattedijke (the Netherlands)	C4 marsh	0-8	0.7	15.5	-21.1	-12.8	-22.5	-1.4	Boschker et al. (1999)
Kattedijke (the Netherlands)	C4 marsh	0-8	0.1	8.3	-20.6	-12.8	-19.6	0.9	Boschker et al. (1999)
Ritthem (the Netherlands)	C4 marsh	0-10	0.1		-21.4	-13.2	-23.1	-1.7	This study
Ritthem (the Netherlands)	C4 marsh	0-10	0.1		-20.5	-13.2	-21.0	-0.5	This study
Ritthem (the Netherlands)	C4 marsh	0-10	1.0	11.3	-18.0	-13.2	-20.7	-2.7	This study
Ritthem (the Netherlands)	C4 marsh	0-10	1.1	18.4	-16.2	-13.2	-21.4	-5.2	This study
Ritthem (the Netherlands)	C4 marsh	0-10	1.9	14.2	-20.4	-13.2	-24.0	-3.6	This study
Valkenisse (the Netherlands)	C4 marsh	0-10	0.1	10.1	-19.6	-13.2	-17.1	2.5	This study
Valkenisse (the Netherlands)	C4 marsh	0-10	0.8	16.8	-23.7	-13.2	-23.2	0.5	This study
Mont Saint Michel (France)	C4 marsh	0-10	1.4		-19.5	-13.2	-18.5	1.0	This study
Mont Saint Michel (France)	C4 marsh	0-10	1.0		-19.3	-13.2	-17.8	1.5	This study
Mont Saint Michel (France)	C4 marsh	0-10	1.3		-19.0	-13.2	-17.1	1.9	This study
Great Marshes (USA)	C4 marsh	0-10	26.2		-14.1	-12.5	-15.9	-1.8	Boschker et al. (1999)
Great Marshes (USA)	C4 marsh	0-10	36.7		-13.4	-12.5	-18.0	-4.5	Boschker et al. (1999)
Great Marshes (USA)	C4 marsh	0-10	21.4		-14.5	-12.5	-16.4	-1.9	Boschker et al. (1999)
North Inlet (USA)	C4 marsh	0-10	4.9	19.3	-17.4	-13.2	-20.3	-2.9	This study
North Inlet (USA)	C4 marsh	0-10	5.1	26.0	-16.1	-13.2	-20.3	-4.2	This study
Canary Creek (USA)	C4 marsh	0-10	25.0	38.9	-13.7	-13.2	-19.8	-6.1	This study
Canary Creek (USA)	C4 marsh	0-10	27.7	28.4	-15.0	-13.2	-20.4	-5.4	This study
Plum Island Sound (USA)	C4 marsh	0-10	6.2	17.9	-17.6	-13.2	-20.9	-3.3	This study
Plum Island Sound (USA)	C4 marsh	0-10	6.9	17.1	-17.0	-13.2	-22.0	-5.0	This study
Plum Island Sound (USA)	C4 marsh	0-10	4.5	16.4	-20.0	-13.5	-24.1	-4.0	This study
Plum Island Sound (USA)	C4 marsh	0-10	4.7	16.5	-20.2	-13.3	-23.3	-3.1	This study
Plum Island Sound (USA)	C4 marsh	0-10	5.4	14.5	-20.2	-13.0	-23.4	-3.2	This study
Plum Island Sound (USA)	C4 marsh	0-10	4.7	14.0	-20.8	-12.5	-22.2	-1.5	This study
Plum Island Sound (USA)	C4 marsh	0-10	4.3	14.9	-17.9	-13.0	-21.6	-3.6	This study
Plum Island Sound (USA)	C4 marsh	0-10	5.1	15.3	-18.5	-12.5	-18.2	0.4	This study
Plum Island Sound (USA)	C4 marsh	0-10	9.6	18.1	-16.6	-13.6	-21.4	-4.8	This study
Plum Island Sound (USA)	C4 marsh	0-10	8.6	15.1	-17.1	-13.8	-19.7	-2.6	This study
Plum Island Sound (USA)	C4 marsh	0-10	0.5	26.1	-19.8	-13.1	-21.7	-1.9	This study
Plum Island Sound (USA)	C4 marsh	0-10	0.5	21.2	-19.1	-13.2	-20.1	-1.0	This study
Plum Island Sound (USA)	C4 marsh	0-10	2.5	15.7	-17.7	-12.9	-22.1	-4.4	This study

Site location	Ecosystem type	Depth range (cm)	% TOC	C/N (molar)	$\delta^{13}\text{C}_{\text{TOC}}$	$\delta^{13}\text{C}_{\text{macrophyte}}$	$\delta^{13}\text{C}_{\text{C}_{15:0}}$	$\delta^{13}\text{C}_{\text{C}_{15:0}} - \delta^{13}\text{C}_{\text{TOC}}$	Data source
Sint Annaland (the Netherlands)	C4 marsh	0-10	1.1	11.1	-21.3	-12.1	-20.6	0.7	This study
Sint Annaland (the Netherlands)	C4 marsh	0-10	0.9	10.0	-22.2	-12.3	-23.9	-1.7	This study
Ritthem (the Netherlands)	C4 marsh	0-2	4.1	11.2	-18.8	-13.2	-21.2	-2.4	This study
Ritthem (the Netherlands)	C4 marsh	8-10	1.0	15.6	-16.6	-13.2	-19.1	-2.6	This study
Ritthem (the Netherlands)	C4 marsh	0-2	4.6	11.7	-18.4	-13.2	-21.8	-3.5	This study
Ritthem (the Netherlands)	C4 marsh	8-10	0.6	15.7	-17.3	-13.2	-17.9	-0.6	This study
Ritthem (the Netherlands)	C4 marsh	0-2	4.9	10.3	-19.7	-13.2	-20.5	-0.8	This study
Ritthem (the Netherlands)	C4 marsh	8-10	5.2	11.3	-20.5	-13.2	-23.2	-2.7	This study
Ritthem (the Netherlands)	C4 marsh	0-2	5.8	13.2	-16.8	-13.2	-21.1	-4.3	This study
Ritthem (the Netherlands)	C4 marsh	8-10	5.1	12.1	-20.2	-13.2	-21.2	-1.0	This study
Waarde marsh (the Netherlands)	C4 marsh	0-2	2.4	19.6	-21.6	-13.2	-24.1	-2.5	This study
Waarde marsh (the Netherlands)	C4 marsh	8-10	1.6	18.3	-25.0	-13.2	-30.0	-5.0	This study
Waarde marsh (the Netherlands)	C4 marsh	0-2	1.4	14.8	-19.8	-13.2	-22.3	-2.5	This study
Waarde marsh (the Netherlands)	C4 marsh	8-10	1.5	16.7	-17.7	-13.2	-25.7	-7.9	This study
Waarde marsh (the Netherlands)	C4 marsh	0-2	2.3	13.7	-22.3	-13.2	-23.2	-1.0	This study
Waarde marsh (the Netherlands)	C4 marsh	8-10	0.3	18.9	-17.4	-13.2	-27.2	-9.8	This study
Waarde marsh (the Netherlands)	C4 marsh	0-2	2.1	13.9	-23.9	-13.2	-24.2	-0.2	This study
Waarde marsh (the Netherlands)	C4 marsh	8-10	2.4	25.3	-19.1	-13.2	-24.3	-5.2	This study
North River (USA)	C4 marsh				-18.1	-13.2	-21.4	-3.3	Cifuentes and Salata (2001)
North River (USA)	C4 marsh				-19.4	-13.2	-21.8	-2.4	Cifuentes and Salata (2001)
Waarde marsh (the Netherlands)	C4 marsh, unvegetated	0-8	0.5	14.7	-25.7		-25.9	-0.2	Boschker et al. (1999)
Waarde marsh (the Netherlands)	C4 marsh, unvegetated	0-8	0.5	13.5	-25.8		-25.5	0.3	Boschker et al. (1999)
Waarde marsh (the Netherlands)	C4 marsh, unvegetated	0-8	0.6	13.7	-26.7		-25.1	1.7	Boschker et al. (1999)
Waarde marsh (the Netherlands)	C4 marsh, unvegetated	0-8	0.7	13.5	-25.7		-23.8	1.9	Boschker et al. (1999)
Waarde marsh (the Netherlands)	C4 marsh, unvegetated	0-8	0.7	13.5	-25.7		-22.8	2.9	Boschker et al. (1999)
Waarde marsh (the Netherlands)	C4 marsh, unvegetated	0-8	0.7	13.5	-25.7		-23.0	2.7	Boschker et al. (1999)
Waarde marsh (the Netherlands)	C4 marsh, unvegetated	0-8	0.7	13.5	-25.7		-22.2	3.5	Boschker et al. (1999)
Waarde marsh (the Netherlands)	C4 marsh, unvegetated	0-8	1.1	13.8	-24.4		-22.9	1.6	Boschker et al. (1999)
Waarde marsh (the Netherlands)	C4 marsh, unvegetated	0-8	0.6	11.8	-24.9		-23.1	1.8	Boschker et al. (1999)
Kattedijke (the Netherlands)	C4 marsh, unvegetated	0-8	0.1	8.6	-20.3		-28.2	-8.0	Boschker et al. (1999)
Kattedijke (the Netherlands)	C4 marsh, unvegetated	0-8	0.2	11.4	-19.7		-27.1	-7.4	Boschker et al. (1999)
Ritthem (the Netherlands)	C4 marsh, unvegetated	0-10	0.1		-22.7		-19.8	2.9	This study
Ritthem (the Netherlands)	C4 marsh, unvegetated	0-10	0.1	8.4	-21.4		-19.1	2.3	This study
Ritthem (the Netherlands)	C4 marsh, unvegetated	0-10	0.2	9.0	-20.0		-19.1	0.9	This study
Ritthem (the Netherlands)	C4 marsh, unvegetated	0-10	0.2	7.7	-20.5		-19.4	1.1	This study
Valkenisse (the Netherlands)	C4 marsh, unvegetated	0-10	0.1	7.6	-24.2		-19.6	4.6	This study
Valkenisse (the Netherlands)	C4 marsh, unvegetated	0-10	0.8	15.9	-26.0		-22.2	3.8	This study
Great Marshes (USA)	C4 marsh, unvegetated	0-10	0.5	10.1	-18.5		-22.8	-4.3	Boschker et al (1999)
Great Marshes (USA)	C4 marsh, unvegetated	0-10	1.9	12.4	-17.1		-19.5	-2.4	Boschker et al (1999)
North Inlet (USA)	C4 marsh, unvegetated	0-10	1.0	13.2	-17.5		-16.9	0.6	This study
North Inlet (USA)	C4 marsh, unvegetated	0-10	0.8	12.7	-19.5		-17.3	2.2	This study
Canary Creek (USA)	C4 marsh, unvegetated	0-10	7.0	15.2	-17.8		-21.4	-3.6	This study
Canary Creek (USA)	C4 marsh, unvegetated	0-10	7.5	17.1	-17.6		-20.2	-2.6	This study
Plum Island Sound (USA)	C4 marsh, unvegetated	0-10	1.9	19.4	-19.8		-19.5	0.3	This study
Plum Island Sound (USA)	C4 marsh, unvegetated	0-10	1.7	18.9	-19.7		-21.4	-1.7	This study
Plum Island Sound (USA)	C4 marsh, unvegetated	0-10	4.1	15.3	-20.9		-25.5	-4.6	This study
Plum Island Sound (USA)	C4 marsh, unvegetated	0-10	3.9	16.6	-20.7		-25.8	-5.1	This study
Plum Island Sound (USA)	C4 marsh, unvegetated	0-10	1.0	20.3	-20.1		-22.3	-2.2	This study

Site location	Ecosystem type	Depth range (cm)	% TOC	C/N (molar)	$\delta^{13}\text{C}_{\text{TOC}}$	$\delta^{13}\text{C}_{\text{macrophyte}}$	$\delta^{13}\text{C}_{\text{C}_{15-0}}$	$\delta^{13}\text{C}_{\text{C}_{15-0}} - \delta^{13}\text{C}_{\text{TOC}}$	Data source
Plum Island Sound (USA)	C4 marsh, unvegetated	0-10	1.1	20.4	-20.3		-21.9	-1.6	This study
Plum Island Sound (USA)	C4 marsh, unvegetated	0-10	2.9	17.1	-18.8		-21.7	-2.9	This study
Plum Island Sound (USA)	C4 marsh, unvegetated	0-10	2.8	16.8	-18.5		-21.5	-3.0	This study
Plum Island Sound (USA)	C4 marsh, unvegetated	0-10	0.2	14.3	-18.1		-22.0	-3.9	This study
Plum Island Sound (USA)	C4 marsh, unvegetated	0-10	0.3	17.2	-19.4		-22.5	-3.1	This study
Ritthem (the Netherlands)	C4 marsh, unvegetated	0-2	0.2	11.1	-17.5		-18.9	-1.4	This study
Ritthem (the Netherlands)	C4 marsh, unvegetated	2-8	0.2	15.8	-24.4		-21.1	3.3	This study
Ritthem (the Netherlands)	C4 marsh, unvegetated	0-2	0.2	10.3	-18.4		-18.3	0.1	This study
Ritthem (the Netherlands)	C4 marsh, unvegetated	2-8	0.1	12.5	-22.8		-21.3	1.5	This study
Waarde marsh (the Netherlands)	C4 marsh, unvegetated	0-2	0.4	15.7	-25.3		-23.0	2.4	This study
Waarde marsh (the Netherlands)	C4 marsh, unvegetated	2-8	0.1	16.9	-24.6		-25.1	-0.5	This study
Waarde marsh (the Netherlands)	C4 marsh, unvegetated	0-2	0.4	18.2	-25.5		-23.8	1.7	This study
Waarde marsh (the Netherlands)	C4 marsh, unvegetated	2-8	0.3	21.4	-26.2		-25.4	0.8	This study
Mallorca (Spain)	Macro alg	0-5	3.7	12.5	-15.2	-13.9	-18.3	-3.1	Holmer et al. (2004)
Mallorca (Spain)	Macro alg	0-5	4.3	15.3	-15.2	-13.9	-19.1	-3.9	Holmer et al. (2004)
Mallorca (Spain)	Macro alg	0-5	4.1	13.7	-15.3		-19.5	-4.2	Holmer et al. (2004)
Mallorca (Spain)	Macro alg	0-5	4.1	12.9	-16.4		-20.3	-3.9	Holmer et al. (2004)
Tana estuary (Kenya)	Mangrove	0-1	2.2	15.2	-25.3	-28.2	-27.0	-1.6	This study
Tana estuary (Kenya)	Mangrove	1-2	2.6	16.6	-25.5	-28.2	-27.9	-2.4	This study
Tana estuary (Kenya)	Mangrove	2-4	2.4	16.3	-25.8	-28.2	-28.7	-2.8	This study
Tana estuary (Kenya)	Mangrove	4-10	2.2	16.0	-25.8	-28.2	-29.0	-3.2	This study
Tana delta (Kenya)	Mangrove	0-1	1.1	14.4	-22.7	-28.2	-24.5	-1.8	This study
Tana delta (Kenya)	Mangrove	1-2	1.0	14.0	-22.5	-28.2	-26.6	-4.1	This study
Tana delta (Kenya)	Mangrove	2-4	1.0	14.0	-22.8	-28.2	-27.2	-4.4	This study
Tana delta (Kenya)	Mangrove	4-10	1.1	13.2	-22.6	-28.2	-27.1	-4.6	This study
Tana delta (Kenya)	Mangrove	0-1	1.2	11.9	-21.7	-28.2	-25.3	-3.7	This study
Tana delta (Kenya)	Mangrove	1-2	1.1	12.3	-21.8	-28.2	-26.0	-4.1	This study
Tana delta (Kenya)	Mangrove	2-4	1.1	11.6	-21.5	-28.2	-25.6	-4.1	This study
Tana delta (Kenya)	Mangrove	4-10	1.1	11.8	-22.2	-28.2	-26.9	-4.7	This study
Tana delta (Kenya)	Mangrove	0-1	1.2	12.3	-20.4	-28.2	-23.3	-2.8	This study
Tana delta (Kenya)	Mangrove	1-2	1.2	12.7	-20.6	-28.2	-24.2	-3.6	This study
Tana delta (Kenya)	Mangrove	2-4	1.1	12.4	-20.8	-28.2	-24.2	-3.4	This study
Gazi bay (Kenya)	Mangrove	0-1	10.3	21.5	-25.2	-29.3	-28.8	-3.6	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	1-2	10.2	19.4	-25.2	-29.3	-28.8	-3.6	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	2-4	8.6	19.6	-25.3	-29.3	-28.8	-3.5	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	4-10	9.2	18.7	-25.1	-29.3	-29.8	-4.7	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	1-2	1.6	19.1	-25.2	-29.3	-27.4	-2.2	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	2-4	2.0	18.8	-25.5	-29.3	-28.4	-3.0	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	4-10	3.2	18.4	-25.6	-29.3	-27.9	-2.3	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	0-1	11.8	18.7	-25.3	-29.3	-26.7	-1.4	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	1-2	14.0	18.1	-24.9	-29.3	-26.9	-2.0	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	2-4	7.5	16.3	-25.1	-29.3	-27.2	-2.1	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	4-10	5.5	19.1	-25.0	-29.3	-27.8	-2.8	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	0-1	0.8	23.1	-25.7	-26.8	-27.3	-1.6	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	1-2	0.6	25.8	-25.6	-26.8	-29.3	-3.7	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	2-4	0.7	17.1	-25.8	-26.8	-28.9	-3.1	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	4-10	1.2	16.2	-25.8	-26.8	-31.9	-6.2	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	0-1	1.4	19.4	-25.2	-27.1	-27.6	-2.4	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	1-2	2.5	22.3	-26.5	-27.1	-27.6	-1.1	Bouillon et al. (2004b)

Site location	Ecosystem type	Depth range (cm)	% TOC	C/N (molar)	$\delta^{13}\text{C}_{\text{TOC}}$	$\delta^{13}\text{C}_{\text{macrophyte}}$	$\delta^{13}\text{C}_{\text{C}_{15-20}}$	$\delta^{13}\text{C}_{\text{C}_{15-20}} - \delta^{13}\text{C}_{\text{TOC}}$	Data source
Gazi bay (Kenya)	Mangrove	2-4	2.2	19.5	-26.2	-27.1	-32.4	-6.2	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	4-10	1.4	22.7	-26.4	-27.1	-32.6	-6.2	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	0-1	13.7	17.0	-24.3	-28.2	-26.1	-1.8	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	1-2	13.0	17.4	-24.6	-28.2	-28.6	-3.9	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	2-4	14.6	19.1	-24.7	-28.2	-29.2	-4.5	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	4-10	13.5	20.2	-25.1	-28.2	-35.6	-10.4	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	0-1	2.9	18.0	-25.6	-28.2	-28.4	-2.8	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	1-2	3.5	20.6	-26.0	-28.2	-33.3	-7.3	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	2-4	2.2	22.7	-26.1	-28.2	-34.0	-8.0	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	4-10	3.9	23.2	-26.4	-28.2	-35.1	-8.7	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	0-1	1.2	15.3	-22.1	-29.1	-27.7	-5.6	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	1-2	2.0	16.8	-24.1	-29.1	-31.7	-7.6	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	2-4	2.3	19.4	-24.8	-29.1	-29.3	-4.5	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	4-10	2.7	19.9	-24.9	-29.1	-33.9	-9.1	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	0-1	4.8	26.6	-24.0	-29.1	-27.5	-3.6	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	1-2	2.6	21.1	-23.9	-29.1	-31.1	-7.2	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	2-4	3.1	19.7	-23.8	-29.1	-28.9	-5.1	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	4-10	2.5	19.7	-24.5	-29.1	-32.9	-8.4	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	0-1	6.1	16.2	-25.2	-29.3	-30.1	-4.9	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	1-2	7.6	19.8	-25.2	-29.3	-30.4	-5.2	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	2-4	8.1	17.8	-25.3	-29.3	-31.3	-6.0	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	4-10	5.9	19.0	-25.5	-29.3	-32.2	-6.7	Bouillon et al. (2004b)
Gazi bay (Kenya)	Mangrove	0-1	12.5	14.4	-24.4	-28.2	-28.1	-3.7	Bouillon et al. (2004b)
Pichavaram (India)	Mangrove	0-1	1.3	15.0	-24.3	-28.2	-26.3	-2.1	Bouillon et al. (2004a)
Pichavaram (India)	Mangrove	1-2	1.2	10.1	-24.4	-28.2	-25.3	-0.9	Bouillon et al. (2004a)
Pichavaram (India)	Mangrove	2-4	1.1	10.1	-22.8	-28.2	-25.6	-2.7	Bouillon et al. (2004a)
Pichavaram (India)	Mangrove	4-10	1.0	12.6	-22.4	-28.2	-26.1	-3.7	Bouillon et al. (2004a)
Pichavaram (India)	Mangrove	0-1	6.9	15.5	-26.2	-28.2	-27.9	-1.7	Bouillon et al. (2004a)
Pichavaram (India)	Mangrove	1-2	5.8	13.8	-26.0	-28.2	-28.5	-2.5	Bouillon et al. (2004a)
Pichavaram (India)	Mangrove	2-4	6.3	18.5	-25.9	-28.2	-28.3	-2.4	Bouillon et al. (2004a)
Pichavaram (India)	Mangrove	4-10	2.5	13.6	-25.1	-28.2	-27.2	-2.1	Bouillon et al. (2004a)
Chunnambar (India)	Mangrove	0-1	0.6	11.1	-23.3	-28.2	-23.6	-0.3	Bouillon et al. (2004a)
Chunnambar (India)	Mangrove	1-3	0.6	12.6	-23.8	-28.2	-27.8	-4.0	Bouillon et al. (2004a)
Chunnambar (India)	Mangrove	3-5	0.4	8.8	-24.5	-28.2	-29.3	-4.8	Bouillon et al. (2004a)
Pambala (Sri Lanka)	Mangrove	0-1	34.8	17.2	-28.4	-30.6	-31.4	-3.0	Bouillon et al. (2004a)
Pambala (Sri Lanka)	Mangrove	1-2	28.2	17.7	-28.6	-30.6	-32.1	-3.5	Bouillon et al. (2004a)
Pambala (Sri Lanka)	Mangrove	2-4	32.3	21.7	-28.1	-30.6	-32.6	-4.5	Bouillon et al. (2004a)
Pambala (Sri Lanka)	Mangrove	4-10	36.6	24.9	-28.1	-30.6	-33.9	-5.8	Bouillon et al. (2004a)
Pambala (Sri Lanka)	Mangrove	0-1	37.2	24.5	-28.2	-30.6	-31.5	-3.3	Bouillon et al. (2004a)
Pambala (Sri Lanka)	Mangrove	1-2	27.1	23.8	-27.9	-30.6	-32.0	-4.1	Bouillon et al. (2004a)
Pambala (Sri Lanka)	Mangrove	2-4	17.5	20.4	-28.0	-30.6	-33.2	-5.2	Bouillon et al. (2004a)
Pambala (Sri Lanka)	Mangrove	4-10	23.0	21.5	-27.9	-30.6	-33.0	-5.1	Bouillon et al. (2004a)
Pambala (Sri Lanka)	Mangrove	0-1	31.0	18.7	-28.5	-30.6	-30.9	-2.4	Bouillon et al. (2004a)
Pambala (Sri Lanka)	Mangrove	1-2	26.9	19.3	-27.4	-30.6	-31.9	-4.5	Bouillon et al. (2004a)
Pambala (Sri Lanka)	Mangrove	2-4	26.7	19.3	-27.7	-30.6	-32.6	-4.9	Bouillon et al. (2004a)
Pambala (Sri Lanka)	Mangrove	4-10	15.9	17.6	-28.0	-30.6	-32.5	-4.5	Bouillon et al. (2004a)

Site location	Ecosystem type	Depth range (cm)	% TOC	C/N (molar)	$\delta^{13}\text{C}_{\text{TOC}}$	$\delta^{13}\text{C}_{\text{macrophyte}}$	$\delta^{13}\text{C}_{\text{total}}$	$\delta^{13}\text{C}_{\text{total}} - \delta^{13}\text{C}_{\text{TOC}}$	Data source
Pambala (Sri Lanka)	Mangrove	0-1	18.5	19.8	-26.8	-30.6	-27.6	-0.8	Bouillon et al. (2004a)
Pambala (Sri Lanka)	Mangrove	1-2	12.5	18.7	-24.8	-30.6	-27.0	-2.2	Bouillon et al. (2004a)
Pambala (Sri Lanka)	Mangrove	2-4	9.2	21.4	-27.0	-30.6	-28.0	-1.0	Bouillon et al. (2004a)
Pambala (Sri Lanka)	Mangrove	4-10	8.9	24.2	-27.7	-30.6	-30.1	-2.3	Bouillon et al. (2004a)
Pambala (Sri Lanka)	Mangrove	0-1	5.2	19.4	-26.1	-30.6	-29.9	-3.8	Bouillon et al. (2004a)
Pambala (Sri Lanka)	Mangrove	1-2	4.6	17.7	-26.1	-30.6	-31.3	-5.2	Bouillon et al. (2004a)
Pambala (Sri Lanka)	Mangrove	2-4	3.7	17.4	-26.2	-30.6	-30.7	-4.5	Bouillon et al. (2004a)
Pambala (Sri Lanka)	Mangrove	4-10	2.2	17.3	-25.8	-30.6	-31.6	-5.8	Bouillon et al. (2004a)
Pambala (Sri Lanka)	Mangrove	0-1	19.3	29.7	-27.8	-30.6	-26.4	1.4	Bouillon et al. (2004a)
Pambala (Sri Lanka)	Mangrove	1-2	6.5	15.2	-27.6	-30.6	-31.6	-4.0	Bouillon et al. (2004a)
Pambala (Sri Lanka)	Mangrove	2-4	7.5	19.2	-27.8	-30.6	-26.4	1.3	Bouillon et al. (2004a)
Pambala (Sri Lanka)	Mangrove	4-10	5.5	18.0	-27.8	-30.6	-31.7	-3.9	Bouillon et al. (2004a)
Nyborg Fjord (Denmark)	Seagrass	0-5	0.1	8.6	-17.3	-7.6	-20.2	-2.9	Boschker et al. (2000)
Nyborg Fjord (Denmark)	Seagrass	0-5	0.3	8.3	-18.0	-7.6	-20.1	-2.1	Boschker et al. (2000)
Nyborg Fjord (Denmark)	Seagrass	0-5	0.1	7.5	-17.7	-7.6	-20.0	-2.3	Boschker et al. (2000)
Nyborg Fjord (Denmark)	Seagrass	0-5	2.2	11.3	-19.9	-9.7	-20.4	-0.5	Boschker et al. (2000)
Nyborg Fjord (Denmark)	Seagrass	0-5	1.9	11.2	-18.7	-9.7	-20.2	-1.5	Boschker et al. (2000)
Nyborg Fjord (Denmark)	Seagrass	0-5	2.2	11.7	-20.4	-9.7	-19.7	0.7	Boschker et al. (2000)
Arcachon Bay (France)	Seagrass	0-5	0.0	14.2	-17.4	-11.0	-19.3	-1.9	Boschker et al. (2000)
Arcachon Bay (France)	Seagrass	0-5	0.0	13.0	-17.5	-10.9	-18.7	-1.2	Boschker et al. (2000)
Arcachon Bay (France)	Seagrass	0-5	0.0	11.8	-17.0	-10.6	-19.1	-2.1	Boschker et al. (2000)
Arcachon Bay (France)	Seagrass	0-5	0.1	12.2	-18.5	-11.1	-19.1	-0.6	Boschker et al. (2000)
Arcachon Bay (France)	Seagrass	0-5	0.2	12.5	-17.4	-10.9	-19.1	-0.6	Boschker et al. (2000)
Arcachon Bay (France)	Seagrass	0-5	0.2	13.8	-17.7	-10.9	-18.4	-0.7	Boschker et al. (2000)
Arcachon Bay (France)	Seagrass	0-5	0.0	10.3	-19.3	-8.4	-18.4	0.9	Boschker et al. (2000)
Arcachon Bay (France)	Seagrass	0-5	0.0	10.0	-19.2	-8.5	-19.2	0.0	Boschker et al. (2000)
Arcachon Bay (France)	Seagrass	0-5	0.0	12.9	-18.8	-8.9	-18.6	0.2	Boschker et al. (2000)
Sint Annaland (the Netherlands)	Seagrass	0-5	0.8	10.3	-22.6	-15.0	-22.7	-0.1	Boschker et al. (2000)
Sint Annaland (the Netherlands)	Seagrass	0-5	0.7	11.7	-23.1	-13.8	-22.1	1.0	Boschker et al. (2000)
Sint Annaland (the Netherlands)	Seagrass	0-5	0.5	10.8	-23.0	-13.8	-22.0	1.0	Boschker et al. (2000)
Ban Pak Klok (Thailand)	Seagrass	0-2	0.3	12.6	-22.6	-12.4	-17.9	4.7	Holmer et al. (2001)
Ban Pak Klok (Thailand)	Seagrass	0-2	0.3	13.8	-21.7	-12.1	-17.9	3.8	Holmer et al. (2001)
Ban Pak Klok (Thailand)	Seagrass	0-2	0.4	13.9	-23.3	-11.3	-18.9	4.4	Holmer et al. (2001)
Ban Pak Klok (Thailand)	Seagrass	0-2	0.6	13.3	-23.0	-10.6	-20.2	2.8	Holmer et al. (2001)
Ban Pak Klok (Thailand)	Seagrass	0-2	0.2	13.5	-22.1	-11.3	-18.2	3.9	Holmer et al. (2001)
Ban Pak Klok (Thailand)	Seagrass	0-2	0.2	11.5	-22.2	-11.8	-18.6	3.6	Holmer et al. (2001)
Mallorca (Spain)	Seagrass	0-5	0.3	10.6	-19.1	-13.5	-18.6	0.5	Holmer et al. (2004)
Mallorca (Spain)	Seagrass	0-5	0.2	8.6	-18.3	-13.5	-18.9	-0.6	Holmer et al. (2004)
Mallorca (Spain)	Seagrass	0-5	0.4	17.3	-16.5	-12.1	-19.8	-3.3	Holmer et al. (2004)
Mallorca (Spain)	Seagrass	0-5	0.5	16.6	-17.4	-12.1	-20.0	-2.6	Holmer et al. (2004)
Mallorca (Spain)	Seagrass	0-5	0.4	11.0	-17.7	-16.2	-19.3	-1.6	Holmer et al. (2004)
Mallorca (Spain)	Seagrass	0-5	0.4	8.2	-17.8	-16.2	-18.7	-0.9	Holmer et al. (2004)
Mallorca (Spain)	Seagrass	0-5	4.8	18.8	-16.3	-13.3	-22.8	-6.5	Holmer et al. (2004)
Mallorca (Spain)	Seagrass	0-5	2.7	14.1	-16.7	-13.3	-23.2	-6.5	Holmer et al. (2004)
Mallorca (Spain)	Seagrass	0-5	3.0	22.4	-14.4	-7.9	-19.1	-4.7	Holmer et al. (2004)
Mallorca (Spain)	Seagrass	0-5	2.6	21.2	-15.1	-7.9	-18.9	-3.7	Holmer et al. (2004)
Gazi bay (Kenya)	Seagrass	0-6	1.3	10.5	-16.0	-10.7	-22.7	-6.8	Bouillon et al. (2004b)
Gazi bay (Kenya)	Seagrass	0-6	1.2	9.7	-16.3	-10.7	-21.2	-4.9	Bouillon et al. (2004b)
Gazi bay (Kenya)	Seagrass	0-6	0.8	15.9	-23.3	-18.6	-30.1	-6.7	Bouillon et al. (2004b)
Gazi bay (Kenya)	Seagrass	0-6	7.0	18.6	-25.5	-17.8	-31.4	-5.8	Bouillon et al. (2004b)

Site location	Ecosystem type	Depth range (cm)	% TOC	C/N (molar)	$\delta^{13}\text{C}_{\text{TOC}}$	$\delta^{13}\text{C}_{\text{macrophyte}}$	$\delta^{13}\text{C}_{\text{+15.0}}$	$\delta^{13}\text{C}_{\text{+15.0}} - \delta^{13}\text{C}_{\text{TOC}}$	Data source
Gazi bay (Kenya)	Seagrass	0-6	3.8	13.9	-24.8	-17.8	-31.5	-6.7	Bouillon et al. (2004b)
Gazi bay (Kenya)	Seagrass	0-6	0.9	12.1	-21.4	-15.7	-24.9	-3.6	Bouillon et al. (2004b)
Gazi bay (Kenya)	Seagrass	0-6	0.8	14.1	-21.3	-15.7	-26.7	-5.4	Bouillon et al. (2004b)
Gazi bay (Kenya)	Seagrass	0-6	0.3	14.5	-21.1	-11.4	-22.8	-1.7	Bouillon et al. (2004b)
Gazi bay (Kenya)	Seagrass	0-6	0.5	12.3	-22.3	-11.4	-24.6	-2.3	Bouillon et al. (2004b)
Gazi bay (Kenya)	Seagrass	0-6	0.5	18.0	-22.3	-14.5	-27.3	-5.0	Bouillon et al. (2004b)
Gazi bay (Kenya)	Seagrass	0-6	0.9	23.2	-23.6	-14.5	-26.8	-3.2	Bouillon et al. (2004b)
Gazi bay (Kenya)	Seagrass	0-6	2.5	10.2	-16.3	-13.0	-21.1	-4.8	Bouillon et al. (2004b)
Laguna Madre (USA)	Seagrass	0			-11.3	-10.8	-14.4	-3.1	Jones et al. (2003)
Laguna Madre (USA)	Seagrass	1			-11.1	-10.8	-17.8	-6.6	Jones et al. (2003)
Laguna Madre (USA)	Seagrass	5			-11.2	-10.8	-17.0	-5.8	Jones et al. (2003)
Laguna Madre (USA)	Seagrass	9			-11.3	-10.8	-17.9	-6.5	Jones et al. (2003)
Laguna Madre (USA)	Seagrass	0			-12.1	-10.8	-17.7	-5.5	Jones et al. (2003)
Laguna Madre (USA)	Seagrass	1			-11.3	-10.8	-17.3	-6.0	Jones et al. (2003)
Laguna Madre (USA)	Seagrass	5			-11.0	-10.8	-17.8	-6.8	Jones et al. (2003)
Laguna Madre (USA)	Seagrass	9			-10.0	-10.8	-19.0	-9.0	Jones et al. (2003)
North River (USA)	Seagrass				-19.7		-23.6	-3.9	Cifuentes and Salata (2001)
North River (USA)	Seagrass				-19.3		-22.0	-2.7	Cifuentes and Salata (2001)
Nyborg Fjord (Denmark)	Seagrass, unvegetated	0-5	0.1	7.0	-17.4		-20.2	-2.8	Boschker et al. (2000)
Nyborg Fjord (Denmark)	Seagrass, unvegetated	0-5	0.1	6.9	-17.8		-19.7	-1.9	Boschker et al. (2000)
Nyborg Fjord (Denmark)	Seagrass, unvegetated	0-5	0.1	7.7	-17.6		-20.0	-2.4	Boschker et al. (2000)
Nyborg Fjord (Denmark)	Seagrass, unvegetated	0-5	2.3	10.9	-20.3		-20.5	-0.2	Boschker et al. (2000)
Nyborg Fjord (Denmark)	Seagrass, unvegetated	0-5	2.2	11.3	-20.0		-21.1	-1.1	Boschker et al. (2000)
Nyborg Fjord (Denmark)	Seagrass, unvegetated	0-5	2.0	11.7	-20.4		-21.6	-1.2	Boschker et al. (2000)
Arcachon Bay (France)	Seagrass, unvegetated	0-5	0.0	12.9	-19.2		-17.2	2.0	Boschker et al. (2000)
Arcachon Bay (France)	Seagrass, unvegetated	0-5	0.1	12.7	-19.8		-24.4	-4.6	Boschker et al. (2000)
Arcachon Bay (France)	Seagrass, unvegetated	0-5	0.0	13.2	-19.8		-24.7	-4.9	Boschker et al. (2000)
Arcachon Bay (France)	Seagrass, unvegetated	0-5	0.2	16.2	-19.3		-19.4	-0.1	Boschker et al. (2000)
Arcachon Bay (France)	Seagrass, unvegetated	0-5	0.1	12.5	-18.2		-19.2	-1.0	Boschker et al. (2000)
Arcachon Bay (France)	Seagrass, unvegetated	0-5	0.2	14.0	-17.9		-19.1	-1.2	Boschker et al. (2000)
Arcachon Bay (France)	Seagrass, unvegetated	0-5	0.0	11.8	-20.6		-19.5	1.1	Boschker et al. (2000)
Arcachon Bay (France)	Seagrass, unvegetated	0-5	0.0	11.4	-19.5		-18.7	0.8	Boschker et al. (2000)
Arcachon Bay (France)	Seagrass, unvegetated	0-5	0.0	12.0	-19.6		-18.8	0.8	Boschker et al. (2000)
Sint Annaland (the Netherlands)	Seagrass, unvegetated	0-5	0.5	12.2	-23.5		-24.2	-0.7	Boschker et al. (2000)
Sint Annaland (the Netherlands)	Seagrass, unvegetated	0-5	0.2	8.5	-23.4		-23.9	-0.5	Boschker et al. (2000)
Sint Annaland (the Netherlands)	Seagrass, unvegetated	0-5	0.3	11.8	-25.0		-23.3	1.7	Boschker et al. (2000)
Laguna Madre (USA)	Seagrass, unvegetated	0			-11.1		-14.8	-3.8	Jones et al. (2003)
Laguna Madre (USA)	Seagrass, unvegetated	1			-11.0		-16.4	-5.4	Jones et al. (2003)
Laguna Madre (USA)	Seagrass, unvegetated	5			-11.2		-17.8	-6.6	Jones et al. (2003)
Laguna Madre (USA)	Seagrass, unvegetated	9			-12.0		-17.2	-5.3	Jones et al. (2003)
Mallorca (Spain)	Seagrass, unvegetated	0-5	0.2	6.7	-21.0		-19.1	1.8	Holmer et al. (2004)
Mallorca (Spain)	Seagrass, unvegetated	0-5	0.2	8.0	-18.4		-19.3	-1.0	Holmer et al. (2004)
Mallorca (Spain)	Seagrass, unvegetated	0-5	0.5	18.1	-15.7		-19.0	-3.3	Holmer et al. (2004)
Mallorca (Spain)	Seagrass, unvegetated	0-5	1.8	23.3	-16.1		-18.6	-2.5	Holmer et al. (2004)
Mallorca (Spain)	Seagrass, unvegetated	0-5	0.2	11.4	-17.7		-19.5	-1.8	Holmer et al. (2004)
Mallorca (Spain)	Seagrass, unvegetated	0-5	1.5		-17.7		-19.1	-1.4	Holmer et al. (2004)
Mallorca (Spain)	Seagrass, unvegetated	0-5	0.2	6.9	-18.2		-18.7	-0.5	Holmer et al. (2004)
Mallorca (Spain)	Seagrass, unvegetated	0-5	0.2	8.4	-17.0		-19.6	-2.5	Holmer et al. (2004)
Mallorca (Spain)	Seagrass, unvegetated	0-5	2.6	27.2	-15.8		-21.0	-5.2	Holmer et al. (2004)
Mallorca (Spain)	Seagrass, unvegetated	0-5	3.1	29.3	-15.1		-20.0	-4.9	Holmer et al. (2004)