

## **ALGAL TURF DOMINATES MOMBASA MARINE PARK AND RESERVE IN KENYA, AN IMPLICATION OF NUTRIENT POLLUTION**

Munga C.N.<sup>1</sup>, A. Vanreusel<sup>2</sup>, D. Obura<sup>3</sup>, F. Dahdouh-Guebas<sup>4</sup>, M.O. Mohamed<sup>5</sup> and N. Amiyo<sup>5</sup>

<sup>1</sup> Kenya Marine and Fisheries Research Institute, PO Box 81651, 1050 Mombasa, Kenya  
E-mail: [ecomama@vub.ac.be](mailto:ecomama@vub.ac.be)

<sup>2</sup> Universiteit Gent, Marine Biology Section, Campus Sterre-S8, Belgium

<sup>3</sup> CORDIO East Africa, PO Box 10135, 80101 Bamburi, Kenya

<sup>4</sup> Université Libre de Bruxelles, Département de Biologie des Organismes, Campus du Solbosch, CP 169, Belgium

<sup>5</sup> Kenya Wildlife Service, PO Box 82144, 80100 Mombasa, Kenya

Mombasa Marine Park and Reserve is one of the Marine Protected Areas in Kenya established more than ten years ago. The Park is a no-take-zone, while the Reserve is a zone where regulated extractive uses such as artisanal fishing are allowed. Since the year 2004, ecological monitoring of the MPA has been going on aimed at generating information on the MPA status with a view to improve its management. Four year monitoring data (2004 to 2007) of percentage benthic cover using the Line Intercept Transect (LIT) method reveal relatively high average percentage benthic cover for algal turf in both the Park and Reserve with sampling sites and seasons. If not checked, algal turf could easily out compete and degrade the hard corals and sea grass beds that are important for MPA ecological and biological health status. The growth of algal turf is triggered by nutrient input mainly from land sources. It is therefore important to identify the nutrient input sources into the MPA so as to address management measures to remedy this ecological threat.

### References

- English S., C. Wilkinson and V. Baker (Eds). 1994. Survey Manual for tropical marine resources. Australian Institute of Marine Science, Townsville, Australia.
- Gorgula S.K. and S.D. Connel. 2004. Expansive covers of turf-forming algae on human-dominated coast: the relative effects of increasing nutrient and sediment loads. *Marine Biology* 145:613-619.