

## MANGROVE SPECIES' ASSOCIATION AND DISTRIBUTION PATTERNS IN TUMPAT, KELANTAN DELTA, EAST COAST OF PENINSULAR MALAYSIA

Satyanarayana Behara<sup>1,2</sup>, Farid Dahdouh-Guebas<sup>1</sup>, Indra Farid Idris<sup>2</sup>, Khairul Azwan Mohamad<sup>2</sup>, Mohd-Lokman Husain<sup>2</sup> and Sulong Ibrahim<sup>2</sup>

<sup>1</sup> Complexité et Dynamique des Systèmes Tropicaux, Département de Biologie des Organismes, Faculté des Sciences, Université libre de Bruxelles, ULB - Campus du Solbosch, CP 169, Avenue Franklin D. Roosevelt 50, 1050 Bruxelles, Belgium  
E-mail: [satyam2149@gmail.com](mailto:satyam2149@gmail.com)

<sup>2</sup> Institute of Oceanography (INOS), University Malaysia Terengganu, 21030 - Kuala Terengganu, Malaysia

We studied mangrove vegetation at Tumpat to estimate different tree structural parameters (e.g., stem density and basal area) at selected sites (PCQ-Method). There were 5 dominant taxa in the vegetation, i.e., *Sonneratia caseolaris*, *Nypa fruticans*, *Avicennia alba*, *Rhizophora mucronata*, and *Bruguiera gymnorrhiza* in order of importance. Total tree density varied between 79 and 132 stems  $0.1\text{ha}^{-1}$ , and basal area from 0.14 to 4.9m<sup>2</sup>  $0.1\text{ha}^{-1}$ . Based on species composition and stem density, the mangrove sites could be separated into two groups (Bray-Curtis similarity: 60%). While Group-1 (sites G9, J5, K4, N6 and O4), dominated by *S. caseolaris* and *N. fruticans*, was distributed throughout the forest, Group-2 (sites C6 and G6), represented largely by *A. alba*, is present close to the bay-mangrove boundary. Elevation measurements indicate that Group-1 species occupied low-lying to elevated grounds (0.87-2.23m above the mean sea level) with a (mean) salinity between 0.38 and 14.6psu, whereas Group-2 occurred preferentially at low to medium elevations (0.86-1.29m) and high salinity (14.6psu). Discrete mangrove associations (=groupings) were discerned, wherein the distribution of species is governed by factors such as proximity of land or sea, freshwater input and elevation.