

FAMENNIAN

Jacques THOREZ¹, Roland DREESSEN² & Maurice STREEL³

(9 figures, 4 tables)

1. Geology Dept, Liège University B18 Sart-Tilman, B-4000 LIEGE, Belgium; E-mail: j.thorez@ulg.ac.be.
2. Flemish Institute for Technological Research (VITO), B-2400 MOL, Belgium; E-mail: roland.dreesen@vito.be.
3. Geology Dept, Liège University B18 Sart-Tilman, B-4000 LIEGE, Belgium; E-mail: Maurice.Streel@ulg.ac.be.

ABSTRACT. The name of the Famennian stage originated in the 19th century from the Famenne region in southern Belgium. After the pioneering work of Dumont, d’Omalius d’Halloy, Gosselet and Murlon, a renewed interest in the 1960’s focused mainly on the lithological, sedimentological, paleontological and palynological characteristics of the Famennian. This resulted in a refined litho- and biostratigraphical framework for the Lower Famennian Famenne Shales and the Upper Famennian Condruz Sandstones in Belgium. The most obvious finding was the distinct diachronous character of the Condruz Group. The present review paper extends this stratigraphical framework into the neighbouring Avesnois (Northern France) and Aachen (Northwestern Germany) areas: it proposes a lithostratigraphical correlation between the units identified within the individual tectonic units. An overview is given of the main lithological-sedimentological characteristics, depositional environments (for both the siliciclastics and the carbonates) and prevailing paleogeographical-paloeclimatological conditions, during the Famennian stage in Belgium and adjacent areas.

KEYWORDS: Famennian, Belgium, Avesnois, Northwestern Germany

Reference to this paper: Thorez, J., Dreesen, R. & Streel, M., 2006. Famennian. In Dejonghe, L., ed., Current status of chronostratigraphic units named from Belgium and adjacent areas, *Geologica Belgica*, Brussels, 9/1-2: 27-45.

1. Introduction

In Belgium, the Famennian is exposed or has been encountered in five distinct tectonic settings (Fig. 1): a) the autochthonous Namur Synclinorium with its prolongation into the Hainaut area; b) the allochthonous Dinant Synclinorium, c) the Theux tectonic window in the Stavelot-Venn Massif; d) the Vesdre Nappe which extends into the Aachen area; e) and the subsurface of the Campine Basin in north-eastern Belgium where the Famennian has been encountered only in boreholes. Both the Namur and Dinant Synclinoria extend into the Avesnois region of northern France. All of these units (and depocenters) surround the London-Brabant Massif as part of the former Old Red Continent. They reached their present-day tectonic configuration during the Asturian orogenic phase. The Dinant and Namur Synclinoria are separated by the Midi-Eifel-Aachen Overthrust, the importance of which still remains a matter of debate: it accounts for 10-15 km of basin shortening according to Michot (1980) versus more than 100 km according to the tectonic nappe concept of Bless *et al.* (1983). Adopting the latter model would have an immediate implication for the palinspastic paleogeographical reconstruction of the Famennian lithostratigraphical units. The Famennian is more complete and best developed in the Dinant Synclinorium.

In contrast, it is rather incomplete or even lacking due to erosion or non-deposition events in the other areas (*e.g.* the northern limb of the Namur Synclinorium, the Hainaut area and especially in the Campine Basin, NE-Belgium), as shown in figure 3.




Figure 1. Location map of the studied areas. B.V.D. = Booze – Le Val Dieu High

2. Name

Famennian (English), Famenniaan (Dutch), Famennium (German), Famennien (French).

3. Age

(Gradstein *et al.* 2004)
 Top: 359.2 ± 2.5 my
 Base: 374.5 ± 2.6 my

4. Authors

The name “Famennian” was first introduced by Dumont (1855, legend of the “Carte géologique de l’Europe”: Faménien ou Dévonien supérieur) when referring to a shaly and sandy (Dumont, 1848) “Condruzian System”. The evolution in concept of subdividing the Famennian Stage is summarized in Table 1 (modified after Dreesen, 1976), starting with the pioneering work of Dumont (1832) and ending with the latest concepts of Bouckaert *et al.* (1968) and Thorez & Dreesen (1986).

5. Historical type area

The name of the Famennian stage originated in the 19th century from the Famenne region in southern Belgium. This geographical region corresponds to a shaly depression located South and East of the Meuse river, in between the Condroz area (folded sandstone-limestone succession) in the North and the limestone ridge (La Calestienne) and the succeeding Lower Paleozoic Ardennes hills in the South (Fig. 2). The name Famenne refers to its Latin name “famina”, pointing to the poor soil conditions of the area.




Figure 2. Location of the Famenne area, the name of which is at the origin of the Famennian Stage.

6. Description

The Famennian Stage (Uppermost Devonian) corresponds in Belgium to a sedimentary sequence of up to 600 m of predominantly siliciclastic shelf sediments inserted in between Upper Devonian (Frasnian) and Lower Carboniferous (Tournaisian) shelf carbonates. With time, a lithofacies shift occurred from predominantly pelitic sediments in the early Famennian, (the Famenne Shales) to predominantly sandy sediments during the late Famennian (the Condroz Sandstones). Fossiliferous nodular shales represent the transitional beds from the underlying Matagne Shales (Frasnian), whereas the Latest Famennian (Strunian) stromatoporoid limestones of the Comblain-au-Pont Fm mark the start of the extensive shallow-marine shelf carbonate sedimentation of the succeeding Lower Carboniferous (Tournaisian) stage, starting with the Hastière Limestone (Figs 2 & 3).

7. Historical background

Dumont (see chapter 4., Authors) correlated his Condruzian system with the Condroz Psammities *s.l.* of d’Omalius d’Halloy (1868). The latter unit then still included the uppermost Frasnian shales at the base, and the Etroeungt “Assise” at the top. The “standard” stratigraphical scheme of the Upper Famennian “Condroz Psammities” was first established by Murlon (1875-1886) in the Ourthe valley, which is south of Liege in the north-eastern corner of the Dinant Synclinorium. Murlon introduced and described several new stratigraphical units, in ascending order: the Esneux, Souverain-Pré, Montfort and Evieux “assises”. These units were then all well exposed in the railroad cuts along the Rivage-Liège track and, in the then, numerous active sandstone quarries. The names of the “assises” were taken from small towns and villages along the Ourthe river, where the lithological units were well exposed. Later on, Murlon (1885) studied coeval lithological units in different geographical areas. He then proposed a lithostratigraphical correlation with the southern part of the Dinant Synclinorium and even with the Avesnois area of northern France, where a shalier facies predominated. Gosselet (1878, 1879, 1880) recognized also important lateral facies changes when he correlated the sandy Condroz Psammities of the Condroz and the Entre-Sambre-et-Meuse regions (northern limb of the Dinant Synclinorium) with the dominantly shaly facies of the Fagne area in northern France. In the latter area, the Sains Shales that occur above the Senzeille and Mariembourg Shales, were then considered as coeval deposits of the Condroz Psammities (= micaceous sandstones) (Fig. 2). The exact lithostratigraphical position of the Souverain-Pré “assise” has been discussed by Dreesen (1976). Since the work by Bouckaert *et al.* (1968) the Souverain-Pré formation has become incorporated into the Condroz Sandstone Group. Figure 3 is a more detailed

UPPER QUARTZO-SHALE SYSTEM										Dumont (1830)	
CONDROZ PSAMMITES s.l.										d' Omalius d' Halloy (1839)	
QUARTZO-SHALY CONDROZIAN SYSTEM										Dumont (1839)	
FAMENNE SHALES					CONDROZ PSAMMITES					d' Omalius d' Halloy (1858)	
FAMENNIAN STAGE										Dumont (1855)	
FRASNIAN SHALE + LIMESTONE		FAGNE SHALES			CONDROZ PSAMMITES			ETROEUNGT LIMESTONE		Gosselet (1860)	
FAMENNE SHALES					CONDROZ PSAMMITES					Mourlon (1873)	
LAYGRED SHALES				PSAMMITES				ETROEUNGT SHALE + LIMESTONE		Gosselet (1879)	
CONDROZ PSAMMITES										Mourlon (1880)	
ESN ASSISE		SVP ASSISE		MFT ASSISE		EVX ASSISE					
LOWER FAMENNIAN			UPPER FAMENNIAN						ETROEUNGT ASSISE	Gosselet (1880)	
SEN ASSISE		MAR ASSISE	SAINS ASSISE								
LOWER FAMENNIAN			Fa 1	Fa 2		Fa 3				Mourlon (1882)	
SEN Fa 1a	MAR Fa 1b	ESN Fa 1c	SVP a	b	MFT a	EVX b	c	d	HAS e	CLP f	
Fa 1			Fa 2								1st Map Legend (1892)
SEN Fa 1a	MAR Fa 1b	ESN Fa 1c	SVP Fa 2a	MFT Fa 2b	EVX Fa 2c		CLP Fa 2d				
LOWER FAMENNIAN				UPPER FAMENNIAN						5th Map Legend (1900)	
SEN Fa 1a	MAR Fa 1b-c			MFT Fa 2a		EVX Fa 2b					
LOWER FAMENNIAN				UPPER FAMENNIAN						Geological Survey of Belgium (1952)	
SEN Fm 1a	MAR Fm 1b	ESN + SVP Fm 1c		MFT Fm 2a	EVX Fm 2b						
LOWER FAMENNIAN				SVP MIDD. FAM	UPPER FAMENNIAN						Sartenaer (1957)
Fm 1			Fm 2								Beugnies (1965)
LOWER ASS. Fm 1a	MIDDLE ASS. Fm 1b	UPPER ASS. Fm 1c	LOWER ASS. Fm 2a	UPPER ASSISE Fm 2b							
LOWER FAMENNIAN			UPPER FAMENNIAN				STRUNIAN		Bouckaert, Streef & Thorez (1968) Thorez & Dreesen (1986)		
Fa 1a	Fa 1b	ESN Fm Fa 2a	SVP Fm Fa 2b	MFT Fm Fa 2c	EVX Fm Fa 2d	CLP Fm Tn 1a					
LOWER FAMENNIAN			MIDDLE FAMENNIAN		UPPER FAMENNIAN		UPPERMOST FAM.		Thorez & Dreesen (1986) Thorez & Dreesen (2002) (Dinant synclinorium, Belgium)		
SEN Fm	MAR Fm	ESN Fm AYE Fm	SVP Fm	(*)	(*)	CLP					
FAMENNE GROUP				CONDROZ GROUP					(*)		this paper

SEN Senzeille
 MAR Mariembourg
 ESN Esneux
 AYE Aye
 SVP Souverain-Pré
 MFT Montfort
 CLT Comblain-La-Tour
 CIN Ciney
 HAV Haversin
 EVX Evieux
 BEV Beverire
 HAS Hastière
 CLP Comblain-Au-Port

Table 1. Historical evolution of the lithostratigraphical subdivision of the Famennian Stage. Asterisks in the lowermost boxes refer to the most recent subdivision depicted in Fig.3.




Figure 3. Detail of the lowermost box in Table 1 showing further lithostratigraphical subdivision of the Famennian Stage within the Dinant Synclinorium. Dashed red lines correspond to successive oolitic ironstone levels (event-stratigraphical marker beds).

correlation scheme for the Famennian formations in the Dinant Synclinorium and represents an enlargement of the inset at the very bottom of Table 1.

Although an international agreement has yet not been reached to define the bases of substages within the Famennian stage, a subdivision into 4 substages has been internationally adopted: from base to top the Lower, Middle, Upper and Uppermost (Strunian?) Famennian (Streel *et al.*, 1998 & 2005). A proposal for a condont-based definition of these substages is still under consideration by the International Subcommittee on Devonian Stratigraphy (Streel, 2005). However, it is adopted here as a chronological subdivision to replace the old subdivisions of 2 or 3 substages and the now obsolete stratigraphic units represented by the Fa or Fm symbols (Table 1, Fig. 3).

8. Lithology and lithostratigraphy

In the Dinant Synclinorium, the Lower Famennian Famenne Group encompasses the shaly Senzeille and Mariembourg Fms (Fig. 4): the latter grade laterally into the Lambermont (*pars*) and Hodimont Fms in the Namur Synclinorium (southern flank), in the Vesdre Nappe and in the Theux Window. Although the Senzeille and Mariembourg Fms can be distinguished on the base of a combination of lithological and macropaleontological criteria (e.g. rhynchonellid brachiopod index species) this distinction has not been made on the published sheets of the new geological map of Wallonia. On the other hand, the Senzeille and Mariembourg Fms merge in a southwestern direction into one single unit, called the Fa-




Figure 4. Lithostratigraphical scheme of the Famenne and Condroz Groups in the different tectonic units of Belgium, incorporating the Avesnois (northern France) and Aachen areas (western Germany). Correlations are based on conodont and miospore zonal markers (left). Black dotted lines correspond to oolitic ironstone levels




Figure 5. Lithostratigraphical scheme (after a palinspastic reconstruction of about 18 km) of the Condroz Sandstone Group in the “classical” Ourthe valley (S of Liège). This section extends from the town of Esneux in the N to the town of Comblain-la-Tour in the S. Black dotted lines represent conspicuous ball-and-pillow levels (“pseudonodules”). Thick red lines correspond to red beds.

menne Shales, typical of the Avesnois area (Fig. 4). Here, the latter Famenne Shales contain nodular and lenticular limestones near their top, yielding conodonts of the *marginifera* zone: hence this upper level can be correlated with the Souverain-Pré Fm in the Ourthe valley (Bouckaert *et al.*, 1977). Moreover, a goniatite-bearing nodular and lenticular limestone interval within the Famenne Shales in the same area, allowed further biostratigraphical correlation with the *Cheiloceras* Limestone of the Aachen region (NW-Germany) (Fig. 4).

The Upper Famennian Condroz Group is best developed, and shows the most complete suite of succeeding formations, in the eastern part of the Dinant Synclinorium (Fig. 3): here, the Montfort Fm grades into the coeval Comblain-la-Tour, Ciney and Haversin Fms, according to a NNE - SSW section. The Haversin Fm in turn passes laterally and in a southwestern direction into the Sains Shales (Avesnois area, northern France). The Evieux Fm is conspicuously interfingering with the Beverire Fm. Its uppermost part is also interrupted by a short transgressive event, the restricted marine or peritidal Fontin Mbr, whereas a thin black shale interval near the boundary of

the Montfort and Evieux Fm (in the Ourthe and Bock valleys) matches another transgressive event, that can probably be correlated with the *annulata* event (Walliser, 1996). Despite the lack of good biostratigraphical control, a thick series of red beds (up to 200 m) provisionally named here as the Huy Citadelle Fm (southern flank of the Namur Synclinorium) (Fig. 4), seems to be coeval of the Evieux Fm. In the Namur Synclinorium, the Samme Fm is partly coeval with the Comblain-au-Pont Fm from the Dinant Synclinorium, but it extends vertically into the Lower Carboniferous Hastière Limestone. In the Campine Basin, the Condroz Sandstone Group (according to borehole data) seems to be represented only by the Evieux Fm. Here, the latter formation is clearly diachronous with respect to the adjacent tectonic units (Laenen, 2003) (Fig. 4). In the Aachen area (NW-Germany), peculiar red-stained goniatite-bearing nodular shales and limestones (the so-called *Cheiloceras* Limestone) occur at or near the transition of the Lower Famenne Shales and the Esneux Fm. This level grades laterally (to the W) into an oolitic ironstone bed and can be traced further westward into the Vesdre and Dinant

Synclinoria (Sartenaer, 1957b; Dreesen, 1987, 1982b). In the Avesnois area, a coeval *Cheiloceratid* goniatite bearing interval is present as well (Bouckaert *et al.*, 1977). Moreover, analogous *Cheiloceras*-bearing marker horizons have been encountered at the same stratigraphical level in different sedimentary basins throughout the Rheno-Hercynian realm (Dreesen, 1989). The Watissart Mbr is a local and rather unusual sandy member of the Esneux Fm in the westernmost part of the Dinant Synclinorium (Silenrieux-Walcourt and Grandrieu-Beaumont area; Dumoulin & Marion, 1997; Dumoulin, 2001) and in the adjacent Avesnois area (Northern France, Waterlot *et al.*, 1967): here, grey quartzitic sandstones ranging in thickness from tens of centimetres to metres were recognized for the first time by Gosselet (1880). They were initially identified as “Grès de Cerfontaine”, and later as “Grès de Watissart” by Beugnies (1965) and Waterlot *et al.* (1967). However, because of the lack of detailed sedimentological and biostratigraphical data, the lithostratigraphical relationships of the Famennian of the Avesnois area with coeval deposits in the adjacent areas in Belgium are still unclear. Figure 5 represents the stratigraphical framework for the Condroz Group in the “classic” Ourthe valley: in this valley Mourlon (1875–1886) made his pioneering geological observations leading to the first detailed (litho)stratigraphical scheme of the Condroz Sandstones, differentiating the succeeding (superposed) lithostratigraphic units or “assises” of Esneux, Souverain-Pré, Montfort and Evieux. Later on, these “assises” became partly replaced by formal formations with corresponding names. More recently, new coeval formations have been introduced, including the Beverire and Comblain-la-Tour Fms (Bouckaert *et al.*, 1968). Of particular interest is the occurrence of thin oolitic ironstone beds (Figs 3 & 4) in the Famennian. These appear at distinct stratigraphical levels in the Lower and Middle Famennian and represent excellent lithostratigraphical marker beds (levels I, II, IIIa, IIIb & IV of Dreesen, 1982a and b). They are interpreted by the authors as transgressive system tracts. Some ironstone beds can be traced over great distances throughout the sedimentary basin, even out of Belgian territory into the Aachen and Avesnois areas. Moreover, they represent good event-stratigraphical markers that can even be correlated with volcano-sedimentary events in the Rhenish basin (Dreesen *et al.*, 1986a and b; Dreesen, 1987). Based on geochemical evidence (REE) the source of the iron has been related to volcanic activity (Laenen *et al.*, 2002).

9. Sedimentology and palaeogeography

The present authors have summarized and graphically highlighted the most relevant lithological, sedimentological and depositional environmental parameters for each of the identified units in tables (Tables 2, 3 & 4). The latter tables give a quick overview of the dominant lithologi-

cal and sedimentological characteristics and subsequent depositional settings, allowing a clear distinction of each formal lithostratigraphical unit of the Belgian Famennian Stage. The Lower Famennian argillaceous sediments represent relatively open marine conditions: they were deposited mainly offshore, on a shallow epicontinental platform, out of the reach of the longshore currents that redistributed the mainly coarser siliciclastics. Paleocological evidence from conodont biofacies analysis supports the above interpretation (Dreesen & Thorez, 1980). The Senzeille and Mariembourg Fms are, respectively, composed of olive-green to greyish green shales and grey to purplish (violet) shales and sandy shales, locally enclosing thin siltstone and fine-grained sandstone beds, as well as nodular or lenticular limestones. Rhynchonellid brachiopods are very common and particular key-species are very helpful in the field for mapping and discriminating between the two shaly formations. Oolitic ironstone marker beds occur within the Famenne Group in most of the investigated tectonic units. The latter Clinton-type oolitic ironstones are composed of lenticular limestones (packstones / grainstones) enriched with various ferruginized allochems. The shaly Hodimont Fm (Laloux *et al.*, 1996) of the Vesdre Nappe is coeval with the Mariembourg Fm of the Dinant and Namur Synclinoria, but it encloses more siltstone beds (decimetric to metric beds). The Hodimont Fm contains numerous nodular and lenticular limestones or coquinas (brachiopod-rich wackestones and packstones). In the Aachen area, a particular red-stained goniatite-bearing nodular mudstone/wackestone, the “*Cheiloceras* Kalk”, occurs near the Lower/Upper Famennian transition. The latter limestone can be traced into the adjacent Vesdre Nappe (Sartenaer, 1957b) and Dinant Synclinorium and grades laterally into ironstone level IIIa (Dreesen, 1982b).

The sediments of the Condroz Group originated during an important progradation of deltaic complexes and a redistribution of the reworked siliciclastic sediments by strong longshore currents onto a shallow epicontinental platform. As a result, a whole spectrum of alluvial, estuarine, restricted marine (lagoonal) and shallow marine siliciclastic depositional settings has developed. The coarser clastics are predominantly well-sorted fine-grained sandstones (mean 45–120 μm) that are chiefly composed of quartz and feldspars. Due to the relative abundance of feldspars, these sandstones can be classified as (micro)arkosic sandstones (Thorez, 1969). The thickness of these (micro-) arkosic sandstones and siltstones varies from a few dm up to several metres, depending on their depositional setting. The sandstones exhibit a large variety of colours as well. Besides a broad spectrum of predominantly siliciclastic sediments (shales, siltstones and sandstones), the Condroz Group contains subordinated carbonate rocks including dolostones and limestones, as well as mixed carbonate-siliciclastic rock types. Their relative frequency depends largely on the type of depositional setting. All of the dolostones (mostly a few cm, excep-

LITHOSTRATIGRAPHY	LITHOLOGY													
	Ss	Si	Sh	Dm	Ds	pAn	SS(D)	SSD	si(Ca)	Lim	As	Vs	Dolc	OI
Comblain-au-Pont Fm	■	■	■							■				
Evieux Fm <i>Fontin Mbr</i>	■	■	■	■					■	■	■	■	■	
Evieux Fm <i>Crupet Mbr</i>	■	■	■	■		■					■	■		
Evieux Fm <i>Royseux Mbr</i>	■	■	■	■		■	■					■	■	
Beverire Fm	■	■	■						■	■				
Montfort Fm <i>Barse Mbr</i>	■	■	■	■	■		■	■			■			
Montfort Fm <i>La Gombe Mbr</i>	■	■	■	■	■		■	■						
Montfort Fm <i>Bon Mariage Mbr</i>	■	■	■	■					■					
Comblain-La-Tour Fm	■	■	■						■	■				
Comblain-La-Tour Fm <i>Poulseur Mbr</i>	■	■	■						■					
Ciney Fm	■	■	■						■					
Ciney Fm <i>Dorinne Mbr</i>	■	■	■						■					
Haversin Fm	■	■	■						■					
Souverain-Pré Fm		■	■						■	■				■
Souverain-Pré Fm <i>Baelen Mbr</i>		■	■						■	■				
Esneux Fm	■	■	■							■				■
Aye Fm	■	■	■							■				■
Mariembourg Fm <i>Senzeille Fm</i>	■	■	■							■				■
Hodimont Fm	■	■							■	■				■

■ abundant ■ frequent ■ rare

Table 2. Occurrence, stratigraphical distribution and relative frequency of characteristic lithological features of the Condroz Sandstone Group, especially in the Dinant Synclinorium. Ss = micro-arkosic sandstone, Si = arkosic siltstone, Sh = shale, Dm = evaporitic dolostone with a massive structure (lack of internal stratification), Ds = straticulated (micaceous) para-autochthonous evaporitic dolomite, SS(D) = dolomitic arkose (contamination by dolomite less than 25%), SSD = sandy dolomite, the dolomite content of which ranges from 25% to 65%, (Ca)Si = calcareous siltstone, Lim = limestones, AS: aridisol, VS = (generally reddened) vertic fluvial paleosol (paleofluvial entisol), Dolc = dolcrete, OI = oolitic ironstones.




Figure 6. Cartoon depicting the major carbonate microfacies characteristics of restricted marine and open marine limestones within the Condroz Sandstone Group (after Dreesen & Thorez, 1994). 1 and 2 refer to exceptional autochthonous limestone deposits: 1. serpulid bioherms of the Fontin Mbr in the Evieux Fm, and 2. mud mounds of the Baelen Mbr of the Souverain-Pré Fm. The upper part of the figure shows the prograding lithofacies and the stratigraphic succession (from right to left) of the corresponding formations with time (*rhomboidea* through *expansa* conodont zones). Arrows point to transport direction of storm surges and tidal currents generating tempestites and allowing mixing of carbonate allochems.

tionally one metre thick) have been interpreted as the result of shallow water evaporation in restricted marine settings or emerging lagoonal ponds. Dolomite is also a characteristic component of paleosols, including dolcretes (dolomitic calcretes) and aridisols: these occur at or near the boundary between the succeeding Montfort and Evieux Fms in the Ourthe and Hoyoux valleys (Thorez, 2002). The progradation of siliciclastic facies during the Famennian is punctuated by the episodic occurrence of bioclastic limestones. Generally the latter occur as nodular or thin lenticular, millimetre to many centimetre thick beds interstratified within shales, siltstones and sandstones. These sometimes densely fossiliferous limestones provide valuable biostratigraphic data and paleoecological information on the prevailing depositional conditions (carbonate microfacies). Not only the host siliclastic sediments but also the enclosed limestones of the Famenne Shales, and especially those of the Condroz Sandstone Group, show a broad spectrum of depositional settings. These are depicted in Figure 6.



















The limestones can be grouped into two major categories, according to the origin of their biogenic components: au-

tochthonous limestones and parautochthonous-allochthonous limestones (Dreesen & Thorez, 1994). Autochthonous limestones are rather exceptional: they are restricted to particular paleogeographical-paleotectonic settings of the Condroz shelf. Three genuine autochthonous carbonate deposits have been described so far in the Condroz sandstone Group: the Fontin Mbr, the Baelen Mbr and the stromatoporoid boundstones (“biostromes”) of the Comblain-au-Pont and coeval Etroeungt Fms. These autochthonous limestones are related to transgressive events, possibly also to eustatic events (Thorez & Dreesen, 1997; Sandberg et al, 2002). The carbonate Fontin Mbr of the Evieux Fm consists of dark-coloured restricted-marine or lagoonal, ostracodal mudstones and algal-peloidal wackestones, locally enclosing small serpulid (microconchid) carbonate build-ups (Dreesen & Jux, 1995). The red- to pink-coloured Baelen Mbr of the Souverain-Pré Fm (in the Vesdre Nappe) consists of relatively deeper water, open-marine cryptalgal-sponge-crinoidal mud mounds displaying characteristic “stromatactis” structures (Dreesen & Flajs, 1984; Dreesen *et al.*, 1985a). The mud mounds developed below wave base but they intermit-

LITHOSTRATIGRAPHY	SEDIMENTARY TEXTURES & STRUCTURES																RGB		
	M		~	~	~	~	~	~	~	~	~	~	~	~	~	~		↑	NGB
Comblain-au-Pont Fm	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Eveux Fm	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Fortin Mbr	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Eveux Fm	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Crupet Mbr	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Eveux Fm	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Royseux Mbr	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Beveire Fm	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Montfort Fm	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Barse Mbr	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Montfort Fm	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
La gombe Mbr	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Montfort Fm	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Bon Mariage Mbr	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Comblain-La-Tour Fm	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Comblain-La-Tour Fm	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Pouiseur Mbr	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Ciney Fm	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Ciney Fm	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Dorinne Mbr	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Haversin Fm	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Souverain-Pré Fm	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Souverain-Pré Fm	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Baelen Mbr	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Esneux Fm	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Aye Fm	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Mariembourg Fm	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Senzeille Fm	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
Hodimont Fm	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■

■ abundant ■ frequent ■ rare

Table 3. Characteristic sedimentary structures of the Condruz Sandstone Group with their relative frequencies (mostly restricted to the Dinant Synclinorium). No clear differentiation can be made for each of the formations. However, some characteristic “finger prints” allow for the identification of some particular depositional environments. See legend of used symbols below.

M	Massive texture (lack of internal bedding due to the lack of mica concentrations or changes in grain size)
	Planar horizontal bedding
	Undulating bedding
	Ripple drift
	Tabular cross-stratification
	Through cross-stratification
	Fluvial channel or creek
	Dune, longitudinal or transversal sand bar, megaripple
	Hummocky cross stratification
	Load cast, slump, ball-and-pillow
	Oscillation ripple-mark (wave activity)
	Asymmetric ripple-mark (current generated)
	Desiccation crack
C	Coquina bed
	Nodular anhydrite (dissolved)
	Dolomite intraclasts
	Nodular limestone or limestone intraclasts
	Geode (dissolved nodular anhydrite)
	Various types of burrowing
	Bioturbated sediment
↑	Plant drift (debris)
NGB	Normal graded bedding (fining-upward)
RGB	Reverse graded bedding (coarsening-upward)

tently reached storm wave base, as indicated by the occurrence of coarse-grained lenticular crinoidal-foraminiferal grainstones. Finally, several stratigraphically distinct stromatoporoid boundstones (“biostromes”) occur interbedded with micaceous shales and siltstones in the latest Famennian (Strunian: Conil, 1968; Conil & Lys, 1980). These boundstones are known from the Comblain-Au-Pont Fm (Dinant Synclinorium), the Dolhain Fm (Vesdre Nappe), the “Strunian” (Aachen area) and from the top of the time-equivalent Etroeungt Fm (Avesnois area). The latter limestones herald the extensive shallow carbonate shelf conditions of the succeeding Lower Carboniferous. For a detailed microfacies analysis of the Etroeungt Fm in the Avesnois area, the reader is referred to Mamet & Pr eat (2003). On the other hand, parautochthonous-allochthonous limestones are more common: they represent the hallmark of the sandy Condruz Group, although they

also occur in the shaly Lower Famennian formations. In contrast with the Famenne Group, the limestone beds in the Condruz Group are more contaminated by siliciclastics. Generally they consist of bioclastic packstones and grainstones with variable mixtures of shallow-marine to restricted-marine skeletal grains and coated grains. Some of the allochthonous carbonates have been interpreted as calcareous tempestites, “seismites” and erosional lags according to sedimentological evidences (Dreesen & Thorez, 1994). An exceptional accumulation (locally exceeding 100 m) of irregular, subrounded, many centimetre-sized clasts of sandy algal-foraminiferal-crinoidal wackestones/ packstones and grainstones, embedded in a siliciclastic matrix, is known as the Souverain-Pr e Fm (Dreesen, 1978). This mixed silicilastic-carbonate facies most probably represents a reworked shallow-marine carbonate ramp that temporarily developed behind sup-

posedly tectonically-controlled submarine crinoidal sand shoals. The carbonate ramp developed during a mid-Famennian eustatic sea-level rise (Thorez & Dreesen, 1997; Sandberg *et al.*, 2002).

Table 3 summarizes the nature, the relative abundance and the association of sedimentary structures characteristic for each formation in the Dinant Synclinorium, where most of the detailed field observations were made (Thorez, 1969, Thorez & Dreesen, 1986 and Thorez *et al.*, 1988). In addition to the above lithological characteristics, sedimentary structures were most helpful for interpreting the various depositional environments of the Condroz Group: these are listed in Table 4. Figure 7 depicts the lateral lithofacies shifts and inferred depositional environments of the Condroz Group within the “classical” Ourthe valley.

During Famennian times, the Condroz shelf corresponded to a relative shallow epicontinental sea, bordering

the south side of the Old Red Continent (Fig. 8; Paproth *et al.*, 1986). The latter continent was intersected by the Caledonian mountain belt. The frequent occurrence of red beds (Evieux Fm), evaporites (Montfort and Evieux Fms), dolcretes and aridisols (Evieux Fm) is symptomatic of prevailing arid to semi-arid climatic conditions within the then tropical trade wind belt (located between paleo-latitude 10° and 20° S). In the then southern hemisphere, the easterly tropical trade winds drove a warm, equatorial sea current westward between the Equator and 20° S (Heckel & Witzke, 1979). This south-equatorial current flowed through the Paleotethys Sea toward Western Europe, before splitting at the east-end side of the Russian Platform generating a southwestwardly moving branch through the basins of Central Europe, known as the West-bound Current (WBC). The latter incoming strong and warm, south-equatorial WBC passed through the Polish Basin and was deflected towards the Ardenno-Rhenish Basin,




Figure 7. Distribution in time and space of the main depositional environments of the Condroz Sandstone in the “classical” Ourthe valley (quarries and outcrops), from the towns of Esneux to Comblain-la-Tour. Syn-sedimentary tectonics directly influence the distribution of depositional environments. Note the (re)activation of listric faults and related pillow levels, the temporary stacking of the barrier system (La Gombe Mbr), the development of distal alluvial facies, containing red beds intercalated with evaporitic dolomite. The Souverain-Pré Fm reaches a maximum thickness on the flank of a listric fault whereas it is wedging out towards Esneux (dolomitized here) and it is thinning towards Comblain-la-Tour (more pelitic matrix). Dotted lines correspond to ball-and-pillow levels. Thick red lines represent red beds.




Table 4. Depositional environments and their stratigraphical distribution in the Lower and Upper Famennian (mainly Dinant Synclinorium). Transition of relative “deep” offshore open-marine settings in the Lower Famennian Senzeille and Mariembourg Fms to progressively more inshore “continentally influenced” depositional settings in the Upper Famennian. This shift represents a “regression” but it is rather the result of prograding pro-delta sands on a shallow epi-continental shelf. The depositional environments can be grouped into 3 main settings: a) distal alluvial to back-barrier setting, with evaporitic lagoons fed by tidal inlets, b) a barrier complex setting including foreshore and shoreface subsettings, and c) offshore settings.

still encroaching the northern shelf areas. This current was sufficiently strong to redistribute the alluvial discharges coming from small prograding deltas on the southern border of the Old Red Continent, where the Netherlands and North-German High and Lowlands were located. Thus, the Condruz Sandstone material represents re-worked pro-delta sands resulting from the destruction of delta lobes located outside Belgium (Paproth *et al.*, 1986). The arid climatic conditions over the surroundings of the London-Brabant High most probably resulted from the fact that the moisture in the trade winds was lost during their passage over the mountain areas of the Bohemian and Carpathian Highs (see Fig. 8).

10. Biostratigraphy

A new lithostratigraphical scheme for the Famennian, mostly applied to the Condruz Group in the Dinant Synclinorium, has been compiled after two decades of detailed and intensive interdisciplinary investigations, involving both sedimentological (Thorez, 1969; Dreesen, 1976; Thorez *et al.*, 1977 & 1988; Dreesen & Thorez, 1980, 1994; Thorez & Dreesen, 2002) and micropaleontological-palynological tools (*i.e.* Bouckaert *et al.*, 1965 & 1968; Dreesen *et al.*, 1986b; Conil *et al.*, 1986; Dusar & Dreesen, 1984). The latter studies lead to the proposition of a detailed biostratigraphical framework, mainly based on conodont and miospore zones (Fig. 9), with accessory input from foraminifera (Bouckaert *et al.*, 1966), ostracodes (Becker *et al.*, 1974) and, for the lower(most) Famennian, brachiopods (*i.e.* Sartenaer, 1957a; Dusar, 1976) and acritarchs (Vanguetaine *et al.*, 1983). The micropaleontological framework provided a powerful tool for intrabasinal biostratigraphical correlations and allowed the first detailed paleogeographical reconstruction of the Condruz Group (Thorez *et al.*, 1977). This was achieved within the area bordered by the Ourthe, Hout-si-Plou, Amblève, Hoyoux, Bocq, Meuse and Lesse rivers, largely thanks to the intensive extraction of sandstone as a building stone during the 1960's-70's. These huge quarries then provided excellent outcrops offering good observation conditions and allowed a bed-by-bed sampling for further petrographical and micropaleontological-palynological investigation. The combined biostratigraphical and lithostratigraphical data support the formal subdivision of the Famennian Stage into four substages, most especially in the thoroughly investigated eastern part of the Dinant Synclinorium. It must be noted however, that some formations are either strongly reduced in thickness, missing, wedging out or are conspicuously diachronous, as shown in Figure 3. This is particularly true along the northern limb of the Namur Synclinorium, in the Hainaut area (lateral extension of the Namur Synclinorium) and especially in the Campine Basin (a good biostratigraphical control is lacking here, hence the correlations are tentative and based on lithostratigraphical criteria only; Laenen, 2003).




Figure 9. Biostratigraphical framework of the Famennian and Lower-Tournaisian Stage and correlations between formal miospore, conodont and ammonoid zones. In the right hand column the former Belgian (litho-)stratigraphical subdivisions are indicated. The latter have now become completely obsolete. Note presence of dotted lines at the base of the VH and VCO miospore biozones: the correlations of the latter miospore zonal boundaries with the conodont zones are correct for the Ardenne-Rhenish realm but they are somewhat different for the deposits on the western side of the Acadian Mountains (*i.e.* in Pennsylvania, USA).

11. Structural setting (syndimentary tectonics)

Block tilting and block faulting (*cf.* listric faults grading into vertical faults, related to the Asturian phase, resulting from compression of the original sedimentary basin) have influenced to a large extent the architecture and paleogeographical evolution of the Condruz Group in the Dinant Synclinorium. At least ten tilted and faulted

blocks have been identified by indirect evidences (Thorez *et al.*, 1988; Thorez & Dreesen, 2002). The latter blocks are bordered by NNW-SSE and SSW-NNE lineaments forming a conspicuous chessboard-like pattern (Thorez & Dreesen 1986, Thorez *et al.*, 1988; Paproth *et al.*, 1986). Ball-and-pillow levels in the Montfort, Comblain-la-Tour and Ciney Fms of the Ourthe, Hoyoux and Bock valleys, were supposedly triggered by seismic shocks, related to the above block tilting: the seismic shocks induced abrupt sea-level changes or they have increased locally the accommodation space. Moreover, the ball-and-pillow levels (12 distinct levels in the Ourthe valley) laterally crosscut the different coeval Members or Formations (Fig.7). The chessboard-like configuration of tilted blocks has also influenced the progradation of the Condroz sandstones in the eastern Dinant Synclinorium. Finally, listric faults and regional irregularities of the basin floor were responsible for the lithostratigraphical architecture of the Famennian in the Campine Basin.

12. Reference sections in Belgium and GSSP

The Global Standard Section and Point (GSSP) for both the Frasnian-Famennian boundary and the Famennian-Tournaisian (Devonian-Carboniferous) boundary are located in the Montagne Noire area of Southern France. The base of the Famennian Stage is defined to coincide with the lower boundary of the Lower *Palmatolepis triangularis* Zone. The GSSP for the Frasnian/Famennian (F/F) stage boundary is drawn in a section exposed near the Upper Coumiac Quarry in the southeastern Montagne Noire (Klapper *et al.*, 1993). The F/F boundary records one of the five largest mass extinction events in the fossil record. Dark shales at the boundary are interpreted to be the equivalent to the upper part of the Kellwasser anoxic event recognized throughout the Paleo-Thetys (Sandberg *et al.*, 1988, 2002). Glass spherules believed to be of impact origin are associated with the F/F boundary in 2 Belgian sections (Senzeille and Hony: Claeys & Casier, 1994; Claeys *et al.*, 1996). The top of the Famennian is defined to coincide with the entry of *Siphonodella sulcata*. The GSSP for the D/C boundary is drawn in an artificial trench ("E") near La Serre, southeastern Montagne Noire (Paproth *et al.*, 1991). Most Belgian reference sections for the formations of the Condroz Group are located in the Ourthe valley, between the towns of Esneux and Comblain-la-Tour (NE-part of Dinant Synclinorium). However, their accessibility is strongly hampered nowadays (abandoned sandstone quarries with badly exposed beds). Incomplete reference sections for the formations of the Famenne Group are located in the Famenne area (former railroad and road cuts near the towns of Senzeille and Mariembourg).

13. Main contributions

Beugnies, 1965; Bouckaert, J. *et al.*, 1968; Bouckaert *et al.*, 1969; Bultynck. & Dejonghe, 2001; Dreesen, 1987; Dreesen *et al.*, 1986a.; Dreesen *et al.*, 1986b; Paproth *et al.*, 1986; Sartenaer, 1957a; Thorez & Dreesen, 1986; Thorez *et al.*, 1977; Thorez *et al.*, 1988.

14. Concluding remarks

The lithostratigraphic framework of the Famennian Stage in Belgium and neighbouring areas conforms with the recommendations of Hedberg's International code (1976). The revised stratigraphical correlation framework of the Famennian presented here is the result of several decades of combined litho- and biostratigraphical research. This work started 125 years ago, with the pioneering work of M. Mourlon, on the lithological units ("assises") in the then well-exposed sections of the Belgian type localities. The PhD thesis of one of the senior authors in 1969 has been the starting point for renewed sedimentology and event-stratigraphical interest in the Belgian Famennian Stage. It was the onset of thorough interdisciplinary research by an extensive and international group of colleagues trying to unravel the complex depositional history of the Condroz Group. In order to reach their goal, all the effects of mutually interplaying paleogeographic, tectonic, paleohydrodynamic and paleoclimatologic conditions had to be taken into account. Even now, many interpretation problems still remain unsolved. However, it is a pity that the quality of exposure of the rocks has dramatically decreased over the years, hampering or even excluding future detailed bed-by-bed sampling in the field.

15. Acknowledgements

The authors are indebted to Dr. John Marshall (Southampton, UK) for reviewing the manuscript and to Katleen Van Baelen (VITO, Mol, Belgium) for the excellent artwork (figures).

References

- BECKER, G., BLESS, M.J.M., STREEL, M. & THOREZ, J., 1974. Palynology and ostracode distribution in the Upper Devonian and basal Dinantian in Belgium, and their dependence on sedimentary facies. *Meded. Rijks. Geol. Dienst, N.S.*, 25-2: 9-99.
- BEUGNIES, A., 1965. Contribution à l'étude du Famennien du bord Nord du Bassin de Dinant. *Ann. Soc. Géol. Belgique*, 88 : 411-450.

- BLESS, M.J.M., BOUCKAERT, J. & PAPROTH, E., 1983. Recent exploration in Pre-Permian rocks around the Brabant Massif in Belgium, The Netherlands and the Federal Republic of Germany. *Geologie en Mijnbouw*, 62: 51-62.
- BOUCKAERT, J., ZIEGLER, W. & THOREZ, J., 1965. Conodont stratigraphy of the Famennian Stage. *Serv. Géol. Belgique*, Mem. 5: 1-62.
- BOUCKAERT, J., CONIL, R. & THOREZ, J., 1966. Position stratigraphique de quelques gîtes famenniens à Foraminifères. *Bull. Soc. belge Géol., Paléont., Hydrol.*, 75: 159-175.
- BOUCKAERT, J., STREEL, M. & THOREZ, J., 1968. Schéma biostratigraphique et coupes de référence du Famennien belge. *Ann. Soc. Géol. Belgique*, 91 (3): 317-336.
- BOUCKAERT, J., STREEL, M., THOREZ, J. & MOUND, M.C., 1969. Biostratigraphic chart of the Famennian Stage (Upper Devonian) in the type localities of Belgium: a preliminary report. *Journal of Paleontology*, 43 (3): 727-734.
- BOUCKAERT, J., DREESEN, R. & DRIJKONINGEN, P., 1977. Recherches biostratigraphiques de quelques coupes de l'Avesnois (Nord de la France). *Soc. Géol. Belgique*, 100: 115-123.
- BULTYNCK, P. & DEJONGHE, L., 2001. Devonian lithostratigraphic units (Belgium). *Geologica Belgica*, 4/1: 39-69.
- CLAEYS, P. & CASIER, J.-G., 1994. Microtektite-like impact glass associated with the Frasnian-Famennian boundary mass extinction. *Earth and Planetary Science Letters*, 122: 303-315.
- CLAEYS, Ph., KYTE, F.T., HERBOSCH, A. & CASIER, J.-G., 1996. Geochemistry of the Frasnian-Famennian boundary in Belgium : mass extinction, anoxic oceans and microtektite layer, but not much Iridium? *Geological Society of America Special Paper*, 307: 491-504.
- CONIL, R., 1968. Le Calcaire carbonifère depuis le Tn1a jusqu'au V2a. *Ann. Soc. géol. Belg.*, 90: 687-726.
- CONIL, R. & LYS, M., 1980. Strunien. In : Cavalier & Roger (coord.) *Les étages français et leurs stratotypes*. Mém. BRGM, 109: 26-35.
- d'OMALIUS d'HALLOY, J.B.J., 1868. *Précis élémentaire de géologie*, 8^{ème} édition, C.Muquardt lib., Bruxelles, 1-636.
- DREESEN, R., 1976. Bijdrage tot de biostratigrafische kennis van het Famenniaan: de Souverain-Pré Formatie in het Bekken van Dinant en het Vesdermassief. *Unpubl. Doct. Thesis*, Leuven Univ. (KUL): 1- 133.
- DREESEN, R., 1978. Position stratigraphique de la Formation de Souverain-Pré dans le Synclinorium de Dinant et dans le Bassin de la Vesdre. *Serv. Géol. Belgique, Prof. Paper 1978/2*: 1-74.
- DREESEN, R., 1982a. Storm-generated oolitic ironstones of the Famennian (Fa1b-Fa2a) in the Dinant and Vesdre Synclinoria (Upper Devonian), Belgium. *Ann. Soc. Géol. Belgique*, 105: 105-129.
- DREESEN, R., 1982b. A propos des niveaux d'oolithes ferrugineuses de l'Ardenne et du volcanisme synsédimentaire dans le Massif Ardenno-Rhénan au Dévonien supérieur. *N.Jb.Geol. Paläont.Abh.* 1982 (1): 1-11.
- DREESEN, R., 1987. Oolitic ironstones as event-stratigraphical marker beds within the Upper Devonian of the Ardenno-Rhenish Massif. In: Young, T. & Taylor, W. (Eds) *Phanerozoic Ironstones. Geol.Soc., Spec. Publ.*, 46: 65-78.
- DREESEN, R., 1989. The « Cheiloceras Limestone » a Famennian (Upper Devonian) event-stratigraphical marker in Hercynian Europe and Northwestern Africa ? *Bull. Soc. belge Géol.*, 98-2: 127-133.
- DREESEN, R. & FLAJS, G., 1984. The « Marbre rouge de Baelen », an important algal-sponge-crinoidal carbonate buildup in the Upper Devonian of the Vesdre Massif (E-Belgium). *C.R. Acad. Sci., Paris*, 299, série II, 10: 639-644.
- DREESEN, R. & JUX, U., 1995. Microconchid buildups from the Late Famennian peritidal-lagoonal settings (Evieux Formation, Ourthe Valley, Belgium). *N.Jb. Geol. Paläont/ Abh.*, 198,1/2 : 107-121.
- DREESEN, R. & THOREZ, J., 1980. Sedimentary environments, conodont biofacies and paleoecology of the Belgian Famennian (Upper Devonian). An Approach. *Ann. Soc. Géol. Belgique*, 103: 97-110.
- DREESEN, R. & THOREZ, J., 1994. Parautochthonous-autochthonous carbonates and conodont in the Late Famennian (Uppermost Devonian) Condros Sandstones of Belgium. *Cour. Forsch. Inst. Senckenberg*, 168: 159-182.
- DREESEN, R., BLESS, M.J.M., CONIL, R., FLAJS, G. & LACHET, C., 1985. Depositional environment, paleoecology and diagenetic history of the "Marbre rouge à crinoïdes de Baelen, Upper Devonian, Verviers Synclinorium, Eastern Belgium). *Ann.Soc. Géol. Belgique*, 108: 311-360.
- DREESEN, R., PAPROTH, E. & THOREZ, J., 1986a. Events documented in the Famennian sediments (Ardenne-Rhenish Massif, Late Devonian, NW Europe). In: McMillan, N.J. et al. (Eds) *Proceedings 2nd Intern. Symp. on the Devonian System.*, Calgary, Canada. *Can. Soc. Petrol. Geol.*, 14, II: 295-308.
- DREESEN, R., SANDBERG, C.A. & ZIEGLER, W., 1986b. Review of Late Devonian and Early Carboniferous conodont biostratigraphy and biofacies models as applied to the Ardenne Shelf. In: Bless, M.J.M. & Streel, M. (Eds) *Late Devonian events around the Old Red Continent*. *Ann. Soc. Géol. Belgique*, 109: 27-42.
- DUMONT, A.H., 1832. Mémoire sur la constitution géologique de la province de Liège. *Nouv. Mém. Acad. royale des sciences et des belles lettres de Bruxelles*, coll. in 4°, VIII: 1-374
- DUMONT, A.H., 1848. Mémoire sur les terrains ardennais et rhénans de l'Ardenne, du Rhin, du Brabant et du Condros. *Mém. Acad. Royale de Belgique*, coll. in 4°, XX and XXII: 1-613.
- DUMONT, A.H., 1855. Carte géologique de l'Europe. Ed E. Noblet, Paris, Liège.

- DUMOULIN, V., 2001. Notice explicative de la carte Géologique de Wallonie - Feuille 52/5-6 Grandrieu-Beaumont, *Ministère de la Région wallonne*, Namur: 1-70.
- DUMOULIN, V. & MARION, J.-M., 1997. Notice explicative de la carte géologique de Wallonie – Feuille 52/7-8 Silenrieux-Walcourt. *Ministère de la Région wallonne*, Namur.
- DUSAR, M., 1976. The Lower Famennian at the southeastern border of the Dinant Basin. *Ann. Soc. Géol. Belgique*, 99, II : 565-570.
- DUSAR, M. & DREESEN, R. (1984): Stratigraphy of the Upper Frasnian and Famennian deposits in the region of Hamoir-sur-Ourthe (Dinant Synclinorium, Belgium). *Belg. Geol. Survey Prof. Paper* 1984/5, 209 :1-55.
- GOSSELET, J., 1878. Quelques documents pour l'étude des schistes de la Famenne. *Ann. Soc. Géol. Nord*, 4: 303-320.
- GOSSELET, J., 1879. Nouveaux documents pour l'étude du Famennien. Tranchées de chemin de fer entre Fléron et Semeries. *Ann. Soc. Géol. Nord*, VI : 389-399.
- GOSSELET, J., 1880: Esquisse géologique du Nord de la France et des contrées voisines. 1er fasc. Terrains Primaires. Lille: 1-35.
- GRADSTEIN, F.M., OGG J.G., BLEEKER W. & LOURENS L.J. 2004: A new Geologic Time Scale with special reference to Precambrian and Neogene. *Episodes*, 27: 83-100.
- HECKEL, P.H. & WITZKE, B.J., 1979. Devonian world paleogeography determined from the distribution of carbonates and related lithic paleoclimatic indicators. In: House, M.R. Scrutton, C. & Basset, M.C. (Eds) "The Devonian System", *Spec. Pap. In Paleontology*, 23: 99-124.
- HEDBERG, H.H., 1976 (Ed.). International stratigraphical guide. A guide to stratigraphic classification, terminology, and procedure. *International Subcommission on stratigraphic classification (ISSC)*, J. Wiley & Sons: 1-200.
- KLAPPER, G., FEIST, R., BECKER, R.T. & HOUSE, M., 1993. Definition of the Frasnian-Famennian Stage boundary. *Episodes*, 16/4: 433-441.
- LAENEN, B., 2003: Lithostratigrafie van het pre-Tertiair in Vlaanderen, Deel II: Dinantiaan & Devoon. *Unpublished Vito-report*, commissioned by the Afdeling Natuurlijke Rijkdommen en Energie (ANRE) of the Flemish Region, 2003/ETE/R/095, July 2003: 1-86.
- LAENEN, B., DREESEN, R. & ROELANDTS, I., 2002. Sequence-stratigraphic significance and comparative REE-fractionation patterns of Rupelian glaucony concentrates and Famennian oolitic ironstones (Belgium). In: Degryse, P. & Sintubin, M. (Eds), *Contributions to the Geology of Belgium and Northwest Europe. Proceeds. 1st Geologica Belgica International Meeting*. Aardk. Mededel. 12: 51-54.
- LALOUX, M., DEJONGHE, L., GHYSEL, P. & HANCE, L., 1996. Notice explicative de la carte géologique de Wallonie – feuille 42/7-8 Fléron-Verviers. *Ministère de la Région wallonne*, Namur: 1-150.
- MAMET, B. & PREAT, A., 2003. Sur les difficultés d'interprétation des hiatus stratigraphiques (exemple tiré de la transition Dévono-Carbonifère, bassin de Dinant). *Geologica Belgica*, 6/1-2: 3-23.
- MICHOT, P., 1980. Belgique- Introduction à la géologie générale. Excursion 211 A, 26^{ème} Congr. Intern. Géol., Paris : 485-576.
- MOURLON, M., 1875-1886. Monographie du Famennien, comprenant les Psammites du Condroz et les Schistes de la Famenne proprement dits (Dévonien supérieur). *Bull. Acad. Royale des Science, des Lettres et des Beaux-Arts de Belgique*, Bruxelles, 1875, 2^{ème} sér., 39 : 602-659 ; 1875, 2^{ème} série, 40 : 761-796 ; 1876, 2^{ème} série, 42 : 854-884 ; 1882, 3^{ème} série, 4 : 504-525 ; 1884, 3^{ème} série, 7 : 295-303 ; 1885, 3^{ème} série, 9 : 238-254 ; 1886, 3^{ème} série, 12 : 369-416 ; 1886, 3^{ème} série, 12 : 613-622.
- MOURLON, M., 1885. Compte rendu d'excursion dans le Famennien de la vallée de l'Ourthe. *Ann. Soc. Géol. Belgique*, 12: 97-107.
- PAPROTH, E., DREESEN, R. & THOREZ, J., 1986. Famennian paleogeography and event stratigraphy in North-western Europe. In: Bless, M.J.M. & Streel, M. (Eds) *Late Devonian events around the Old Red Continent*. *Ann. Soc. Géol. Belgique*, 109: 175-186.
- PAPROTH, E., FEIST, R. & FLAJS, G., 1991. Decision on the Devonian-carboniferous boundary stratotype. *Episodes*, 14/4: 331-336.
- SANDBERG, C.A., ZIEGLER, W., DREESEN, R., & BUTLER, J.L., 1988. Late Frasnian mass extinction: condont event stratigraphy, global changes and possible causes. *Courrier Forschungsinstitut Senckenberg*, 102: 263-307.
- SANDBERG, C.A., MORROW, J.R. & ZIEGLER, W., 2002. Late Devonian sea-level changes, catastrophic events and mass extinctions. In: Koeberl C. & MacLeod, K.G., (Eds), *Catastrophic Events and Mass Extinctions: Impacts and Beyond*. *Geological Society of America Special Paper* 356: 473-487.
- SARTENAER, P., 1957a. Esquisse d'une division stratigraphique nouvelle des dépôts du Famennien inférieur du Bassin de Dinant. *Bull. Soc. Belg. Géol.*, LXV, 3: 421-446.
- SARTENAER, P., 1957b. La découverte en Belgique de la Zone à Cheiloceras. *Ann. Soc. Géol. Belgique*, LXXX, 1956-1957 : B 463-470.
- STREEL, M., 2005. Subdivision of the Famennian Stage into four substages and correlation with the neritic and continental miospore zonation (SDS Business Meeting, Florence, August 23, 2004). *IUGS Subcomm. Devonian Stratigraphy, Newsletter* 21: 14, 16-17..
- STREEL, M., BRICE, D., DEGARDIN, J.-M., DERYCKE, Cl., DREESEN, R., GROESSENS, E., HANCE, L., LEGRAND-BLAIN, M., LETHIERS, F., LOBOZIAK, S., MAZIANE N., MILHAU, B., MISTIAEN, B., POTY, E., ROHART, J.-C., SARTENAER, P., THOREZ, J., VACHARD, D., BLIECK, A. 1998.

- Proposal for a Strunian Substage and a division of the Famennian Stage into four Substages. *IUGS Subcomm. Devonian Stratigraphy, Newsletter*, 15: 47-49.
- STREEL, M., BELKA, Z., DREESEN, R., DURKINA, A.V., GROOS-UFFENORDE, H., HANCE, L., HARTKOPF-FRÖDER, C., HAYDUKIEWICZ, J., KORN, D., PERRI, M.-C., PIECHA, M., SPALLETTA, C. 2005: Relation of the neritic microfaunas and continental microfloras with the conodont and other pelagic faunas of the latest part of the Famennian (SDS Business Meeting, Florence, August 23, 2004), *IUGS Subcomm. Devonian Stratigraphy, Newsletter 2005* : 17-20.
- THOREZ, J., 1969. Sédimentologie du Famennien supérieur dans le Synclinorium de Dinant. *Unpubl. Doct. Thesis (available on CD-Rom from the Geol. Dept. Library) Liège Univ.*: 1-225.
- THOREZ, J., 2002. Paleosols and inferred paleoclimatology in the Condroz Sandstone (Upper Famennian, northeastern part of the Dinant Synclinorium, Belgium). *In: Degryse, P. & Sintubin, M. (Eds) Contributions to the Geology of Belgium and Northwest Europe, Proceeds. 1st Geologica Belgica International Meeting. Aardk. Mededel.*, 12: 145-148.
- THOREZ, J. & DREESEN, R., 1986. A model of a regressive depositional system around the Old Red Continent as Exemplified by a Field Trip in the Upper Famennian "Psammities du Condroz" in Belgium. *In: Bless, M.J.M. & Streel, M. (Eds), Symposium on the Old Red Continent, Aachen, 1986. Ann. Soc. Géol. Belgique*, 109 : 285-323.
- THOREZ, J. & DREESEN, R., 1997. Sequence- and (bio)event stratigraphy: a tool for global correlation in Euramerica, as exemplified by the late Upper Devonian in Belgium. *1st Intern. Conf. on North Gondwana Mid-Paleozoic biodynamics (IGCP 421), Vienna, 17-21 September, Abstract*: 56-61.
- THOREZ, J. & DREESEN, R., 2002. The Condroz Sandstone Group (Upper Famennian) Revisited. Leuven 2002 "On the crossroads...", *Geologica Belgica Intern. Meeting*, 11-15 September 2002, Field Trip Guide book.
- THOREZ, J., STREEL, M., BOUCKAERT, J. & BLESS, M.J.M., 1977. Stratigraphie et paléogéographie de la partie orientale du Synclinorium de Dinant (Belgique) au Famennien supérieur : un modèle de bassin sédimentaire reconstitué par analyse pluridisciplinaire sédimentologique et micropaléontologique. *Med. Rijks. Geol. Dienst (The Netherlands), Nrw. Ser.*, 28: 17-32.
- THOREZ, J., GOEMAERE, E. & DREESEN, R., 1988. Tide- and Wave-influenced depositional environments in the Psammities du Condroz (Upper Famennian) in Belgium. *In: de Boer, P.L. et al. (Eds) Tide-Influenced Sedimentary Environments and Facies*, D.Reidel Publ., The Netherlands: 389-415.
- VANGUESTAINE, M., DECLAIRFAYT, Th., ROUHART, A. & SMEESTERS, A., 1983. Zonation par acritarches du Frasnien Supérieur – Famennien Inférieur dans les Bassins de Dinant, Namur, Herve et Campine (Dévonien Supérieur de Belgique). *Ann. Soc. Géol. Belgique*, 106: 121-171.
- WALLISER, O.H., 1996. Global events in the Devonian and Carboniferous, *In: Walliser, O.H. (ed.): Global Events and Event Stratigraphy in the Phanerozoic*. Springer, Berlin, 225-250.
- WATERLOT, G., BEUGNIES, A. & GODFRIAUX, Y., 1967. Carte géologique détaillée de la France à l'échelle de 1/50.000: Maubeuge – XXVIII-6, *Ed. Bureau de Recherches Géologiques et Minières*.

Manuscript received on 09.11.2004 and accepted for publication on 17.10.2005.