

WARNANTIAN

Edouard POTY¹ & Luc HANCE²

(4 figures)

1. *Paléontologie animale et humaine, Département de Géologie, Université de Liège, Bât. B18, Allée du 6 Août, Sart Tilman, 4000-Liège, Belgium; E-mail: e.poty@ulg.ac.be.*
2. *Carmeuse Coordination Center, Bd de Lauzelle, 65, 1348 Louvain-la-Neuve, Belgium; E-mail: Luc.hance@skynet.be.*

ABSTRACT. The Warnantian Substage is about 8 to 12 Ma long and was defined as corresponding to the Cf6 Foraminifer Zone by Conil *et al.*, 1977 (= MFZ13 to MFZ15 Zones of Devuyst & Hance (in Poty *et al.*, in press). Its base is taken at the base of the Bonne River Formation (Fm); its top is defined by the base of the Namurian stage, but its upper part is almost completely missing in the Namur-Dinant Basin. The substage is covered by the RC7 and RC8 Coral Zones. It comprises the Bonne River and the Anhée Fms, which are mainly composed of shallowing-upward parasequences which were deposited by aggradation on the shelf, as suggested by their lateral regularity. It corresponds to the HST of the third-order sequence 8 (Thon-Samson Member of the Bonne River Fm) and to the third-order sequences 9 and 10 (Hance *et al.*, 2001; Poty *et al.*, 2002). The lower (MFZ13-MFZ14, RC7) and the upper Warnantian (MFZ15, RC8) can be correlated respectively with the British Asbian and Brigantian Substages.

KEYWORDS: Warnantian, Viséan, Carboniferous, Belgium, lithostratigraphy, biostratigraphy, palaeogeography

Reference to this paper: Poty, E. & Hance, L., 2006. Warnantian. In Dejonghe, L., ed., Current status of chronostratigraphic units named from Belgium and adjacent areas, *Geologica Belgica*, Brussels, 9/1-2: 139-144.

1. Name

Warnantian (English), Warnantiaan (Dutch), Warnantium (German), Warnantien (French).

2. Age

From 331 to 319 Ma (i.e. 12 Ma long) in a time scale with minimum ages and from 334.5 to 326.5 (i.e. 8 Ma long) in a time scale with maximum ages (Menning *et al.*, 2001)

3. Authors

The base of the substage is defined by Conil, R., Groessens, E. & Pirlet, H. (1977. Nouvelle charte stratigraphique du Dinantien type de la Belgique. *Annales de la Société géologique du Nord*, XCVI : 363-371) at the « base of the thick bed of dark bioclastic limestone overlying the beige algal limestones of the top of the Livian » (free translation from French), i.e. the base of the Thon-Samson Member (Mbr) of the Bonne River Formation (Fm). The substage corresponds to the *Neoarchaediscus* Zone (Cf6) of Conil *et al.* (1977). Its top is defined by the base of the Namurian Stage.




Figure 1. Location of the Warnantian stratotype.

4. Historical type area

The Warnantian Substage is named after Warnant, a locality in the Mollignée valley, near Dinant (Dinant sedimentation area), where the substage is well exposed (De Jaiffé quarry and railway cutting at Warnant; Demanet, 1938; Conil & Pirlet, 1970; Conil & Pirlet, 1974; Laloux *et al.*, 1988). However, the « Camp de César » old quarry, at Thon-Samson (Figs 1, 2; Pirlet, 1963; Pirlet, 1968; Conil & Pirlet, 1974) in the Namur sedimentation

area (NSA), was chosen as boundary stratotype by Conil *et al.* (1977). In the Samson valley, the Transcar quarry at Maizeret and the old quarry (Fig. 3) in front of the junction between the valley road and the road to Maizeret (Pirlet, 1963, 1968) are good boundary parastratotypes. Geological maps n° 145 (Andenne-Couthuin, 1893) and n° 155 (Gesves-Ohey, 1901), under revision by Delcambre & Pingot (in preparation).

5. Description

In the stratotype area, only the lowest part of the Warnantian, including the Thon-Samson Mbr and the lower part of the Poilvache Mbr, is exposed (Fig. 2, 4). This part corresponds to the Cf6 α , β Subzones of Conil *et al.* (1991). The rest of the substage lacks and the Namurian siliciclastics directly top the Poilvache Mbr.

The Thon-Samson Mbr rests on the stromatolitic limestones of the Bay-Bonnet Mbr (top of the Livian Substage), and comprises at its base a shallowing-upward parasequence composed of 3 m of grey bioclastic limestone with cherts, capped by a 30 cm-thick bed of stromatolitic limestone. It is topped by 5 to 6 m of massive grey crinoidal limestone.

The Poilvache Mbr is composed of 6,5 m of thin-bedded stromatolitic limestone interbedded with some argillaceous layers (« plates escailles ») overlain by 14 m of thick-bedded pale stromatolitic to bioclastic limestones.

The most complete sections of the substage are in the Condroz (CSA) and in the Dinant (DSA) sedimentation areas. But the upper Warnantian (corresponding to the Cf6 δ Foraminiferal Subzone of Conil *et al.*, 1991, ~ Zone MFZ15 – *Janischewskina typica* Interval Zone of Devuyt & Hance (in Poty *et al.*, in press) and to the RC8 Rugose Coral Zone of Poty, 1985), is always poorly developed and often almost totally absent.



Figure 2. Base of the Warnantian Substage in its stratotype (“Camp de César” old quarry at Thon-Samson).

6. Historical background

The Viséan was formerly considered as a series and was divided into three stages by Conil *et al.* (1977), from base to top: Moliniacian (lower Viséan), Livian (middle Viséan) and Warnantian (upper Viséan). The Viséan was recently redefined as a stage by the IUGS Subcommission on Carboniferous Stratigraphy (Heckel, 2004) and, as a consequence, the Warnantian is now a substage. The British Asbian and Brigantian Substages correspond respectively to the lower and the upper Warnantian.




Figure 3. Base of the Warnantian in the old quarry in front of the junction between the valley road and the road to Maizeret at Thon-Samson (parastratotype).

7. Lithology

The lithology and the lithostratigraphy (Fig. 4) of the substage are relatively constant in the Namur-Dinant basin, but deposits are locally missing: the gap includes the uppermost Warnantian in the DSA and progressively extends stratigraphically downwards up to the uppermost Livian in the NSA. A deep palaeokarst filled with Namurian siliciclastic deposits is sometimes developed at the top of the substage.

In the CSA and DSA, where the substage is best developed, it is about 120 m-thick and comprises two formations: the Bonne River Fm and the Anhée Fm.

The Bonne River Fm comprises the Thon-Samson and the Poilvache Mbrs. The Thon-Samson Mbr (sensu Poty *et al.*, 2002, is 8 m-thick and composed of massive well-bedded, pale to dark limestone (grainstones-rudstones), usually crinoidal, sometimes with corals and brachiopods. The Poilvache Mbr (ibid.) is about 80 m-thick and comprises pale to dark limestones, sometimes cherty, organized in shallowing-upward parasequences dominated by lime mudstones and stromatolitic limestones. It is poor in fossils.

The Anhée Fm (ibid.) is composed of two members: (1) the Lower Mbr is about 25 m-thick and comprises parasequences of dark limestones dominated by wackestones and packstones, locally very fossiliferous (Chabôfosse Facies); (2) the Upper Mbr, a few metres to 8 m-thick, comprises argillaceous limestones, shales and siliceous shales. The upper part of the Lower Mbr and the Upper Mbr were previously attributed respectively to the lower and to the upper parts of the Warnant Fm (Paproth *et al.*, 1983).

The Bonne River Fm and most of the Lower Mbr of the Anhée Fm correspond to the lower Warnantian. The top of the Lower Mbr and the Upper Mbr of the Anhée Fm correspond to the upper Warnantian that is very poorly developed and often missing.

Note that collapse brecciation triggered by the solution of the underlying Livian evaporites (Grands-Malades Fm) can affect the Warnantian limestones ("Grande Brèche Viséenne").

In the Visé-Maastricht sedimentation area (VSA), the Substage corresponds to the upper part of the Visé Limestone Fm (Pirlet, 1967; Poty *et al.*, 2002), which is the historical type unit of the Viséan Stage, and to the upper part of the Berneau Fm (Poty *et al.*, 2002).

The Visé Limestone comprises pale to grey sedimentary limestone breccias, turbidites and massive algal and bioclastic boundstones forming buildups very rich in fossils. The Berneau Fm is mainly composed of dark limestone turbidites and breccias. Most of the upper part of the Substage is also missing in the area. The thickness of the Substage varies from 0 m in the Souvré tectonic block to some hundreds metres in the Maastricht block system (Poty, 1991).

In the Avesnois (Mansy *et al.*, 1989), the substage corresponds to the Saint-Hilaire and Saint-Rémy-Chaussée Limestones (lower Warnantian) and to the Queue Noir-Jean shales and silicified shales (upper Warnantian). In the Boulonnais, it comprises the Joinville and the Réty Fm; its uppermost part is also missing (Poty, 1994).

8. Sedimentology and palaeogeography

The Thon-Samson Mbr corresponds to the HST of the third-order sequence 8 of Hance *et al.* (2001), while the Poilvache Mbr and the Anhée Fm correspond respectively to the TST (and LST?) and to the HST (and FSST?) of sequence 9. The sequence 10 is not developed in the basin. During the Warnantian, the Visé-Maastricht sedimentation area belonged to the Campine Basin which was separated from the Namur-Dinant Basin by the emergent Brabant Massif. Therefore, the Warnantian deposits in the Campine Basin are not similar to those known in the Namur-Dinant Basin (see above). The Warnantian is better developed in Great-Britain and Ireland where it is locally more than 1000 m-thick (George *et al.*, 1976).

9. Palaeontology

9.1. Foraminifers

The Warnantian is covered by the MFZ13 to MFZ15 foraminifer zones of Devuyt & Hance (in Poty *et al.*, in press). It is equivalent to the Cf6 Zone of Conil *et al.* (1977, 1991), which is well documented by Laloux (1988) and Laloux *et al.* (1988).

The MFZ13 Zone starts with the entry of *Neoarchaediscus* and *Planospirodiscus* in the lowermost part of the Thon-Samson Mbr in its type area (NSA), in sequence 1 of Pirlet (1968; see also Laloux, 1988). *Vissariotaxis* sp., *Palaeotextularia* with a bilaminar wall and typical *Endothyra spira* enter at this level. *Chomatemediocris* sp., *Neoarchaediscus incertus*, *Vissariotaxis compressa* and primitive *Howchinia* are other typical elements. *Pseudoendothyra* sp. appears higher in the zone.

The MFZ14 Zone, equivalent to the Cf6γ Subzone and the lower part of the Cf6δ Subzone of Conil *et al.* (1977, 1991), covers the uppermost part of the Poilvache Mbr and most of the Lower Member of the Anhée Fm (sensu Poty *et al.* 2002). *Howchinia bradyana* and *Bradyina rotula*




Figure 4. Stratigraphical pattern of the Warnantian in southern Belgium.

enter in the uppermost part of the Poilvache Mbr (DSA), together with cribrate *Palaeotextulariidae* (*Cribrostomum* and *Koskinobigenarina*). The successive incomings of *Cribrospira panderi*, *Bibradya* sp., *Asteroarchaediscus* sp., *Globispiroplectammina* sp. and *Plectostaffella* sp. complete the association. *Archaediscus* ex gr. *karreri*, *Endothyranopsis crassa*, large *Omphalotis* and *Eostaffella* aff. *mosquensis* are also present. Archaeodiscids at the 'tenuis stage' appear in the uppermost part of the Zone. *Loeblichia paraammonoides* enters in the upper part of the lower member of the Anhéé Fm in the DSA and CSA, at the same level or at a short distance above the entry of *Warnantella* (Laloux, 1988; Laloux, in Poty *et al.* 1988). The base of the MFZ15 Zone is defined at the entry of *Janischewskina typica* a few metres above *L. paraammonoides*. It correlates with the upper part of the Cf6 δ Subzone and with the lower part of the Cf7 Zone that does not yield significant new taxa. Most of the key taxa typical of Zone MFZ14 are still present. *Asteroarchaediscus* and archaeodiscids at the 'tenuis stage' are abundant. The first *Monotaxinoides* are reported by Conil *et al.* (1991) in the upper part of the zone.

9.2. Corals

The Warnantian is covered by the rugose coral Zones RC7 and RC8 (Poty, 1985; Conil *et al.*, 1991; Poty *et al.*, in press).

In the type area, the first coral assemblage of the Zone RC7 is recorded in the massive crinoidal limestone of the upper part of the Thon-Samson Mbr. The new characteristic taxa are *Siphonodendron scaleberense*, *S. pauciradiale*, *Siphonophyllia samsonensis* (= *S. benburbensis*) and *Dibunophyllum* sp. The genus *Diphyphyllum* (*D. furcatum*) appears in the overlying Poilvache Mbr (sequence "i" of Pirlet, 1968).

The Anhéé Fm is characterized by the appearance of *Lithostrotion maccoyanum*, *Siphonodendron junceum*, *Diphyphyllum lateseptatum*, *D. fasciculatum*, and later the genus *Aulophyllum* (*A. fungites*). Other common taxa appearing in the RC7 Zone are: *Lithostrotion decipiens*, *Siphonodendron intermedium*, *Clisiophyllum keyserlingi*, *Dibunophyllum bipartitum* and *Pseudozaphrentoides juddi*. The genera *Lonsdaleia sensu lato* (including fasciculate, subcerioid and cerioid species) and *Palastraea* appear in the upper part of the Lower Member of the Anhéé Fm and mark the base of the RC8 Zone.

RC7 and RC8 zones are easily correlatable everywhere in Europe.

9.3. Other fossils

Brachiopods are common, but they need to be revised to be used for the stratigraphy of the substage. The occurrence of gigantoproductids in the substage is worth mentioning. The Warnantian corresponds to the top of the *Gnathodus homopunctatus* – *Taphrognathus transatlanticus* and to the *Gnathodus bilineatus* Conodont Zones

(Paproth *et al.*, 1983), and to the goniatite faunas GF10 to GF16 of Ramsbottom (in Paproth *et al.*, 1983). However conodonts and goniatites are very uncommon and without practical biostratigraphical value in Belgium.

10. Chronostratigraphy

Fig. 4 gives the Foraminiferal and Coral zonations used in Western Europe. More details about the chronostratigraphy and the stratigraphical correlations outside Western Europe are given in Poty *et al.* (in press).

11. Geochronology

No radiometric dates are available for the Warnantian in Belgium. The time scale used is based on datations obtained from other Carboniferous basins in Europe and in Australia (Menning *et al.*, 2001).

12. Structural setting

Warnantian deposits are part of the Variscan orogenic cycle. Those of the Namur-Dinant Basin were deposited by aggradation on the shelf as suggested by their lateral regularity, while those of the Campine Basin were affected by tectonic block faulting. In southern Belgium, Warnantian deposits are distributed on both sides of the Midi-Eifel fault zone and occur in two main structural units: the Brabant Parautochton and the Ardenne Allochton. These structural units were part of the Namur-Dinant Basin during the time of deposition. Warnantian deposits are largely folded except for those situated in the northern part of the Brabant Parautochton. The Warnantian stratotype is located on the southern flank of the Namur Synclinorium, at the front of the Variscan folding and the beds are gently folded. In the Campine Basin, Warnantian deposits are faulted but not folded, excepted for those situated in the Visé area which suffered Variscan folding.

13. Reference sections in Belgium

In the DSA, three sections are important reference sections: The old "De Jaiffe" quarry and the railway cutting at Warnant (Demanet, 1938; Pirlet, 1968; Conil & Pirlet, 1970; Conil & Pirlet, 1974; Laloux *et al.*, 1988), a section situated in front of the Yvoir bridge at Anhéé and the disused Watrisse quarry (Pirlet, 1968; Conil & Pirlet, 1970) south of Anhéé. In the CSA, at Royseux (Hoyoux valley), sections along the road (Pirlet, 1964) and on the slopes of both sides of the valley (Poty *et al.*, 1988; Aretz, 2001) expose very fossiliferous limestones of the Anhéé Fm ("Chabôfosse Facies") overlying the Bonne River Fm.

The latter formation is well exposed on the east bank of the river Bonne at Modave (Pirlet, 1968). In the NSA, the lower part of the substage is well exposed in the Samson valley, in the Transcar quarry at Maizeret (fig. 1) and in an old quarry in front of the junction between the valley road and the road to Maizeret (right side of the valley, fig. 1, 3; Pirlet, 1963, 1968). The Warnantian part of the Visé Limestone (historical Viséan stratotype) is exposed in old quarries situated south of Visé (Pirlet, 1967), but access is now very difficult.

14. Main contributions

Conil, Groessens & Pirlet (1977); Conil & Pirlet (1970); Devuyt, Hance & Poty, 2005; Hance, Poty & Devuyt (2001, 2002); Laloux *et al.* (1988); Paproth *et al.* (1983); Pirlet (1963, 1964, 1967, 1968); Poty *et al.* (2002).

15. Remarks

The base of the Warnantian Substage is placed at the bioclastic limestone bed which marks the base of the Bonne River Fm (Thon-Samson Mbr). The underlying Bay-Bonnet Mbr is composed of stromatolitic limestones and is almost totally devoid of fossils. Therefore, the appearance of the characteristic taxa of the substage is due to ecological factors allowing their presence, and a lectostratotype has to be searched in a succession developed in open marine facies.

16. Acknowledgements

M. Aretz (Universität zu Köln) and F.X. Devuyt (Trinity College, Dublin) kindly corrected the English and commented on an early version of this paper. Very pertinent comments of D. Vachard on the foraminifer associations were appreciated.

References

- ARETZ, M., 2001. The upper Viséan coral-horizons of Royseux – the development of an unusual facies in Belgian Early Carboniferous. *Tohoku University Museum Bulletin*, 1: 86-95.
- CONIL, R., GROESSENS, E., LALOUX, M., POTY, E. & TOURNEUR, F., 1991. Carboniferous Guide Foraminifera, Corals and Conodonts in the Franco-Belgian and Campine Basins: their potential for widespread correlations. *Courier Forschungsinstitut Senckenberg*, 130: 15-30.
- CONIL, R., GROESSENS, E. & PIRLET, H., 1977. Nouvelle charte stratigraphique du Dinantien type de la Belgique. *Annales de la Société géologique du Nord*, XCVI: 363-371.
- CONIL, R. & PIRLET, H., 1970. Le Calcaire Carbonifère du Synclinorium de Dinant et le sommet du Famenien. In: Colloque sur la stratigraphie du carbonifère, STREEL, M. & WAGNER, R.H., édité., Les Congrès et Colloques de l'Université de Liège, 55: 47-63.
- CONIL, R. & PIRLET, H., 1974. Excursion A; In BOUCKAERT, J. & STREEL, M., International Symposium on Belgian Micropaleontological Limits, Namur 1974, Guidebook; Ministry of Economic Affairs, Geological Survey of Belgium: 1-11.
- DEMANET, F., 1938. La faune des couches de passage du Dinantien au Namurien. *Mémoire de l'Institut Royal des Sciences Naturelles de Belgique*, 84: 1-201.
- DEVUYST, F.-X., HANCE, L. & POTY, E., 2005. The Dinantian of Southern Belgium revisited: sedimentary history and biostratigraphy. A guidebook of key sections. I.U.G.S. Subcommittee on Carboniferous Stratigraphy field trip, May 2005: 1-79.
- GEORGE, T.N., JOHNSON, G.A.L., MITCHELL, M., PRENTICE, J.E., RAMSBOTTOM, W.H.C., SEVASTOPULO, G.D. & WILSON, R.B., 1976. A correlation of Dinantian rocks in the British Isles. *Geological Society of London, special report 7*: 1-87.
- HANCE, L., POTY, E. & DEVUYST, F.-X., 2001. Stratigraphie séquentielle du Dinantien type (Belgique) et corrélation avec le Nord de la France (Boulonnais, Avesnois). *Bulletin de la Société Géologique de France*, 172, 4: 411-426.
- HANCE, L., POTY, E. & DEVUYST, F.X., 2002. Sequence stratigraphy of the Belgian Lower Carboniferous – tentative correlation with the British Isles. In HILLS, L.V., HENDERSON, C.M. & BAMBER, W. (eds.) Carboniferous and Permian of the World. *Canadian Society of Petroleum Geologists Memoirs*, 19: 41-51.
- HECKEL, P.H., 2004. Chairman's column. *Newsletter on Carboniferous stratigraphy, I.U.G.S. Subcommittee on Carboniferous Stratigraphy*, 22: 1-3.
- LALOUX, M., 1988. Foraminifères du Viséen supérieur et du Namurien du bassin franco-belge. In LALOUX, M., BOUCKAERT, J., CONIL, R., GROESSENS, E., LAURENT, S., OVERLAU, P., PIRLET, H., POTY, E., SCHILTZ, M. & VANDENBERGHE, N., Pre Congress excursion to the Carboniferous stratotypes in Belgium. *Bulletin de la Société belge de Géologie*, 96 (3): 205-220.
- LALOUX, M., GROESSENS, E., OVERLAU, P. & PIRLET, H., 1988. Warnant. In: LALOUX, M., BOUCKAERT, J., CONIL, R., GROESSENS, E., LAURENT, S., OVERLAU, P., PIRLET, H., POTY, E., SCHILTZ, M. & VANDENBERGHE, N., Pre Congress excursion to the Carboniferous stratotypes in Belgium. *Bulletin de la Société belge de Géologie*, 96 (3): 221-226.
- MANSY, J.L., CONIL, R., MEILLIEZ, F., KHATIR, A., DELCAMBRE, B., GROESSENS, E., LYS, M., POTY, E., SWENNEN, R., TRENTESAUX, A. & WEYANT, M., 1989. Nouvelles données stratigraphiques

- et structurales sur le Dinantien de l'Avesnois. *Annales de la Société Géologique du Nord*, CVIII: 125-142.
- MENNING, M., WEYER, D., DROZDZEWSKI, G. & WENDT, I., 2001. More radiometric ages for the Carboniferous time scale. *Newsletter on Carboniferous Stratigraphy*, IUGS Subcommission on Carboniferous Stratigraphy, 19: 16-18.
- PAPROTH, E., CONIL, R., BLESS, M. J. M., BOONEN, P., BOUCKAERT, J., CARPENTIER, N., COEN, M., DELCAMBRE, B., DEPRIJCK, CH., DEUZON, S., DREESEN R., GROESSENS, E., HANCE, L., HENNEBERT, M., HIBO, D., HAHN, G., HAHN, R., HISLAIRE, O., KASIG, W., LALOUX, M., LAUWERS, A., LEES, A., LYS, M., OP DE BEEK K., OVERLAU, P., PIRLET H., POTY E., RAMSBOTTOM, W., STREEL M., SWENNEN, R., THOREZ, J., VANGUESTAINE, M., VAN STEENWINKEL, M. & VIESLET, J. L. 1983. Bio- and lithostratigraphic subdivisions of the Dinantian in Belgium, a review. *Annales de la Société Géologique de Belgique*, 106: 185-239.
- PIRLET, H., 1963. Sédimentologie des formations du Viséen supérieur, V3b dans la vallée du Samson (Bassin de Namur, Belgique). *Annales de la Société Géologique de Belgique*, 86, 1: 1-41.
- PIRLET, H., 1964. Lithologie, stratigraphie et tectonique du Viséen supérieur de Royseux (bord sud du Synclinorium de Dinant). *Annales de la Société Géologique de Belgique*, 86, 8: 397-404.
- PIRLET, H., 1967. Nouvelle interprétation des carrières de Richelle : le Viséen de Visé (2^e note sur les calcaires de Visé). *Annales de la Société Géologique de Belgique*, 90, 4-6: 299-328.
- PIRLET, H., 1968. La sédimentation rythmique et la stratigraphie du Viséen supérieur V3b, V3c inférieur dans les synclinoriums de Namur et de Dinant. *Académie Royale de Belgique, classe des sciences, mémoires in-4^e, 2^e série*, XVII: 1-98.
- POTY, E., 1985. A rugose coral biozonation for the Dinantian of Belgium as a basis for a coral biozonation of the Dinantian of Eurasia. *Compte-rendu X Congrès International de stratigraphie et de Géologie du Carbonifère*, Madrid 1983, 4: 29-31.
- POTY, E., 1991. Tectonique de blocs dans le prolongement oriental du Massif du Brabant. *Annales de la Société Géologique de Belgique*, 114, 1: 265-275.
- POTY, E., 1994. Nouvelles précisions sur les corrélations stratigraphiques du Dinantien du Boulonnais et de la Belgique : application de la biozonation corallienne. *Comptes Rendus de l'Académie des Sciences, Paris*, 319, s. II: 467-473.
- POTY, E., ARETZ, M. & BARCHY, L., 2002. Stratigraphie et sédimentologie des « Calcaires à *Productus* » du Carbonifère inférieur de la Montagne Noire (Massif Central, France). *Comptes Rendus Géoscience*, 334: 843-848.
- POTY, E., CONIL, R., GROESSENS, E., LALOUX, M. & LAURENT, S., 1988. Royseux. In LALOUX, M., BOUCKAERT, J., CONIL, R., GROESSENS, E., LAURENT, S., OVERLAU, P., PIRLET, H., POTY, E., SCHILTZ, M. & VANDENBERGHE, N., Pre Congress excursion to the Carboniferous stratotypes in Belgium. *Bulletin de la Société belge de Géologie*, 96 (3): 243-47.
- POTY, E., DEVUYST, F.-X. & HANCE, L., in press. Upper Devonian and Mississippian foraminiferal and rugose coral zonation of Belgium and Northern France: a tool for Eurasian correlations. *Geological Magazine*.
- POTY, E., HANCE, L., LEES, A. & HENNEBERT, M., 2002. Dinantian lithostratigraphic units (Belgium) . In BULTYNCK, P. & DEJONGHE, L., eds., Guide to a revised lithostratigraphic scale of Belgium. *Geologica Belgica*, 4/1-2: 69-94.

Manuscript received on 14.05.2005 and accepted for publication on 17.10.2005.