

Emerging and recurring diseases in cetaceans worldwide and the role of environmental stressors

Marie-Françoise Van Bressem¹, Juan Antonio Raga², Giovanni di Guardo³, Paul Jepson⁴, Pdraig Duignan⁵, Ursula Siebert⁶, Tom Barrett⁷, Marcos César de Oliveira Santos⁸, Ignacio Moreno⁹, Salvatore Siciliano¹⁰, Alex Aguilar¹¹ and Koen Van Waerebeek¹

¹ Cetacean Conservation Medicine Group (CMED), CEPEC, Museo de Delfines, Pucusana, Peru; ² Marine Zoology Unit, Cavanilles Institute of Biodiversity and Evolutionary Biology, University of Valencia, P.O. Box 22085, 46071 Valencia, Spain; ³ University of Teramo, Faculty of Veterinary Medicine, Department of Comparative Biomedical Sciences, Piazza Aldo Moro, 45 - 64100 - Teramo, Italy; ⁴ Institute of Zoology, Regent's Park, London NW1 4RY, United Kingdom; ⁵ Veterinary Laboratory Agency, Luddington, Warwickshire, CV37 9SJ, U.K.; ⁶ Christian Albrechts University, D-25761 Buesum, Germany; ⁷ Institute for Animal Health, Pirbright, UK; ⁸ Projeto Atlantis, Laboratório de Biologia da Conservação de Cetáceos, Departamento de Zoologia, Instituto de Biociências, Universidade Estadual Paulista "Júlio de Mesquita Filho" (UNESP), Campus Rio Claro, Brasil, 13506-900; ⁹ Grupo de Estudos de Mamíferos Aquáticos do Rio Grande do Sul (GEMARS) & Centro de Estudos Costeiros, Limnológicos e Marinhos da Universidade Federal do Rio Grande do Sul (CECLIMAR/UFRGS), Rua Felipe Néri, 382/203, 90440-150 Porto Alegre RS, Brazil; ¹⁰ Grupo de Estudos de Mamíferos Marinhos da Região dos Lagos (GEMM-Lagos) & Laboratório de Ecologia, Departamento de Endemias Samuel Pessoa, Escola Nacional de Saúde Pública/FIOCRUZ, Rua Leopoldo Bulhões, 1480-térreo, Manguinhos, Rio de Janeiro, RJ 21041-210, Brazil; ¹¹ Department of Animal Biology, Faculty of Biology, University of Barcelona, 08071 Barcelona, Spain

ABSTRACT

Emerging and recurring infectious diseases known or suspected to have the potential to significantly impact cetacean populations, and possible synergistic effects of environmental factors are reviewed. Cetacean morbilliviruses and papillomaviruses and brucellosis may affect population densities through high mortality rates or interference with reproduction. Evidence is available for the role of environmental factors in the emergence/recurrence and severity of at least six infectious conditions i.e. lobomycosis, toxoplasmosis, tattoo skin disease, generalized bacterial infections, miscellaneous skin diseases and morbillivirus epizootics. Other micro-parasites of potential importance include rhabdo-, herpes- and parainfluenza-viruses as well as *Helicobacter* spp., *Streptococcus* spp., *Salmonella* spp. and *Mycobacterium marinum*. The population impact and aetiology of newly emerging skin diseases in South America are unknown and represent a cause of concern.

INTRODUCTION

Microparasites, including viruses, bacteria, and protozoans, may constrain the growth of wild animal populations (Anderson and May 1979; Anderson, 1982; Fenner, 1983; Gulland, 1995). The intensity of this effect is a function of the heightened mortality rate and/or reduction in reproductive capacity. Thus, microparasites may increase the risk of extinction of small populations in combination with other factors (May, 1986; Thorne and Williams, 1988; Gulland, 1995, Raga *et al.*, 1997). Here we review some emerging and recurring infectious diseases known or suspected to have the potential to significantly impact cetacean populations, and possible synergistic effects of environmental factors.

CETACEAN MORBILLIVIRUS

The dolphin, porpoise and pilot whale morbilliviruses (DMV, PMV and PWMV) are strains of a recently recognized member of the genus *Morbivirus* (single, negative-strand RNA viruses of family *Paramyxoviridae*) called cetacean morbillivirus (CeMV) (Barrett *et al.*, 1993; Blixenkrone-Möller *et al.*, 1996; Kennedy, 1998; Taubenberger *et al.*, 2000; Di Guardo *et al.*, 2005). They trigger highly lethal epizootic diseases in cetaceans causing bronchiolo-interstitial pneumonia and interstitial pneumonia, non-suppurative encephalitis and lymphoid depletion (Kennedy *et al.*,

1991, Domingo *et al.*, 1992, Duignan *et al.*, 1992; Schulman *et al.*, 1997). CeMV caused epizootics in bottlenose dolphins (*Tursiops truncatus*) along the Atlantic USA coast in 1982, 1987-1988 and 1993-1994 (Lipscomb *et al.*, 1994; Krafft *et al.*, 1995, Duignan *et al.*, 1996). Over 50% of the inshore bottlenose dolphins off New Jersey and south to Canaveral National Seashore, Florida, USA, may have died during the 1987-1988 epizootic (Lipscomb *et al.*, 1994). PMV caused mortalities of an unknown number of harbour porpoises (*Phocoena phocoena*) along the coasts of the United Kingdom and the Netherlands in 1988-1990 (Kennedy *et al.*, 1988; Visser *et al.*, 1993). DMV was responsible for lethal epizootics in at least two small cetacean species from the Mediterranean Sea in 1990-1992 and again in 2006-2007 (Domingo *et al.*, 1990; Van Bresseem *et al.*, 1991, 1993a; Di Guardo *et al.*, 1992, 2005; Fernandez *et al.*, 2008; Raga *et al.*, 2008). The epizootic started along the central coast of Spain in 1990 and ended in Turkey and the Greek Islands in 1992 (Aguilar and Raga, 1993). It affected predominantly striped dolphins (*Stenella coeruleoalba*). Sexually mature individuals suffered the highest mortality though dependent calves also represented a significant portion of the toll probably because of the death of their mothers (Calzada *et al.*, 1994). DMV apparently did not persist as an endemic infection in Mediterranean striped dolphins after the epidemic was over (Van Bresseem *et al.*, 2001a). At the end of October 2006, an unusual high number of long-finned pilot whales (*Globicephala melas*) stranded along the coast of the Strait of Gibraltar (Fernandez *et al.*, 2008). The epizootic spread eastward and northeast till April 2007. The freshly dead *G. melas* examined by immuno-histology were positive for morbillivirus infection and a virus sequence very closely related to the DMV isolated in 1990-1992 was amplified by reverse-transcriptase polymerase chain reaction (RT-PCR) (Fernandez *et al.*, 2008). In early July 2007 dead or moribund striped dolphins and *G. melas* were detected in the Gulf of Valencia (Raga *et al.*, 2008). A DMV strain closely related to those isolated during the 1990-1992 epidemic was detected in seven of 10 *S. coeruleoalba* as well as in a juvenile *G. melas* by immunohistochemistry and RT-PCR (0.4% difference in the complete H gene) (Raga *et al.*, 2008; Tiwari, Banyard, Raga, and Barrett, unpublished work). In summer-autumn 2007, over two hundred striped dolphins were found dead along the coasts of Spain. Juvenile striped dolphins were significantly more frequently affected than adults in the current epizootic, likely because older dolphins were still protected by the immunity developed during the 1990-1992 epidemic (Raga *et al.*, 2008). The virus apparently reached the coasts of France in August (F. Dupraz, MARMAM message, day November 2007) and those of Italy (Ligurian Sea) in August-November 2007 (Garibaldi *et al.*, 2008). An isolated case was reported in Sardinia in 2007 (S. Appino, MARMAM message, day December 2007).

Taking into account that both 1990-1992 and 2006-2007 DMV epidemics started close to, or in the Gibraltar Strait (Raga *et al.*, 2008) and that DMV was circulating in the North Sea earlier that year (Wohlstein *et al.*, 2007), we believe that DMV-infected cetaceans, possibly *G. melas*, entered the Strait of Gibraltar and propagated the infection to striped dolphins, currently the most abundant odontocete in the Mediterranean (Notarbartolo di Sciarra *et al.*, 1993; Aguilar, 2000). *G. melas* has been proposed as the reservoir and vector of infection to other cetacean species of the North Atlantic (Duignan *et al.*, 1995; Van Bresseem *et al.*, 1998). Long-finned pilot whales are commonly seen in the western Mediterranean including the Strait of Gibraltar and may form mixed groups with striped dolphins (Duguay, 1989; Raga *et al.*, 1991, pers. observations). Interestingly, the outbreak of PMV mortalities in the NE Atlantic and North Sea was also contemporaneous with the 1990-1992 DMV epidemic. Whether these facts indicate that some environmental factor(s) triggered or influenced these outbreaks remains to be clarified. The dolphins that succumbed to the 1990-1992 epizootic were in a poor nutritional state with their lipid reserves estimated to be only about 60% of the usual values for this population, an energy depletion that could not be explained by the effect of the disease alone (Aguilar *et al.*, 1992; Aguilar and Raga, 1993). Also, the prevalence of ectoparasites and epizoids was much higher than reference values for this population, suggesting that the dolphins had gone through a period of restricted mobility (Aguilar and Raga, 1993; Aznar *et al.*, 1994). These unusual factors were associated with abnormally high temperatures and low rainfall conditions that prevailed in the western Mediterranean in the winter preceding the outbreak, and which apparently depressed the peak of productivity that occurs each year in early spring and determines the regional abundance of fish (Aguilar and Raga, 1993). Polychlorinated biphenyls (PCBs) loads of striped dolphins that died during the 1990-92 epidemic were higher than in the individuals that survived it (Kannan *et al.*, 1993; Aguilar and Borrell, 1994), and given their well-known immunosuppressive effect in mammals (Busbee *et al.*, 1999), it was suggested that the high PCB loads may have compromised immune response and increased the severity of the outbreak (Aguilar and Borrell, 1994). Though the role of pollutants in the 2007 epidemic remains to be investigated, recent pollutant data obtained through analyses of biopsies from apparently healthy striped dolphins in 1987-2002 suggested that PCB and DDT concentrations have gradually decreased in this species (Aguilar and Borrell, 2005). Recurrent epizootics with high mortalities among all age classes may have a profound impact on the population dynamics of Mediterranean *S. coeruleoalba*.

Morbilliviruses may also have been involved in the mass strandings of Fraser's dolphins (*Lagenodelphis hosei*) observed along the Atlantic coast of South America in 1997-1999 (Moreno *et al.*, 2003). Indeed, three of four *L. hosei* that died during these events had serum antibodies against DMV (Van Bresseem *et al.*, 2001a). The role of these viruses in cetacean mass-stranding worldwide should be considered.

BRUCELLA SPP.

Brucellosis is a globally distributed zoonotic disease of mammals that i.a. is pathogenic for the reproductive system and may cause abortion. The aetiologic agent is a Gram-negative, facultative intracellular bacteria of the genus *Brucella*. In the 1990s, previously unknown strains of *Brucella* were detected in free-ranging pinnipeds and cetaceans from the Americas, Europe, the Antarctic and western North Pacific as well as in captive bottlenose dolphins (Ewalt *et al.*, 1994; Ross *et al.*, 1994; Jahans *et al.*, 1997; Clavareau *et al.*, 1998; Miller *et al.*, 1999; Bricker *et al.*, 2000; Van Bresseem *et al.*, 2001b; Foster *et al.*, 2002; Prenger-Berninghoff *et al.*, 2003; Ohishi *et al.*, 2003, 2004; Davison *et al.*, 2008). Phylogenetic analyses of the marine *Brucella* isolates show that these form a new group that originate very near the *B. ovis* branch and that the seal, porpoise and dolphin brucellae have diverged a very long time ago, concurrent with the evolution of their hosts (Bourg *et al.*, 2007). On the basis of biological and molecular characteristics, Foster *et al.* (2007) proposed two *Brucella* species in marine mammals: *Brucella ceti* with cetaceans as preferred hosts and *Brucella pinnipedialis* with seals as the preferred hosts.

Disorders associated with brucellosis in cetaceans include placentitis, orchitis, abortion, lung infection and non-suppurative meningo-encephalitis (Miller *et al.*, 1999; Gonzalez *et al.*, 2002; Prenger-Berninghoff *et al.*, 2003; Ohishi *et al.*, 2004; Davison *et al.*, 2008). Brucellosis is endemic in several species and populations worldwide and, in Peruvian small cetaceans affects significantly more sexually mature than immature individuals (Van Bresseem *et al.*, 2001b, 2007a). When enzootic *B. ceti* infection may constitute a measurable limiting factor among environmental variables reducing population growth through negatively influencing reproduction (Van Bresseem *et al.*, 2001b, 2007a).

TOXOPLASMOSIS

The protozoan *Toxoplasma gondii* is widespread in marine mammals including odontocetes (reviewed in Dubey *et al.*, 2003). In cetaceans, toxoplasmosis was first reported in a Guiana dolphin (*Sotalia guianensis*) from Brazil (Bandoli and Oliveira, 1977). Infected free-ranging dolphins have been found in Europe, the Americas and the Caribbean (Bandoli and Oliveira, 1977; Domingo *et al.*, 1990; Cruickshank *et al.*, 1990; Inskoop *et al.*, 1990; Di Guardo *et al.*, 1995a,b; Resendes *et al.*, 2002; Cabezón *et al.*, 2004; Dubey *et al.*, 2007). They presented lympho-adenitis, lesions in the adrenal glands, myocarditis, acute interstitial pneumonia and non-suppurative encephalitis (Inskoop *et al.*, 1990; Dubey *et al.*, 2003; 2007). Disseminated toxoplasmosis was reported in a female *T. truncatus* and her calf from Florida, in a female Risso's dolphin (*Grampus griseus*) from the Mediterranean and in two beluga whales (*Delphinapterus leucas*) from the St Lawrence estuary, Quebec (Inskoop *et al.*, 1990; De Guise *et al.*, 1995; Resendes *et al.*, 2000). Two pregnant infected females (a captive *Tursiops aduncus* and the Mediterranean *Grampus griseus*) aborted fetuses with generalized toxoplasmosis (Jardine and Dubey, 2002; Resendes *et al.*, 2002). Clinical toxoplasmosis was generally associated with immunosuppression following a morbillivirus infection (Domingo *et al.*, 1990; Di Guardo *et al.*, 1995b; Schulman *et al.*, 1997) and/or high concentrations of environmental contaminants including PCBs (Mikaelian *et al.*, 2000). Serological data demonstrated that *T. gondii* circulates in Mediterranean *T. truncatus*, *D. delphis* and *S. coerulealba* (Cabezón *et al.*, 2004), coastal bottlenose dolphins from the USA (Dubey *et al.*, 2003, 2005) and belugas from Quebec (Mikaelian *et al.*, 2000).

T. gondii is the sole recognized species in the genus and, until recently, was composed of three major genotypes designed as types I-III that have emerged as dominant strains worldwide (Conrad *et al.*, 2005). Two new clades of *T. gondii*, named type A and X, were recently detected in the southern sea otter (*Enhydra lutris nereis*) from California and *Enhydra lutris kenyoni* from Washington State (Conrad *et al.*, 2005; Sundar *et al.*, 2008). Members of the *Felidae* family are the only known definitive hosts for *T. gondii* genotypes I-III but many mammals can be intermediate hosts if infected through the ingestion of contaminated food or water, or transplacentally. The origin and transmission of types A and X are not yet known. Some *T. gondii* infections of marine mammals, especially in coastal species, have been linked to freshwater runoffs contaminated with cat excrements containing oocysts (Miller *et al.*, 2002). Oocysts are very resistant and likely to survive in the sea and may be concentrated by molluscs (Dubey *et al.*, 2007). The infection of offshore species like *G. griseus* and *S. coeruleoalba* may be linked to ship run-off waters when hygienic conditions are low and when rodents, cats or contaminated soil are present onboard. The global maritime trade is thought to be responsible for the dissemination of *T. gondii* from its original niche in South America to the other continents (Lehman *et al.*, 2006).

Though the potential of *T. gondii* to impact cetacean populations has not yet been investigated, its ability to cause abortions and lethal systemic diseases renders it highly suspect. The possible reactivation of latent *T. gondii* infection during morbillivirus outbreaks may synergistically increase the severity of the viral disease and death rate. *T. gondii* is thought to contribute to the slow rate recovery of the southern sea otter population (Conrad *et al.*, 2005).

LOBOMYCOSIS AND LOBOMYCOSIS-LIKE DISEASE

Caused by a yeast-like organism known as *Lacazia loboi* (Taborda *et al.*, 1999) (syn. *Loboa loboi*; Caldwell *et al.*, 1975), lobomycosis (or lacaziosis) naturally affects humans, *T. truncatus* and *S. guianensis* (de Vries and Laarman, 1973; Caldwell *et al.*, 1975; Simões-Lopes *et al.*, 1993; Reif *et al.*, 2006; Van Bresseem *et al.*, 2007b). *Lacazia loboi* cells found in *T. truncatus* infected tissues are significantly smaller than those found in humans, suggesting that the organism may not be identical in the two hosts (Haubold *et al.*, 2000). In humans, lobomycosis is a self-limited, chronic fungal infection of the skin, endemic in rural regions in South and Central America. Water, earth and vegetation are considered ecological habitats of the fungus and the agent accesses the skin by penetration or accidental trauma (Paniz-Mondolfi *et al.*, 2007). Patients with lobomycosis may have immunoregulatory disturbances possibly responsible for the lack of pathogen containment (Vilani-Moreno *et al.*, 2005; Honda *et al.*, 2007). Lobomycosis in dolphins is characterized by grayish, whitish to slightly pink, verrucous lesions, often in pronounced relief that may ulcerate (Migaki *et al.*, 1971). In *T. truncatus* from Florida the disease was associated with an impaired immune function possibly caused by anthropogenic factors (Bossart, 1984; Reif *et al.*, 2006). Variation in salinity and water temperature may also play a role (Reif *et al.*, 2006). The disease has been reported in dolphins from the Atlantic coast of the USA, Europe and South America (De Vries and Laarman, 1973; Caldwell *et al.*, 1975; Symmers, 1983; Simões-Lopes *et al.*, 1993; Reif *et al.*, 2006; Van Bresseem *et al.*, 2007b). In South America, several cases highly reminiscent of lobomycosis were observed in free-ranging coastal *T. truncatus* and *S. guianensis*. In the absence of a histological diagnosis, the disease was called lobomycosis-like disease (LLD; Van Bresseem *et al.*, 2007b). LLD seem to have been emerging recently over a relatively short span of time in Colombia, Ecuador, Peru, Brazil and Venezuela (Van Bresseem *et al.*, 2007b, 2008a; Bermúdez *et al.*, 2008). Prevalence in inshore *T. truncatus* varied from 1.6% (Gulf of Guayaquil, Ecuador) to 20% (Tramandai estuary, Brazil) (Van Bresseem *et al.*, 2007b; Moreno *et al.*, 2008). Most of the populations affected inhabited polluted waters harbouring ports or cities and, in the case of Guayaquil, intense shrimp farming activities (CPPS, 2000; CVC, 2003; UNEP, 2006; WHO/UNICEF/WSSCC, 2001). The origin of the fungus in coastal waters is unknown. The role of increased shipping in introducing *Lacazia loboi* to new ecological niches should be explored. Indeed, the discharge of water, sediments and biofilms from ships' ballast water tanks is a prominent vector of aquatic invasive species (Ruiz *et al.*, 2000; Drake *et al.*, 2007) and several cases of LLD have recently been diagnosed in small cetaceans in the vicinity of large ports in Colombia, Ecuador and Brazil (Van Bresseem *et al.*, 2007b; 2008a). The appearance of new cases of LLD in the lagoon of Mayotte, southwest Indian Ocean and possibly off Amakusa-Shimoshima Island, East China Sea and gives weight to this hypothesis (Kiska and Van Bresseem, 2008; Shirakihara *et al.*, 2008).

Lobomycosis and LLD may eventually contribute to the death of severely affected dolphins (Simões-Lopes *et al.*, 1993; Van Bresseem *et al.*, 2007b; Bermúdez-Villapol *et al.*, 2008). All dolphins that died with extensive lobomycosis or LLD were adult, and all LLD affected individuals sighted have been large, presumably adult, individuals, except for a calf *S. guianensis* from the Lagamar estuary, Brazil (Van Bresseem *et al.*, 2008a). As the rates of increase of dolphin population size appears to be more sensitive to the non-calf survival rate than to the calf survival rate (Reilly and Barlow, 1986), lobomycosis and LLD are likely to lower the growth of coastal populations, and may have a significant impact on small, isolated ones.

GENITAL PAPILOMAVIRUSES

Papillomaviruses (PVs), small, non-enveloped, double-stranded DNA viruses (family *Papillomaviridae*) are epitheliotropic pathogens which may induce proliferation of the stratified squamous epithelia of the skin and mucosa and cause lesions known as warts, papillomas and condylomas in mammals and birds (Antonsson and Hansson, 2002; Terai *et al.*, 2002, Bernard, 2005).

Genital papillomatosis was observed in sperm whales (*Physeter macrocephalus*) caught off Iceland (Lambertsen *et al.*, 1987), in dusky dolphins (*Lagenorhynchus obscurus*), long-beaked common dolphins (*Delphinus capensis*), common bottlenose dolphins and Burmeister's porpoises (*Phocoena spinipinnis*) from Peru (Van Bresseem *et al.*, 1996), in *T. truncatus* from Florida (Bossart *et al.*, 2005) and Cuba (Cruz *et al.* 2006) and in a Guiana dolphin from Brazil (M. Marcondes, pers. commn. May 2007) as well as in *T. truncatus* kept in captivity in Europe and the USA (Bossart *et al.*, 2005; Rector *et al.*, 2006). Genital lesions macroscopically and microscopically consistent with PV-induced papillomas have also been seen in *P. phocoena*, *D. delphis* and *S. coeruleoalba* stranded along the coasts of the United Kingdom (Jepson, P.D., pers. obs.). Prevalence of the disease ranged from high to medium in cetaceans from Peru (33-66.7%; Van Bresseem *et al.*, 1996), Cuba (28.7%; Cruz *et al.*, 2006) and Iceland (10%; Lambertsen *et al.*, 1987). Sexual variation in wart prevalence was found in *L. obscurus* and *P. spinipinnis* with males being two and three times more infected than females, respectively (Van Bresseem *et al.*, 1996). The same possibly occurs in *T. truncatus* from Cuba (Cruz *et al.*, 2006).

Phocoena spinipinnis papillomavirus type 1 and another still uncharacterized PV cause genital warts in *P. spinipinnis* (Van Bresseem *et al.*, 2007c). At least three PVs are associated with genital papillomatosis in *T. truncatus*

from Europe and Atlantic USA (Chiers *et al.*, 2004; Rector *et al.*, 2006; Rehtanz *et al.*, 2006;). Other still uncharacterised PVs were detected in papillomas from Peruvian *T. truncatus* and *L. obscurus* (Cassonnet *et al.*, 1998). The high prevalence (48.5%) of genital warts in 33 Burmeister's porpoises examined in 1993-1995 (Van Bresseem *et al.*, 1996) and the detection of PV sequences in 5 of 7 genital warts sampled in this study indicate that PV infection is frequent in this species. Porpoises may become infected early in life through vertical and horizontal transmission (Van Bresseem *et al.*, 1996, 2007c). Genital lesions of sufficient severity that may impede, or at least, hamper copulation were seen in two of 20 male *P. spinipinnis* but not in other species from Peru. One sperm whale also suffered extensive genital papillomatosis (Lambertsen *et al.*, 1987). PVs in some circumstances (especially if non-randomly distributed) may exert an indirect impact on the dynamics of this population (Lambertsen *et al.*, 1987; Van Bresseem *et al.*, 1999).

TATTOO SKIN DISEASE

Tattoo skin disease (TSD) is characterised by very typical, irregular, grey, black or yellowish, stippled lesions that may occur on any part of the body but show a preferential corporal distribution depending on the species (Van Bresseem and Van Waerebeek, 1996). It has been observed in several species of free-ranging Delphinidae, Phocoenidae and Ziphiidae worldwide as well as in captive bottlenose dolphins (*Tursiops truncatus*) and a bowhead whale (*Balaena mysticetus*) from Alaska (see Van Bresseem *et al.*, 1999, 2008b; Bracht *et al.*, 2006). TSD is caused by poxviruses (Flom and Houk, 1979, Geraci *et al.*, 1979, Van Bresseem *et al.*, 1993, 2006) that may belong to a new genus of *Chordopoxvirinae*, but have a common, most immediate ancestor with terrestrial poxviruses of the genus *Orthopoxvirus* (Bracht *et al.*, 2006; Pearce *et al.*, 2008). These viruses are thought to induce humoral immunity that protects neonates and young calves from the disease (Smith *et al.*, 1983, Van Bresseem and Van Waerebeek, 1996; Van Bresseem *et al.*, 2006). Individual tattoo lesions may persist for months or years and recurrence may occur (Van Bresseem *et al.*, 2003).

A still unpublished study (Van Bresseem *et al.*, 2008b) suggests that the epidemiological pattern of TSD may be an indicator of a degrading and stressful aquatic environment. In odontocetes that had died a traumatic death (TD) as well as in free-ranging *T. truncatus* from the Sado estuary, Portugal, prevalence varied with the age class, being generally null in neonates and young calves, peaking among juveniles and decreasing in adults. In contrast, prevalence increased or remained high in adult dolphins and porpoises that had died in poor health (PH) as well as in coastal *P. spinipinnis* from Peru. Prevalence of TSD was significantly higher in adult dolphins and porpoises from the PH category than in adults from the TD category. In coastal dolphins and porpoises, prevalence did not decrease significantly in adults while it did in adult neritic and oceanic cetaceans. These findings suggest that adult odontocetes that died in PH and Peruvian *P. spinipinnis* had a depressed immune system with reduced capacity to fight infections.

Clinical and epizootiological data do not indicate that poxvirus infection induces a high mortality rate when enzootic (Van Bresseem and Van Waerebeek, 1996; Van Bresseem *et al.*, 2003). However, it may kill neonates and calves without protective immunity and thus interfere with host population dynamics. An apparent, severe, generalized poxvirus infection in a calf *S. guianensis* stranded alive along the coast of Pará state, Brazil in March 2008 gives weight to this hypothesis. Super-infection by fungi or bacteria may exacerbate the disease (Smith *et al.*, 1983; Van Bresseem *et al.*, 2003, 2007b).

INFECTIOUS DISEASES IN *PHOCOENA PHOCOENA* FROM THE NORTH AND BALTIC SEAS

Recent studies have demonstrated a significant association between chronic PCB exposure and infectious diseases (including parasitic, bacterial and mycotic pneumonia and generalized bacterial infections) in harbour porpoises from the North and Baltic Seas (Jepson *et al.*, 1999, Siebert *et al.*, 1999; Das *et al.*, 2004; Jepson *et al.*, 2005; Hall *et al.*, 2006; Valderrama *et al.*, 2008). The infectious disease group of harbour porpoises from the British Isles had a significantly higher sum of the concentrations of 25 individual chlorobiphenyl congeners ($\Sigma 25\text{CBs}$) than the TD group. The mean $\Sigma 25\text{CBs}$ level in TD animals was 13.6mg/kg lipid, whereas the mean level in the PH group was 27.6mg/kg lipid. Adult females in both groups had significantly lower $\Sigma 25\text{CBs}$ levels than adult males because of off-loading of organochlorines during gestation and lactation (Jepson *et al.*, 2005). The full impact of PCBs on harbour porpoise populations in European waters is unknown.

EMERGING SKIN DISEASES IN SOUTH AMERICA

Miscellaneous skin diseases of unknown aetiologies have been reported from several regions in South America (Flach, 2006; Brownell *et al.*, 2007; Van Bresseem *et al.*, 2007b, 2008a; Flach *et al.*, 2008; S. Siciliano, unpublished data). Many are associated with unrelated wounds, scars and tooth rakes. Though these lesions may eventually heal, their effect on the overall health, fitness and reproductive success of individuals is unknown. A possible relation with poor water quality is suggested as lesions were observed in individuals living in waters heavily contaminated by organic material

originating from fishmeal factories, municipal sewage and aquaculture as well as ballast water from cargo ships (Van Bresseem *et al.*, 2007b).

Several bacteria including the common opportunistic pathogens *Vibrio* spp., *Aeromonas hydrophila* and *Pseudomonas aeruginosa* that thrive in the aquatic environment have been isolated from skin lesions in cetaceans worldwide (reviewed in Van Bresseem *et al.*, 2008c). In the lower Chesapeake Bay, USA, coastal ecosystems are frequently invaded by micro-organisms from ballast waters with an estimated 10^{20} bacteria (including *V. cholerae*) and viruses annually discharged (Drake *et al.*, 2007). Salmon and shrimp farms heavily use prophylactic antibiotics resulting in the increase of antibiotic resistance in fish and human pathogens including *Vibrio cholerae* and *Aeromonas hydrophila* (Weber *et al.*, 1994; Rhodes *et al.*, 2000; Cabello, 2004, 2006). The combination of aquaculture, intense ship traffic and the release of untreated run-off waters likely favour a high density of pathogenic bacteria, some of which are antibiotic-resistant, that may infect wounds. In this context, we believe that combined microbial and organic contamination is possibly at the origin of the emergence of frequent miscellaneous skin lesions in coastal cetaceans from South America. The worldwide dissemination of fungi and antibiotics-resistant marine bacteria through water ballast and sea product transportation may favour the emergence and spread of skin diseases to other areas of the world.

CONCLUSIONS

As for any other mammals, cetaceans are normally infected by a wide variety of pathogens that are order or family-specific (i.e. morbilli-, papilloma- and poxviruses as well as *Brucella ceti*) or opportunistic (bacteria, fungi, protozoans). Infection by specific pathogens are likely to have occurred for thousands of years with some equilibrium between populations and pathogens as in other species (Begon *et al.*, 1996; Blaser and Kirschner, 2007). Environmental degradation including biological and chemical pollution, climate change, fisheries and heavy boat traffic, may have changed this pattern by lowering the immune response of the populations, shortening the food supply, increasing stress and facilitating the introduction of alien pathogens, among others. The number and severity of diseases have increased, in part because of more dedicated research in this field (Gulland and Hall, 2007). However, taking this bias into account, there has been an increase in the frequency and severity of some diseases as for example those resulting from exposure to harmful algal blooms (HABs) as well as lobomycosis and whitish velvety cutaneous lesions in cetaceans from the Americas to cite a few (Reif *et al.*, 2006; Gulland and Hall, 2007; Van Bresseem *et al.*, 2007b; Flach *et al.*, 2008; Siciliano *et al.*, 2008).

There is conclusive evidence that chemical pollution has increased the emergence and severity of several diseases (Martineau *et al.*, 1988; Aguilar and Borrell, 1994; De Guise *et al.*, 1998; Jepson *et al.*, 1999, 2005; Ross, 2002; Hall *et al.*, 2006; Van Bresseem *et al.*, 2008b). Biological pollution is an emerging issue (Mos *et al.*, 2006; Pereira *et al.*, 2007; Van Bresseem *et al.*, 2007b). Coastal cetaceans seem particularly more at risk because, besides living in an environment that is prone to be more polluted, they may face higher concentrations of pathogenic micro-organisms in the waters they inhabit and may also be more often wounded and stressed by interactions with boats and fisheries. With a depressed immune system and a ruptured skin, these animals would make ideal targets for opportunistic pathogens. The continued degradation of the marine environment and increased pressure on its inhabitants will possibly provoke more frequent epizootics, help disseminate fungal and bacterial diseases and increase the prevalence and severity of infectious illnesses in cetaceans worldwide.

ACKNOWLEDGEMENTS

The work of M-F Van Bresseem was funded by WDSCS and CSI. CEPEC research has benefitted from sustained IFAW support. Van Waerebeek attends the IWC/60 meeting as head of scientific delegation for Federal Public Service, Public health, Food Chain security and Environment, Brussels, Belgium. S. Siciliano is supported by fellowships of the FIOCRUZ/FAPERJ and PROJETO PIATAM OCEANO – CENPES/PETROBRAS, UFF, UFPA and MPEG.

REFERENCES

- Aguilar, A. 2000. Population biology, conservation threats, and status of Mediterranean striped dolphins (*Stenella coeruleoalba*). *J Cet Res Manag* 2: 17-26.
- Aguilar, A. and Raga, J.A. 1993. The striped dolphin epizootic in the Mediterranean Sea. *Ambio* 22(8): 524-528.
- Aguilar, A. and Borrell, A. 1994. Abnormally high polychlorinated biphenyl levels in striped dolphins (*Stenella coeruleoalba*) affected by the 1990-1992 Mediterranean epizootic. *Sc Tot Environm* 154: 237-247.
- Aguilar, A. and Borrell, A. 2005. DDT and PCB reduction in the western Mediterranean in 1987-2002, as shown by levels in dolphins. *Mar Environm Res* 59: 391-404.

- Aguilar, A., Borrell, A., Calzada, N. and Grau, E. 1992. Body fat reserves in striped dolphins examined during the western Mediterranean die-off. *Proceedings of the Mediterranean Striped Dolphin Mortality International Workshop*, 47-52.
- Anderson, R.M. 1982. Population dynamics of infectious disease agents: theory and applications. Chapman and Hall, London
- Anderson, R.M. and May, R.M. 1979. Population biology of infectious diseases: Part I. *Nature* 280:361-367
- Antonsson, A. and Hansson, B. G. 2002. Healthy skin of many animal species harbors papillomaviruses which are closely related to their human counterparts. *J Virol* 76: 12537-12542.
- Aznar, J., Balbuena, J.A. and Raga, J.A. 1994. Are epizootics biological indicators of a Western Mediterranean striped dolphin die-off? *Dis Aquat Org* 18:59-163
- Bandoli, J.G. and de Oliveira, C.A.B. 1977. Toxoplasmose em *Sotalia guianensis* (Van Beneden, 1863), Cetacea, Delphinidae. *Fol Méd* 75: 459-468
- Barrett, T., Visser, I.K.G., Mamaev, L.V., Goatley, L., Van Bresse, M.-F., Osterhaus, A.D.M.E. 1993. Dolphin and porpoise morbilliviruses are genetically distinct from phocine distemper virus. *Virol* 193, 1010-1012.
- Begon, M., Harper, J.L. and Townsend, C.R. 1996. Ecology, 3rd edn. Blackwell Science Oxford
- Beineke, A., Siebert, U., Mac Lachlan, M., Bruhn R., Thron K., Failing K., Müller G. and Baumgärtner W. 2005. Investigations of the potential influence of environmental contaminants on the thymus and spleen of harbor porpoises (*Phocoena phocoena*). *Environm Sc Technol* 39, 3933-3938.
- Bermúdez-Villapol, L.A., Sayegh, A.J. and Van Bresse, M-F. 2008. Lobomycosis-like disease in a bottlenose dolphin (*Tursiops truncatus*) from Venezuela. Paper SC/60/DW7 presented to the IWC Scientific Committee, May 2008 (unpublished) [Available from the Office of this Journal]
- Bernard, H-U. 2005. The clinical importance of the nomenclature, evolution and taxonomy of human papillomaviruses. *J Clin Virol* 32S: S1-S6.
- Blaser, M.J. and Kirschner, D. 2007. The equilibria that allows bacterial persistence in human hosts. *Nature* 449: 843-849.
- Blixenkron-Möller, M., Bolt, G., Jensen, T.D., Harder, T. and Svansson, V. 1996. Comparative analysis of the attachment protein gene (H) of dolphin morbillivirus. *Virus Res* 40:47-56.
- Bossart, G. 1984. Suspected acquired immunodeficiency in an Atlantic bottlenosed dolphin with chronic-active hepatitis and lobomycosis. *JAVMA* 185 1413-1414
- Bossart, G.D., Ghim, S.J., Rehtanz, M., Goldstein, J., Varela, R., Ewing, R.Y., Fair, P.A., Lenzi, R., Joseph, B., Hicks, C.L., Schneider, L.S., McKinnie, C., Reif, J.S., Sanchez, R., Lopez, A., Novoa, S., Bernal, J., Goretti, M., Rodriguez, M., Defran, R.H. and Bennett Jenson, A. 2005 Orogenital neoplasia in Atlantic bottlenose dolphins (*Tursiops truncatus*). *Aquat Mamm* 31: 473-480.
- Bourg, G., O'Callaghan, D. and Boschioli, M.L. 2007. The genomic structure of *Brucella* strains isolated from marine mammals gives clues to evolutionary history within the genus. *Vet Microb* 125: 375-380
- Bracht, A.J., Brudek, R.L., Ewing, R.Y., Manire, C.A., Burek, K.A., Rosa, C., Beckmen, K.B., Maruniak, J.E. and Romero, C.H. 2006. Genetic identification of novel poxviruses of cetaceans and pinnipeds. *Arch Virol* 151: 423-438
- Bricker, B.J., Ewalt, D.R., Macmillan, A.P., Foster, G. and Brew, S. 2000. Molecular characterization of *Brucella* strains isolated from marine mammals. *J Clin Microb* 38: 1258-1262.
- Bross, M.H., Soch, K., Morales, R. and Mitchell, R.B. 2007. *Vibrio vulnificus* infection: diagnosis and treatment. *Am Fam Phys* 15: 540-544.
- Brownell, R. L. Jr., Carlson, C.A., Galletti Vernazzani, B. and Cabrera, E. 2007. Skin lesions on blue whales off Southern Chile: possible conservation implications? Paper SC/59/SH21 presented to the IWC Scientific Committee, May 2007 (unpublished).
- Busbee, D., Tizard, I., Scott, J., Ferrick, D. and Ott-Reeves, E. 1999. Environmental pollutants and marine mammal health: the potential impact of hydrocarbons and halogenated hydrocarbons on immune system dysfunction. *J Cet Res Manag (Special Issue 1)*: 223-248.
- Cabello, F.C. 2004 Antibióticos y acuicultura en Chile: consecuencias para la salud humana y animal. *Rev Med Chile* 132: 1001-1006.
- Cabello, F.C. 2006. Heavy use of prophylactic antibiotics in aquaculture: a growing problem for human and animal health and for the environment. *Environm Microb* 8: 1137-1144.
- Cabezón, O., Resendes, A.R., Raga, J.A., Agusti, C., Alegre, F., Mons, J.L., Dubey, J.P. and S. Almeria. 2004. Seroprevalence of *Toxoplasma gondii* antibodies in wild dolphins from the Spanish Mediterranean coast. *J Parasitol* 90: 643-644.
- Caldwell, D.K., Caldwell, M.C., Woodard, J.C., Ajello, L., Kaplan, W. and McLure, H.M. 1975. Lobomycosis as a disease of the Atlantic bottle-nosed dolphin (*Tursiops truncatus* Montagu, 1821). *Am J Trop Med* 24: 105-114

- Calzada, N., Lockyer, C. H. and Aguilar, A. 1994. Age and sex composition of the striped dolphin die-off in the western Mediterranean. *Mar Mamm Sc* 10: 299-310.
- Cassonnet, P., Van Bresseem, M-F., Desaintes, C., Van Waerebeek, K. and Orth G. 1998. Papillomaviruses cause genital warts in small cetaceans from Peru. The World Marine Mammal Science Conference, Monaco, January 1998.
- Chiers, K. Lacave, G., Salbany, A., Roque, L. and Bossart, G.D. 2004. Detecteion of a novel papillomavirus in genital papillomatous lesions of bottlenose dolphins. 22nd meeting of the European Society of Veterinary pathology and 6th meeting of the European Society of Veterinary Clinical Pathology/ European College of Veterinary Clinical Pathology and Continuing Education Day, Olsztyn- Poland 15-18 September 2004.
- Clavareau, C., Wellemans, V., Walravens, K., Tryland, M., Verger, J.M., Grayon, M., Cloeckaert, A., Letesson, J.J., Godfroid, J. 1998. Phenotypic and molecular characterization of a *Brucella* strain isolated from a minke whale (*Balaenoptera Acutorostrata*). *Microbiol* 144: 3267-3273.
- Conrad, P.A., Miller, M.A., Kreuder, C., James, E.R., Mazet, J., Dabritz, H., Jessup, D.A., Gulland, F.M. and Grigg, M.E. 2005. Transmission of *Toxoplasma*: Clues from the study of sea otters as sentinels of *Toxoplasma gondii* flow into the marine environment. *Int J Parasitol* 35: 1155-1168.
- Cruickshank, J.J, Haines, D.M., Palmer, N.C. and St Aubin, D. 1990. Cysts of *Toxoplasma*-like organism in an Atlantic bottlenose dolphin. *Can Vet J* 31: 213-215.
- Cruz, D., Guevara, C., Blanco, M., Sánchez, L. and Chamizo, E.G. 2006. Papilomatosis en genitales de delfines (*Tursiops truncatus*) de zonas costeras del archipelago cubano. VI congreso Internacional de Ciencias Veterinarias. La Habana, Cuba, 13 April 2006.
- Comision Permanente Del Pacifico Sur – CPPS. 2000. Estado del Medio Ambiente Marino y Costero del Pacífico Sudeste. Quito, Ecuador. 165 pp
- Corporacion Autonoma Regional Del Valle Del Cauca-CVC. 2003. Plan de gestión ambiental regional del Valle del Cauca 2002 –2012: Participación con Compromiso. Grupo de Planificación Ambiental y Grupo Archivo Corporativo. Segunda Edición. Cali, Colombia. 264 pp.
- Das, K., Siebert, U., Fontaine, M., Jauniaux, T., Holsbeek, L. and Bouquegneau J.M. 2004. Ecological and pathological factors related to trace metal concentrations in harbour porpoises *Phocoena phocoena* from the North Sea and adjacent areas. *Mar Ecol Progr Series* 281: 283-295.
- Davison, N.J., Cranwell, M.P., Perrett, L.L., Dawson, C.E., Deaville, R., Stubberfield, E.J., Jarvis, D.S. and Jepson, P.D. 2008. First case of meningoencephalitis associated with *Brucella* species in a live-stranded striped dolphin (*Stenella coeruleoalba*) in England. *Vet Rec* (in press)
- De Guise, S., Lagacé, A., Béland, P., Girard, C. and Higgins, R. 1995. Non-neoplastic lesions in beluga whales (*Delphinapterus leucas*) and other marine mammals from the St Lawrence Estuary. *J Comp Pathol* 112: 257-271.
- De Guise, S., Martineau, D., Beland, P. and Fournier, M. 1998. Effects of in vitro exposure of beluga whale leukocytes to selected organochlorines. *J Toxicol Environ Health A* 55:479-493.
- De Vries, G.A and Laarman, J.J. 1973. A case of Lobo's disease in the dolphin *Sotalia guianensis*. *Aquat Mamm* 1: 26-33.
- Di Guardo, G., Agrimi, U., Amaddeo, D., McAliskey, M. and Kennedy, S. 1992. Morbillivirus infection in a striped dolphin (*Stenella coeruleoalba*) from the coast of Italy. *Vet Rec* 130: 579-580.
- Di Guardo, G., Agrimi, U., Morelli, L., Cardeti, G., Terracciano, G. and Kennedy, S. 1995a. Post mortem investigations on cetaceans found stranded on the coasts of Italy between 1990 and 1993. *Vet Rec* 136:439-442
- Di Guardo, G., Corradi, A., Agrimi, U., Zizzo, N., Morelli, L., Perillo, A., Kramer, L., Cabassi, E. and Kennedy, S. 1995b. Neuropathological lesions in cetaceans found stranded from 1991 to 1993 on the coasts of Italy. *European J Vet Pathol* 1: 47-51.
- Di Guardo, G., Marruchela, G., Agrimi, M. And Kennedy, S. 2005. Morbillivirus infections in aquatic mammals: a brief overview. *J Vet Med A* 52:88-93.
- Domingo, M., Ferrer, L., Pumarola, M., Marco, A., Plana, J., Kennedy, S., Mcalisky, M., Rima, B.K. 1990. Morbillivirus in dolphins. *Nature* 348, 21-21.
- Domingo, M., Visa, J., Pumarola, M., Marco, A.J., Ferrer, L., Rabanal, R. and Kennedy S. 1992. Pathologic and immunocytochemical studies of morbillivirus infection in striped dolphins (*Stenella coeruleoalba*). *Vet Pathol* 29:1-10
- Domingo, M., Vilafranca, M., Visa, J., Prats, N., Trudgett, A. and Visser, I. 1995. Evidence for chronic morbillivirus infection in the Mediterranean striped dolphin (*Stenella coeruleoalba*). *Vet Microb* 44: 229-239.
- Drake, L., Doblin, M.A, Dobbs, F.C. 2007. Potential microbial bioinvasions via ships' ballast water, sediment, and biofilm. *Mar Poll Bull* 55: 333-341

- Dubey, J.P., Zarnke, R., Thomas, N.J., Wong, S.K., Van Bonn, W., Briggs, M., Davis, J.W., Ewing, R., Mense, M., Kwok, O.C.H., Romand, S. and Thulliez, P. 2003. *Toxoplasma gondii*, *Neospora caninum*, *Sarcocystis canis*-like infections in marine mammals. *Vet Parasitol* 116: 275-296.
- Dubey, J.P., Fair, P.A., Bossart, G.A., Hill, D., Fayer, R., Sreekumar, C., Kwok, O.C.H. and Thulliez, P. 2005. A comparison of several serologic tests to detect antibodies to *Toxoplasma gondii* in naturally exposed bottlenose dolphins (*Tursiops truncatus*). *J Parasitol* 91: 1074-1081.
- Dubey, J.P., Morales, J.A., Sundar, N., Velmurugan, G.V., González-Barrientos, Hernández-Mora and Su, C. 2007. Isolation and genetic characterization of *Toxoplasma gondii* from striped dolphin (*Stenella coeruleoalba*) from Costa Rica. *J Parasitol* 93: 710-711.
- Duguy, R. 1989. Les mammifères marins de la Méditerranée occidentale. *Bull Soc Zool France* 114: 89-96.
- Duignan, P.J., Geraci, J.R., Raga, J.A. and Calzada, N. 1992. Pathology of morbillivirus infection in striped dolphins (*Stenella coeruleoalba*) from Valencia and Murcia, Spain. *Can J Vet Res* 56:242-248
- Duignan, P.J., House, C., Geraci, J.R., Early, G., Copland, H., Walsh, M.T., Bossart, G.D., Cray, C., Sadove, S., St Aubin, D.J. and Moore, M. 1995. Morbillivirus infection in two species of pilot whales (*Globicephala* sp.) from the western Atlantic. *Mar Mamm Sci* 11, 150-162.
- Duignan, P. J., House, C., Odell, D. K., Wells, R. S., Hansen, L. J., Walsh, M. T., Aubin, D. J. S., Rima, B. K. and Geraci, J. R. 1996. Morbillivirus infection in bottlenose dolphins: evidence for recurrent epizootics in the western Atlantic and Gulf of Mexico. *Mar Mamm Sci* 12: 499-515.
- Ewalt, D.R., Payeur, J.B., Martin, B.M., Cummins, D.R. and Miller, W.G. 1994. Characteristics of a *Brucella* species from a bottlenose dolphin (*Tursiops Truncatus*). *J Vet Diagn Invest* 6: 448-452
- Gonzalez, L., Patterson, I.A., Reid, R.J., Foster, G., Barberan, M., Blasco, J.M., Kennedy, S., Howie, F.E., Godfroid, J., MacMillan, A.P., Shock, A. and Buxton, D. 2002. Chronic meningoencephalitis associated with *Brucella* sp. infection in live-stranded striped dolphins (*Stenella coeruleoalba*). *J Comp Pathol* 126:147-52.
- Fenner, F. 1983. The Florey Lecture. 1983: biological control, as exemplified by smallpox eradication and myxomatosis. *Proc R Soc Lond B* 218:259-285
- Fernández A, Esperón F, Herraéz P, Espinosa de los Monteros A, Clavel C, Bernabé A, et al. 2008. Morbillivirus and pilot whale deaths, Mediterranean Sea. *Emerg Infect Dis* 2008 May [Epub ahead of print]
- Flach, L. 2006. Photo-identification study reveals human threats towards estuarine dolphins in southeast Brazil. In: Siciliano S, Borobia M, Barros NB, Marques FC, Trujillo T, Flores PAC. (eds) Workshop on research and conservation of the genus *Sotalia*, Armação dos Búzios, Rio de Janeiro, Brazil.
- Flach, L., Van Bresse, M-F., Reyes, J.C., Echegaray, M., Siciliano, S., Santos, M., Vididi, F., Crespo, E., Klaich, J., Moreno, I., Tavares, M., Felix, F. and Van Waerebeek, K. 2008. Miscellaneous skin lesions of unknown aetiology in small cetaceans from South America. Paper SC/60/DW4 presented to the IWC Scientific Committee, May 2008 (unpublished) [Available from the Office of this Journal].
- Flom, J.O. and Houk, E.J. 1979. Morphologic evidence of poxvirus in "tattoo" lesions from captive bottlenosed dolphins. *J Wildl Dis* 15: 593-596.
- Foster, G., Macmillan, A.P., Godfroid, J., Howie, F., Ross, H.M., Cloeckert, A., Reid, R.J., Brew, S. And Patterson, I.A.P. 2002. A review of *Brucella* sp. infection of sea mammals with particular emphasis on isolates from Scotland. *Vet Microb* 90: 563-580.
- Foster, G., Osterman, B.S., Godfroid, J., Jacques, I. and Cloeckert, A. 2007. *Brucella ceti* sp. nov. and *Brucella pinnipedialis* sp. nov for *Brucella* strains with cetaceans and seals as their preferred hosts. *Int J Syst Evol Microb* 57: 2688-2693.
- Forcada, J., Aguilar, A., Hammond, P.S., Pastor, X. and Aguilar, R. 1994. Distribution and numbers of striped dolphins in the western Mediterranean Sea after the 1990 epizootic outbreak. *Mar Mamm Sc* 10(2): 137-150.
- Garibaldi, F., Mignone, W., Caroggio, P., Ballardini, M., Podestà, M., Bozzetta, E., Casalone, C., Marsilio, F., Di Francesco, C.E., Proietto, U., Colangelo, P., Scaravelli, D. and Di Guardo, G. 2008. Serological evidence of *Morbillivirus* infection in striped dolphins (*Stenella coeruleoalba*) found stranded on the Ligurian Sea coast of Italy. *Proceedings of 22nd ECS Conference*, Egmond aan Zee, The Netherlands, 10-12. March. 2008, pp. 192-193.
- Geraci, J.R., Hicks, B.D. and St Aubin, D.J. 1979. Dolphin pox: a skin disease of cetaceans. *Can J Comp Med* 43: 399-404
- Gonzalez, L., Patterson, I.A., Reid, R.J., Foster, G., Barberan, M., Blasco, J.M., Kennedy, S., Howie, F.E., Godfroid, J., MacMillan, A.P., Shock, A. and Buxton, D. 2002. Chronic meningoencephalitis associated with *Brucella* sp. infection in live-stranded striped dolphins (*Stenella coeruleoalba*). *J. Comp. Pathol.* 126:147-52.
- Gulland, F. 1995. The impact of infectious diseases on wild animal populations-a review. In: Grenfeld B.T., Dobson, A.P. (eds) Ecology of infectious diseases in natural population. Cambridge University Press, Cambridge, p 20-51
- Gulland, F. M. and A. J. Hall. 2007. Is marine mammal health deteriorating? Trends in the global reporting of marine mammal disease. *EcoHealth* 4: 135-150.

- Hall, A., Hugunin, K., Deaville, R., Law, R.J., Allchin, C.R., and Jepson P. 2006. The risk of infection from polychlorinated biphenyl exposure in the harbor porpoise (*Phocoena phocoena*): a case-control approach. *Environ Health Perspect* 114: 704-711.
- Haubold, E.M., Cooper, C.R. Jr, Wen, J.W., McGinnis, M.R. and Cowan, D.F. 2000. Comparative morphology of *Lacazia loboi* (syn. *Loboa loboi*) in dolphins and humans. *Med Mycol* 38: 9-14.
- Honda, K., Horner, K. and Raugi, G.J. 2007. Lobomycosis. in: Fivenson, D.P., Butler, D.F., Callen, J.P, Quirk, C. and Elston, D.M. (Eds) e-Medicine World Medical Library online: <http://www.emedicine.com/derm/topic832.htm>
- Inskip, W., Gardiner, C.H., Harris, R.K., J.P.Dubey and Goldston, R.T. 1990. Toxoplasmosis in Atlantic bottle-nosed dolphins (*Tursiops truncatus*). *J Wildl Dis* 26: 377-382.
- Jahans, K.L., Foster, G. and Broughton, E.S. 1997. The characterization of *Brucella* strains isolated from marine mammals. *Vet Microb* 57: 373-382.
- Jardine, J.E. and Dubey, J.P. 2002. Congenital toxoplasmosis in an Indo-Pacific bottlenose dolphin (*Tursiops aduncus*). *J Parasitol* 88: 197-199
- Jepson, P.D., Bennett, P.M., Allchin, C.R., Law, R.J., Kuiken, T., Baker, J.R., Rogan, E. and Kirkwood, J.K. 1999. Investigating potential associations between chronic exposure to polychlorinated biphenyls and infectious disease mortality in harbour porpoises from England and Wales. *Sci Total Environ* 243/244: 339-348.
- Jepson, P.D., Bennett, P.M., Deaville, R., Allchin, C.R., Baker, J.R. and Law, R.J. 2005. Relationships between polychlorinated biphenyls and health status in harbor porpoises (*Phocoena phocoena*) stranded in the United Kingdom. *Environ Toxicol Chem* 24: 238-348.
- Kannan, K., Tanabe, S., Borrell, A., Aguilar, A., Focardi, S. and Tatsukawa, R. 1993. Isomer-specific analysis and toxic evaluation of polychlorinated biphenyls in striped dolphins affected by an epizootic in the western Mediterranean Sea. *Arch Environm Cont Toxicol* 25: 227-233.
- Kennedy, S., Smyth, J.A., Cush, P.F., McCullough, S.J., Allan, G.M. and McQuaid, S. 1988. Viral distemper now found in porpoises. *Nature* 336: 21
- Kennedy, S., Smyth, J.A., Cush, P.F., McAliskey, M., McCullough, S.J. and Rima, B.K. 1991. Histopathologic and immunocytochemical studies of distemper in harbour porpoises. *Vet Pathol* 28:1-7
- Kennedy, S. 1998. Morbillivirus infections in aquatic mammals. *J Comp Path* 119:201-225
- Kiska, J. and Van Bresse, M-F. 2008. Lobomycosis-like disease in Indo-Pacific bottlenose dolphins (*Tursiops aduncus*) from the southwest Indian Ocean. Paper SC/60/DW10 presented to the IWC Scientific Committee, May 2008 (unpublished) [Available from the Office of this Journal]
- Krafft, A., Lichy, J.H., Lipscomb, T.P., Klaunberg, B.A., Kennedy, S. and Taunbenberger, J.K. 1995. Postmortem diagnosis of morbillivirus infection in bottlenose dolphins (*Tursiops truncatus*) In the Atlantic and Gulf of Mexico epizootics by polymerase chain reaction-based assay. *J Wildl Dis* 31:410-415
- Lambertsen, R.H., Kohn, B.A., Sundberg, J.P. and Buergelt, C.D. 1987. Genital papillomatosis in sperm whale bulls. *J Wildl Dis* 23:361-367
- Lehanne, L. and Rawlin, G.T. 2001. Topically acquired bacterial zoonoses from fish: a review. *Med J Austral* 7: 480-481.
- Lehmann, T., Marcet, P. L., Graham, D. H., Dahl, E. R. and Dubey, J.P. 2006. Globalization and the population structure of *Toxoplasma gondii*. *PNAS* 103: 11423-11428.
- Lipscomb, T.P., Schulman, F.Y., Moffett, D. and Kennedy, S. 1994. Morbilliviral disease in Atlantic bottlenose dolphins (*Tursiops truncatus*) from the 1987-1988 epizootic. *J Wildl Dis* 30:567-57
- May, R.M. 1986. The cautionary tale of the black-footed ferret. *Nature* 320:13-14
- Martineau, D., Lagace, A., Beland, P., Higgins, R., Armstrong, D. and Shugart, L.R. 1988. Pathology of stranded beluga whales (*Delphinapterus leucas*) from the St. Lawrence Estuary. Quebec. *Can J Comp Pathol* 98:287-
- Migaki, G., Valerio, M.G., Irvine, B.A and Graner, F.M. 1971. Lobo's disease in an Atlantic bottle-nosed dolphin. *JAVMA* 159: 578-582.
- Mikaelian, I., Boisclair, J., Dubey, J.P., Kennedy, S. and Martineau, D. 2000. Toxoplasmosis in beluga whales (*Delphinapterus leucas*) from the St Lawrence estuary: two cases reports and a serological survey. *J Comp Pathol* 122: 73-76.
- Miller, W.G., Adams, L.G., Ficht, T.A., Cheville, N.F., Payeur, J.P., Harley, D.R., House, C. and Ridgway, S.H. 1999. Brucella-induced abortions and infection in bottlenose dolphins (*Tursiops truncatus*). *J Zoo Wildl Med* 30: 100-110.
- Miller, M.A., Gardner, I.A., Kreuder, C., Paradies, D.M., Worcester, K.R., Jessup, D.A., Dodd, E., Harris, M.D., Ames, J.A., Packham, A.E. and Conrad, P.A. 2002. Coastal freshwater runoff is a risk factor for *Toxoplasma gondii* infection of southern sea otters (*Enhydra lutris nereis*). *Int J Parasitol* 32: 997-1006.
- Moreno, I.B., Schiavon, D.D., Martins, M.B., Ott, P.H., Caon, G. and Oliveira, L.R. 2003. Fraser's dolphin (*Lagenodelphis hosei* Fraser, 1956) in southern Brazil. *LAJAM* 2: 39-46.
- Moreno, I. B., Ott, P. H., Tavares, M., Oliveira, L. R., Borba, M. R., Driemeier, D., Nakashima, S. B., Heinzemann, L. S., Siciliano, S. and Van Bresse, M- F. 2008. Mycotic dermatitis in common bottlenose dolphins (*Tursiops*

- truncatus*) from southern Brazil, with a confirmed record of lobomycosis disease. Paper SC/60/DW1 presented to the IWC Scientific Committee, May 2008 (unpublished). 9pp. [Available from the Office of this Journal].
- Mos, L., Morsey, B., Jeffries, S.J., Yunker, M.B., Raverty, S., De Guise, S. and Ross, P.S.R. 2006. Chemical and biological pollution contribute to the immunological profiles of free-ranging harbour seals. *Environm Toxicol Chemist* 25: 310-317.
- Notarbartolo di Sciarra, G., Venturino, M.C., Zanardelli, M., Bearzi, G., Borsani, F.J. and Cavalloni, B. 1993. Cetaceans in the central Mediterranean Sea: distribution and sighting frequencies. *Boll Soc Zool d'Italia*, 60:131-3
- Ohishi, K., Takishita, K., Kawato, M., Zenitani, R., Bando, T., Fujise, Y., Goto, Y., Yamamoto, S., Maruyama, T. 2004. Molecular evidence of new variant *Brucella* in North Pacific common minke whales. *Microb Infect* 6: 1199-2204.
- Ohishi, K., Zenitani, R., Bando, T., Goto, Y., Uchida, K., Maruyama, T., Yamamoto, S., Miyazaki, N., Fujise, Y. 2003. Pathological and serological evidence of *Brucella*-infection in baleen whales (Mysticeti) in the western North Pacific. *Comp Immunol Microb Infect Dis* 26: 125-136.
- Paniz-Mondolfi, A.E., Reyes-Jaimes, O., Dávila-Jones, L. 2007. Lobomycosis in Venezuela. *Int J Dermatol* 46: 180-185.
- Pearce, G., Blacklaws, B.A., Gajda, A.M., Jepson, P., Deaville, R. and Van Bresseem, M-F. 2008. Molecular identification and phylogenetic relationships in poxviruses from cetacean skin lesions. 2008 Conference of the European Cetacean Society, The Netherlands, 10-12 March 2008.
- Pereira, C.S., Amorim, S.D. Santos, A.F.M., Siciliano, S., Moreno, I.M.B., Ott, P.H. and Rodrigues, D.P. 2007. *Vibrio* spp. isolados de mamíferos marinhos capturados na região litorânea do sudeste ao sul do Brasil. *Pesq Vet Bras* 27:81-83.
- Prenger-Berninghoff E., Thoele A., Siebert U., Stede M. and Weiss R. 2003. Phenotypic and molecular characterization of *Brucella* strains isolated from marine mammals of the German North Sea. *Int J Med Microbiol* 293 (36): 320.
- Raga, J.A., Radúan, A., Balbuena, J.A., Aguilar, A., Grau, E. and Borrell, A. 1991. Varamientos de cetáceos en las costas españolas del Mediterraneo durante el período 1982-1988. *Misc Zool* 15: 215-226.
- Raga, J.A., Balbuena, J.A., Aznar, F.J. and Fernandez, M. 1997. The impact of parasites on marine mammals: a review. *Parassit* 39:293-296.
- Raga, J.A., Banyard, A., Domingo, M., Corteyn, M., Van Bresseem, M-F, Fernández, M., Aznar, F.J. and Barrett, T. 2008. Dolphin morbillivirus epizootic resurges in the Mediterranean. *Emerg Infect Dis* 14: 471-473
- Rector, A., Lacave, G., Mostmans, S., Van Doorslaer, K., Rehtanz, M., Salbany, A., Roque, L., Ghim, S-J., Bennett Jenson, A. and Van Ranst, M. 2006. The genetic characterization of a novel close-to-root papillomavirus in a bottlenose dolphin: *Tursiops truncatus* papillomavirus type 1 (TtPV-1). March 17th - 20th, 2006 34 Annual Symposium of EAAM.
- Rehtanz, M., Ghim, S. J., Rector, A., Van Ranst, M., Fair, P., Bossart, G. D. and Jenson, A.B. 2006. Isolation and characterization of the first American bottlenose dolphin papillomavirus: *Tursiops truncatus* papillomavirus type 2. *J Gen Virol* 87, 3559-3565.
- Reif, J.S., Mazzoil, M.S., McCulloch, S.D., Varela, R., Goldstein, J.D., Fair, P. and Bossart, G.D. 2006. Lobomycosis in Atlantic bottlenose dolphins from the Indian River Lagoon, Florida. *JAVMA* 228: 104-108.
- Reilly, S.B. and Barlow, J. 1986. Rates of increase in dolphin population size. *Fish Bull* 84:527-533
- Resendes, A.R., Almería, S., J.P., Dubey, Obón, E., Juan-Sallés, C., Degollada, E., Alegre, E., Cabezón, O., Pont, S., and Domingo, M. 2002. Disseminated toxoplasmosis in a pregnant Risso's dolphin (*Grampus griseus*) with transplacental fetal infection. *J. Parasitol* 88: 1029-1032.
- Rhodes, G., Huys, G., Swings, J., Mc Gann, P., Hiney, M., Smith P et al. 2000. Distribution of oxytetracycline resistance plasmids between aeromonads in hospital and aquaculture environments: Implication of Tn1721 in dissemination of the tetracycline resistance determinant Tet A. *Appl Environ Microbiol* 66: 3883-90.
- Ross, P.S.R. 2002. The role of immunotoxic environmental contaminants in facilitating the emergence of infectious diseases in marine mammals. *HERA* 8: 277-292
- Ross, H.M., Foster, G., Reid, R.J., Jahans, K.L. and Macmillan, A.P. 1994. *Brucella* species infection in sea-mammals. *Vet Rec* 134: 359.
- Ruiz, G.M., Rawlings, T.K., Dobbs, F.C., Drake, L.A., Mullady, T., Huq, A., and Colwell, R.R. 2000. Global spread of microorganisms by ships. *Nature* 408: 49-50.
- Schulman, F.Y., Lipscomb, T.P., Moffett, D., Krafft, A.E., Lichy, J.H., Tsai, M.M., Taubenberger, J.K. and Kennedy, S. 1997. Histologic, immunohistochemical, and polymerase chain reaction studies of bottlenose dolphins from the 1987-88 United States Atlantic coast epizootic. *Vet Pathol* 34 288-
- Shirakihara, M., Amano, M. and Van Bresseem, M-F. 2008. Skin lesion in a resident population of Indo-Pacific bottlenose dolphins in Japan. Paper SC/60/DW6 presented to the IWC Scientific Committee, May 2008 (unpublished) [Available from the Office of this Journal]

- Siciliano, S., Van Bresseem, M-F., Moreno, I.B., Ott, P.H., Tavares, M., Flores, P.A.C., Flach, L., Cesar Reyes, J., Echegaray, M., de Oliveira Santos, M.C., Vididi, F., Crespo, E., Klaich, J., Felix, F., Sanino, G.P. and Van Waerebeek, K. 2008. Review of lobomycosis and lobomycosis-like disease in Cetacea from South America. Paper SC/60/DW13 presented to the IWC Scientific Committee, May 2008 (unpublished) [Available from the Office of this Journal]
- Siebert, U., Joiris, C., Holsbeek, L., Benke, H., Failing, K., Frese, K. and Petzinger, E. 1999. Potential relation between mercury concentrations and necropsy findings in cetaceans from German waters of the North and Baltic Seas. *Mar Poll Bull* 38, 285-295.
- Simões-Lopes, P.C., Paula, G.S. and Xavier, F.M., Scaramelo, A.C. 1993. First case of lobomycosis in bottlenose dolphin on Southern Brazil. *Mar Mamm Sc* 9: 329 - 331.
- Smith, A.W., Skilling, D.E. and Ridgway, S.H. 1983. Calicivirus-induced vesicular disease in cetaceans and probable interspecies transmission. *JAVMA* 11: 1223-1225
- Sundar, N., Cole, R.A., Thomas, N.J., Majumbar, D., Dubey, J.P. and Su, C. 2008. Genetic diversity among sea otter isolates of *Toxoplasma gondii*. *Vet Parasitol* 151:125-132
- Symmers, W.S. 1983. A possible case of Lobo's disease acquired in Europe from a bottle-nosed dolphin (*Tursiops truncatus*). *Bull So Pathol Exot Fil* 76: 777-784.
- Taubenberger, J.K., Tsai, M., Krafft, A.E., Lichy, J.H., Reid, A.H., Schulman, F.Y., Lipscomb, T.P. 1996. Two morbilliviruses implicated in bottlenose dolphin epizootics. *Emerg Infect Dis* 2: 213-216.
- Taborda, P.R., Taborda, V.A. and McGinnins, M.R. 1999. *Lacazia loboi* gen. Nov., comb., the etiologic agent of Lobomycosis. *J Clin Microb* 37: 2031-2033
- Terai, M., DeSalle, R. and Burk, R. D. 2002. Lack of canonical E6 and E7 open reading frames in bird papillomaviruses: *Fringilla coelebs* papillomavirus and *Psittacus erithracus timneh* papillomavirus. *J Virol* 76, 10020-10023.
- Thorne, E.T. and Williams, E. 1988. Disease and endangered species: the black- lack-footed ferret as a recent example *Conserv Biol* 2:66-74
- Tsai, Y.H., Hsu, R.W., Huang, T.J., Hsu, W.H., Huang, K.C., Li, Y.Y. and Peng, K.T. 2000. Necrotizing soft-tissue infections and sepsis caused by *Vibrio vulnificus* compared with those caused by *Aeromonas* species. *Med J Austral* 173: 256-259.
- Tsur, I., Goffman, O., Yakobsen, B., Moffet, D. and Kennedy, S. 1997. Morbillivirus infection in a bottlenose dolphin (*Tursiops truncatus*) from the Mediterranean Sea. *Eur J Vet Pathol* 2:83-85
- UNEP. 2006. Permanent Commission for the South Pacific (CPPS). Humboldt Current, GIWA Regional Assessment 64. University of Kalmar, Kalmar, Sweden. 98 pp.
- Valderrama, C.A., Macgregor, S.K., Rowcliffe, M and Jepson, P.D. 2008. Review of monophasic strain of *Salmonella* group B isolated from cetaceans in England and Wales between 1990 and 2002. *Environm Microbiol* (in press)
- Van Bresseem, M-F. and Van Waerebeek, K. 1996. Epidemiology of poxvirus in small cetaceans from the Eastern South Pacific. *Mar Mam Sci* 12: 371-382
- Van Bresseem, M-F., Visser, I.K.G., Van De Bildt, M.W.G., Teppema, J.S., Raga, J.A. and Osterhaus, A.D.M.E. 1991. Morbillivirus infection in Mediterranean striped dolphins (*Stenella coeruleoalba*). *Vet Rec* 129: 471-472.
- Van Bresseem, M-F., Visser, I.K.G., De Swart, R.L., Örvell, C., Stanzani, L., Androukaki, E., Siakavara, K. and Osterhaus, A.D.M.E. 1993a. Dolphin morbillivirus in different parts of the Mediterranean Sea. *Arch Virol* 129:235-242.
- Van Bresseem, M-F., Van Waerebeek, K., Reyes, J-C., Dekegel, D. and Pastoret P-P. 1993b. Evidence of poxvirus in dusky dolphin (*Lagenorhynchus obscurus*) and Burmeister's porpoise (*Phocoena spinipinnis*) from coastal Peru. *J Wildl Dis* 29: 109-113.
- Van Bresseem, M-F., Van Waerebeek, K and Bennett, M. 2006. Orthopoxvirus neutralising antibodies in small cetaceans from the Southeast Pacific. *LAJAM* 51: 49-54.
- Van Bresseem, M-F., Van Waerebeek, K., Piérard, G. and Desaintes, C. 1996. Genital and lingual warts in small cetaceans from coastal Peru. *Dis Aquat Org* 26: 1-10
- Van Bresseem, M-F., Van Waerebeek, K., Flemming, M. and Barrett, T. 1998. Serological evidence of morbillivirus infection in small cetaceans from the Southeast Pacific. *Vet Microb* 2:89-98.
- Van Bresseem, M-F., Van Waerebeek, K. and Raga, J.A. 1999. A review of virus infections of cetaceans and the potential impact of morbilliviruses, poxviruses and papillomaviruses on host population dynamics. *Dis Aquat Org* 38: 53-65.
- Van Bresseem, M-F., Van Waerebeek, K., Jepson, P.D., Raga, J.A., Duignan, P.J., Nielsen, O., Di Benedetto, A.P., Siciliano, S., Ramos, R., Kant, W., Peddemors, V., Kinoshita, R., Ross, P.S., Lopez-Fernandez, A., Evans, K., Crespo, E. and Barrett, T. 2001a. An insight into the epidemiology of dolphin morbillivirus worldwide. *Vet Microb* 81: 287-304.

- Van Bresseem, M.-F., Van Waerebeek, K., Raga, J.A., Godfroid, J., Brew, S.D. and MacMillan, A.P. 2001b. Serological evidence of *Brucella* species infection in odontocetes from the South Pacific and the Mediterranean. *Vet Rec* 148: 657-661.
- Van Bresseem, M.-F., F., Gaspar, R. and Aznar, J. 2003. Epidemiology of tattoo skin disease in bottlenose dolphins (*Tursiops truncatus*) from the Sado estuary, Portugal. *Dis Aquat Org* 56:171-179.
- Van Bresseem, M.-F., Van Waerebeek, K. and Bennett, M. 2006. Orthopoxvirus neutralising antibodies in small cetaceans from the Southeast Pacific. *LAJAM* 5:49-54.
- Van Bresseem, M.-F., Raga, A.J., Barrett, T., Siciliano, S., Di Benedetto, A.P., Crespo, E. and Van Waerebeek, K. 2007a. A brief review of micro-parasites that may have a potential impact on the population dynamics of small cetaceans from South America. Paper SC/59/DW8 presented to the IWC Scientific Committee, May 2007 (unpublished).
- Van Bresseem, M.-F., Van Waerebeek, K., Reyes, J.C., Félix, F., Echegaray, M., Siciliano, S., Di Benedetto, A.P., Flach, L., Viddi, F., Avila, I.C., Herrera, J.C., Tobón, I.C., Bolaños, J., Moreno, I.B., Ott, P.H., Sanino, G.P., Castineira, E., Montes, D., Crespo, E., Flores, PAC, Haase, B., Mendonça de Souza, S.M.F., Laeta, M. and Frago, A.B. 2007b. A preliminary overview of skin and skeletal diseases and traumata in small cetaceans from South American waters. *LAJAM* 6 (in press)
- Van Bresseem, M.-F., Cassonnet, P., Rector, A., Desaintes, C., Van Waerebeek, K., Alfaro-Shigueto, J., Van Ranst, M and Orth, G. 2007c. Genital warts in Burmeister's porpoises: characterization of *Phocoena spinipinnis* papillomavirus type 1 (PsPV-1) and evidence for a second, distantly related PsPV. *J Gen Virol* 88, 1928-1933.
- Van Bresseem, M.-F., Santos, M. O., Oshima, J.E. 2008a (submitted). Skin diseases in marine tucuxis (*Sotalia guianensis*) from the Parana estuary, Brazil: a possible indicator of a compromised marine environment. *Mar Environm Res*.
- Van Bresseem, M.-F., Van Waerebeek, K., Raga, J.A., Jepson, P.D., Duignan, P., Deaville, R., Flach, L., Viddi, F., Baker, J.R., Di Benedetto, A.P., Echegaray, M., Genov, T., Reyes, J.C., Felix, F., Gaspar, R., Ramos, R., Peddemors, V. and Siebert, U. 2008b (submitted). Epidemiology of tattoo skin disease worldwide: a general health indicator for cetaceans and their environment? *Dis Aquat Org*.
- Van Vresseem, M.-F., Van Waerebeek, K., Flach, L., Reyes, J.C., de Oliveira Santos, M.C., Siciliano, S., Echegaray, M., Viddi, F., Felix, F., Crespo, E., Sanino, G.P., Avila, I.C., Freijja, N. and Castro, C. 2008c. Skin diseases in cetaceans. Paper SC/60/DW8 presented to the IWC Scientific Committee, May 2008 (unpublished) [Available from the Office of this Journal]
- Vilani-Moreno, F.R., Lauris, J.R. and Opromolla, D.V., 2005. Cytokine quantification in the supernatant of mononuclear cell cultures and in blood serum from patients with Jorge Lobo's disease. *Mycopathol* 158: 17-24.
- Visser, I.K.G., Van Bresseem, M.-F., De Swart, R.L., van de Bildt, M.W.G., Vos, H.W., Van der Heijden, R.W.J., Saliki, J.T., Örvell, C., Kitching, P., Kuiken, T., Barrett, T. and Osterhaus, A.D.M.E. 1993. Characterization of morbilliviruses isolated from dolphins and porpoises in Europe. *J Gen Virol* 74:631-641.
- Weber, J.T., Mintz, E.D., Canizares, R., Semiglia, A., Gomez, I., Sempertegui, R., Davila, A., Greene, K.D., Pühr, N.D., Cameron, D.N., Tenover, F.C., Barrett, T.J., Bean, N.H., Ivey, C., Tauxe, R.V., and Blake, P.A. 1994. Epidemic cholera in Ecuador: multidrug-resistance and transmission by water and seafood. *Epid Infect* 112:1-11.
- WHO/UNICEF/WSSCC. 2001. Global Water Supply and Sanitation Assessment 2001. Report World Health Organization and United Nations Children's Found.
- Wohlsein, P., Puff, C., Kreutzer, M., Siebert, U and Baumgärtner, W. 2007. Distemper in a dolphin. *Emer Infect Dis* 13: 1959-1961.