

The effect of salinity on the acute toxicity of copper and cadmium in European sea bass *Dicentrarchus labrax* L.

Loaiza Iván^{1,2}, Marjan Diricx² and Gudrun De Boeck²

¹ Marine and Lacustrine Science and Management (Oceans and Lakes), Faculty of Science and Bioengineering Science, Vrije Universiteit Brussel (VUB), Pleinlaan 2, 1050 Brussels, Belgium
E-mail: ivan.loaiza.alamo@vub.ac.be

² Systematic Physiological and Ecotoxicological Research (SPHERE), University of Antwerp, Groenenborgerlaan 171, 2020 Antwerp, Belgium
E-mail: marjan.diricx@ua.ac.be; gudrun.deboeck@ua.ac.be

Contamination by heavy metals in aquatic environments is a serious concern due to increasing human activities over the last decades. Moreover, productive areas such as estuaries have even been more affected because of the complex interaction between fluvial and marine processes, creating a stressful environment with gradients of salinity and other important parameters which can intensify negative effects of pollutants on biota such as euryhaline fish (Mieiro *et al.*, 2012; Oliva *et al.*, 2012). European sea bass (*Dicentrarchus labrax* L.) is an euryhaline species able to withstand a large variation in salinity and is considered as a bioindicator due to both economic and ecological importance (Loizeau, 2001; Schnitzler *et al.*, 2011). Copper as essential micronutrient has important roles in cellular, enzymatical and protein mechanisms but is potentially toxic when its concentration increases above a certain threshold (Blanchard and Grosell, 2005; Grosell *et al.*, 2007), and can even be more toxic than cadmium, a non-essential metal (Roméo *et al.*, 2000) which is ubiquitous in environment and one of the most deleterious heavy metal pollutants (Nath *et al.*, 1984 cited by Loaiza, 2011). Therefore, the effect of salinity on the acute toxicity of copper and cadmium in European sea bass *D. labrax* L. was examined to determine 96h-LC50 values (the lethal concentration where 50% of the population dies) at different exposure concentration of both metals (at 0 to 5100 $\mu\text{gCu.l}^{-1}$ and at 0 to 20000 $\mu\text{gCd.l}^{-1}$) over a broad salinity range (1, 2.5, 5, 10, 20, 28, 35‰) in juvenile sea bass during a period of 10 days. The results showed an important influence of the salinity in the toxicity in both elements, where for Cu mortality first decreased with increasing salinity but increased again at higher salinities and for Cd, mortality decreased continuously with increasing salinity, to a point where no reliable LC50 could be determined at the 2 highest salinities. In addition, it was observed that the individuals were affected in different ways. For instance Cu exposed fish at low salinity showed opened gills at die-off. Copper is readily taken up by the gill, where it inhibits the action of Na⁺/K⁺ adenosine triphosphatase, resulting in an osmoregulatory disturbance as Na⁺ is lost to the dilute external environment (Blanchard and Grosell, 2006). In conclusion, the salinity influences the toxicity differently for Cu and Cd. This not only occurs due to the fact that salinity plays an important role in the speciation of pollutants and competition by cations in uptake processes, but as can be clearly seen in the case of Cu, also by the physiology of the fish that responds to the changes in the environment. Further research on the relationship between bio-accumulation of the metals and the mortality at different salinities is planned.

Keywords: *Dicentrarchus labrax* L.; salinity; acute exposure.

References

- Blanchard J. and M. Grosell. 2005. Effects of salinity on copper accumulation in the common killifish (*Fundulus heteroclitus*). *Environmental toxicology and chemistry* 24(6):1403-1413.
- Blanchard J. and M. Grosell. 2006. Copper toxicity across salinities from freshwater to seawater in the euryhaline fish *Fundulus heteroclitus*: Is copper an ionoregulatory toxicant in high salinities? *Aquatic Toxicology* 80:131-139.
- Grosell M., J. Blanchard, K.V. Brix and R. Gerdes. 2007. Physiology is pivotal for interactions between salinity and acute copper toxicity to fish and invertebrates. *Aquatic Toxicology* 84:162-172.
- Loizeau V. 2001. A steady-state model of PCB bioaccumulation in the sea bass (*Dicentrarchus labrax*) food web from the Seine Estuary, France. *Estuaries* 24:1074-87.

- Loaiza I. 2011. Bioacumulación de cadmio y plomo en concha de abanico *Argopecten purpuratus* (Lamarck, 1819) en cultivo de fondo, en Parachique - Bahía de Sechura. Tesis. Facultad de Pesquería. UNALM
- Mieiro C.L., J.P. Coelho, M. Pacheco, A.C. Duarte and M.E. Pereira. 2012. Trace elements in two marine fish species during estuarine residency: Non-essential versus essential . *Marine Pollution Bulletin* (in press).
- Oliva M., J.J. Vicente, C. Gravato, L. Guilhermino and M.D. Galindo-Riaño. 2012. Oxidative stress biomarkers in Senegal sole, *Solea senegalensis*, to assess the impact of heavy metal pollution in a Huelva estuary (SW Spain) seasonal and spatial variation. *Ecotoxicol. Environ. Saf.* 75:151-162.
- Roméo M., N. Bennani, M. Gnassia-Barelli, M. Lafaurie and J.P. Girard. 2000. Cadmium and copper display different responses towards oxidative stress in the kidney of the sea bass *Dicentrarchus labrax*. *Aquatic Toxicology* 48:185-194.
- Schnitzler J.G., J.P. Thomé, M. Lepage and K. Das. 2011. Organochlorine pesticides, polychlorinated biphenyls and trace elements in wild European sea bass (*Dicentrarchus labrax*) off European estuaries. *Science of the Total Environment* 409:3680-3686.