

International Council for the
Exploration of the Sea

4274

CM 1982/L:40
Biological
Oceanography
Committee

BIBLIOGRAPHY OF NORTH SEA
BENTHIC LITERATURE

by

A Eleftheriou
DAFS Marine Laboratory
Aberdeen
SCOTLAND

INTRODUCTION

At the first meeting of the ICES North Sea Benthos Working Group (NSBWG) in Texel (8-12 March, 1982) it was recognised that there is a need for a comprehensive bibliography whose aim would be the provision of a list of works concerning qualitative and quantitative aspects of the North Sea level bottom communities. Thus the prepared list comprises mainly published works although an attempt to include as many unpublished documents (reports, data sheets, theses etc.) has also been made.

This compilation has been limited to ecological works concerning the zoobenthos from the subtidal and offshore areas of the North Sea while studies of estuarine, intertidal, fjordic or shallow sea areas have been excluded. Such an exclusion will inevitably restrict the selection of the relevant works which appear in this list. Moreover, works concerning studies of population dynamics of commercial species of molluscs and decapod crustaceans are adequately covered by existing works and they are therefore not included in this list.

It should be stressed that although care was taken to make this bibliography complete and accurate, it is possible that a number of works on faunistic aspects of a localised nature may have been overlooked. Nevertheless, it is hoped that this bibliography will form the basis for a more comprehensive compilation which will subsequently be produced for all those who are interested in the level bottom communities in the North Sea in connection with fisheries or pollution matters.

LITERATURE

Addy, J.M., 1979

Some studies of benthic communities
in areas of oil industry activity.
PhD Thesis, University College of
Swansea.

Addy, J.M., Levell, D. and
Hartley, J.P., 1978

Biological monitoring of sediments
in the Ekofisk oil field. pp 515-539.
In: Proceedings of Conference on Assess-
ment of ecological impacts of oil
spills. Am. Inst. Biol. Sci., June 1978
Keystone Colorado. 514-539.

Allen, J.A., 1960

On the biology of Crangon allmani Kinahan
in Northumberland waters. J. mar. biol.
Ass. UK 39, 481-508

Allen, J.A., 1966

The dynamics and inter-relationships of
mixed populations of Caridea found off
the north-east coast of England.
In: Some contemporary studies in Marine
Science, H. Barnes (ed.), Allen and
Unwin, London, 45-56.

Bakke, T., 1981

The 1980 survey for benthic fauna in the
"Statfjord" area. Institute for Marine
Research, Norway, unpublished report,
7pp.

Barel, C.D.N., Kramers, P.G.N.,
1977

A survey of the echinoderm associates
of the north-east Atlantic area. Zool.
Verhandel. No. 156, 3-159.

Birkett, L., 1953

Changes in the composition of the bottom
fauna of the Dogger Bank area. Nature,
London, 171, 265.

Birkett, L., 1954

Standing crop, mortality and yield of a
Mactra patch. ICES North Sea Committee,
CM 1954, No. 72, 4pp.

Bisschop, G., 1977

Bijdrage tot de studie van de Nematodenfauna
van de Noordzee en de Westerschelde ter
hoogte van haar monding. MSc Thesis,
State University of Ghent. 129pp.

Blegvad, H., 1922

Animal communities in the southern North
Sea. Proc. Zool. Soc. Lond., 27-32.

Blohm, A., 1915

Die Dekapoden der Nord-und Ostsee (mit
Ausnahme der Natantia Boas). Wiss.
Meeresunters. Abt. Kiel, 17, 1-114.

- Caspers, H., 1978 Ecological effects of sewage sludge on benthic fauna off the German North Sea coast. In: Water pollution research. Proceedings of the 8th international conference, Sydney 1976. S.H. Jenkins (ed.), Pergamon Press, Oxford, 951-956.
- Creutzberg, F., 1979 "Aurelia" - cruise reports on the benthic fauna of the Southern North Sea. Introductory Report, Internal Report of the Netherlands Institute for Sea Research, Texel 1979 (4), 31p.
- Creutzberg, F. and van Leeuwen, F., 1980 The life cycle of Crangon allmani Kinahan in the Southern North Sea. ICES Biological Committee CM 1980/L:71, 7pp.
- Creutzberg, F., Wapenaar, P., Duineweld, G. and Lopez, N.L., 1982 Distribution and density of the benthic fauna in the Southern North Sea in relation to bottom characteristics and hydrographic conditions. In: Symposium on Biological Productivity of Continental Shelves in the temperate zone of the North Atlantic, Kiel, 1982, No. 6, 17pp.
- Davis, F.M., 1923. Quantitative studies on the fauna of the sea bottom. No. 1. Preliminary investigations in the Dogger Bank. Fishery Invest. Lond. Ser. 2, 6, 1-54.
- Davis, F.M., 1925 Quantitative studies on the fauna of the sea bottom. No. 2. Results of the investigations on the Southern North Sea 1921-1924. Fishery Invest. Lond. Ser. 2, 8, 1-50.
- De Coninck, L., 1973 Macrobenthos and meiobenthos. In: Math. Modelsea (1973). ICES Fisheries improvement and Hydrography Committee, CM 1973/E:19, 385-398.
- Dicks, B., 1976 Offshore biological monitoring. In: Marine Ecology and Oil Pollution. J.M. Baker (ed.), Applied Science Publs. Ltd., 325-440.
- Dicks, B., Hartley, J., 1975 Biological survey of the benthic fauna in the Forties oil field, June 1975. Unpublished Report, BP Petroleum Development (UK) Ltd.
- Dorjes, J., 1968 Das Makrobenthos. In: Sedimentologie, Faunenzonierung und Faziesabfolge vor der Ostküste der inneren Deutschen Bucht. Reineck, H.E., Dorjes, J., Gadaw, S., and Hertwerk, G. Seckenberg. leth. 49, 272-284.

- Dorjes, J., 1977
Uber die Bodenfauna des Borkum Riffgrundes (Nordsee). Seckenberg. marit. 9 (1/2), 1-17.

Dorjes, J., Gadow, S., Reineck, H.E. and Singh, I.B., 1969
Die Rinnen der Jade (Sudliche Nordsee). Sedimente und Macrofauna. Seckenberg. marit. 1 (50), 5-62.

Dorjes, J., Gadow, S., Reineck, H.E. and Singh, I.B., 1970
Sedimentologie und Macrofauna der Nordergrunde und der Aussenjade (Nordsee). Seckenberg. marit. 2, 31-59.

Dyer, M.F., 1982
Series of North Sea benthos surveys with trawl and headline camera. J. mar. biol. Ass. UK 62, 297-313.

Eagle, R.A., Hardiman, P.A., Norton, M.G., Nunny, R.S. and Rolfe, M.S., 1979
The field assessment of effects of dumping wastes at sea: 5. The disposal of solid wastes off the north east coast of England. Fish. Res. Techn. Rep. MAFF-Lowestoft, No. 51, 34pp.

Ehrenbaum, E., 1896
Die Cumaceen und Schizopoden von Helgoland nebst neueren Beobachtungen über ihr Vorkommen in der deutschen Bucht und in der Nordsee. Wiss. Meeresunters. Abt. Kiel, 2, 401-435.

Eisma, D., 1966
The distribution of benthic marine molluscs off the main Dutch coast. Neth. J. Sea Res. 3 (1), 107-163.

Eleftheriou, A., 1970
Notes on the polychaete Pseudopolydora pulchra (Carazzi) from British waters. Cah. Biol. mar. 11, 459-474.

Eliason, A., 1962
Die Polychaeten der Skagerrak-Expedition 1933. Zool. Bidr. Uppsala, 33, 207-293.

Enequist, P., 1950
Studies on the soft-bottom amphipods of the Skagerrak. Zool. Bidr. Uppsala, 28, 1-492.

Faubel, A., 1978
New Nerillidae (Archiannelida) from the sublittoral of the North Sea and the mid-Atlantic (NW Africa). Zool. Scr., 7 (4), 257-262.

Fauchald, K., 1963
Nephtyidae (Polychaeta) from Norwegian waters. Sarsia, 13, 1-32.

Fauvel, P., 1914
Annelides polychetes non-pelagiques provenant des campagnes de l'HIRONDELLE et de la PRINCESS-ALICE (1885-1910). Result. Camp. sci. Monaco 46, 1-362.

- Ferbrache, J., 1982
Forties field benthic surveys. A review with recommendations for future requirements. Unpublished Report, 14pp., BP Petroleum Development (UK) Ltd.

Forsman, B., 1940
Untersuchungen über die Cumaceen des Skageraks. Zool. Bidr. Uppsala, 18, 1-162.

Gerdes, D., 1977
The re-establishment of an Amphiura filiformis (O.F. Müller) population in the inner part of the German Bight. In: Biology of benthic organisms. Proc. 11th Europ. Symp. Mar. Biol., Galway, October 1976. B.F. Keegan, P.O' Ceidigh and P.J.S. Boaden (eds.), Pergamon Press, 277-284.

Gerlach, S.A., 1978
Food-chain relationships in subtidal silty sand marine sediments and the role of meiofauna in stimulating bacterial productivity. Oecologia, 33, 55-69.

Glemarec, M., 1973
The benthic communities of the European North Atlantic continental shelf. Oceanogr. mar. Biol. Ann. Rev., 11, 263-289.

Govaere, J.C.R., 1976
On Stauronereis caecus (Dorvilleidae, Polychaeta) new for the Southern North Sea. Cah. Biol. mar., 17, 31-36.

Govaere, J.C.R., 1978
Numerieke analyse van het macrobenthos in de Southern Bight (Noordzee). PhD Thesis, State University Ghent, 220pp.

Govaere, J.C.R., Thielemans, L.K.H. and De Boever, R., 1977
Studie van het macrobenthos in de zuidelijke Noordzee. In: Inventaris van de fauna en flora. J.C.J. Nihoul, L. De Coninck (eds.), Projekt Zee, National Onderzoeks-en Ontwikkelingsprogramma, 7, Brussels, 115-165.

Govaere, J.C.R., Van Damme, D., Heip, C. and De Coninck, L.A.P., 1980
Benthic communities in the Southern Bight of the North Sea and their use in ecological monitoring. Helgolander Meeresunters., 33, 507-521.

Hagmeier, A., 1925
Vorlaufiger Bericht über die vorbereitenden Untersuchungen der Bodenfauna der Deutschen Bucht mit dem Petersen-Bodengreifer. Ber. dt. wiss. Kommn. Meeresforsch. NF 1, 247-272.

- Hagmeier, A., 1930
Eine Fluktuation von Mactra (Spisula)
subtruncata Da Costa an der ostfriesischen
Kuste. Ber. dt. wiss. Kommn. Meeres-
forsch. NF 5, 126-155.
- Hartley, J.P., 1977
Survey of the sublittoral macrobenthos
in the Moray Firth, with particular
reference to the Beatrice Oil Field.
June 1977. Monitoring Report. Sphere
Environmental Consultants Ltd., 9pp.
- Hartley, J.P., 1979
Biological monitoring of the sea-bed in
the Forties Oil Field. Paper presented
at the Soc. Petrol. Ind. Biologists
Conference on Ecological Damage Assess-
ment. Arlington, Virginia, 12-14 Nov.
1979, 215-253.
- Hartley, J.P., 1981
The family Paraonidae (Polychaeta) in
British waters: a new species and
new records with a key to species. J.
mar. biol. Ass. UK 61, 133-149.
- Hartley, J.P., 1982
Benthic studies in two North Sea oil
fields. PhD Thesis, University of
Wales, 191pp.
- Hartley, J.P. and
Cartlidge, D.M., 1981
Third biological survey of the benthic
sediments in the Forties Oil Field,
June 1981. Unpublished Report to
BP Petroleum Development (UK) Ltd.
- Hartmann-Schroder, G., 1974
Polychaeten von Expeditionen der
"Anton Dohrn" in Nordsee und Skagerrak.
Veroff. Inst. Meeresforsch. Bremerh.
14, 169-274.
- Hartmann-Schroder, G. and
Stripp, K., 1968
Beitrag zur Polychaetenfauna der
Deutschen Bucht. Veroff. Inst.
Meeresforsch. Bremerh. 11, 1-24.
- Heincke, F., 1894
Die Mollusken Helgolands. Wiss.
Meeresunters. Abt. Kiel 1, 121-153.
- Heinen, A., 1911
Die Nephthydeen und Lycoriden der
Nord-und Ostsee. Wiss. Meeresunters.
Abt. Kiel, 13, 1-88.
- Heip, C. and
Decraemer, W., 1974
The diversity of nematode communities
in the Southern North Sea. J. mar.
biol. Ass. UK 54, 251-255.
- Heip, C., Herman, R.,
Bisschop, G., Govaere, J.C.R.,
Holvoet, M., van Damme, D.,
Vanosmael, C., Willems, K.R.
and De Coninck, L.A.P., 1979
Benthic studies of the Southern Bight of the
North Sea and its adjacent continental
estuaries. Progress Report I. ICES
Biological Oceanography Committee, CM 1979/
L:9, 133-163.

- Hilbig, B. and
Dittmer, J.D., 1979 *Gyptis helgolandica* sp. n. (Hesionidae, Annelida) a new Polychaete from the sub-littoral zone of the German Bight. Veroff. Inst. Meeresforsch. Bremerh. 18, 101-110.
- Høglund, H., 1947 Foraminifera in the Gullmar Fjord and the Skagerak. Zool. Bidr. Uppsala, 26, 1-328.
- Howard, F.G., 1977 Some notes on the biology of Pandalus borealis Krøyer on Fladen. ICES Shellfish and Benthos Committee CM 1977/K:30, 2pp.
- Hult, J., 1941 On the soft bottom isopods of the Skager Rak. Zool. Bidr. Uppsala, 21, 1-234.
- Institute of Environmental Engineering, 1979 Murchison Field. Environmental base-line study. August 1978 survey. Unpublished Report to CONOCO, 257pp.
- Institute of Environmental Engineering, 1980 Hutton and Murchison Oil Fields. Macrofauna assessment, 1980 survey. Unpublished Report to CONOCO.
- Institute of Environmental Engineering, 1982 West Sole Oil Field environmental study. Unpublished Report to BP(UK).
- Institute of Environmental Engineering, 1982 Brae Oil Field environmental study. 1981 survey. Unpublished Report to MARATHON, 300pp.
- Jensen, P., 1976 Free-living marine nematodes from a sublittoral region in the North Sea off the Belgian Coast. Biol. Jaarb. Dodonaea, 44, 231-255.
- Jones, R., 1982 Some observations on energy transfer through the North Sea and Georges Bank food webs. In: Symposium on Biological Productivity of Continental Shelves in the temperate zone of the North Atlantic, Kiel 1982, No. 30, 15pp.
- Josefson, A.B., 1975 Ophyotrocha longidentata sp. n. and Dorvillea erucaeformis (Malmgren) (Polychaeta, Dorvilleidae) from the West Coast of Scandinavia. Zool. Scr. 4, 49-54.
- Josefson, A.B., 1981 Persistence and structure of two deep macrobenthic communities in the Skagerrak (west coast of Sweden). J. exp. mar. Biol. Ecol. 50 (1), 63-98.

- Josefson, A.B., 1982
Regulation of population size, growth
and production of a deposit feeding
bivalve: a long term field study
of three deep water populations off
the Swedish west coast. J. exp. mar.
Biol. Ecol. 59, 125-150
- Juario, J.V., 1972
Cyartonema germanicum sp. n. (Nematoda:
Siphonolaimidae) aus dem Sublitoral der
Deutschen Bucht und Bemerkungen zur
Gattung Cyartonema Cobb 1920. Veröff.
Inst. Meeresforsch. Bremerh. 14, 81-86.
- Juario, J.V., 1974
Neue freilebende Nematoden aus dem
Sublitoral der Deutschen Bucht. Veröff.
Inst. Meeresforsch. Bremerh. 14, 275-303.
- Juario, J.V., 1975
Nematode species composition and seasonal
fluctuation of a sublittoral meiofauna
community on the German Bight. Veröff.
Inst. Meeresforsch. Bremerh. 15, 283-337.
- Kingston, P.F., 1980
Firth of Forth benthic faunal survey
1976-1977. Unpublished data sheets,
Heriot-Watt University, Edinburgh.
- Kingston, P.F. and
Mackie, A.S.Y., 1980
Octobranchus floriceps sp. nov.
(Polychaeta: Trichobranchidae) from
the northern North Sea with a
re-examination of O. antarcticus Monro.
Sarsia, 65, (3/4), 249-254.
- Kirkegaard, J.B., 1958
Records of the group Pogonophora in the
Skagerrak. Nature, Lond. 181, 1086-1087.
- Kirkegaard, J.B., 1969
A quantitative investigation of the
central North Sea Polychaeta. Spolia
Zool. Mus. haun. 29, 1-284.
- Kirkegaard, J.B., 1978
Production by polychaetes on the Dogger
Bank in the North Sea. Meddr Danm. Fisk.
-og Havunders. N.S., 7, 497-509.
- Klein, G., 1969
Amphipoden ans der Wesermündung und der
Helgoländer Bucht, mit Beschreibung von
Talorchestia frisiae n. sp. Veröff.
Inst. Meeresforsch. Bremerh. 11, 173-194.
- Klein, G., 1973
Populationsdynamik und Produktionsleistung
des benthischen röhrenbauenden Amphipoden
Ampelisca brevicornis an zwei stationen
in der Helgoländer Bucht. Unpublished
Diplomarbeit, Ruhr Universität, Bochum,
28pp.

- Klein, G., Rachor, E.
and Gerlach, S.A., 1975 Dynamics and productivity of two populations of the benthic tube-dwelling amphipod Ampelisca brevicornis in Helgoland Bight. *Ophelia* 14, 139-159.
- Lorenzen, S., 1972 Die Nematodenfauna im Verklappungsgebiet für Industrie-abwässer nordwestlich von Helgoland. I. Araeolaimida und Monhysterida. *Zool. Anz.* 187, 223-248.
- Lorenzen, S., 1972 Die Nematodenfauna im Verklappungsgebiet für Industrie-abwässer nordwestlich von Helgoland. II. Desmodorida und Chromadorida. *Zool. Anz.* 187, 283-302.
- Lorenzen, S., 1972 Die Nematodenfauna im Verklappungsgebiet für Industrie-abwässer nordwestlich von Helgoland. III. Cyatholaimidae, mit einer Revision von Pomponema Cobb 1917. *Veröff. Inst. Meeresforsch. Bremerh.* 13, 285-300.
- Lorenzen, S., 1972 Demoscolex-Arten (freilebende Nematoden) von der Nord- und Ostsee. *Veröff. Inst. Meeresforsch. Bremerh.* 13, 307-316.
- Lorenzen, S., 1973 Freilebende Meeresnematoden aus dem Sublitoral der Nordsee und der Kieler Bucht. *Veröff. Inst. Meeresforsch. Bremerh.* 14, 103-130.
- Lorenzen, S., 1974 Die Nematodenfauna der sublitoralen Region der Deutschen Bucht, insbesondere im Titan-Abwassergebiet bei Helgoland. *Veröff. Inst. Meeresforsch. Bremerh.* 14, 305-327.
- Lorenzen, S., 1977 Revision der Xyalidae (freilebende Nematoden) auf der Grundlage einer kritischen Analyse von 56 Arten aus Nord- und Ostsee. *Veröff. Inst. Meeresforsch. Bremerh.* 16, 197-261.
- McIntyre, A.D., 1958 The Ecology of Scottish inshore fishing grounds. I. The bottom fauna of east coast grounds. *Mar. Res.* 1, 1-24.
- McIntyre, A.D., 1961 New records of polychaetes from Scottish coastal and offshore waters. *Proc. Roy. Soc. Edinb., Sect. B*, 67, No. 14, 351-362.
- McIntyre, A.D., 1961 Quantitative differences in the fauna of boreal mud associations. *J. mar. Biol. Ass. UK* 41, 599-616.

- Moore, D.C., 1981
Studies of environmental effects of sewage sludge dumping off the Firth of Forth. ICES, Marine Environmental Quality Committee, CM 1981/E:42, 3pp.
- Muhlenhardt-Siegel, U., 1981
Die Biomasse mariner Macrobenthos-Gesellschaften im Einflussbereich der Klärschlammverklappung von der Elbmündung. Helgolander Meeresunters. 34 (4), 427-437.
- Nihoul, J.C.J. and Polk, P., 1977
Chaines trophiques et cycles des nutrients. Projekt Zee, 8. Programme National de Recherche et de developpement, Bruxelles, 1-339.
- Nolte, W., 1913
Zur Kenntnis der Maldaniden der Nord-und Ostsee. Wiss. Meeresunters. Abt. Kiel, 15, 1-94.
- Norman, A.M., 1869
Shetland final dredging report. Part II. On the Crustacea, Tunicata, Polyzoa, Echinodermata, Actinozoa and Porifera. Rep. Br. Ass. Advmt Sci., 1868, 247-336.
- Norton, P.E.P., 1973
The history and future of the North Sea benthos. Proc. Roy. Soc. Edinb., 76B, 193-200.
- Petersen, C.G.J., 1913
Valuation of the sea. II. The animal communities of the sea bottom and their importance for marine zoogeography. Rep. Dan. biol. Stn., 21, 1-44.
- Petersen, C.G.J., 1915
On the animal communities of the sea bottom in the Skagerrak, the Christiania Fjord and the Danish waters. Rep. Dan. biol. Stn., 23, 3-28.
- Petersen, C.G.J., 1918
The sea bottom and its production of fish-food. Rep. Dan. biol. Stn., 25, 1-62.
- Petersen, C.G.J. and Boysen Jensen, P., 1911
Valuation of the Sea. I. Animal life of the sea bottom, its food and quantity. Rep. Dan. biol. Stn., 20, 1-76.
- Petersen, G.H., 1977
The density, biomass and origin of the bivalves of the central North Sea. Meddr. Danm. Fisk-og Havunders. 7, 221-273.
- Poizat, C., 1980
Gastéropodes opistobranches mesopsammiques du Skagerrak (Suède Occidentale). Distribution et dynamique de population. Vie Milieu 30 (3/4), 209-223.

- Poizat, C. and Vicente, N., 1977 Donnees preliminaires sur l'ecologie et la biologie de quelques Gasteropodes opisthobranches du Skagerrak (Suede Occidentale). Bull. Mus. Hist. nat. (Ser. 3), 439 (Ecol. gen., 34).., 3-18.
- Por, F.D., 1964 Les Harpacticoides (Crustacea, Copepoda) des fonds meubles du Skagerak. Cah. Biol. mar. 5 (3), 233-270.
- Poulsen, E.M., 1946 Investigations on the Danish Fishery for and the biology of the Norway lobster and the Deep-Sea Prawn. Rep. Dan. biol. Stn. 48 (1943-1945), 27-50.
- Rachor, E., 1972 On the influence of industrial waste containing H_2SO_4 on the bottom fauna off Helgoland (German Bight). In: Marine Pollution and sea life. M. Ruivo (ed.), Fishing News (Books) Ltd, London, 390-392.
- Rachor, E., 1976 Structure, dynamics and productivity of a population of Nucula nitidosa (Bivalvia: Protobranchiata) in the German Bight. Ber. dt. wiss. Kommn. Meeresforsch., 24, 296-331.
- Rachor, E., 1977 Faunenverarmung in einem Schlickgebiet in der Nahe Helgolands. Helgolander wiss. Meeresunters. 30, 633-651.
- Rachor, E., 1980 The inner German Bight - an ecologically sensitive area as indicated by bottom fauna. Helgolander Meeresunters. 33, 522-530.
- Rachor, E. and Bartel, S., 1981 Occurrence and ecological significance of the spoon-worm Echiurus echiurus in the German Bight. Veröff. Inst. Meeresforsch. Bremerh. 19, 71-88.
- Rachor, E. and Dethlefsen, V., 1974 Effects of acid-iron waste disposal in the marine environment. ICES Fishery Improvement Committee, CM 1974/E:27, 14pp.
- Rachor, E. and Gerlach, S.A., 1975 Variations in macrobenthos in the German Bight. In: ICES Symp. Changes in North Sea Fish Stocks, and their causes. Aarhus July 9-12, 1975, 11.
- Rachor, E. and Gerlach, S.A., 1978 Changes of macrobenthos in a sublittoral sand area of the German Bight, 1967 to 1975. Rapp. P.-v. Reun. Cons. int. Explor. Mer, 172, 418-431.

- Rachor, E. and Salzwedel, H., 1976
- Studies on population dynamics and productivity of some bivalves in the German Bight. Proc. 10th Europ. Symp. Mar. Biol. Ostend, September 1975. G. Persoone and E. Jaspers (eds.), Universe Press, 575-588.
- Rachor, E., Salzwedel, H., Gerdes, D. and Krause, H., 1981
- Boden-tiergemeinschaften im Sublitoral der Deutschen Bucht und ihre zeitliche Variabilität. Verhandlungen dtsch. zool. Ges. Bremen, (in press).
- Raiett, D.S., 1937
- The benthic amphipoda of the North-Western North Sea and adjacent waters. Proc. Roy. Soc. Edin. 57, (No. 18), 241-254.
- Redant, F., 1974
- The benthic and epibenthic fauna of the Thornton Bank area. ICES Shellfish and Benthos Committee, CM 1974/K:22, 5pp.
- Redant, F., 1977
- Epibenthos. In: Inventaris van de fauna en flora. J.C.J. Nihoul and L. De Coninck (eds.), Projekt Zee, 7, Nationaal Onderzoeks-en Ontwikkelingsprogramma, Brussels, 167-202.
- Reibisch, J., 1905
- Faunistisch-biologische Untersuchungen über Amphipoden der Nordsee. Wiss. Meeresunters. Abt. Kiel, 8, 145-188.
- Reineck, H.-E., Dorjes, J., Gadow, S. and Hertweck, G., 1968
- Sedimentologie, Faunenzonierung und Faziesabfolge vor der Ostküste der inneren Deutschen Bucht. Seckenberg. leth. 49, 261-309.
- Remane, A., 1940
- Einführung in die Zoologische Ökologie der Nord- und Ostsee. In: G. Grimpe and E. Wagler (eds.): Die Tierwelt der Nord- und Oestsee, Ia: 1-238, Leipzig (Becker and Erler).
- Salvini-Plawen, L. von, 1971
- Die nordatlantischen Caudofoveata des zoologischen Museums zu Kopenhagen (Mollusca, Aculifera). Steenstrupia 2, 27-34.
- Salzwedel, H., 1979
- Reproduction, growth, mortality and variations in abundance and biomass of Tellina fabula (Bivalvia) in the German Bight in 1975/76. Veröff. Inst. Meeresforsch. Bremerh. 18, 111-202.

- Salzwedel, H., 1980 Energy budgets for two populations of the bivalve Tellina fabula in the German Bight. Veröff. Inst. Meeresforsch. Bremerh. 18, 257-287.
- Schrader, E., 1911 Lamellibranchiaten der Nordsee. Wiss. Meeresunters. Abt. Kiel, 12, 1-71.
- Schuster, O., 1952 Die vareler Rinne im Jadebusen. Abh. seckenberg. naturforsch. Ges. 486, 1-38.
- Shelton, R.G.J., 1973 Some effects of dumped, solid wastes on marine life and fisheries. In: North Sea Science, E.D. Goldberg (ed.), Massachusetts Institute of Technology, 415-436.
- Small, W., 1912 Report on the Annelida Polychaeta collected in the North Sea and adjacent parts by the Scottish Fishery Board vessel 'Goldseeker'. Ann. Mag. Nat. Hist. 10, 165-185.
- Souplet, A. and Dewarumez, J.M., 1980 Les peuplements benthiques du littoral de la region de Dunkerque. Cah. Biol. mar. 21, 23-39.
- Souplet, A., Glacon, R., Dewarumez, J.M. and Smigelski, F., 1980 Distribution des peuplements benthiques littoraux en Mer du Nord, du Cap Blanc-Nez a la frontiere de Belgique. C.R. Acad. Sci. Paris Ser. D, 290, 627-630.
- Steele, J.H., 1974 The structure of marine ecosystems. Blackwell Scient. Publs. Oxford, 128pp.
- Stephen, A.C., 1923 Preliminary survey of the Scottish waters of the North Sea by the Petersen grab. Sci. Invest. Fish. Scot., 1922, No. 3, 21pp.
- Stephen, A.C., 1933 Studies on the Scottish marine fauna: The natural faunistic divisions of the North Sea as shown by the quantitative distribution of the molluscs. Trans. Roy. Soc. Edinb. 57 (pt III). (No. 22), 601-616.
- Stephen, A.C., 1934 Studies on the Scottish marine fauna: Quantitative distribution of the echinoderms and the natural faunistic divisions of the North Sea. Trans. Roy. Soc. Edinb. 57 (pt III). (No. 32), 777-787.

- Stephensen, K., 1923 Crustacea Malacostraca. V: Amphipoda I. Dan. Ingolf Expedition, 3 (8), 1-100.
- Stripp, K., 1969 Die Assoziationen des Benthos in der Helgolander Bucht. Veröff. Inst. Meeresforsch. Bremerh. 12, 95-142.
- Stripp, K., 1969 Jahrezeitliche Fluktuationen von Macrofauna und Meiofauna in der Helgolander Bucht. Veröff. Inst. Meeresforsch. Bremerh. 12, 65-94.
- Stripp, K., 1969 Das Verhältnis von Macrofauna und Meiofauna in den Sedimenten der Helgoländer Bucht. Veröff. Inst. Meeresforsch. Bremerh. 12, 143-148.
- Stripp, K. and Gerlach, S.A., 1969 Die Bodenfauna im Verklappungsgebiet von Industrieabwässern nordwestlich von Helgoland. Veröff. Inst. Meeresforsch. Bremerh. 12, 149-156.
- Sussbach, S. and Breckner, A., 1911 Die Seigel, Seesterne und Schlangensterne der Nord- und Ostsee. Wiss. Meeresunters. Abt. Kiel 12, 167-300.
- Thorson, G., 1957 Bottom communities. In: Treatise on Marine Ecology and Paleoecology. J.W. Hedgpeth (ed.). Geol. Soc. America, Memoir 67, 1, 461-534.
- Ursin, E., 1952 Change in the composition of the bottom fauna of the Dogger Bank area. Nature, Lond. 170, 324.
- Ursin, E., 1960 A quantitative investigation of the echinoderm fauna of the central North Sea. Meddr. Danm. Fisk-og. Havunders. N.S. 2 (24), 1-204.
- Van Damme, D. and Heip, C., 1977 Het meiobenthos in de Zuidelijke Noordzee. In: Inventaris van de fauna en flora. J.C.J. Nihoul and L. De Coninck (eds.). Projekt Zee, 7, Nationaal Onderzoeks-en Ontwikkelingsprogramma, Brussels, 1-113.
- Van Noort, G.J. and Creutzberg, F., 1981 "Aurelia" - cruise reports on the benthic fauna of the Southern North Sea. Report 5: Trawl Survey October-November 1974. Internal Report of the Netherlands Institute for Sea Research, Texel 1981, (1), 116pp.

- Van Noort, G.J.,
Van Leeuwen, F. and
Creutzberg, F., 1979
- "Aurelia" - cruise reports on the benthic fauna of the Southern North Sea.
Report 1: Trawl survey April-May 1972.
Internal Report of the Netherlands Institute for Sea Research, Texel 1979, (5), 104pp.
- Van Noort, G.J.,
Van Leeuwen, F. and
Creutzberg, F., 1979
- "Aurelia" - cruise reports on the benthic fauna of the Southern North Sea.
Report 2: Trawl Survey June-July 1972.
Internal Report of the Netherlands Institute for Sea Research, Texel, 1979, (6), 107pp.
- Van Noort, G.J.,
Van Leeuwen, F. and
Creutzberg, F., 1979
- "Aurelia" - cruise reports on the benthic fauna of the Southern North Sea. Report 3: Trawl Survey October-November 1972.
Internal Report of the Netherlands Institute for Sea Research, Texel 1979, (7), 115pp.
- Van Noort, G.J.,
Van Leeuwen, F. and
Creutzberg, F., 1979
- "Aurelia" - cruise reports on the benthic fauna of the Southern North Sea.
Report 4: Trawl Survey January-February 1973. Internal Report of the Netherlands Institute for Sea Research, Texel, 1979, (8), 100pp.
- Van Noort, G.J. and
Creutzberg, F., 1981
- "Aurelia" - cruise reports on the benthic fauna of the Southern North Sea.
Report 5: Trawl Survey October-November 1974. Internal Report of the Netherlands Institute for Sea Research, Texel 1981, (1), 116pp.
- Vincx, M., 1981
- New and little-known nematodes from the North Sea. Cah. Biol. mar. 22, 431-451.
- Voit, M., 1911
- Die Glyceriden der Nordsee. Wiss. Meeresunters. Abt. Kiel, NF 13, 89-125.
- Warwick, R.M. and
Buchanan, J.B., 1970
- The meiofauna off the coast of Northumberland. I. The structure of the nematode population. J. mar. biol. Ass. UK, 50, 129-146.
- Ziegelmeier, E., 1953
- Kurze Mitteilung über zwei für die Deutsche Bucht neue Polychatenarten der Gattungen Capitomastus und Goniada. Helgolander wiss. Meeresunters. 4, 253-259.
- Ziegelmeier, E., 1954
- Investigations of the bottom fauna. Ann. Biol. 11 (1956), 63.

