

Deciphering Mega-Ripple variability in an anthropogenically-steered environment: implications for mine burial studies.

Sonia Papili^(1,2), Matthias Baeye⁽³⁾, Vera Van Lancker⁽³⁾

1. Belgian Navy, Marinebasis Zeebrugge, Graaf Jansdijk 1, 8380 Zeebrugge, Belgium Email: sonia.papili@mil.be,

2. RCMG (Renard Centre of Marine Geology), Department Geology and Soil Science, Ghent University, Krijgslaan 281-S8, 9000 Gent. Belgium. Email: sonia.papili@ugent.be

3. Royal Belgian Institute of Natural Sciences. Management Unit of the North Sea Mathematical Models. Gulledele 100, 1200 Brussels, Belgium. Email: Vera.vanlancker@mumm.ac.be, Matthias.Baeye@mumm.ac.be

ABSTRACT

In 2007 the Ministry of Defence, in collaboration with Ghent University, developed a project on the understanding of mega-ripple variability in view of improving mine burial prediction models in sandbank areas. Results will assist in the monitoring of sea-mines, heritage of two World Wars, nowadays partially or totally buried by sandy bedforms.

In this paper we will present results from ‘The Vlakte Van De Raan – Akkaert Bank’ region, a sandbank area on the Belgian Continental Shelf where intensive disposal of dredged material takes place. Five years time series of multibeam echosoundings (Kongsberg Simrad EM1002-EM3002) were built-up.

Migration of dunes (2-3m in height) has been observed of up to 20m in a South West direction opposite to the dominant North East current direction. Magnitudes of migration are largest near to the disposal ground (Van Lancker et al., 2012). Megaripples has been observed too, but their variability was not quantified. During this study a series of parallel profiles, aligned transversally to the dunes, were investigated in time. Profiles with ebb-dominance were analyzed separately from those with a flood-dominance. Differences are highlighted between those closely settled to the disposal ground and those further located.

From the bedform migration, a volume quantification was performed following the results of Duffy (2012). Results are being analysed against hydro-meteorological influences and volumes of dumped material.

REFERENCES

- V. Van Lancker, M. Baeye, I. Du Four, R. Janssens, S. Degraer, M. Fettweis, F. Francken, J.S. Houziaux, P. Luytan, D. Van den Eynde, M. Devolder, K. De Cauwer, J. Monbaliu, E. Toorman, J. Portilla, A. Ullman, M. Liste Munõz, L. Fernandez, H. Komijani, T. Verwaest, R. Delgado, J. De Schutter, J. Janssens, Y. Levy, J. Vanlede, M. Vincx, M. Rabaut, N. Vandenberghe, E. Zeelmaekers, A. Goffin. (2012). *Quantification of Erosion/Sedimentation patterns to trace the natural versus anthropogenic sediment dynamics ‘QUEST4D’*. Final report Brussels : Belgian Science Policy Office.103 p. (Research Program Science for a Sustainable Development).
- Garret P; Duffy, John E. Hughes Clarke. (2012). Measurement of bedload transport in a coastal sea using repeat swath bathymetry surveys: assessing bedload formulae using sand dune migration. International Association of Sedimentologists and published for them by Blackwell Publishing Ltd. 44, pag 249-272