

Yoshida, N.
Osaka J. Math.
39 (2002), 89–96

A SEQUENCE IN THE CLASSICAL SCHOTTKY SPACE

NAOSHI YOSHIDA

(Received June 5, 2000)

1. Introduction

Let \mathbb{M} be the topological group of all linear fractional transformations. Its multiplication is the composition of mappings and its topology is the uniform convergence topology on the extended complex plane $\widehat{\mathbb{C}}$.

Let r be a positive integer. We denote the free group with basis $\{1, \dots, r\}$ by F_r . The mapping from $\theta \in \text{Hom}(F_r, \mathbb{M})$ to $(\theta(1), \dots, \theta(r)) \in \mathbb{M}^r$ is bijective. We give $\text{Hom}(F_r, \mathbb{M})$ the topology such that this bijection is a homeomorphism. When $\theta \in \text{Hom}(F_r, \mathbb{M})$ is a monomorphism, θ^{-1} is the inverse of the isomorphism θ whose range is restricted to $\text{Im } \theta$. For $\varphi \in \mathbb{M}$ and $\theta \in \text{Hom}(F_r, \mathbb{M})$, we define $\varphi\theta \in \text{Hom}(F_r, \mathbb{M})$ to be $(\varphi\theta)(x) = \varphi \circ \theta(x) \circ \varphi^{-1}$ for every x in F_r . In this way, \mathbb{M} acts on $\text{Hom}(F_r, \mathbb{M})$.

Let r be a positive integer greater than one. Define the *Schottky space* \mathbb{S}_r of rank r to be

$$\mathbb{S}_r = \{\theta \in \text{Hom}(F_r, \mathbb{M}) \mid \text{Im } \theta \text{ is a Schottky group of rank } r\}.$$

\mathbb{S}_r is \mathbb{M} -invariant. The Schottky space of rank r defined in Chuckrow [2] is \mathbb{S}_r/\mathbb{M} . But the results of Chuckrow [2] which we use also hold for the Schottky space in our sense. We denote by $\partial\mathbb{S}_r$ the boundary of \mathbb{S}_r in $\text{Hom}(F_r, \mathbb{M})$. An element of $\partial\mathbb{S}_r$ is called a *cusp* if its image has parabolic transformations. The following results are shown in Chuckrow [2]:

- (1) \mathbb{S}_r is open and connected in $\text{Hom}(F_r, \mathbb{M})$ (Chuckrow [2, Lemma 5]).
- (2) Every element of $\partial\mathbb{S}_r$ is a monomorphism and has an image without elliptic transformations (Chuckrow [2, Theorem 4]).
- (3) If $\theta \in \partial\mathbb{S}_r$ is not a cusp, then $\text{Im } \theta$ does not act discontinuously on any open subset of $\widehat{\mathbb{C}}$ (Chuckrow [2, Theorem 5]).

Define the *classical Schottky space* \mathbb{S}_r^0 of rank r to be

$$\mathbb{S}_r^0 = \{\theta \in \text{Hom}(F_r, \mathbb{M}) \mid \text{Im } \theta \text{ is a classical Schottky group of rank } r\}.$$

Let $\overline{\mathbb{S}_r^0}$ be the closure of \mathbb{S}_r^0 in $\text{Hom}(F_r, \mathbb{M})$. If θ belongs to $\partial\mathbb{S}_r \cap \overline{\mathbb{S}_r^0}$, then $\text{Im } \theta$ acts

discontinuously on some open subset of $\widehat{\mathbb{C}}$ (Marden [4, Proposition 3.1]). Thus every element of $\partial\mathbb{S}_r \cap \overline{\mathbb{S}_r^0}$ is a cusp.

For each loxodromic transformation f , we denote the multiplier of f by $\lambda(f)$ ($|\lambda(f)| > 1$). The main result of this paper is as follows:

Theorem. *Let r be an integer greater than one. If a sequence $\{\theta_n\}_{n=1}^{+\infty}$ in \mathbb{S}_r^0 converges to θ in $\partial\mathbb{S}_r \cap \overline{\mathbb{S}_r^0}$ as n tends to $+\infty$, then for each parabolic transformation φ of $\text{Im } \theta$, $\lambda(\theta_n \circ \theta^{-1}(\varphi))$ converges to 1 conically as n tends to $+\infty$. Namely, $\lambda(\theta_n \circ \theta^{-1}(\varphi))$ converges to 1 and*

$$\left\{ \frac{|\lambda(\theta_n \circ \theta^{-1}(\varphi)) - 1|}{|\lambda(\theta_n \circ \theta^{-1}(\varphi))| - 1} \right\}_{n=1}^{+\infty}$$

is bounded.

Using McMullen [7, Theorem 7.3], we obtain the following:

Corollary. *Let r be an integer greater than one. If a sequence $\{\theta_n\}_{n=1}^{+\infty}$ in \mathbb{S}_r^0 converges to θ in $\partial\mathbb{S}_r \cap \overline{\mathbb{S}_r^0}$ as n tends to $+\infty$, then*

- (1) $\text{Im } \theta_n$ converges to $\text{Im } \theta$ geometrically;
- (2) the limit set of $\text{Im } \theta_n$ converges to the limit set of $\text{Im } \theta$ in the sense of Hausdorff convergence;
- (3) the Patterson-Sullivan measure of $\text{Im } \theta_n$ converges to the measure of $\text{Im } \theta$ weakly;
- (4) the critical exponent of $\text{Im } \theta_n$ converges to the critical exponent of $\text{Im } \theta$, as n tends to $+\infty$.

In section 2, we will recall the definition of a Schottky group, and we will also prove a lemma. In section 3, we will prove our theorem. In section 4, we will show that \mathbb{S}_r^0 in our theorem cannot be replaced with \mathbb{S}_r even if θ belongs to $\partial\mathbb{S}_r \cap \overline{\mathbb{S}_r^0}$.

The author is deeply grateful to Professor Hiroki Sato for his valuable suggestions and encouragement. Also he is grateful to Professors Katsuhiko Matsuzaki and Yoshihide Okumura. Thanks are due to the referees for their careful reading and valuable suggestions.

2. Schottky Groups

Let r be an integer greater than one. A subgroup G of \mathbb{M} is a *Schottky group of rank r* if there exist a set of generators h_1, \dots, h_r of G and $2r$ mutually disjoint Jordan curves $C_1, C_{-1}, \dots, C_r, C_{-r}$ on $\widehat{\mathbb{C}}$ which satisfy the following conditions:

- (1) $C_1, C_{-1}, \dots, C_r, C_{-r}$ bound a $2r$ -ply connected region R .
- (2) For each i in $\{1, \dots, r\}$, h_i maps C_i onto C_{-i} .
- (3) For each i in $\{1, \dots, r\}$, $h_i(R)$ and R are mutually disjoint.

In the above definition, if Jordan curves can be replaced with circles, then G is called a *classical Schottky group of rank r* . A Schottky group of rank r is free of rank r , purely loxodromic and acts discontinuously on some open subset of $\widehat{\mathbb{C}}$.

EXAMPLE (cf. McMullen [6, Theorem 3.1]). For each positive integer n , let C_{1n}, \dots, C_{r+1n} be circles on $\widehat{\mathbb{C}}$ which bound an $(r+1)$ -ply connected region ($r \geq 2$). Suppose that C_{1n}, \dots, C_{r+1n} converge to circles C_1, \dots, C_{r+1} as n tends to $+\infty$, respectively; C_1, \dots, C_{r+1} may be tangent but cannot intersect. Define $\theta_n, \theta \in \text{Hom}(F_r, \mathbb{M})$ to be

$$\theta_n(i) = \rho_{r+1n} \circ \rho_{in}, \quad \theta(i) = \rho_{r+1} \circ \rho_i \quad \text{for every } i \text{ in } \{1, \dots, r\},$$

respectively, where ρ_{jn} and ρ_j are the reflections in C_{jn} and C_j on $\widehat{\mathbb{C}}$, respectively ($j = 1, \dots, r+1$). It is shown that $\{\theta_n\}_{n=1}^{+\infty}$ is contained in \mathbb{S}_r^0 and converges to θ as n tends to $+\infty$. If $\varphi \in \text{Im } \theta$ is parabolic, then there exist $k, l \in \{1, \dots, r+1\}$ such that φ and $\rho_k \circ \rho_l$ are conjugate in the group generated by $\rho_1, \dots, \rho_{r+1}$ (in this case, C_k and C_l are tangent). Since the composite of two reflections in two mutually disjoint circles is hyperbolic, $\lambda(\theta_n \circ \theta^{-1}(\varphi))$ is real for every n . Therefore, $\lambda(\theta_n \circ \theta^{-1}(\varphi))$ converges to 1 conically as n tends to $+\infty$: this is a special case of our theorem.

We notice the following:

Lemma 1 (Marden [4, Lemma 4.1]). *Suppose that G is a Schottky group and that u, v and w are three distinct limit points of G . Fix a region R as in the above definition of a Schottky group. Then there exists one and only one $\varphi \in G$ such that u, v and w belong to three distinct components of $\widehat{\mathbb{C}} - \varphi(R)$.*

In order to prove our theorem, we will prove the following lemma.

Lemma 2. *Let G be a classical Schottky group. Suppose that f and g belong to G and have no common fixed points. Let u, v and w be the repulsive fixed point of f , the attractive fixed point of f and the attractive fixed point of g , respectively. Then there exist two closed disks P and Q in $\widehat{\mathbb{C}}$ which have the following properties:*

- (1) P and Q contain u and w , respectively and they do not intersect each other.
- (2) $f(P)$ contains P and Q and it does not contain v .
- (3) Q contains at least one of $g(u)$ and $g(v)$.

Proof. Let r be the rank of G . Suppose that R is a region as in the above definition of a Schottky group. Since G is classical, we may assume that every component of ∂R is a circle. Note that u, v and w are limit points of G . By Lemma 1, there exists $\varphi \in G$ such that u, v and w belong to three distinct components of $\widehat{\mathbb{C}} - \varphi(R)$. Let U, V and W be components of $\widehat{\mathbb{C}} - \varphi(R)$ which contain u, v and w , respectively. By

the definitions of U and V , we can show that $f(U)$ contains $\widehat{\mathbb{C}} - V$ and does not contain v . In particular, $f(U)$ contains U and W . If the repulsive fixed point of g does not belong to U (or V), then $g(u)$ (or $g(v)$) belongs to W . Thus we can put $P = U$ and $Q = W$. \square

3. Proof of Theorem

Choose a loxodromic transformation ψ of $\text{Im } \theta$ which does not fix the fixed point of φ . We define $\varphi_n = \theta_n \circ \theta^{-1}(\varphi)$ and $\psi_n = \theta_n \circ \theta^{-1}(\psi)$ for each n . Note that φ_n and ψ_n have no common fixed points. Let p_n and q_n be the repulsive fixed point of φ_n and the attractive fixed point of φ_n , respectively. We write k_n for $\lambda(\varphi_n)$. Clearly, k_n converges to 1.

Choose an element γ of \mathbb{M} such that $\gamma \circ \varphi \circ \gamma^{-1}(z) = z/(z+1)$. Both $\gamma(p_n)$ and $\gamma(q_n)$ converge to 0 as n tends to $+\infty$. We assume that n is sufficiently large such that neither $\gamma(p_n)$ nor $\gamma(q_n)$ is ∞ . For each n , define $\gamma_n \in \mathbb{M}$ to be

$$\gamma_n(z) = \frac{1 - k_n}{\gamma(p_n) - \gamma(q_n)}(\gamma(z) - \gamma(q_n)).$$

We write

$$\gamma \circ \varphi_n \circ \gamma^{-1}(z) = \frac{a_n z + b_n}{c_n z + d_n}, \quad (a_n d_n - b_n c_n = 1),$$

for each n . Note that $c_n \neq 0$ and that c_n^2 converges to 1. Since $\gamma(p_n)$ and $\gamma(q_n)$ are the solutions of the quadratic equation $c_n x^2 - (a_n - d_n)x - b_n = 0$,

$$(\gamma(p_n) - \gamma(q_n))^2 = (\gamma(p_n) + \gamma(q_n))^2 - 4\gamma(p_n)\gamma(q_n) = \frac{(a_n + d_n)^2 - 4}{c_n^2}.$$

Using $(a_n + d_n)^2 = k_n + k_n^{-1} + 2$, we have

$$(\gamma(p_n) - \gamma(q_n))^2 = \frac{(k_n - 1)^2}{k_n c_n^2}.$$

Since both k_n and c_n^2 converge to 1,

$$\left(\frac{1 - k_n}{\gamma(p_n) - \gamma(q_n)} \right)^2 = k_n c_n^2 \longrightarrow 1 \quad (n \longrightarrow +\infty).$$

Thus γ_n converges to γ , or some subsequence of $\{\gamma_n\}$ converges to $-\gamma$, where $(-\gamma)(z) = -(\gamma(z))$. Considering fixed points and multipliers, we can show $\gamma_n \circ \varphi_n \circ \gamma_n^{-1}(z) = z/(z+k_n)$. Since $\gamma \circ \varphi \circ \gamma^{-1}(z) = z/(z+1)$ and k_n converges to 1, γ_n converges to γ as n tends to $+\infty$.

Let $\sigma \in \mathbb{M}$ map z to $1/z$. Define f_n and f to be

$$f_n = \sigma \circ \gamma_n \circ \varphi_n \circ \gamma_n^{-1} \circ \sigma^{-1} \text{ and } f = \sigma \circ \gamma \circ \varphi \circ \gamma^{-1} \circ \sigma^{-1},$$

respectively. Then $f_n(z) = k_n z + 1$ and $f(z) = z + 1$. Note that $1/(1 - k_n)$ is the repulsive fixed point of f_n . Define g_n and g to be

$$g_n = \sigma \circ \gamma_n \circ \psi_n \circ \gamma_n^{-1} \circ \sigma^{-1} \text{ and } g = \sigma \circ \gamma \circ \psi \circ \gamma^{-1} \circ \sigma^{-1},$$

respectively. Clearly, g_n converges to g as n tends to $+\infty$. Let w_n and w be the attractive fixed points of g_n and g , respectively. Note that neither w_n nor w is ∞ . By Lemma 2, there exist two closed disks P_n and Q_n in $\widehat{\mathbb{C}}$ for each n which have the following properties:

- (1) P_n and Q_n contain $1/(1 - k_n)$ and w_n , respectively and they do not intersect each other.
- (2) $f_n(P_n)$ contains P_n and Q_n and it does not contain ∞ .
- (3) Q_n contains at least one of $g_n(\infty)$ and $g_n(1/(1 - k_n))$.

From (2), both P_n and Q_n are contained in \mathbb{C} . We put

$$P_n = \{z \in \mathbb{C} \mid |z - \alpha_n| \leq \rho_n\}.$$

We easily obtain

$$f_n(P_n) = \{z \in \mathbb{C} \mid |z - (k_n \alpha_n + 1)| \leq \rho_n |k_n|\}.$$

From $P_n \subset f_n(P_n)$, we deduce that

$$|\alpha_n(k_n - 1) + 1| \leq \rho_n(|k_n| - 1).$$

Let l_n be the ray which has α_n as its initial point and which passes through the center (in the Euclidean sense) of Q_n . Suppose that l_n crosses ∂P_n at u'_n , ∂Q_n at u_n and v_n , and $f_n(\partial P_n)$ at v'_n (u_n lies between u'_n and v_n). Under this condition,

$$|u_n - v_n| \leq |u'_n - v'_n| = |v'_n - \alpha_n| - \rho_n \leq |\alpha_n(k_n - 1) + 1| + \rho_n |k_n| - \rho_n.$$

Using $|\alpha_n(k_n - 1) + 1| \leq \rho_n(|k_n| - 1)$, we have

$$|u_n - v_n| \leq 2\rho_n(|k_n| - 1).$$

We assume that n is sufficiently large such that the following inequalities are satisfied:

$$\begin{aligned} |w - w_n| &< \frac{|w - g(\infty)|}{4}, \\ |g(\infty) - g_n(\infty)| &< \frac{|w - g(\infty)|}{4}. \end{aligned}$$

$$\left| g(\infty) - g_n \left(\frac{1}{1 - k_n} \right) \right| < \frac{|w - g(\infty)|}{4}.$$

From these inequalities, we obtain

$$\begin{aligned} \frac{|w - g(\infty)|}{2} &< |w_n - g_n(\infty)|, \\ \frac{|w - g(\infty)|}{2} &< \left| w_n - g_n \left(\frac{1}{1 - k_n} \right) \right|. \end{aligned}$$

Since Q_n contains w_n and at least one of $g_n(\infty)$ and $g_n(1/(1 - k_n))$, and $|u_n - v_n|$ is the diameter (in the Euclidean sense) of Q_n ,

$$\frac{|w - g(\infty)|}{2} < |u_n - v_n|.$$

Since $|u_n - v_n| \leq 2\rho_n(|k_n| - 1)$,

$$|w - g(\infty)| < 4\rho_n(|k_n| - 1).$$

Using this inequality and $|\alpha_n(k_n - 1) + 1| \leq \rho_n(|k_n| - 1)$, we have

$$\begin{aligned} 1 &\geq \frac{|\alpha_n(k_n - 1) + 1|}{\rho_n(|k_n| - 1)} \\ &\geq \frac{|\alpha_n|}{\rho_n} \frac{|k_n| - 1}{|k_n| - 1} - \frac{1}{\rho_n(|k_n| - 1)} \\ &> \frac{|\alpha_n|}{\rho_n} \frac{|k_n| - 1}{|k_n| - 1} - \frac{4}{|w - g(\infty)|}. \end{aligned}$$

Since w_n does not belong to P_n ,

$$1 < \frac{|w_n - \alpha_n|}{\rho_n} \leq \frac{|w_n|}{\rho_n} + \frac{|\alpha_n|}{\rho_n} < \frac{4|w_n|(|k_n| - 1)}{|w - g(\infty)|} + \frac{|\alpha_n|}{\rho_n}.$$

Since $|w_n|(|k_n| - 1)$ converges to 0 as n tends to $+\infty$, $|\alpha_n|/\rho_n$ is greater than $1/2$ for sufficiently large n . Therefore,

$$\frac{|k_n - 1|}{|k_n| - 1} < \frac{\rho_n}{|\alpha_n|} \left(1 + \frac{4}{|w - g(\infty)|} \right) < 2 \left(1 + \frac{4}{|w - g(\infty)|} \right)$$

for sufficiently large n . This completes the proof.

4. Convergence of critical exponents

Let B^3 be the unit ball model of three-dimensional hyperbolic space, and let ∂B^3 be the sphere at infinity of B^3 . \mathbb{M} acts naturally on both of B^3 and ∂B^3 . A discrete

subgroup of \mathbb{M} acts on B^3 discontinuously. A discrete subgroup of \mathbb{M} is called *geometrically finite* if there exists a finite-sided fundamental polyhedron for its action on B^3 and *geometrically infinite* otherwise. A Schottky group is geometrically finite.

Let G be a discrete subgroup of \mathbb{M} . Define the *critical exponent* $\delta(G)$ of G to be

$$\delta(G) = \inf \left\{ \alpha \geq 0 \mid \sum_{g \in G} \exp(-\alpha \rho(\mathbf{o}, g(\mathbf{o}))) < +\infty \right\},$$

where $\mathbf{o} = (0, 0, 0)$ and $\rho(\mathbf{o}, g(\mathbf{o}))$ is the hyperbolic distance between \mathbf{o} and $g(\mathbf{o})$. Furthermore, suppose that G is geometrically finite. Then, there exists one and only one Borel probability measure μ on ∂B^3 such that it is supported on the limit set of G and that for every g in G and every Borel subset E of ∂B^3 , the following equality holds:

$$\mu(g(E)) = \int_E |g'(x)|^{\delta(G)} d\mu(x),$$

where $|g'(x)|$ is the linear distortion of g at x in the spherical metric on ∂B^3 (Sullivan [8, Theorem 1]). We call this μ the *Patterson-Sullivan measure of G* .

Let r be an integer greater than one. For every θ in $\partial \mathbb{S}_r$, $\text{Im } \theta$ is discrete (Marden [4, Lemma 2.2]). Using McMullen [7, Theorem 7.3], we obtain the following:

Proposition. *Suppose that a sequence $\{\theta_n\}_{n=1}^{+\infty}$ in \mathbb{S}_r converges to a cusp θ as n tends to $+\infty$ and that $\text{Im } \theta$ is geometrically finite. If for each parabolic transformation φ of $\text{Im } \theta$, $\lambda(\theta_n \circ \theta^{-1}(\varphi))$ converges to 1 conically as n tends to $+\infty$, then*

- (1) $\text{Im } \theta_n$ converges to $\text{Im } \theta$ geometrically;
- (2) the limit set of $\text{Im } \theta_n$ converges to the limit set of $\text{Im } \theta$ in the sense of Hausdorff convergence;
- (3) the Patterson-Sullivan measure of $\text{Im } \theta_n$ converges to the measure of $\text{Im } \theta$ weakly;
- (4) the critical exponent of $\text{Im } \theta_n$ converges to the critical exponent of $\text{Im } \theta$, as n tends to $+\infty$.

For every θ in $\partial \mathbb{S}_r \cap \overline{\mathbb{S}_r^0}$, $\text{Im } \theta$ is geometrically finite (Jørgensen, Marden and Maskit [3]). Hence from this we obtain the corollary stated in the introduction.

Finally, we will show that \mathbb{S}_r^0 in our theorem cannot be replaced with \mathbb{S}_r . If $\theta \in \partial \mathbb{S}_r$ is not a cusp, then $\text{Im } \theta$ is geometrically infinite. Using Mostow rigidity, we can prove this claim (see, for example, Matsuzaki and Taniguchi [5, Theorem 4.25]). If a sequence $\{\eta_m\}_{m=1}^{+\infty}$ in \mathbb{S}_r converges to η and if $\text{Im } \eta$ is geometrically infinite, then $\delta(\text{Im } \eta_m)$ converges to 2 as m tends to $+\infty$ (Bishop and Jones [1, Theorem 6.2]). It is essentially proved in Chuckrow [2] that $\partial \mathbb{S}_r$ removed all cusps is dense in $\partial \mathbb{S}_r$. Consequently, by diagonal method, for each θ in $\partial \mathbb{S}_r$, there exists a sequence $\{\theta_n\}_{n=1}^{+\infty}$ in \mathbb{S}_r such that θ_n converges to θ and $\delta(\text{Im } \theta_n)$ converges to 2 as n tends to $+\infty$. On the other hand, if a discrete subgroup G of \mathbb{M} is geometrically finite and if the limit

set of G does not coincide with $\widehat{\mathbb{C}}$, then $\delta(G)$ is less than 2 (Sullivan [8, Theorem 1]). Therefore, \mathbb{S}_r^0 in our theorem cannot be replaced with \mathbb{S}_r even if θ belongs to $\partial\mathbb{S}_r \cap \overline{\mathbb{S}_r^0}$.

References

- [1] C.J. Bishop and P.W. Jones: *Hausdorff dimension and Kleinian groups*, Acta Math. **179** (1997), 1–39.
- [2] V. Chuckrow: *On Schottky groups with applications to kleinian groups*, Ann. of Math. **88** (1968), 47–61.
- [3] T. Jørgensen, A. Marden and B. Maskit: *The boundary of classical Schottky space*, Duke Math. J. **46** (1979), 441–446.
- [4] A. Marden: *Schottky groups and circles*, Contributions to Analysis, Academic Press, New York and London, 273–278, 1974.
- [5] K. Matsuzaki and M. Taniguchi: *Hyperbolic Manifolds and Kleinian Groups*, Oxford Univ. Press, Oxford, 1998.
- [6] C.T. McMullen: *Hausdorff dimension and conformal dynamics III: Computation of dimension*, Amer. J. Math. **120** (1998), 691–721.
- [7] C.T. McMullen: *Hausdorff dimension and conformal dynamics I: Strong convergence of Kleinian groups*, J. Diff. Geom. **51** (1999), 471–515.
- [8] D. Sullivan: *Entropy, Hausdorff measures old and new, and limit sets of geometrically finite Kleinian groups*, Acta Math. **153** (1984), 259–277.

Graduate School of Science and Engineering
Shizuoka University
836 Ohya, Shizuoka 422–8529, Japan