

Sadek, M. and El-Sissi, N.
Osaka J. Math.
52 (2015), 515–525

PARTITIONS WITH EQUAL PRODUCTS AND ELLIPTIC CURVES

MOHAMMAD SADEK and NERMINE EL-SISSI

(Received February 27, 2013, revised January 21, 2014)

Abstract

Let a, b, c be distinct positive integers. Set $M = a + b + c$ and $N = abc$. We give an explicit description of the Mordell–Weil group of the elliptic curve $E_{(M,N)}: y^2 - Mxy - Ny = x^3$ over \mathbb{Q} . In particular we determine the torsion subgroup of $E_{(M,N)}(\mathbb{Q})$ and show that its rank is positive. Furthermore there are infinitely many positive integers M that can be written in n different ways, $n \in \{2, 3\}$, as the sum of three distinct positive integers with the same product N and $E_{(M,N)}(\mathbb{Q})$ has rank at least n .

1. Introduction

The subject of partitioning integers has been used to construct infinite families of elliptic curves with positive rank. Many authors attack questions linking partitioning integers and elliptic curves arising from these integers. For example in [1] the elliptic curve $E_n: y^2 = x^3 - nx$ where $n = a^4 + b^4$ is proved to be of rank at least 2 over $\mathbb{Q}(a, b)$. If n can be written as a sum of two biquadrates in two different ways, then [4] indicates the existence of an infinite number of integers n such that $E_n(\mathbb{Q})$ is of rank at least 3, and this lower bound is improved in [1] to be 4.

In this note we study the partitions of a positive integer into three positive integers with the same product. A triple of positive integers (x, y, z) is said to be a *partition* of a positive integer M if $M = x + y + z$. The integers x, y, z are the *parts* of the partition. We set $N = xyz$. If we are looking for all integer triples (x, y, z) with sum M and product N , then we are trying to solve two Diophantine equations. Eliminating z will yield the elliptic curve

$$E_{(M,N)}: y^2 - Mxy - Ny = x^3.$$

In [5], the elliptic curve $E_{(M,N)}$ was shown to be of positive rank over \mathbb{Q} under the mild condition that if $M = d_1 + d_2 + d_3$, $N = d_1d_2d_3$ where $d_1 > d_2 > d_3$ then $d_1(d_2 - d_3)^3 \neq d_3(d_1 - d_2)^3$. We reprove this fact and give an explicit description of the torsion part $E_{(M,N)}^{\text{tor}}(\mathbb{Q})$ of the Mordell–Weil group $E_{(M,N)}(\mathbb{Q})$. More precisely we show that $E_{(M,N)}^{\text{tor}}(\mathbb{Q})$ is one of the following subgroups $\mathbb{Z}/3\mathbb{Z}$, $\mathbb{Z}/6\mathbb{Z}$, or $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$.

If the curve $E_{(M,N)}$ has positive rank over \mathbb{Q} , then this means that there is an infinite number of triples of nonzero rational numbers adding up to M and having product N . We investigate the rank of this elliptic curve if M can be written as a sum of three nonzero positive integers with product N in at least two different ways.

We parametrize positive integers M that have two different partitions into triples with product N . The parametrization of M is given in terms of four parameters p, q, r, s . Consequently we show that $E_{(M,N)}$ has rank at least 2 over $\mathbb{Q}(p, q, r, s)$. Thus there exists an infinite number of pairs of integers (M, N) such that $E_{(M,N)}(\mathbb{Q})$ has rank at least 2.

A weaker result is presented when M has three different partitions into triples with product N . An infinite parametric family of such pairs (M, N) is constructed. The parametrization depends on three parameters p, q, r . We prove that $E_{(M,N)}$ has rank at least 3 over $\mathbb{Q}(p, q, r)$, and hence the existence of an infinite number of pairs of integers (M, N) for which $E_{(M,N)}(\mathbb{Q})$ has rank at least 3.

In this note, a partitioning question is used to exhibit infinite families of elliptic curves with positive rank. This partitioning question corresponds to an interesting geometric problem. Namely, if (M, N) is a pair of positive integers such that $4M$ is the perimeter of a rectangular box \mathcal{R} with integer side lengths, and N is the volume of \mathcal{R} , then how many different rectangular boxes with integer sides have the same perimeter and volume? Indeed, that $\text{rank } E_{(M,N)}(\mathbb{Q}) \geq 1$ means that there are an infinite number of rectangular boxes with rational sides, and the same perimeter and volume. We parametrize the pairs (M, N) for which there exist two rectangular boxes with integer sides, perimeter $4M$, and volume N . Moreover we give infinite parametric pairs (M, N) for which there exists three rectangular boxes with integer sides, perimeter $4M$, and volume N . These pairs (M, N) give rise to elliptic curves of rank at least 2 and 3 respectively.

All calculations throughout this note were performed using Sage [8], and Mathematica [6]. We would like to thank the anonymous referee for several comments and suggestions.

2. Partitions

In this section we collect elementary properties about partitions with equal products.

Lemma 2.1. *Let M be a positive integer that has at least two distinct partitions into triples with equal product N . The following statements are true:*

- a) *There is no common entry between any of the triples. In particular, $N \notin \{p, pq\}$ where p, q are primes.*
- b) *N is not a prime power.*
- c) *N is a product of at least four (not necessarily distinct) primes.*

Proof. a) Assume that $M = a + b + c = a + d + e$ and $abc = ade$. One has $b + c = d + (bc/d)$, i.e., $bd + cd = d^2 + bc$ or $b(d - c) = d(d - c)$, in other words, $b = d$ or $d = c$. In both cases, this contradicts the fact that the partitions are distinct.

b) If $N = p^r$, then $M = p^{r_1} + p^{r_2} + p^{r_3} = p^{s_1} + p^{s_2} + p^{s_3}$ where $r_1 \geq r_2 \geq r_3$, $s_1 \geq s_2 \geq s_3$. Dividing by $\min(p^{r_3}, p^{s_3})$, one sum will be divisible by p while the other is not.

c) Using parts a) and b), it is enough to show that N cannot be a product of exactly three primes. So we assume on the contrary that $N = pqr$ where $p \geq q \geq r$ are primes. According to a) the only possible partitions for M are given by $M = 1 + 1 + pqr = p + q + r$. The contradiction follows from $pqr \geq p^3 > 3p > p + q + r$. \square

Let M, N, x, y, z be nonzero integers satisfying the following relations:

$$x + y + z = M \quad \text{and} \quad xyz = N.$$

These two equations are equivalent to the following cubic equation $Mxy - x^2y - xy^2 = N$. We homogenize the above cubic equation to obtain the following equation

$$NZ^3 + XY^2 + X^2Y - MXYZ = 0$$

describing a planar curve $C_{(M,N)}$ in $\mathbb{P}_{\mathbb{Q}}^2$ with $(X : Y : Z) = (0 : 1 : 0) \in C_{(M,N)}(\mathbb{Q})$. Therefore given that $\Delta = N^3(M^3 - 27N) \neq 0$, the Jacobian $E_{(M,N)} := \text{Jac}(C_{(M,N)})$ of the planar curve $C_{(M,N)}$ is an elliptic curve defined by the following Weierstrass equation.

$$E_{(M,N)}: Y^2 - MXY - NY = X^3.$$

In fact, $C_{(M,N)}$ is isomorphic to $E_{(M,N)}$ via the following transformation:

$$\begin{aligned} \alpha_{(M,N)}: C_{(M,N)} &\xrightarrow{\cong} E_{(M,N)}, \\ (X : Y : Z) &\mapsto (-NZ : -NY : X). \end{aligned}$$

An ordered triple of nonzero integers (d_1, d_2, d_3) such that $d_1 + d_2 + d_3 = M$ and $d_1d_2d_3 = N$ is sent to a point in $C_{(M,N)}(\mathbb{Q})$ and hence a point in $E_{(M,N)}(\mathbb{Q})$.

We will not treat triples of the form (d, d, d) . The reason is that the corresponding cubic curve $y^2 - 3dxy - d^3y = x^3$ is singular. Therefore we are going to assume throughout that the pairs $(M, N) = (3d, d^3)$ are excluded.

From now on we will assume that if $N = d^3a$, where a is cube-free, then $\gcd(M, d) = 1$. Otherwise the Weierstrass equation describing $E_{(M,N)}$ is not minimal. Moreover this allows us to assume that the parts of each partition of M with product N are coprime.

Corollary 2.2. *Let M, N be nonzero integers such that $N^3(M^3 - 27N) \neq 0$. There is a one-to-one correspondence between the set of ordered triples*

$$\Gamma_{(M,N)}(\mathbb{Q}) = \{(d_1, d_2, d_3) : d_1 + d_2 + d_3 = M, d_1 d_2 d_3 = N, d_i \in \mathbb{Q}\}$$

and

$$E_{(M,N)}(\mathbb{Q}) = \{(x : y : z) \in \mathbb{P}^2(\mathbb{Q}) : y^2 z - Mxyz - Nyz^2 = x^3, xyz \neq 0\}.$$

Proof. The bijection map is as follows:

$$\begin{aligned} \Gamma_{(M,N)}(\mathbb{Q}) &\rightarrow C_{(M,N)}(\mathbb{Q}) \rightarrow E_{(M,N)}(\mathbb{Q}), \\ (d_1, d_2, d_3) &\mapsto (d_1 : d_2 : 1) \mapsto (-N : -Nd_2 : d_1), \\ (-Nc/a, b/a, -a^2/bc) &\leftarrow (Nc : -b : -a) \leftarrow (a : b : c). \end{aligned}$$

We restrict x, y, z to be non-zero since we divide by them in the inverse map. \square

Before we proceed with investigating the torsion subgroup of the Mordell–Weil group of $E_{(M,N)}(\mathbb{Q})$, we need to recall the classification of torsion points of elliptic curves over \mathbb{Q} , see (Chapter VIII, §8, Theorem 7.5) in [7].

Lemma 2.3. *Let E be an elliptic curve over \mathbb{Q} . Then the torsion subgroup of $E(\mathbb{Q})$ is one of the following fifteen groups:*

$$\mathbb{Z}/N\mathbb{Z}, \quad 1 \leq N \leq 10 \quad \text{or} \quad N = 12; \quad \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2N\mathbb{Z}, \quad 1 \leq N \leq 4.$$

Lemma 2.4. *Let $E_{(M,N)}$ be the elliptic curve described above. One has $\mathbb{Z}/3\mathbb{Z} \subseteq E_{(M,N)}^{\text{tor}}(\mathbb{Q})$.*

Proof. The point $(0 : 0 : 1)$ is a rational point on

$$E_{(M,N)} : y^2 z - Mxyz - Nyz^2 = x^3.$$

Indeed the subgroup generated by $(0 : 0 : 1)$ is $\{(0 : 0 : 1), (0 : N : 1), O\}$. \square

We define the following set $S_{(M,N)}$ of classes of triples of nonzero rational numbers as follows:

$$S_{(M,N)} = \{(a, a, b) : a \neq b, 2a + b = M, a^2 b = N\} / \sim$$

where $(x_1, x_2, x_3) \sim (y_1, y_2, y_3)$ if and only if $x_i = y_j$ for some i, j .

Proposition 2.5. *Let a, b be nonzero rational numbers such that $2a + b = M$ and $a^2 b = N$. Then the point $(-N : -Na : a)$ is a torsion point of order 6 in $E_{(M,N)}(\mathbb{Q})$. In particular, if $\#S_{(M,N)} = 2$, then $E_{(M,N)}^{\text{tor}}(\mathbb{Q}) = \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$.*

Proof. This is direct calculation using the formulas for addition on elliptic curves, see p. 58 in [7]. More precisely, the subgroup generated by the point $(-ab, -a^2b)$ is:

$$\langle(-ab, -a^2b)\rangle = \{(-ab, -a^2b), (0, 0), (-a^2, -a^3), (0, a^2b), (-ab, -ab^2), O\}.$$

We observe that $(-a^2, -a^3)$ is the image of the partition (b, a, a) and $(-ab, -ab^2)$ is the image of the partition (a, b, a) in $E_{(M,N)}(\mathbb{Q})$. □

REMARK 2.6. In Proposition 2.5, the case $\#S_{(M,N)} > 2$ cannot occur, since this will contradict the fact that the number of torsion points on $E_{(M,N)}$ must be less than 16, see Lemma 2.3. This implies that no positive integer M can be written in more than two different ways as $M = x_1 + 2x_2$ where $x_1x_2^2 = N$ and $x_1, x_2 > 0$.

Proposition 2.7. *Let M, N be integers such that M can be written as a sum of three distinct positive rational numbers $d_1 > d_2 > d_3$ whose product is N . We assume that $N^3(M^3 - 27N) \neq 0$. Assume moreover that $d_1(d_2 - d_3)^3 \neq d_3(d_1 - d_2)^3$. Then $\text{rank } E_{(M,N)}(\mathbb{Q}) \geq 1$.*

Proof. We recall that $\Delta = N^3(M^3 - 27N)$ is the discriminant of $E_{(M,N)}$. Hence $\Delta \neq 0$ implies that $E_{(M,N)}$ is an elliptic curve.

We observe that the six rational points

$$P_{ij} = (-d_i d_j, -d_i d_j^2) \in E_{(M,N)}(\mathbb{Q}), \quad i \neq j,$$

satisfy the following identities:

$$P_{ij} + P_{ji} = O, \quad P_{ij} + P_{ik} = (0, 0), \quad j \neq k,$$

$$P_{ij} + P_{kj} = (0, N), \quad i \neq k, \quad 2P_{ij} = \left(\frac{d_i d_j (d_i - d_k)(d_j - d_k)}{(d_i - d_j)^2}, \frac{d_i d_j^2 (d_i - d_k)^3}{(d_i - d_j)^3} \right).$$

We claim that P_{ij} is of infinite order for every i, j . Assume on the contrary that P_{ij} is of finite order. First of all we see that P_{ij} is not a 2-torsion, since otherwise $P_{ij} = -P_{ij} = P_{ji}$ and $d_i d_j^2 = d_j^2 d_i$ which implies $d_i = d_j$, a contradiction. Therefore we have 12 points of finite order $P_{ij}, 2P_{ij}, i \neq j$. Moreover, the points P_{ij} and $2P_{ij}$ are distinct. For the latter statement, it is easy to show that the six points P_{ij} are distinct. However, if $P_{ij} \in 2S$ where $S = \{P_{ij}, 1 \leq i, j \leq 3\}$, then we have one of the following possibilities: if $P_{ij} = 2P_{ki}$ then $P_{ki} = (0, 0)$; if $P_{ij} = 2P_{jk}$ then $P_{jk} = (0, N)$; if $P_{ij} = 2P_{ji}$ then $3P_{ij} = (0, 0)$; if $P_{ij} = 2P_{ik}$ then $-d_j = d_k(d_i - d_j)(d_k - d_j)/(d_i - d_k)^2$ and observing the signs of both sides of the equality implies $(i, j, k) \in \{(1, 2, 3), (3, 2, 1)\}$, moreover $-d_j^2 = d_k^2(d_i - d_j)^3/(d_i - d_k)^3$ but this implies $(i, j, k) \in \{(2, 1, 3), (2, 3, 1)\}$, a contradiction; if $P_{ij} = 2P_{kj}$, then $-d_i = d_k(d_k - d_i)(d_j - d_i)/(d_k - d_j)^2$ hence $(i, j, k) \in \{(2, 3, 1), (2, 1, 3)\}$, moreover $-d_i = d_k(d_k - d_i)^3/(d_k - d_j)^3$ but this implies $(i, j, k) \in \{(3, 1, 2), (1, 3, 2)\}$, a contradiction.

Now if $2P_{ij} = 2P_{ki}$ then $2(P_{ij} + P_{ik}) = O$, a contradiction; if $2P_{ij} = 2P_{jk}$ then $2(P_{ij} + P_{kj}) = O$; if $2P_{ij} \in \{2P_{ji}, 2P_{ik}, 2P_{kj}\}$ then $d_1(d_2 - d_3)^3 = d_3(d_1 - d_2)^3$ which is ruled out by our assumption. None of these 12 points is a point of the subgroup $\{O, (0, 0), (0, N)\}$. Therefore we have fifteen points of finite order, three of them make up a subgroup of order three. According to the classification of torsion points on elliptic curves, Lemma 2.3, this is a contradiction. \square

Proposition 2.7 provides an easy method to construct elliptic curves with positive rank.

Theorem 2.8. *Let M, N be integers such that M can be written as a sum of three positive rational numbers $d_1 > d_2 > d_3$ whose product is N . Assume moreover that $N^3(M^3 - 27N) \neq 0$ and $d_1(d_2 - d_3)^3 \neq d_3(d_1 - d_2)^3$. The Mordell–Weil group of the elliptic curve $E_{(M,N)}: y^2 - Mxy - Ny = x^3$ satisfies $E_{(M,N)}(\mathbb{Q}) \cong \mathbb{Z}^r \times E_{(M,N)}^{\text{tor}}(\mathbb{Q})$, $r \geq 1$, where*

$$E_{(M,N)}^{\text{tor}}(\mathbb{Q}) \cong \begin{cases} \mathbb{Z}/3\mathbb{Z} & \text{if } \#S_{(M,N)} = 0, \\ \mathbb{Z}/6\mathbb{Z} & \text{if } \#S_{(M,N)} = 1, \\ \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z} & \text{if } \#S_{(M,N)} = 2. \end{cases}$$

Proof. We recall that there is a bijection between ordered triples of nonzero rational numbers adding up to M and having product N , and rational points on $E_{(M,N)}$, see Corollary 2.2. Moreover Proposition 2.7 implies that if $d_1 + d_2 + d_3 = M$ and $d_1d_2d_3 = N$ where d_1, d_2, d_3 are distinct nonzero rational numbers, then $(-d_id_j, -d_id_j^2)$ is a point of infinite order in $E_{(M,N)}(\mathbb{Q})$. Therefore according to Corollary 2.2 and Proposition 2.7, the only points of finite order $(x : y : z)$ are the ones corresponding to triples of the form (a, a, b) , $a \neq b$, where $2a + b = M$ and $a^2b = N$, or the ones with at least one of the entries being zero. In other words, the finite points are lying either in $S_{(M,N)}$ or the subgroup $\{O, (0, 0), (0, N)\}$ respectively. Now the statement of the theorem follows from Lemma 2.4 and Proposition 2.5. \square

3. A family of elliptic curves with rank at least 2

In Theorem 2.8, we proved that if an integer M is the sum of three distinct positive rational numbers whose product is N , then there is a corresponding elliptic curve $E_{(M,N)}$ of positive rank. In what follows, we study the arithmetic of these elliptic curves and introduce further conditions on the given integers to increase the rank of the corresponding elliptic curves. For this purpose, we use positive integers which have more than one partition into three distinct parts such that the product of these parts are equal in each partition.

We consider the following system of Diophantine equations:

$$(1) \quad \begin{aligned} X + Y + Z &= U + V + W, \\ XYZ &= UVW. \end{aligned}$$

In §2 of [2], a complete solution of the following system is found.

$$(2) \quad \begin{aligned} x + y + z &= u + v + w, \\ x^3 + y^3 + z^3 &= u^3 + v^3 + w^3. \end{aligned}$$

Using the following transformations

$$\begin{array}{l} x = -X + Y + Z, \quad u = -U + V + W, \\ y = X - Y + Z, \quad v = U - V + W, \\ z = X + Y - Z, \quad w = U + V - W, \end{array} \left| \begin{array}{l} X = \frac{y+z}{2}, \quad U = \frac{v+w}{2}, \\ Y = \frac{x+z}{2}, \quad V = \frac{u+w}{2}, \\ Z = \frac{x+y}{2}, \quad W = \frac{u+v}{2}, \end{array} \right.$$

we transform the latter system (2) of equations into the following system

$$(3) \quad X + Y + Z = U + V + W,$$

$$(4) \quad (X + Y + Z)^3 - 24XYZ = (U + V + W)^3 - 24UVW.$$

Therefore, equation (3) reduces equation (4) to $XYZ = UVW$.

We obtain a complete solution of system (1) using the transformations above and the complete solution of (2) found in Theorem 1 of [2]. The solution is given in terms of quadratic polynomials in four parameters such that each parameter appearing in the solution is of first degree, more explicitly, the solution of (1) is

$$(5) \quad \begin{aligned} X &= p(r + s), & Y &= q(p + s), & Z &= r(q + s), \\ U &= q(r + s), & V &= r(p + s), & W &= p(q + s) \end{aligned}$$

where p, q, r, s are parameters.

Theorem 3.1. *Let M and N be given by parametrization (5). Then the points $P = (-N : -NX : Y)$ and $Q = (-N : -NU : V)$ are two independent points in $E_{(M,N)}$ over $\mathbb{Q}(p, q, r, s)$. In particular, $E_{(M,N)}$ has rank at least 2 over $\mathbb{Q}(p, q, r, s)$.*

Proof. To show that $E_{(M,N)}$ has rank at least 2, we need to specialize p, q, r, s in the above parametrization so that the specialization of P, Q are independent over \mathbb{Q} . This holds because the specialization map is a homomorphism. Putting $p = 1$,

$q = 2$, $r = 3$, $s = 4$ yields $x = 7$, $y = 10$, $z = 18$, $u = 14$, $v = 15$, $w = 6$, and the points P , Q are specialized to $(-126, -882)$ and $(-84, -1176)$ on the elliptic curve $E_{(35,1260)}$. The height-pairing matrix associated to these points has non-zero determinant $1.70464760105805 \dots$. This means that P, Q are independent in $E_{(M,N)}$ over $\mathbb{Q}(p, q, r, s)$. \square

In the following example a different family of parametric pairs (M, N) given in terms of one parameter q is introduced such that $E_{(M,N)}$ has rank at least 2 over $\mathbb{Q}(q)$.

EXAMPLE 3.2. Let $k \geq 2$ be an integer. We define p as follows:

$$p = \frac{q^{2k-1} + 1}{q + 1}.$$

Now we define M_k as follows:

$$\begin{aligned} M_k &= 1 + pq^2 + q^{2k-1} \\ &= p + q + q^{2k} \end{aligned}$$

where the product of the parts in each of the above partitions of M_k is $N_k = pq^{2k+1}$. A specialization argument yields that the points $(-pq^{2k+1}, -p^2q^{2k+3})$, $(-q^{2k+1}, -q^{2k+2})$ are independent in $E_{(M_k, N_k)}$ over $\mathbb{Q}(q)$, and the curves $E_{(M_k, N_k)}$ have rank at least 2 over $\mathbb{Q}(q)$.

4. A family of elliptic curves with rank at least 3

As it has been illustrated, there is an infinite number of pairs of positive integers (M, N) , where M can be written in two different ways as a sum of three distinct positive integers with the same product N , and such that for each such pair the corresponding elliptic curve has rank at least two over \mathbb{Q} . The latter statement holds due to the fact that the specialization homomorphism is injective at infinitely many families of parameters. This suggests that if the number of partitions with the same product increases, then the rank of the corresponding elliptic curve might get larger.

We start by finding infinite number of integer solutions to the following Diophantine system:

$$\begin{aligned} (6) \quad & x_1 + y_1 + z_1 = x_2 + y_2 + z_2 = x_3 + y_3 + z_3, \\ & x_1 y_1 z_1 = x_2 y_2 z_2 = x_3 y_3 z_3. \end{aligned}$$

Given positive integers p, q, r, s , we are looking for pairs of positive integers (M, N) such that M has at least three partitions into three parts with equal product N and $pqr s \mid N$. More accurately we find w, z which make the triples (pw, qs, rz) ,

$(pqrw, s, z)$ and (w, qrs, pz) solutions to the system. It is an elementary linear algebra exercise to show that

$$w = \frac{qr^2 - qp - qr + p + q - r}{pqr^2 - p^2qr + p^2 - p - r + 1}s,$$

$$z = -\frac{p(1 - qr)w + (q - 1)s}{r - 1}.$$

Thus we can generate infinite number of integer solutions to the system (6) by clearing denominators. More precisely the following parametrizations solve system (6):

$$(7) \quad \begin{aligned} p, \quad q, \quad r, \\ s &= pqr^2 - p^2qr + p^2 - p - r + 1, \\ w &= qr^2 - qp - qr + p + q - r, \\ z &= pq^2r^2 - pq^2r - pqr + p + q - 1. \end{aligned}$$

Theorem 4.1. *Let M and N be given by parametrization (7). The points $P_1 = (-prwz, -p^2rw^2z)$, $P_2 = (-sz, -s^2z)$ and $P_3 = (-pwz, -pw^2z)$ are three independent points in $E_{(M,N)}$ over $\mathbb{Q}(p, q, r)$. In particular, $E_{(M,N)}$ has rank at least 3 over $\mathbb{Q}(p, q, r)$.*

Proof. By specializing $p = 2, q = 2, r = 3, s = 12, w = 9, z = 39$, we get the following partition of 159:

$$159 = 18 + 24 + 117 = 108 + 12 + 39 = 9 + 72 + 78.$$

The points P_i are specialized to $(-2106, -37908)$, $(-468, -5616)$ and $(-702, -6318)$ on the elliptic curve $y^2 - 159xy - 50544y = x^3$. The determinant of the height matrix associated to these points is $4.55758994382846 \dots$. This means that P_1, P_2, P_3 are independent. \square

We can obtain parametric solutions to the system (6) by solving two homogeneous linear equations, and consequently we reach an infinite family of positive integers that have three distinct partitions with the same product. These integers provide us with a family of elliptic curves of rank at least 3.

5. Elliptic curves with higher rank

The task of finding positive integers with large number of partitions into triples whose parts have equal products seems a hard problem. In fact the largest number of such partitions of a given integer up to the knowledge of the authors is 13. The integer 17116 has 13 different partitions with product $2^{10}3^35^27^211.13.19$, see D16 in [3].

We showed that positive integers that have two or three partitions with equal products can yield elliptic curves with rank at least two or three respectively. We may expect that the higher the number of partitions we can produce with equal products, the

higher the rank of the corresponding elliptic curve that we can construct. Therefore if we manage to find integers with arbitrary number of such partitions, then we should predict that the corresponding elliptic curves will have arbitrary large ranks.

For example the partitions (x_i, y_i, z_i) of $M = 17116$ with product $N = 2^{10} \cdot 3^3 \cdot 2 \cdot 5^2 \cdot 7^2 \cdot 11 \cdot 13 \cdot 19$ are:

(1512, 7700, 7904), (1520, 7280, 8316), (1540, 6840, 8736),
 (1596, 6160, 9360), (1716, 5320, 10080), (1755, 5120, 10241),
 (1760, 5096, 10260), (1792, 4950, 10374), (2016, 4180, 10920),
 (2128, 3900, 11088), (2200, 3744, 11172), (2548, 3168, 11400),
 (2736, 2940, 11440).

The elliptic curve

$$E: y^2 - 17116xy - 2^{10}3^35^27^211.13.19y = x^3$$

has Mordell–Weil group $E(\mathbb{Q}) \cong \mathbb{Z}/3\mathbb{Z} \times \mathbb{Z}^n$, $n \geq 6$, where the points $(-N : -Nx_i : y_i)$, $i \leq 6$, are independent.

References

- [1] J. Aguirre and J.C. Peral: *Sums of biquadrates and elliptic curves*, Glas. Mat. Ser. III **48** (2013), 49–58.
- [2] A. Choudhry: *Some Diophantine problems concerning equal sums of integers and their cubes*, Hardy–Ramanujan J. **33** (2010), 59–70.
- [3] R.K. Guy: *Unsolved Problems in Number Theory*, second edition, Springer, New York, 1994.
- [4] F.A. Izadi, F. Khoshnam and K. Nabardi: *Sums of two biquadrates and elliptic curves of rank ≥ 4* , Math. J. Okayama Univ. **56** (2014), 51–63.
- [5] J.B. Kelly: *Partitions with equal products*, II, Proc. Amer. Math. Soc. **107** (1989), 887–893.
- [6] Wolfram Research: *Mathematica Edition*, Version 5.0, Wolfram Research.
- [7] J.H. Silverman: *The Arithmetic of Elliptic Curves*, Graduate Texts in Mathematics **106**, Springer, New York, 1986.
- [8] W.A. Stein et al.: *Sage Mathematics Software (Version 4.3.5)*, The Sage Development Team, <http://www.sagemath.org>.

Mohammad Sadek
Department of Mathematics and Actuarial Science
American University in Cairo
Egypt
e-mail: mmsadek@aucegypt.edu

Nermine El-Sissi
Department of Mathematics and Actuarial Science
American University in Cairo
Egypt
e-mail: nelsissi@aucegypt.edu