

Ji, Q., Tan, G. and Yu, G.
Osaka J. Math.
52 (2015), 1–14

L^2 -ESTIMATES ON WEAKLY q -CONVEX DOMAINS

QINGCHUN JI, GUO TAN and GUANGSHENG YU

(Received June 6, 2013)

Abstract

We establish an estimate on weakly q -convex domains in \mathbb{C}^n which provides a unified approach to various existence results for the $\bar{\partial}$ -problem. We also prove a Diederich–Fornaess type result for weakly q -convex domains.

1. Introduction

Let $\Omega \subseteq \mathbb{C}^n$ be a pseudoconvex domain and let $\phi \in C^2(\Omega)$ be a strictly plurisubharmonic function. A variant of Hörmander’s theorem ([10]) states that for any $\bar{\partial}$ -closed $(0, 1)$ -form $f = f_j d\bar{z}_j \in L^2_{0,1}(\Omega, \text{loc})$ there exists a solution of $\bar{\partial}u = f$ satisfying

$$\int_{\Omega} |u|^2 e^{-\phi} \leq \int_{\Omega} |f|^2_{\sqrt{-1} \partial \bar{\partial} \phi} e^{-\phi}$$

where $|f|^2_{\sqrt{-1} \partial \bar{\partial} \phi} := \phi^{j\bar{k}} f_j \bar{f}_k$ and $(\phi^{j\bar{k}}) := (\phi_{j\bar{k}})^{-1}$. A geometric observation is that $\sqrt{-1} \partial \bar{\partial} \phi$ is the curvature form of the Hermitian metric $e^{-\phi}$ on the trivial line bundle. As proved in [9], the length of the $(0, 1)$ -form could be calculated w.r.t. another curvature form. The pointwise norm $|f|^2_{\sqrt{-1} \partial \bar{\partial} \psi}$ is used in [9] instead of $|f|^2_{\sqrt{-1} \partial \bar{\partial} \phi}$ where ψ is any strictly plurisubharmonic function such that $-e^{-\psi}$ is plurisubharmonic. The latter result was then further generalized to non-plurisubharmonic weights ([7], [8], [2], [3]), i.e., the curvature of the Hermitian metric on trivial bundle is not necessarily positive. Berndtsson–Blocki–Donnelly–Fefferman type results are closely related to the Ohsawa–Takegoshi extension theorem and Bergman metric (see [4], [5], [2], [8]).

We will consider, in the present paper, the $\bar{\partial}$ -problem on q -convex domains. We follow [11] in defining the notions of q -convexity and q -subharmonicity. We begin by recalling some basic notions and related preliminaries on exterior algebra. We prove a Diederich–Fornaess type result for weakly q -convex domains (Theorem 1). Let $\varphi \in C^\infty(\bar{\Omega})$ be a q -subharmonic function and let $\psi \in C^\infty(\bar{\Omega})$ be a function such that the real $(1, 1)$ -form $\delta \sqrt{-1} \partial \bar{\partial} \varphi - \sqrt{-1} \partial \psi \wedge \bar{\partial} \psi$ is q -positive semi-definite (see Definition 3)

for some constant $\delta \in [0, 4)$, we will establish the following a priori estimate

$$(*) \quad \|\bar{\partial}_{\varphi-(1/2)\psi}^* g\|_{\varphi}^2 + \|\bar{\partial} g\|_{\varphi}^2 \geq \frac{(2-\sqrt{\delta})^2}{4} \int_{\Omega} \langle F_{\varphi} g, g \rangle e^{-\varphi},$$

for any (p, q) -form $g \in \text{Dom}(\bar{\partial}^*) \cap C_{p,q}^{\infty}(\bar{\Omega})$ on weakly q -convex domains with smooth boundary. Here we have used the notation $F_{\varphi} = \varphi_{j\bar{k}} d\bar{z}_{\bar{k}} \wedge \partial/\partial\bar{z}_{j\bar{1}}$. When $\psi = 0$ we can choose $\delta = 0$, so $(*)$ generalizes Hörmander's estimate to q -convex domains and q -subharmonic weight functions. Actually, $(*)$ also implies the following Donnelly–Fefferman type estimate.

$$(**) \quad \|\bar{\partial}_{\varphi+\tau\psi}^* g\|_{\varphi+\psi}^2 + \|\bar{\partial} g\|_{\varphi+\psi}^2 \geq \tau^2 \int_{\Omega} \langle F_{\psi} g, g \rangle e^{-\varphi-\psi},$$

for any $g \in \text{Dom}(\bar{\partial}^*) \cap C_{p,q}^{\infty}(\bar{\Omega})$ where $\varphi \in C^{\infty}(\bar{\Omega})$ is a q -subharmonic function, $\psi \in C^{\infty}(\bar{\Omega})$ with $-e^{-\psi}$ being q -subharmonic and $\tau \in (0, 1/2]$ is a constant. This estimate implies an existence theorem of Berndtsson–Błocki–Donnelly–Fefferman type (see Corollary 2 below). This kind of theorems may help produce a desired curvature term without the contribution of the metric which has important applications (e.g., Ohsawa–Takegoshi type extension theorems). The curvature operator F_{φ} of a certain Hermitian metric will play an important role in our formulation of main results. Applications for p -convex Riemannian manifolds can be found in [12].

Here are the main results of the present paper:

Theorem 1. *Let $\Omega \Subset \mathbb{C}^n$ be a weakly q -convex domain with smooth boundary and let $r \in C^{\infty}(\bar{\Omega})$ be a defining function for Ω . Then for any strictly q -subharmonic function $\phi \in C^{\infty}(\bar{\Omega})$, there exist constants $K > 0$, $\eta_0 \in (0, 1)$ such that for any $\eta \in (0, \eta_0)$ the function $\rho := -(-re^{-K\phi})^{\eta}$ is strictly q -subharmonic on Ω .*

Theorem 2. *Let Ω be a weakly q -convex domain in \mathbb{C}^n ($1 \leq q \leq n$) and let $\varphi \in C^2(\Omega)$ be a q -subharmonic function on Ω and $\psi \in C^1(\Omega)$. Assume that the real $(1, 1)$ -form $\delta\sqrt{-1}\bar{\partial}\bar{\partial}\varphi - \sqrt{-1}\partial\psi \wedge \bar{\partial}\psi$ is q -positive semi-definite for some constant $\delta \in [0, 4)$. Then for any $\bar{\partial}$ -closed (p, q) -form $f \in L_{p,q}^2(\Omega, \text{loc})$ ($0 \leq p \leq n$), if*

$$\int_{\Omega} \langle F_{\varphi}^{-1} f, f \rangle e^{-\varphi+\psi} < \infty,$$

there exists a $(p, q-1)$ -form $u \in L_{p,q-1}^2(\Omega, \varphi - \psi)$ such that

$$\bar{\partial}u = f, \quad \|u\|_{\varphi-\psi}^2 \leq \frac{4}{(2-\sqrt{\delta})^2} \int_{\Omega} \langle F_{\varphi}^{-1} f, f \rangle e^{-\varphi+\psi},$$

where F_{φ}^{-1} is defined by (8) and it is required implicitly that $F_{\varphi}^{-1} f$ is defined almost everywhere in Ω .

Corollary 1. *Let Ω be a weakly q -convex domain in \mathbb{C}^n ($1 \leq q \leq n$) and let φ be a q -subharmonic function on Ω . Then for any $\bar{\partial}$ -closed (p, q) -form $f \in L^2_{p,q}(\Omega, \text{loc})$ ($0 \leq p \leq n$), if*

$$\int_{\Omega} \langle F_{\varphi}^{-1} f, f \rangle e^{-\varphi} < \infty,$$

there exists a $(p, q-1)$ -form $u \in L^2_{p,q-1}(\Omega, \varphi - \psi)$ such that

$$\bar{\partial}u = f, \quad \|u\|_{\varphi}^2 \leq \int_{\Omega} \langle F_{\varphi}^{-1} f, f \rangle e^{-\varphi}.$$

Corollary 2. *Let Ω be a weakly q -convex domain in \mathbb{C}^n ($1 \leq q \leq n$) and let φ be a q -subharmonic function on Ω , $\psi \in C^2(\Omega)$ be a function such that $-e^{-\psi}$ is q -subharmonic. For any constant $\delta \in [0, 1)$ and $\bar{\partial}$ -closed (p, q) -form $f \in L^2_{p,q}(\Omega, \text{loc})$ ($0 \leq p \leq n$), if*

$$\int_{\Omega} \langle F_{\psi}^{-1} f, f \rangle e^{-\varphi + \delta\psi} < \infty$$

then there exists a $(p, q-1)$ -form $u \in L^2_{p,q-1}(\Omega, \varphi - \delta\psi)$ such that

$$\bar{\partial}u = f, \quad \|u\|_{\varphi - \delta\psi}^2 \leq \frac{4}{(1-\delta)^2} \int_{\Omega} \langle F_{\psi}^{-1} f, f \rangle e^{-\varphi + \delta\psi}.$$

Corollary 3. *Let Ω be a weakly q -convex domain in \mathbb{C}^n ($1 \leq q \leq n$) and let φ be a q -subharmonic function on Ω , $\psi \in C^2(\Omega)$ be a strictly plurisubharmonic function such that $-e^{-\psi}$ is q -subharmonic. For any constant $\delta \in [0, 1)$ and $\bar{\partial}$ -closed (p, q) -form $f \in L^2_{p,q}(\Omega, \text{loc})$ ($0 \leq p \leq n$), if*

$$\int_{\Omega} \psi^{\bar{j}k} f_{I,\bar{j}\bar{k}} \overline{f_{I,\bar{k}\bar{k}}} e^{-\varphi + \delta\psi} < \infty$$

then there exists a $(p, q-1)$ -form $u \in L^2_{p,q-1}(\Omega, \varphi - \delta\psi)$ such that

$$\bar{\partial}u = f, \quad \|u\|_{\varphi - \delta\psi}^2 \leq \frac{4}{q^2(1-\delta)^2} \int_{\Omega} \psi^{\bar{j}k} f_{I,\bar{j}\bar{k}} \overline{f_{I,\bar{k}\bar{k}}} e^{-\varphi + \delta\psi}$$

where $(\psi^{\bar{j}k}) := (\psi_{j\bar{k}})^{-1}$.

Corollary 4. *Let Ω be a weakly q -convex domain in \mathbb{C}^n ($1 \leq q \leq n$) and let φ be a q -subharmonic function on Ω , $\psi \in C^2(\Omega)$ be a q -subharmonic function such that $-e^{-\psi}$ is q -subharmonic. For any $\bar{\partial}$ -closed (p, q) -form $f \in L^2_{p,q}(\Omega, \text{loc})$ ($0 \leq p \leq n$), if*

$$\int_{\Omega} \langle F_{\psi}^{-1} f, f \rangle e^{-\varphi} < \infty$$

then there exists a $(p, q - 1)$ -form $u \in L^2_{p, q-1}(\Omega, \varphi)$ such that

$$\bar{\partial}u = f, \quad \|u\|_\varphi^2 \leq 4 \int_\Omega \langle F_\psi^{-1} f, f \rangle e^{-\varphi}.$$

Corollary 5. *Let Ω be a bounded weakly q -convex domain in \mathbb{C}^n ($1 \leq q \leq n$) and let φ be a q -subharmonic function on Ω . For any $\bar{\partial}$ -closed (p, q) -form $f \in L^2_{p, q}(\Omega, \varphi)$ ($0 \leq p \leq n$), there exists a $(p, q - 1)$ -form $u \in L^2_{p, q-1}(\Omega, \varphi)$ such that*

$$\bar{\partial}u = f, \quad \|u\|_\varphi \leq \frac{2d}{q} \|f\|_\varphi$$

where d is the diameter of Ω .

Since there are plenty of q -subharmonic functions which are not plurisubharmonic when $q \geq 2$, our results provide more flexibility in choosing weights for L^2 -estimates. Such flexibility may help us make generalizations and improvements on existence results for the $\bar{\partial}$ -problem. Let ρ be the function in Theorem 1 above, then it is easy to see that $-e^{-\psi}$ is strictly q -subharmonic on Ω where $\psi := -\log(-\rho)$, as a consequence, we obtain Theorem 2.4 in [11]. Theorem 1 was originally proved by Diederich and Fornæss ([6]) for pseudoconvex domains, i.e. the case of $q = 1$. Theorem 2 was obtained by Błocki ([5]) for $(0, 1)$ -forms on pseudoconvex domains. Corollary 1 is a strengthened version of Theorem 3.1 in [11]. In the case of $q = 1$, Corollary 2 recovers a result due to Błocki ([3]). The arguments used in [3] and [5] do not indicate the estimates (*), (**). Corollary 3 above improves the main result in [1] and our Corollary 5 improves slightly a result due to Hörmander (Theorem 2.2.3 in [10]) when $q \geq 2$.

2. Weakly q -convex domains

We begin by establishing the basic notation.

We will adhere to the summation convention that sum is performed over strictly increasing multi-indices. The coordinates of \mathbb{C}^n are chosen such that the standard Kähler form of \mathbb{C}^n is given by $\sqrt{-1} dz_j \wedge d\bar{z}_j$. Let Ω be a domain in \mathbb{C}^n and let $\phi \in C^\infty(\Omega)$, we denote by $\nabla^{0,1}\phi$ the $(0, 1)$ -part of the gradient $\nabla\phi$ of ϕ w.r.t. the standard Kähler metric, i.e. $\nabla^{0,1}\phi = \phi_j \partial/\partial\bar{z}_j$. We use $\langle \cdot, \cdot \rangle$ to denote the induced (pointwise) Hermitian inner product of (p, q) -forms on Ω . Following [10], the weighted L^2 Hermitian inner product of (p, q) -forms will be denoted by $(\cdot, \cdot)_\varphi$ and the corresponding Hilbert space will be denoted by $L^2_{p, q}(\Omega, \varphi)$.

Let Ω be a domain in \mathbb{C}^n , $1 \leq q \leq n$, we recall the notion of q -subharmonicity ([11], [14] and [13]).

DEFINITION 1. Let φ be an upper semi-continuous function on Ω , we say φ is q -subharmonic on Ω if the restriction of φ to any q dimensional complex submanifold of Ω is subharmonic w.r.t. the induced metric.

REMARK 1. It is easy to show (see [1], [14]) that for any $\varphi \in C^2(\Omega)$, φ is q -subharmonic if and only if any sum of q eigenvalues of the complex Hessian

$$\varphi_{i\bar{j}} dz_i \otimes d\bar{z}_j$$

of φ is nonnegative. If any sum of q eigenvalues of the complex Hessian is positive, φ is then called strictly q -subharmonic. Moreover, every q -subharmonic function could be approximated by a decreasing sequence of smooth q -subharmonic functions. It is easy to see that 1-subharmonicity is equivalent to plurisubharmonicity.

Following [11] and [13], we introduce the notion of q -convexity.

DEFINITION 2. Assume Ω is a smooth domain, and r a defining function for Ω , then we say that Ω is weakly q -convex if at every point $b \in \partial\Omega$ we have

$$r_{i\bar{j}}(b)g_{i\bar{k}}\overline{g_{j\bar{k}}} \geq 0$$

for every $(0, q)$ -form $g = g_{j\bar{j}} d\bar{z}_j$ such that

$$r_i g_{i\bar{k}} = 0$$

for all multi-indices K with $|K| = q - 1$. For a general domain $\Omega \subseteq \mathbb{C}^n$, we call it weakly q -convex if it could be exhausted by smooth weakly q -convex domains.

REMARK 2. It is easy to see that q -subharmonicity (convexity) implies $(q + 1)$ -subharmonicity (convexity). The notions of q -subharmonicity and q -convexity are both invariant under a unitary change of coordinates, but not preserved by biholomorphic transformations.

Assume $\Omega \subseteq \mathbb{C}^n$ is a smooth domain, and $r \in C^\infty(\overline{\Omega})$ is a defining function for Ω . Let $\phi \in C^\infty(\overline{\Omega})$ and $g \in C_{p,q}^\infty(\overline{\Omega})$ satisfy

$$r_i g_{i\bar{i}K} = 0$$

for all multi-indices K with $|I| = p$, $|K| = q - 1$, then we have the standard Kohn–Morrey–Hörmander identity

$$\begin{aligned} \|\bar{\partial}g\|_\phi^2 + \|\bar{\partial}_\phi^*g\|_\phi^2 &= \int_\Omega \partial_j \partial_{\bar{k}} \phi g_{i\bar{j}K} \overline{g_{i\bar{k}K}} e^{-\phi} \\ &\quad + \int_{\partial\Omega} \partial_j \partial_{\bar{k}} r g_{i\bar{j}K} \overline{g_{i\bar{k}K}} \frac{1}{|\nabla r|} e^{-\phi} \\ &\quad + \int_\Omega |\partial_{\bar{j}} g_{i\bar{j}}|^2 e^{-\phi}. \end{aligned}$$

When Ω is q -convex, we obtain the following inequality

$$(1) \quad \|\bar{\partial}g\|_{\phi}^2 + \|\bar{\partial}_{\phi}^*g\|_{\phi}^2 \geq \int_{\Omega} \partial_j \bar{\partial}_{\bar{k}} \phi g_{I, \bar{J} \bar{K}} \overline{g_{I, \bar{K} \bar{K}}} e^{-\phi}.$$

We denote by $\bigwedge^{p,q}$ the linear space of (p, q) -forms, i.e. $\bigwedge^{p,q} = \text{span}_{\mathbb{C}}\{dz_I \wedge d\bar{z}_J \mid |I| = p, |J| = q\}$. For any real $(1, 1)$ -form $\theta = \sqrt{-1} \theta_{j\bar{k}} dz_j \wedge d\bar{z}_{\bar{k}}$, we introduce a self-adjoint linear operator on $\bigwedge^{p,q}$ by setting

$$(2) \quad F_{\theta} = \theta_{j\bar{k}} d\bar{z}_{\bar{k}} \wedge \frac{\partial}{\partial z_j} \lrcorner$$

where \lrcorner means the interior product. We also set the notation $F_{\phi} := F_{\sqrt{-1} \partial \bar{\partial} \phi}$ for a smooth function ϕ .

With the linear operator F_{ϕ} , we can rewrite the integrand on the right hand side of (1) as follows

$$(3) \quad \begin{aligned} \partial_j \bar{\partial}_{\bar{k}} \phi g_{I, \bar{J} \bar{K}} \overline{g_{I, \bar{K} \bar{K}}} &= \left(\phi_{j\bar{k}} \frac{\partial}{\partial z_j} \lrcorner g \right)_{I, \bar{K}} \cdot \overline{\left(\frac{\partial}{\partial z_k} \lrcorner g \right)_{I, \bar{K}}} \\ &= \left\langle \phi_{j\bar{k}} \frac{\partial}{\partial z_j} \lrcorner g, \frac{\partial}{\partial z_k} \lrcorner g \right\rangle \\ &= \langle F_{\phi} g, g \rangle. \end{aligned}$$

Consequently, we obtain by the Kohn–Morrey–Hörmander identity and (3)

$$(4) \quad \|\bar{\partial}g\|_{\phi}^2 + \|\bar{\partial}_{\phi}^*g\|_{\phi}^2 \geq \int_{\Omega} \langle F_{\phi} g, g \rangle e^{-\phi} := (F_{\phi} g, g)_{\phi}.$$

Denote the eigenvalues of the matrix $(\theta_{j\bar{k}})$ by

$$\lambda_1 \leq \dots \leq \lambda_n,$$

after a unitary change of coordinates, we have $F_{\theta} = \lambda_j d\bar{z}_{\bar{j}} \wedge \partial/\partial z_j \lrcorner$. For any multi-indices I, J with $|I| = p, |J| = q$, set

$$(5) \quad \lambda_{I, J} := \sum_{j \in J} \lambda_j,$$

it holds that

$$\begin{aligned} F_{\theta} dz_I \wedge d\bar{z}_J &= \lambda_j dz_I \wedge d\bar{z}_j \wedge \frac{\partial}{\partial z_j} \lrcorner d\bar{z}_J \\ &= \lambda_j dz_I \wedge d\bar{z}_j \wedge \sum_{a=1}^q (-1)^{a-1} \delta_{j j_a} d\bar{z}_{j_1} \wedge \dots \wedge \widehat{d\bar{z}_{j_a}} \wedge \dots \wedge d\bar{z}_{j_q} \\ &= \sum_{j \in J} \lambda_j dz_I \wedge d\bar{z}_j = \lambda_{I, J} dz_I \wedge d\bar{z}_J \end{aligned}$$

where the circumflex over a term means that it is to be omitted. Hence eigenvalues of the map F_θ are given by

$$(6) \quad \lambda_{I,J}, \quad |I| = p, |J| = q.$$

DEFINITION 3. Let $\theta = \sqrt{-1}\theta_{j\bar{k}} dz_j \wedge d\bar{z}_k$ be a real $(1,1)$ -form on \mathbb{C}^n , $1 \leq q \leq n$. θ is said to be q -positive semi-definite (q -positive) if $\lambda_1 + \cdots + \lambda_q \geq 0$ (> 0) where $\lambda_1 \leq \cdots \leq \lambda_n$ are the eigenvalues of the matrix $(\theta_{j\bar{k}})$.

REMARK 3. By formula (6), θ is q -positive semi-definite if and only if the operator $F_\theta: \bigwedge^{p,q} \rightarrow \bigwedge^{p,q}$ is a positive semi-definite for any $0 \leq p \leq n$. We have the following criterion for q -subharmonicity of a smooth function ϕ .

ϕ is q -subharmonic (strictly q -subharmonic) on Ω if and only if F_ϕ is q -positive semi-definite (definite) at each point of Ω .

Since $F_\theta: \bigwedge^{p,q} \rightarrow \bigwedge^{p,q}$ is self-adjoint, we have the following orthogonal decomposition

$$(7) \quad \bigwedge^{p,q} = \text{Ker } F_\theta \oplus \text{Im } F_\theta,$$

which implies that F_θ induces an isomorphism $F_\theta|_{\text{Im } F_\theta}: \text{Im } F_\theta \rightarrow \text{Im } F_\theta$. We can therefore define

$$(8) \quad F_\theta^{-1} := (F_\theta|_{\text{Im } F_\theta})^{-1}: \text{Im } F_\theta \rightarrow \text{Im } F_\theta$$

for any real $(1,1)$ -form θ . Notice that F_θ itself is not required to be invertible in the above definition.

When θ is q -positive, we know by (6)

$$(9) \quad (F_\theta^{-1}g)_{I,J} = \lambda_{I,J}^{-1} g_{I,J}$$

holds for any $g = g_{I,J} dz_I \wedge d\bar{z}_J \in \bigwedge^{p,q}$ and any given multi-indices I, J satisfying $|I| = p, |J| = q$.

If the function ϕ is further assumed to be strictly plurisubharmonic, we denote by $(\phi^{\bar{j}k})$ the inverse matrix of the complex Hessian matrix $(\phi_{j\bar{k}})$, then we have

$$(10) \quad \begin{aligned} \langle F_\phi^{-1}g, g \rangle &= \lambda_{I,J}^{-1} |g_{I,J}|^2 \\ &= \left(\sum_{j \in J} \lambda_j \right)^{-1} |g_{I,J}|^2 \\ &\leq \frac{1}{q^2} \sum_{j \in I} \lambda_j^{-1} |g_{I,J}|^2 \\ &\stackrel{(3)(5)}{=} \frac{1}{q^2} \phi^{\bar{j}k} g_{I,\bar{j}\bar{k}} \overline{g_{I,\bar{k}\bar{k}}} \end{aligned}$$

for arbitrary $g = g_{I,\bar{J}} dz_I \wedge d\bar{z}_{\bar{J}} \in \bigwedge^{p,q}$.

We conclude this section by proving a Diederich–Fornaess type result for smooth bounded weakly q -convex domains.

Proof of Theorem 1. By Remark 3, it suffices to show that $\langle F_\rho g, g \rangle > 0$ for any $0 \neq g \in \bigwedge^{0,q}$. A direct computation gives

$$(11) \quad \begin{aligned} \langle F_\rho g, g \rangle &= \eta(-r)^{\eta-2} e^{-\eta K \phi} \cdot [K r^2 (\langle F_\phi g, g \rangle - \eta K |\nabla^{0,1} \phi \lrcorner g|^2) \\ &\quad - r (\langle F_r g, g \rangle - 2\eta K \Re \langle \nabla^{0,1} \phi \lrcorner g, \nabla^{0,1} r \lrcorner g \rangle) \\ &\quad + (1 - \eta) |\nabla^{0,1} r \lrcorner g|^2]. \end{aligned}$$

Throughout the proof, we denote by A_1, A_2, \dots various constants which are independent of η, K .

Since the boundary of Ω is assumed to be smooth, for any sufficiently small $\varepsilon > 0$ there is a smooth map $\pi : N_\varepsilon \rightarrow \partial\Omega$ such that

$$(12) \quad \text{dist}(z, \partial\Omega) = |z - \pi(z)|, \quad z \in N_\varepsilon$$

where $N_\varepsilon := \{z \in \Omega \mid r(z) > -\varepsilon\}$. As the function $r \in C^\infty(\bar{\Omega})$ is a defining function for Ω , there exists a constant $A_1 > 0$ which only depends on ε such that

$$(13) \quad \text{dist}(z, \partial\Omega) \leq -A_1 r(z), \quad A_1 \leq |\nabla r(z)|, \quad z \in N_\varepsilon.$$

For any $g \in \bigwedge^{0,q}$, $z \in N_\varepsilon$, set

$$g_1(z) = \frac{1}{|\nabla^{0,1} r(z)|^2} \nabla^{0,1} r(z) \lrcorner \bar{\partial} r(z) \wedge g, \quad g_2(z) = \frac{1}{|\nabla^{0,1} r(z)|^2} \bar{\partial} r(z) \wedge \nabla^{0,1} r(z) \lrcorner g,$$

then we have $g = g_1(z) + g_2(z)$, $|g|^2 = |g_1(z)|^2 + |g_2(z)|^2$ and

$$(14) \quad \nabla^{0,1} r(z) \lrcorner g_1(z) = 0, \quad |g_2(z)| \leq \frac{1}{|\nabla^{0,1} r(z)|} |\nabla^{0,1} r(z) \lrcorner g|$$

for every $z \in N_\varepsilon$. From (12) and the first inequality in (13), there is a constant $A_2 > 0$ such that

$$(15) \quad \begin{aligned} |\langle F_r g_1, g_1 \rangle(z) - \langle F_r g_1, g_1 \rangle(\pi(z))| &= \left| \int_0^1 \frac{d}{dt} \langle F_r g_1, g_1 \rangle(tz + (1-t)\pi(z)) dt \right| \\ &\leq -A_2 r(z) |g|^2 \end{aligned}$$

holds for any $z \in N_\varepsilon$. By (3), the identity in (14) and Definition 2, we get

$$\langle F_r g_1, g_1 \rangle(\pi(z)) \geq 0, \quad z \in N_\varepsilon.$$

Therefore, for any $z \in N_\varepsilon$, the following estimate follows from (15)

$$\langle F_r g_1, g_1 \rangle(z) \geq A_2 r(z) |g|^2.$$

Taking into account of the inequality in (14) and $|g_1(z)| \leq |g|$, the above estimate implies that

$$(16) \quad \langle F_r g, g \rangle(z) \geq A_2 r(z) |g|^2 - \frac{A_3}{|\nabla^{0,1} r(z)|} |\nabla^{0,1} r(z) \lrcorner g| |g|$$

holds for any $z \in N_\varepsilon$ where $A_3 > 0$ is another constant.

Since ϕ is strictly q -subharmonic on $\bar{\Omega}$, there is a constant $\sigma > 0$ such that

$$(17) \quad \langle F_\phi g, g \rangle(z) - \eta K |\nabla^{0,1} \phi(z) \lrcorner g|^2 \geq (\sigma - A_4 \eta K) |g|^2$$

holds for any $z \in \Omega$ where $A_4 := \sup_\Omega |\nabla^{0,1} \phi|^2$. From (11) and (17), there exists a constant $A_5 > 0$ such that

$$(18) \quad \langle F_\rho g, g \rangle(z) \geq \eta (-r)^{\eta-2} e^{-\eta K \phi} \left[K r^2(z) \left(\sigma - \frac{\eta}{1-\eta} A_4 K \right) - A_5 \right] |g|^2$$

holds for any $z \in \Omega$.

When $K > 4A_5/(\sigma \varepsilon^2)$ and $\eta \in (0, \sigma/(2A_4 K + \sigma))$, (18) implies that

$$(19) \quad \langle F_\rho g, g \rangle \geq \frac{1}{4} \eta (-r)^{\eta-2} e^{-\eta K \phi} K \varepsilon^2 \sigma |g|^2$$

holds on $\Omega \setminus N_\varepsilon$.

Similarly, for any constants $\eta \in (0, \sigma/(2A_4 K))$ and $K > (4/\sigma)(A_2 + (\sigma^2/(4A_4)) + 2A_6^2 + \sigma^2)$, $A_6 := A_3/(2A_1)$, from (11), (16) and (17) it follows that the following inequality holds on N_ε

$$(20) \quad \begin{aligned} \langle F_\rho g, g \rangle &\geq \eta (-r)^{\eta-2} e^{-\eta K \phi} [K(\sigma - A_4 \eta K) - A_2] r^2 |g|^2 \\ &\quad + 2(A_6 + A_4 \eta K) |\nabla^{0,1} r \lrcorner g| r |g| \\ &\quad + (1-\eta) |\nabla^{0,1} r \lrcorner g|^2 \\ &\geq \eta (-r)^{\eta-2} e^{-\eta K \phi} \left[K(\sigma - A_4 \eta K) - A_2 - \frac{2A_6^2 + 2A_4^2 \eta^2 K^2}{1-\eta} \right] r^2 |g|^2 \\ &\geq \eta (-r)^{\eta-2} e^{-\eta K \phi} \left(\frac{K\sigma}{2} - A_2 - 4A_6^2 - \sigma^2 \right) r^2 |g|^2 \\ &\geq \frac{1}{4} \eta (-r)^{\eta-2} e^{-\eta K \phi} K r^2 \sigma |g|^2. \end{aligned}$$

By combining (19) and (20), we know Theorem 1 is true for any constant $K > (4/\sigma)(A_2 + \sigma^2/(4A_4) + A_5/\varepsilon^2 + 2A_6^2 + \sigma^2)$ and $\eta_0 := \sigma/(2A_4 K + \sigma)$. \square

3. Donnelly–Fefferman type estimate

We will prove, in this section, the existence results in the present paper. The key for our proofs is to establish an a priori estimate of Donnelly–Fefferman type from which we get the existence theorem 1. Since the constant δ involved in this estimate would be allowed to have value zero, we also obtain an existence result of Donnelly–Fefferman type and Hörmander type (with one weight function). We first recall a basic lemma from functional analysis which is due to Hörmander (see [10]).

Lemma. *Let $H_1 \xrightarrow{T} H_2 \xrightarrow{S} H_3$ be a complex of closed and densely defined operators between Hilbert spaces. For any $f \in \text{Ker } S$ and any constant $C > 0$, the following conditions are equivalent.*

1. *There exists some $u \in H_1$ such that $Tu = f$ and $\|u\|_{H_1} \leq C$.*
2. *$|(f, g)_{H_2}|^2 \leq C^2(\|T^*g\|_{H_1}^2 + \|Sg\|_{H_3}^2)$ holds for each $g \in \text{Dom}(T^*) \cap \text{Dom}(S)$.*

Proof of Theorem 2. We consider first the case where Ω is a bounded domain in \mathbb{C}^n with smooth boundary and $\varphi, \psi \in C^\infty(\overline{\Omega})$

We will apply the above lemma to following weighted L^2 -spaces of differential forms

$$H_1 = L^2_{p,q-1}\left(\Omega, \varphi - \frac{1}{2}\psi\right), \quad H_2 = L^2_{p,q}\left(\Omega, \varphi - \frac{1}{2}\psi\right), \quad H_3 = L^2_{p,q+1}\left(\Omega, \varphi - \frac{1}{2}\psi\right)$$

and the operators

$$T = \bar{\partial} \circ e^{-(1/4)\psi}, \quad S = e^{-(1/4)\psi} \circ \bar{\partial}.$$

In order to use the above lemma, we need to show that the following estimate

$$(21) \quad \begin{aligned} & |(f, g)_{\varphi-(1/2)\psi}|^2 \\ & \leq \frac{4(F_\varphi^{-1}f, f)_{\varphi-\psi}}{(2-\sqrt{\delta})^2} (\|e^{-(1/4)\psi} \bar{\partial}_\varphi^* g\|_{\varphi-(1/2)\psi}^2 + \|e^{-(1/4)\psi} \bar{\partial} g\|_{\varphi-(1/2)\psi}^2) \end{aligned}$$

holds for arbitrary $g \in \text{Dom}(\bar{\partial}^*) \cap C^\infty_{p,q}(\overline{\Omega})$.

Let $g \in \text{Dom} \bar{\partial}^* \cap C^\infty_{p,q}(\overline{\Omega})$, from

$$\bar{\partial}_\varphi^* g = \bar{\partial}_{\varphi-(1/2)\psi}^* g + \frac{1}{2} \nabla^{0,1} \psi \lrcorner g,$$

by using Cauchy's inequality with ε , it follows that

$$\|\bar{\partial}_\varphi^* g\|_\varphi^2 \leq \frac{1+\varepsilon}{\varepsilon} \|\bar{\partial}_{\varphi-(1/2)\psi}^* g\|_\varphi^2 + \frac{1+\varepsilon}{4} \|\nabla^{0,1} \psi \lrcorner g\|_\varphi^2$$

for any positive constant ϵ . For any $\varepsilon \in [0, 4)$, let

$$\epsilon = \frac{2}{\sqrt{\delta}} - 1,$$

then the above inequality becomes

$$(22) \quad \|\bar{\partial}_\varphi^* g\|_\varphi^2 \leq \frac{2}{2 - \sqrt{\delta}} \|\bar{\partial}_{\varphi - (1/2)\psi}^* g\|_\varphi^2 + \frac{1}{2\sqrt{\delta}} \|\nabla^{0,1} \psi \lrcorner g\|_\varphi^2.$$

Since $\delta\sqrt{-1}\partial\bar{\partial}\varphi - \sqrt{-1}\partial\psi \wedge \bar{\partial}\psi$ is q -positive semi-definite, we get the following inequality

$$(23) \quad \delta\langle F_\varphi g, g \rangle \geq |\nabla^{0,1} \psi \lrcorner g|^2.$$

Substituting (22) and (23) into Hörmander's estimate (4), the q -subharmonicity of φ gives

$$\begin{aligned} \frac{2}{2 - \sqrt{\delta}} \|\bar{\partial}_{\varphi - \frac{1}{2}\psi}^* g\|_\varphi^2 + \|\bar{\partial}g\|_\varphi^2 &\geq \|\bar{\partial}_\varphi^* g\|_\varphi^2 + \|\bar{\partial}g\|_\varphi^2 - \frac{1}{2\sqrt{\delta}} \|\nabla^{0,1} \psi \lrcorner g\|_\varphi^2 \\ &\geq \frac{2 - \sqrt{\delta}}{2} \int_\Omega \langle F_\psi g, g \rangle e^{-\varphi} \end{aligned}$$

which further implies the desired estimate (*) as follows

$$\begin{aligned} \|e^{-(1/4)\psi} \bar{\partial}_{\varphi - (1/2)\psi}^* g\|_{\varphi - (1/2)\psi}^2 + \|e^{-(1/4)\psi} \bar{\partial}g\|_{\varphi - (1/2)\psi}^2 &= \|\bar{\partial}_{\varphi - (1/2)\psi}^* g\|_\varphi^2 + \|\bar{\partial}g\|_\varphi^2 \\ &\geq \|\bar{\partial}_{\varphi - (1/2)\psi}^* g\|_\varphi^2 + \frac{2 - \sqrt{\delta}}{2} \|\bar{\partial}g\|_\varphi^2 \\ &\geq \frac{(2 - \sqrt{\delta})^2}{4} \int_\Omega \langle F_\varphi g, g \rangle e^{-\varphi}. \end{aligned}$$

Since φ is q -subharmonic, the Cauchy–Schwarz inequality applied to the positive semi-definite Hermitian form $(F_\varphi \cdot, \cdot)_\varphi$ gives

$$\begin{aligned} |(f, g)_{\varphi - \frac{1}{2}\psi}|^2 &= |(F_\varphi \circ F_\varphi^{-1} e^{(1/2)\psi} f, g)_\varphi|^2 \\ &\leq (e^{(1/2)\psi} f, e^{(1/2)\psi} F_\varphi^{-1} f)_\varphi (F_\varphi g, g)_\varphi \\ &\leq \frac{4(F_\varphi^{-1} f, f)_{\varphi - \psi}}{(2 - \sqrt{\delta})^2} (\|e^{-(1/4)\psi} \bar{\partial}_{\varphi - (1/2)\psi}^* g\|_{\varphi - \frac{1}{2}\psi}^2 + \|e^{-(1/4)\psi} \bar{\partial}g\|_{\varphi - (1/2)\psi}^2) \end{aligned}$$

where F_φ^{-1} is defined by (8). Thus the estimate (21) has been proved for $g \in \text{Dom} \bar{\partial}^* \cap C_{p,q}^\infty(\bar{\Omega})$. By using the density lemma (Proposition 1.2.4 in [10]), we know that (22) holds for any $g \in \text{Dom}(T^*) \cap \text{Dom}(S)$. Consequently, by the lemma we mentioned at

the beginning of this section, there exists some $v \in L^2_{p,q-1}(\Omega, \varphi - (1/2)\psi)$ such that

$$Tv = f, \quad \|v\|_{\varphi - (1/2)\psi}^2 \leq \frac{4}{(2 - \sqrt{\delta})^2} (F_\varphi^{-1} f, f)_{\varphi - \psi}.$$

Set $u = e^{-(1/4)\psi} v$, then we get $u \in L^2_{p,q-1}(\Omega, \varphi - \psi)$ and

$$(24) \quad \bar{\partial}u = f, \quad \|u\|_{\varphi - \psi}^2 = \|v\|_{\varphi - (1/2)\psi}^2 \leq \frac{4}{(2 - \sqrt{\delta})^2} (F_\varphi^{-1} f, f)_{\varphi - \psi}.$$

Theorem 2 now follows, in its full generality, from (24), the standard argument of smooth approximation and taking weak limit (see e.g. [10]). \square

Proof of Corollary 1. Corollary 1 follows from Theorem 2 by choosing $\delta = 0$ and $\psi = 0$. \square

Proof of Corollary 2. Let $\varphi_1 = \varphi + \psi$ and $\psi_1 = (1 + \delta)\psi$, then φ_1 is q -subharmonic. Since

$$(1 + \delta)^2 \sqrt{-1} \partial \bar{\partial} \varphi_1 - \sqrt{-1} \partial \psi_1 \wedge \bar{\partial} \psi_1 = (1 + \delta)^2 [\sqrt{-1} \partial \bar{\partial} \varphi + \sqrt{-1} e^\psi \partial \bar{\partial} (-e^{-\psi})],$$

the assumption that φ and $-e^{-\psi}$ are both q -subharmonic functions implies that $(1 + \delta)^2 \sqrt{-1} \partial \bar{\partial} \varphi_1 - \sqrt{-1} \partial \psi_1 \wedge \bar{\partial} \psi_1$ is q -positive semi-definite. Applying Theorem 2 to the weights φ_1 and ψ_1 , we obtain Corollary 2. \square

Proof of Corollary 3. Corollary 3 follows directly from Corollary 2 and the point-wise inequality (10). \square

Proof of Corollary 4. Corollary 4 follows directly from Corollary 2 by choosing the constant δ to be 0. \square

Proof of Corollary 5. Without loss of generality, we assume that Ω contains the origin of \mathbb{C}^n . Let $\psi = q|z|^2/d^2$, then (9) implies that $F_\psi^{-1} = (d^2/q^2)\text{Id}$ on (p, q) -forms. Since the complex Hessian of $-e^{-\psi}$ is given by

$$\frac{q}{d^2} e^{-\psi} \left(dz_i \otimes d\bar{z}_i - \frac{q}{d^2} z_i dz_i \otimes z_j d\bar{z}_j \right),$$

we know that any sum of q eigenvalues of the complex Hessian of $-e^{-\psi}$ is not less than

$$\frac{q}{d^2} e^{-\psi} \left[\left(1 - \frac{q}{d^2} |z|^2 \right) + q - 1 \right] = \frac{q^2}{d^2} e^{-\psi} \left(1 - \frac{|z|^2}{d^2} \right) \geq 0.$$

So $-e^{-\psi}$ is, by definition, a q -subharmonic function on Ω (but not plurisubharmonic). Applying Corollary 4 with the weight $\psi = q|z|^2/d^2$, we obtain the following estimate for the solution u

$$\|u\|_{\varphi}^2 \leq \frac{4d^2}{q^2} \|f\|_{\varphi}^2.$$

This completes the proof of Corollary 5. □

References

- [1] H. Ahn and N.Q. Dieu: *The Donnelly–Fefferman theorem on q -pseudoconvex domains*, Osaka J. Math. **46** (2009), 599–610.
- [2] B. Berndtsson: *The extension theorem of Ohsawa–Takegoshi and the theorem of Donnelly–Fefferman*, Ann. Inst. Fourier (Grenoble) **46** (1996), 1083–1094.
- [3] Z. Błocki: *A note on the Hörmander, Donnelly–Fefferman, and Berndtsson L^2 -estimates for the $\bar{\partial}$ -operator*, Ann. Polon. Math. **84** (2004), 87–91.
- [4] Z. Błocki: *The Bergman metric and the pluricomplex Green function*, Trans. Amer. Math. Soc. **357** (2005), 2613–2625.
- [5] Z. Błocki: *Suita conjecture and the Ohsawa–Takegoshi extension theorem*, to appear in Invent. Math.
- [6] K. Diederich and J.E. Forneaess: *Pseudoconvex domains: bounded strictly plurisubharmonic exhaustion functions*, Invent. Math. **39** (1977), 129–141.
- [7] K. Diederich and G. Herbot: *Extension of holomorphic L^2 -functions with weighted growth conditions*, Nagoya Math. J. **126** (1992), 141–157.
- [8] K. Diederich and T. Ohsawa: *An estimate for the Bergman distance on pseudoconvex domains*, Ann. of Math. (2) **141** (1995), 181–190.
- [9] H. Donnelly and C. Fefferman: *L^2 -cohomology and index theorem for the Bergman metric*, Ann. of Math. (2) **118** (1983), 593–618.
- [10] L. Hörmander: *L^2 estimates and existence theorems for the $\bar{\partial}$ operator*, Acta Math. **113** (1965), 89–152.
- [11] L.-H. Ho: *$\bar{\partial}$ -problem on weakly q -convex domains*, Math. Ann. **290** (1991), 3–18.
- [12] Q. Ji, X. Liu and G. Yu: *L^2 -estimates on p -convex Riemannian manifolds*, Adv. Math. **253** (2014), 234–280.
- [13] T. Napier and M. Ramachandran: *The Bochner–Hartogs dichotomy for weakly 1-complete Kähler manifolds*, Ann. Inst. Fourier (Grenoble) **47** (1997), 1345–1365.
- [14] H. Wu: *On certain Kähler manifolds which are q -complete*; in Complex Analysis of Several Variables (Madison, Wis., 1982), Proc. Sympos. Pure Math. **41**, Amer. Math. Soc., Providence, RI, 1984, 253–276.

Qingchun Ji
School of Mathematics
Key Laboratory of Mathematics for Nonlinear Sciences
Fudan University
Shanghai 200433
China
e-mail: qingchunji@fudan.edu.cn

Guo Tan
School of Mathematics
Key Laboratory of Mathematics for Nonlinear Sciences
Fudan University
Shanghai 200433
China
e-mail: 10210180016@fudan.edu.cn

Guangsheng Yu
School of Mathematics
Key Laboratory of Mathematics for Nonlinear Sciences
Fudan University
Shanghai 200433
China
e-mail: 12110180009@fudan.edu.cn