

COASTAL BUT NOT LITTORAL: MARINE RESOURCES IN NASCA DIET

Patrick H. Carmichael, Brenda V. Kennedy, and Lauren Cadwallader

Carmichael and Kennedy were members of the Department of Archaeology, University of Calgary, when this study was originally undertaken (1990). Patrick Carmichael is now in the Department of Sociology and Anthropology at Mount Royal University (pcarmichael@mtroyal.ca). Brenda Kennedy has left academe to work as an international program consultant. Lauren Cadwallader is a Research Associate at the McDonald Institute for Archaeological Research, University of Cambridge (lc340@cam.ac.uk).

Abstract

We examine the contribution of marine resources to the Nasca dietary economy (Early Intermediate Period, c. 100 BC – AD 600, Peruvian south coast) through ceramic iconography, settlement patterns, maritime subsistence technology, fish and shell remains, and stable isotope analysis. Each data set has limitations, but combined a consistent pattern emerges. Though the rich marine biomass of the Peru Current offers potential for huge food surpluses, we conclude the Nasca use of their littoral zone was minor. This result contrasts with earlier and later subsistence patterns in the same area, and with contemporary dietary systems elsewhere along the Andean coast. This challenge to conventional wisdom on coastal economies highlights the need for new research to understand the full range of Andean adaptations, especially those which appear counter-intuitive. This study also questions the notion that percentage frequencies of motifs in the iconography reflect daily realities.

The cold upwelling of the Peru Current endows Peru with one of the world's richest marine biomasses. Archaeology has long recognized the importance of fishing and littoral gathering

throughout Peruvian prehistory (Rostworoski 1981; Marcus 1987; Sandweiss et al. 1989), and maritime resources figure prominently in models of cultural transformation (Moseley 1975; Quilter and Stocker 1983; Quilter et al 1991; Sandweiss 2009; Feldman 2009). As coastal Peru is a hyperarid desert in which agriculture depends on unpredictable water supplies from seasonal river flooding, an emphasis or partial reliance on marine resources provides an obvious means of augmenting protein and coping with uncertain conditions.

In this study we examine the role of marine resources in Early Intermediate Period (EIP) Nasca culture (c. 100 BC – AD 600) of the Peruvian south coast (Figure 1). While the Nasca are considered a coastal people – in that they inhabited a coastal desert – they are known primarily from inland valley sites situated 40-60 kilometers from the ocean (Kroeber and Collier 1998; Schreiber 1999, Silverman 2002; Proulx 2007; Reindel 2009; Orefici 2012). Elsewhere on the Peruvian coast, rivers running down from the Andes have floodplains funneling outward to the sea. In such locales fishing and farming are easily integrated pursuits. In a pattern going back to the Late Preceramic and Initial Period in the Casma Valley, Pozorski and Pozorski refer to a “complementary coastal-inland subsistence pattern” or “economic symbiosis” between settlements on the coastline and those in riparian ecological zones (2012:367-369). This makes good sense, and for their examples they have good evidence. Our question is whether the Nasca engaged in a similar economic symbiosis, and if so, to what extent? For some archaeologists the answer is obvious. The Nasca provide us with iconographic depictions of sea creatures and fishermen, and many researchers have inferred that fishing was an important part of the economy (Blasco and Ramos 1980:128; Allen 1981:43; Lumbreras 1983:125; Proulx 1983:100; Townsend 1985:125; Silverman 1986:11; Carmichael 1988:34). We present several lines of evidence which converge to challenge this traditional view.

In southern Peru, the coastal desert reaches its maximum width in the regions of Ica and Nazca. Here, Andean rivers are separated from the ocean by a wide desert plain and range of seafront hills. In the Río Grande de Nazca Basin (hereafter Nazca Basin) all ten principal rivers merge inland, and only the Río Grande itself continues to the sea (Figure 1). These rivers have erratic seasonal flows, and their channels are dry most of the year, though sub-surface water may be present four meters or more below the surface. The region also experiences the most extreme sun and wind regimes on the entire Peruvian coast (on hydrology and climate see Schreiber and Lancho 2003:24-30; Beresford-Jones 2011:9-28). The lower reaches of the Ríos Ica and Grande trace much of their route through deep, narrow gorges which occasionally widen into small basins like verdant oases in the midst of an arid desert landscape. The river mouths are meager, anticlimactic effluents. Back from the oceanfront hills and away from the rivers the terrain is open, rolling country, characterized by stony pampas with rock outcrops punctuating the horizon. In this environment agriculture is highly circumscribed, being confined to the river margins and oases, and entirely dependent on seasonal flooding. The annual flood does not always arrive. Even the Río Grande, the largest river in the Nazca Basin, is considered one of the driest and most irregular rivers on the entire Pacific coast (ONERN 1971:197). Ironically, dry periods can be followed by destructive floods (Beresford-Jones et al. 2009:245). This is a tenuous environment for agriculturalists relying primarily on varieties of maize, beans, root crops, fruit, cotton, and gourds (Silverman 1993:289-292; Kellner and Schoeninger 2008:232; Vaughn 2009:58; Chiou et al. 2013:43). It is only logical for the Nasca to have supplemented their diet with nearby marine protein. The forty kilometers of desert separating the coastline from inland valleys was not a serious obstacle, being only a one-day walk for local inhabitants. At the time this study was first conceived (1988), some researchers assumed Nasca fishing towns subject to

inland urban centers contributed substantial food surpluses, which buffered the effects of periodic drought and supported dense populations including craft and military specialists, and a noble class ruling a state-level society. It was a model based on logical assumptions, but what did the archaeology say?

This study summarizes the evidence for Nasca marine exploitation from iconography, settlement patterns, subsistence technology, fish and shell remains, and stable isotope analysis of human remains. In the past, isotopic studies have concentrated on the importance of maize in the pre-Columbian diet (e.g., Ericson et al. 1989; Burger and van der Merwe 1990), with more recent isotope studies of Nasca diet focusing on identifying foreigners in the population (Knudson et al. 2009; Conlee et al. 2009), dietary shifts between time periods (Kellner and Schoeninger 2008), residence mobility (Webb et al. 2013), and general isotopic signatures for a region (Horn et al. 2009; Cadwallader et al. 2012). Results from these studies pertaining to the current topic are considered below: our work is focused exclusively on the issue of marine resource exploitation.

Preface and Update

This work was originally presented by Carmichael and Kennedy in 1991 at the Institute for Andean Studies Annual Conference, Berkeley, and later prepared for a monograph as yet unpublished (Carmichael 1998). The current, updated version has benefitted from a much expanded literature in the intervening years, and from the input of our co-author Lauren Cadwallader, who brought the isotope section up to date. However, in spite of numerous additional independent studies, and enhanced methods of analysis, our basic conclusions remain unchanged.

The chronology used in this article is the traditional South Coast framework set out by Rowe and Menzel (1973; see Figure 2). For comparison, the recent chronology advocated by Carmichael (2013) for the Southern Nasca Region is shown in Figure 3. In the new chronology Carmichael moved the original Nasca style phase 8 (EIP Epoch 8) to the Middle Horizon (MH), and Nasca 1 (EIP 1) to the transition between the Early Horizon (EH) and EIP, while the estimated time allotments are moved down a century. These adjustments do not seriously impinge on the findings reported herein which, as stated, follow the traditional Rowe and Menzel (1973) chronology. To date, our work is still the only study designed specifically to address the question of maritime input to Nasca diet.

In the interests of clarity, Nasca will be spelled with a “z” when referring to toponyms and geography, and with an “s” when referencing the archaeological culture.

Iconography

Depictions of sea creatures on Nasca pottery have been used to infer economic importance for maritime resources (Blasco and Ramos 1980:128; Proulx 1983:100; Silverman 1986:11; Carmichael 1988:34). Fishermen with nets, fish, shellfish, aquatic plants, and littoral birds are depicted, and some Nasca deities are based on marine creatures (Proulx 2006:83-85, 134-135, 149-156, 169). Piscine geoglyphs and petroglyphs are also documented in interior regions (Proulx 2007:67-68; Nieves 2010:28). It is evident the Nasca were well acquainted with marine life and, given the tremendous food potential of ocean resources, it is reasonable to hypothesize the littoral zone played an important role in the regional dietary economy.

A perusal of museum collections and the illustrated literature gives the impression that marine motifs are extremely common. However, the majority of collections lack secure

provenience, and the question of collector bias remains an unknown factor. In order to objectively assess motif frequencies, Carmichael (1992) assembled a large sample of excavated Nasca pottery (N=447 painted vessels). The pots came from 138 documented burials which could be confidently assigned to specific style phases according to the Dawson Seriation (Carmichael 1988:461-464), and derive from sites along the Nazca River (see Figure 4). Following Roark's (1965:57) motif classification system (still widely used today) the sample was divided into three groups based on primary motifs as follows: Referential – subject matter depicting life forms or objects in the real world (birds, plants, animals, fish and shellfish, humans) and assumed to represent secular or daily reality; Conceptual – subject matter that is not identifiable with beings or objects in the real world (spirits or deities) and therefore sacred; and Abstract – designs for which no referent can be identified (geometric, abstract, decorative designs – meaning, if any, unknown). Results are shown in Table 1, where the frequency of marine themes is noted under the referential category (adapted from Carmichael 1992:189). As shown in Table 1 marine themes account for 10.6% of the EIP 2-4 sample, and only 0.7% of the EIP 5-8 sample (see chronology in Figure 2, compare with Figure 3). If it is assumed that motif frequencies reflect economic importance, then it must be concluded fishing was virtually abandoned in the later epochs. This seems unlikely as climate studies indicate that beginning in the 5th century AD (EIP 5, Figure 2; Middle Nasca, Figure 3) the south coast experienced a period of protracted desiccation (Kevin Vaughn provides an excellent summary of several recent climate studies in Kanter and Vaughn 2012:75-76; also see Mächtle and Eitel 2013). At the same time riparian woodlands were significantly reduced to increase farmland, thereby altering soil and wind conditions which created an environment of ever-increasing aridity and fragility,

and culminated in the collapse of agricultural production (Beresford-Jones et al. 2011:290). These conditions do not favor a reduction in fishing activity during Late Nasca times.

If marine iconography is not documenting economic concerns, what is being expressed? A more useful approach frames the question within an indigenous Andean context. Previously, Carmichael noted that in traditional Andean thought the ocean was believed to be the source of all water (1992:188-189). The female sea deity, Mamacocha, was also “mother of lakes and water,” and springs were worshipped as her daughters (Cobo 1990:33, 117 [1653]). On a conceptual level no distinction was drawn between fresh water and salt water; rather, all water originated from the sea. Symbols of the sea are still used in rain ceremonies near Nazca today. In times of drought a man is sent to the coast at night to collect foamy sea water in a jug from a place where the waves crash against the rocks. The water is later sprinkled on the summit of a hill above Nazca in the belief this offering will cause rain in the mountains within two weeks (Urton 1982:11). In another account Antunez de Moyolo observed sea water being poured over seaweed near Nazca in order to bring rain (Reinhard 1988:51). In addition to sea water and seaweed, seashells were also used as offerings to ensure continued water supplies (Cobo 1990:117 [1653]), and any objects or images associated with the sea may have been perceived in a similar manner. In this context Nasca marine motifs need not be read as specific references to sea life, but rather, as general references to water/fertility concepts – understandable preoccupations for farmers dependant on inconsistent water supplies in a rainless, hyperarid desert (Carmichael 1992:188-189). The later reduction in depictive marine themes need not be read as a lack of concern for water; as Table 1 shows, in EIP 5-8 there is a greater than two-fold increase in conceptual (supernatural) themes, which include piscine motifs such as the “Mythical

Killer Whale” and water-related forms of the “Anthropomorphic Mythical Being” (Proulx 2006: Figs. 5.3, 5.9-10, 5.47-52, 5.54; and see Carmichael 1992: Figs. 7-8, 18).

Aside from marine themes, our isotope study (see below) provides another curious finding in regard to motif frequencies: approximately 50% of bone carbon was ultimately derived from C₄ plants (either directly or through animals eating them) with maize being the most probable candidate, yet as a primary motif on pottery, maize was entirely absent in the plant category (Table 1). A similar study done on Proulx’s (1968) Ica pottery samples for EIP 3-4 found maize comprised 4% of plant motifs (Carmichael 1992:188). The critical question here is whether depictions which can be identified with objects in the real world are “mundane” or “secular” images of daily life as stated by Roark (1965:57), or have interpretations fallen subject to classification? Carmichael has argued that Nasca iconography is a symbolic, interrelated system from which components cannot be isolated and treated as accurate reflections of ordinary reality (1992, 1994). A different view is presented by Donald Proulx (2006: 18, 140, 205). We should remain cautious in using iconography to reconstruct subsistence. Nonetheless, iconography unequivocally demonstrates the Nasca engaged in some fishing involving nets. The question is the extent to which this activity underpinned the regional dietary economy.

Settlement Patterns

Nasca sites are well documented in the inland valley areas of Ica and the Nazca Basin (Massey 1986; Browne and Baraybar 1988; Cook 1994; Schreiber 1998; Silverman 2002; Proulx 2007; Reindell 2009). These inland regions are separated from the ocean by 40-60 kilometers of desert plain, transected by the Ica and Grande rivers which trace their routes through coastal hills to the shores of the Pacific. Little systematic research had been done in the littoral zone before

Carmichael's 1989-1990 survey (Carmichael 1998 [and see Engel 1966, 1981, 1991]).

Carmichael's study area encompassed the coastal region opposite the inland centers of Ica and Nazca (Figure 1). While sites from all time periods were recorded, a main objective was to document Nasca fishing villages, which informants insisted were abundant and huge.

After six months of field work no oceanfront Nasca sites were found (EIP 2-8). Settlements dating to the Early Horizon (EH) and Late Intermediate Period (LIP) are present in the littoral zone, but EIP and Middle Horizon (MH) sites were not documented on the seafront. They occur in the oases several kilometers inland from the mouths of the Ica and Grande Rivers (both called Monte Grande), and it would be surprising if, eventually, a few sherds are not found on the shores, but remains indicative of permanent oceanfront habitation from these periods are thus far lacking.

Here, and hereafter in this paper, a qualification is required: Carmichael's survey findings reflect surface evidence as it appeared in 1989-1990. Recently disturbed areas, or deeply buried deposits, may yet modify his results. As previously noted, regional chronologies are currently in flux, and researchers' definitions can vary by area. The reader is again directed to the chronology used for Carmichael's survey, and in this article, shown in Figure 2 (Carmichael 1998:7).

At the sites of El Chucho (Cerro Chucho; Chucchio) and Karwa (Karwas; Carhua) in the Bahía de la Independencia at the northern end of the study area (Figure 1), occupation continued from the EH through the Necrópolis phase (EH 10 - EIP 2) according to Engel (1981:28-29) and Paul (1991:18) (also see Garcia and Pinilla 1995 for an excellent discussion of the Bahía during this time). The smaller site of Morro Quemado, located at the south end of the Bahía, also corresponds to this phase (Garcia and Pinilla 1995; Carmichael 1998:82). Carmichael did not find pottery dating to EIP 1-2 at these locales. His evidence suggests El Chucho and Karwa were

reoccupied in the LIP (Carmichael 1998: 73, 75). Evidently, earlier and later peoples fared quite well in this littoral environment.

Large and small villages from the LIP were encountered throughout the study area, and represent a more extensive use of the shoreline than any previous period in prehistory (Carmichael 1998:29). Here we deal only with the LIP site of La Yerba, located at the mouth of the Rio Ica (Figure 1), which provides a good example of what a permanent fishing village looks like (Carmichael 1998:87-88). The site has been known since the beginning of the last century when it was first described (colorfully) by Max Uhle¹; Engel refers to it as 15b VII-30 and 15b VII-35 (1981:20), or Max Uhle I and Max Uhle II (1991:157), and Cook designates it as PV-62-L-3 (1994:223). La Yerba covers two large, adjacent, shell-strewn hills facing the ocean on the south side of the Rio Ica, about 700 meters in from the beach line at the north end of La Yerba dune field (for which Carmichael named the site). Surface remains cover an area approximately 200 X 120 meters, and mounded to a height of 15 meters or more. In and around these hills are exposed sections of reed walls, foundation segments of silt-stone blocks, and adobes with mud plaster. Note that these more permanent materials are not available on the beachfront, and must have been hauled from some distance. Midden deposits contain thick, rich organic layers full of charcoal and ash. In addition to a great variety of shellfish, maize and gourd fragments are plentiful. Sea lion, whale, camelid, and canine bones are also present, as are pieces of coarse cloth, string, and netting. Fragments of large cooking and storage vessels are found over the entire site, and painted fineware sherds are plentiful. A nearby cemetery dates to the same period. At La Yerba, we have clear evidence of year-round occupation demonstrated not just by the amount of debris, but by the energy investment in durable construction materials and abundance of both utilitarian and fineware pottery, which had to be carried across the desert. A

corresponding cemetery demonstrates territorial permanency. It is fair to suggest that La Yerba participated in an economic symbiosis with its inland neighbors.

The EH-Necropolis sites of El Chucho and Karwa contain similar midden deposits, but very little pottery was visible on the surface in 1989-1990, and this mainly eroded or undecorated utilitarian ware. However, where LIP La Yerba consists of two, roughly circular mounds, the EH mounds are linear. El Chucho has 18 linear mounds, the larger ones estimated at 100-150 meters in length by 20 meters wide and 2-4 meters high (Carmichael 1998:73). The survey by Cordy-Collins at Karwa in 1980 (1998:37) identified 9 major mounds ranging from 31-234 meters long, 10-47 meters wide, and 1-6 meters high. Modest human burials were observed at both sites. These villages were important fishing communities of their era, and possibly year-round occupations. They may have engaged in economic symbiosis with agrarian settlements, probably in the Ica Valley rather than the Paracas Peninsula according to Garcia and Pinilla (1995:65).

Since we have major examples of oceanfront villages in the EH and LIP, why is the record blank for the EIP and MH? Here we will concentrate on the EIP, though the same considerations apply to the MH. First, the littoral zone is not a uniform environment. The coastline includes sand beaches, gravel beaches, rocky beaches, and sheer rock cliffs, with shoreline depths ranging from waist deep to a 10 meter plunge. Varieties of shellfish, fish, birds, and sea mammals inhabit these various microenvironments, so that particular resources are concentrated in given stretches. Access to potable water is a major problem today, although a few seeps are reported along the Bahía de la Independencia, and brackish water can be found below the dry river beds of the Ica and Grande rivers behind the shore. Seasonal storms lash the coast making cliffs dangerous for deep water fishing and egg gathering. Unpredictable events such as El Niños and

red tides (toxic algal blooms) shut down fisheries. In spite of such challenges EH and LIP villages flourished on this coastline for centuries.

The common factor shared by the large EH sites of El Chucho and Karwa with LIP La Yerba is location – they are situated on (or immediately adjacent to) extensive sandy beaches. The principal constituents of the midden (mound) deposits at all three sites are species of clams harvested in abundance from the sandy sublittoral. These sites also contain gastropods and bivalves from the rocky sublittoral in addition to sea mammal bones and domesticated plant foods, but the bulk of the deposits are composed of clam shells. Why were El Chucho and Karwa abandoned at the beginning of the EIP? Were the clam beds exhausted due to environmental disasters or over-harvesting, did the fresh water seeps fail, or fish migrations shift? Why did a permanent village the size of La Yerba not appear at the mouth of the Río Ica before the LIP?² Whatever factors may have been involved (and human agency must be considered among them), the settlement pattern data demonstrate that, for the Nasca, the return on harvesting marine resources did not warrant the energy investment required to maintain permanent shoreline occupations. Nonetheless, a considerable amount of protein can be obtained and preserved during short-term, seasonal visits.

Marine Subsistence Technology

The apparent absence of permanent, year-round Nasca fishing villages along the oceanfront suggests fishing and littoral gathering activities – both of which undeniably occurred – were conducted during temporary visits to the shore. Another approach to estimating the importance of these pursuits is to examine the archaeological evidence for marine subsistence technology. Fishing gear requires maintenance, which is best undertaken where repair materials are plentiful,

and when free time allows. Prized implements are more likely to be kept with the owner rather than abandoned in a distant cache. Such considerations, combined with the fact that the ocean is a one to two day walk from most Nasca villages, suggests the possibility that gear accompanied its owners back to their villages after a fishing trip.

On the Peruvian coast from Preceramic times onward, fishing technology included cotton nets, stone weights, gourd floats, cotton line, and fishhooks fashioned from shell, bone, and cactus-spines (Quilter and Stocker 1983:548; Bird and Hislop 1985:224-225; Pozorski and Pozorski 1987:14-15). If fishing was an important activity for the EIP Nasca, some evidence of such artifacts might be expected. However, none has been reported from excavated habitation sites, and no fishing gear (nets, sinkers, gourd floats, hooks, fish line, or harpoons) was present among the contents of 213 Nasca tombs documented by Carmichael (1995). Nasca graves often contain utilitarian items such as spear-throwers and darts, obsidian knives and points, cactus-spine needles, yarn, spindles and spindle whorls, combs, gourd bowls, baskets, and clothing (Carmichael 1988:483-499). The absence of artifacts associated with fishing is notable. Fragments of netting have been reported as surface finds (Proulx 2007:9), including knotted and looped netting (Proulx 2006: 176), but it is not clear whether these are hand-sized pieces of net bags or parts of full-size fishing nets. Ceramic iconography graphically illustrates the presence of fish nets (Lapiner 1976: 214; Proulx 2006: 177), perhaps as large as the traditional Peruvian *atarraya* or circular cast net (Marcus 1987:16-17). Marcus (ibid: 17) provides an example of one found in a Late Intermediate Period (LIP) tomb. We anticipate similar nets will be found eventually in secure Nasca contexts. However, there is no suggestion in the iconography of larger hanging nets (*red de cortina*) which require floats on the surface and weights on the bottom. Historically, hanging nets were strung between two rafts (ibid: 18). Nasca water-craft

have never been identified archaeologically. Iconography suggests the use of one-man floats, perhaps made from an inflated animal skin (Proulx 2006:123).

Shell and Fish Remains

Shell remains from crabs, chitons, sea urchins, barnacles, mussels, clams, and many other mollusks from both sandy and rocky littoral habitats frequently dot the surface of inland sites in the Ica-Nazca region. One might assume shellfish were a constant and important dietary element throughout prehistory, but recent field work documents variations in shell densities through time, with EIP Nasca sites containing sparse remains compared to deposits from earlier and later periods (Cook and Parrish 2005:140; Beresford-Jones 2011:92). Controlled excavations at EIP sites consistently reveal the shell component to be minor and scattered (Van Gijsegheem 2004:298; Cook and Parrish 2005:140; Vaughn and Linares 2006:602; Vaughn 2009:133; Beresford-Jones et al. 2011:280). Some specimens are found to be edge-worn, cut, perforated, or with traces of red pigment on exterior and interior surfaces (Rodríguez de Sandweiss 1993:295). Mussel shells appear to have been used as spoons. Mollusks were also used to make beads for necklaces and bracelets, which were worn by men, women, and children (Carmichael 1988:485-486). Rodríguez de Sandweiss concluded that, in some features excavated at Cahuachi, shell remains were not a byproduct of subsistence, but rather represent ritual offerings (1993:298). While shell remains are present at many inland sites, there is no evidence to suggest shellfish played a significant role in the Nasca diet.

Tangible evidence of fish at inland sites is more equivocal, as fish bones are delicate, may be boiled down, and on small species like anchovies be entirely consumed. Fish remains are sometimes mentioned in the literature as present at excavated sites, but seldom quantified. Van

Gijseghem (2004:297) included fish bone in a list of faunal recoveries from La Puntilla (Late Paracas – Nasca 1), and Rodríguez de Sandweiss (1993:296) lists several species found at Cahuachi. Neither researcher calculated MNIs, indicating minor representation. Valdez alone reports abundant fish heads in some excavations at Cahuachi, which he identifies as *Odontesthes regia regia* (*pejerry*), a small fish known to school periodically at river mouths (1988:148, 150). Apparently these finds were localized, as Silverman's excavations (1993) did not encounter significant fish remains. Cahuachi is well known as the preeminent centre for Early Nasca ceremonial feasting (Silverman 1993; Valdez 1994; Orefici 2012), but fish bone is not mentioned at village sites of the period such as Upanca (Vaughn and Linares 2006) or Marcaya (Vaughn 2009). We are left pondering the extent to which fish may or may not have contributed to the Nasca diet.

The evidence summarized thus far does not support a strong reliance on marine resources. It may be argued the paucity of fishing gear and limited shellfish and fish remains at inland sites is due to sampling error or differential preservation, and that large food surpluses can be generated from temporary sea shore encampments. If only the meat from mollusks and fish was transported inland (having been sun-and-wind dried on the beach) there would be few tangible traces in the archaeological record. For these reasons we have investigated another avenue of inquiry, the stable carbon, nitrogen and sulfur isotope values of the Nasca people themselves.

Isotope Evidence

Stable isotope analysis can provide extremely useful dietary information, but application of the technique is limited to situations in which subsistence issues involve isotopically distinct food groups. Stable light isotopes such as carbon, nitrogen and sulfur have been used in the past to

explore two main issues surrounding dietary choice: 1) relative reliance on C₃ versus C₄ plants, especially maize and 2) relative reliance on marine versus terrestrial resources.

Carbon isotopes can be used to distinguish between three groups of plants because of the different photosynthetic pathways used. C₃ plants include most trees, shrubs, temperate-zone grasses and the majority of domesticated Andean plants (potatoes, squash, quinoa, beans) and have $\delta^{13}\text{C}$ values in the range of -33‰ to -23‰ (O'Leary 1988; Tieszen and Boutton 1989; Tieszen 1991; Sharp 2007). C₄ plants are characterized by plants that grow in arid, warm conditions and include many grasses as well as maize and kiwicha (a pseudo-cereal similar to quinoa) with $\delta^{13}\text{C}$ values ranging typically from -16‰ to -9‰, although the values can be as low as -21‰ (Hatch and Slack 1966; O'Leary 1988; Sharp 2007; Tieszen 1991). Crassulacean Acid Metabolism (CAM) plants, notably cacti and succulents, can have $\delta^{13}\text{C}$ values that span the range of C₃ and C₄ plants, although their isotope value will often reflect the type of environment that they grow in, for example CAM plants in arid hot environments will have a $\delta^{13}\text{C}$ value similar to C₄ plants (Eickmeier and Bender 1976; Cadwallader et al. 2012). There is a slight fractionation of carbon isotopes as the food chain is ascended of approximately +1‰ (DeNiro and Epstein 1978).

Nitrogen isotope values are useful in distinguishing between different trophic levels, with the $\delta^{15}\text{N}$ value increasing due to fractionation as the food chain is ascended by approximately 3‰-5‰ at each step (Bocherens and Drucker 2003; O'Connell et al. 2012). Thus herbivores, omnivores and carnivores can be distinguished, and the degree of carnivory estimated (Ambrose 1993, 2000; O'Connell and Hedges 1999). It is important to note that $\delta^{15}\text{N}$ values can be elevated in plants as a response to aridity or can be artificially raised due to the addition of fertilizer

(Ambrose 1991). Thus it is important to have faunal isotope values to act as a baseline when interpreting human data in order to account for these possible effects.

Together, carbon and nitrogen are useful in distinguishing between the consumption of marine and freshwater resources. Whilst the $\delta^{15}\text{N}$ value will be elevated for both of these diets (allowing its identification from terrestrial resources) a more enriched $\delta^{13}\text{C}$ value (similar to those of a C_4 diet) will identify it as a marine diet as opposed to a freshwater diet which has much lower values (Schoeninger and DeNiro 1984).

Sulfur isotope ratios ($\delta^{34}\text{S}$) are used as another potential indicator of marine resource consumption. $\delta^{34}\text{S}$ values of the marine environment are typically in the region of 15‰ to 20‰ (Zhao et al. 2003; Hoefs 2004) whilst terrestrial sources are ^{34}S depleted and in the range of 2‰ to 8‰ (Schwarcz 1991; Macko et al. 1999; Privat et al. 2007). Sulfur isotopes do not fractionate as they ascend the food chain and thus the values in human tissues should be directly equivalent of the foods consumed (Kennedy and Krouse 1989; González-Martín et al. 2001; Richards et al. 2003). However, the sea-spray effect means that terrestrial plant grown near the coast can have sulfur isotope values similar to marine resources (Richards et al. 2003). Thus sulfur isotopes can also indicate proximity to the sea. There have been few Andean archaeological studies which have incorporated $\delta^{34}\text{S}$ analyses (Kelley et al. 1989; Aufderheide et al. 1994; Macko et al. 1999; Horn et al. 2009), largely due to the fact that the sample of choice is hair or soft tissue, since the sulfur content of bone is very low.

Materials and Methods. Carbon, nitrogen and sulfur isotope values were determined for a large sample of human remains (N=75) derived from the Kroeber collection, which was assembled in 1926 and is housed at the Field Museum of Natural History in Chicago. Well-provenienced

specimens of bone, hair and soft tissue (skin) were selected from gravelots recovered from seven sites along the Río Nazca: Agua Santa, Aja, Cahuachi, Cantayo, Majoro Chico, Ocongalla, and Soisongo (Figure 4; and see Kroeber and Collier 1998; Schreiber 1998). The sample spans the EIP and includes a small number of MH and LIP burials (Table 2). It is restricted to adults, but incorporates both males and females and persons of different social status.³ While bone and soft tissue specimens represent interred individuals, the hair sample also incorporates wigs worn by the deceased and hair bundles included in the grave goods. On the basis of forensic examination, we believe that multiple hair samples found in a single grave can be attributed to different individuals (Carmichael et al. n.d.).

Sample preparation techniques used at the University of Calgary followed accepted methods for the various materials analyzed and the isotopes involved. Bone collagen was extracted as per the methods of Sealy and Van der Merwe (1986). Hair and skin specimens were washed in acetone (to remove any lipids), followed by distilled water, freeze-dried and analyzed as bulk samples. The hair can be regarded as pure keratin once it has been cleaned in this manner whereas the skin will be a mixture of collagen, keratin, elastin and other minor components (Odland 1991; Wenstrup et al. 1991). Due to the different turnover times of the tissues, bone collagen gives approximately a lifetime dietary average (Hedges et al. 2007; Rummel et al. 2007), whereas hair represents the last few months of life (Saitoh et al. 1969; Tobin 2005). The different components in skin turnover at different rates and some have been shown to not turnover at all (El-Harake et al. 1998; Ritz-Timme et al. 2003; Babraj et al. 2005), but as collagen is the predominant component the diet represented by the isotope signal is approximately that consumed in the last few months of life.

Specimens for carbon and nitrogen analysis (all tissues types) were combusted in a Carlo-Erba carbon-nitrogen analyzer, and their isotope ratios assessed in a VG Isogas Prism mass spectrometer. Samples for sulfur isotope analysis were Parr-bombed, sulfur retrieved as barium sulfate, and analyzed in a VG 602 mass spectrometer. Isotope ratios are expressed using the delta (δ) notation and are expressed in ‰ relative to international standards, such as VPDB for carbon and AIR for nitrogen (Hoefs 2004). The original data is now unavailable and as such machine precision cannot be given.

During the original analysis in 1990, atomic carbon to nitrogen ratios were used to assess the integrity of the specimens. Carbon and nitrogen values for all tissues were rejected if the C/N ratio of the specimen fell outside the acceptable range of 2.9-3.6 (DeNiro 1985). This value is based on bone collagen and subsequently an acceptable C/N ratio for hair keratin has been published as 3.0-3.8 (O'Connell and Hedges 1999). Given the similarity in the ranges for bone collagen and keratin we assume here that the keratin results are valid. The C/N ratio of skin collagen, the major component of skin, should be theoretically the same as bone collagen due to the similarities in the collagen composition (Odland 1991; Wenstrup et al. 1991) thus the use of 2.9-3.6 as a quality control assessment here is regarded as appropriate in the circumstances. Not all of the human remains yielded useable results. It is unknown whether there are any paired results in the data set.

Results

Only mean stable isotope results and their standard deviations are available and are summarized in Table 3.

Carbon and Nitrogen. The three tissue types are largely comparable, based on the standard deviations and the spread of the data, although the carbon isotope values from the bone collagen are narrower. The means for the bone and hair carbon and nitrogen values are in good agreement when the tissue spacing between collagen and keratin is taken into consideration (O'Connell et al. 2001). The skin (n=5) results are broadly similar to the other tissues and as expected based on tissue spacing studies (DeNiro and Epstein 1978, 1981; Lyon and Baxter 1978; O'Connell et al. 2001; Tieszen et al. 1983; White and Schwarcz 1994). There is no indication that there is a change in diet towards the end of life in general for the population analyzed.

The carbon isotope values suggest that a mixed C₃/C₄ diet was consumed, with a significant reliance on C₄ foods. Without faunal remains it is difficult to know whether this C₄ signal in humans is the result of animals consuming either wild C₄ foods, such as grasses which are abundant on the coast, or domesticated C₄ foods, such as maize or kiwicha (Cadwallader et al. 2012). Again, without a faunal baseline, interpretation of the $\delta^{15}\text{N}$ values is slightly problematic in terms of defining the types and relative quantities of meat consumed. However, comparing this data to published faunal data from similar sites (Horn et al. 2009; Cadwallader 2013) and to modern studies of marine isotope values (Tieszen and Chapman 1992), our results suggest that terrestrial meat was the main source of protein and that marine foods were not consumed in any significant quantity (see Figure 5).

Sulfur. The hair and the skin sulfur results (n=10 for both) are broadly similar. These low values can be interpreted as stemming from a terrestrial diet with little or no input from marine resources, especially when viewed in light of resources with known sulfur values from the coast (Horn et al. 2009). They also indicate that no foods were consumed that had been subject to the sea-spray effect (unsurprising given the distance from the sea) nor had any terrestrial meat

sources been foddered on marine plants such as seaweed (see Figure 6). Thus these results fully support the carbon and nitrogen data.

To assess the possibility of temporal variation in types of foods consumed by the Nasca, we divided our sample into two groups: specimens from EIP 2-4 and EIP 5-8 (Table 4). Carbon and sulfur isotope values for the early material are slightly lower than for later material, while nitrogen values are essentially the same. The difference in carbon collagen values is statistically significant for the early versus late EIP ($t(12)=-2.8$ $p < .05$)⁴, although actually this increase could be attributed to natural variation within the background isotopic resources, especially as the early material derives from different sites than the later sample.

Several MH and LIP specimens were also included in our Field Museum sample. MH values are in the same range as those from Nasca sites, while LIP values are slightly enriched suggesting a somewhat greater importance of maize and possibly marine resources as has been seen elsewhere (Cadwallader 2013), although this result is based on only one sample (Table 4).

Finally, we can ask how the Nazca Valley data compare with those from other south coast valleys during the EIP. Figure 7 plots our bone collagen data against other contemporaneous populations from the south coast region. There is a striking similarity between all the data, although the Ica Valley population is less enriched in $\delta^{13}\text{C}$ (Cadwallader 2013). Importantly for this study none of the populations show any sign of a reliance on marine resources based on the isotopic data.

Discussion

Isotope data demonstrates the Nasca subsistence economy was firmly based on agriculture with maize probably the principal crop. Marine resources played a minor role in the regular diet, a

finding confirmed in other isotope studies (Kellner and Schoeninger 2008:236; Horn et al. 2009:192; Cadwallader 2013:187; Webb, et al. 2013:133-135). However, seafood could have provided relief during prolonged droughts, and mollusks may have been a seasonal source of protein (Erlandson 1988), which would not be identified in bulk analyses of bone collagen. The isotopic analysis suggests that terrestrial animal protein was somewhat frequently eaten although there is a paucity of faunal isotopic evidence for this period and region. This is in contrast to faunal analyses which suggest it played a minor dietary role.

The results of the isotope study complement other lines of evidence. The absence of Nasca sites on the oceanfront, the thin and scattered nature of marine remains at inland sites, and the paucity of fishing gear beyond fragments of netting can no longer be assumed to reflect preservation or sampling biases, but accurately indicate the limited dietary importance of marine resources. While the frequency of marine motifs in the iconography parallels the seafood component of the diet in EIP 2-4, isotope data indicate marine input remained relatively constant throughout the EIP (Table 4). In EIP 5-8, depictive, primary motifs showing fish and fishermen are rare (Table 1). Hence, the iconographic and isotopic data do not agree. In addition, isotope values suggest that maize composed a significant portion of the Nasca diet, although an indirect dietary route (i.e. humans eating camelids with a C₄ signal) cannot be ruled out without future faunal isotopic analyses (Cadwallader et al. 2012). Horn et al. (2009:192) also found that maize played a major role in subsistence, while Kellner and Schoeninger refer to it as the main plant staple (2008:236). Maize has also been found in significant quantities in the archaeobotanical record (Silverman 1993; Valdez 1994; Piacenza 2005; Beresford-Jones et al. 2011). However, as primary motifs on pottery, depictions of maize were absent from the plant category in Carmichael's sample of 447 vessels (Table 1 and see Carmichael 1992:188). In Proulx's (1968)

Ica sample of EIP 3-4 pottery they represent only 4% of plant motifs. Other plants such as beans, peppers and fruits are more than five times as common. We argue motif frequencies have no relation to economic reality, and do not reflect mundane or daily experience.⁵

The Nasca were familiar with the ocean. They certainly gathered mollusks and engaged in some form of net fishing. How might these activities be envisaged? While encamped at the mouth of the Rio Ica in October and November 1989, Carmichael observed trucks dropping off groups of fisherman from inland areas (Ocucaje, the cities of Ica and Pisco, and even a few from the sierra). They set up small, reed mat huts, or ramadas. In mid-October there were about 20 of these ramadas, sheltering 50-60 people, mostly adult males, but a few wives and children among them. There were a few commercial teams of six or eight men, but most were independents, and spent more time enjoying the beach than working it. These people brought the barest necessities – a sweater, blanket, and a plastic bowl, cup, and spoon for each, along with a few tin cooking pots (unlike the ancient inhabitants of nearby La Yerba who brought their best serving ware and large ceramic storage vessels – investments in permanent residency). The modern fishermen stayed for several weeks, harvesting macha (*Mesodema donacium*) on the sandy shores, steaming open the shells, and sun-drying the meat for transport inland. Many sacks of dried clams were prepared and loaded on returning trucks. Farther north in the Bahía de la Independencia, Carmichael occasionally encountered teams of net fishermen from Ica who camped for a week or two, patiently waiting for schooling fish to come close to sandy shores. Carmichael's survey study concluded that the Nasca likely favored small, temporary sea shore encampments of a similar nature to those erected by fishermen today (1998).

Earlier studies which assumed fishing to be an integral part of the Nasca economy relied heavily on the maritime potential argument. But proximity to the ocean (or any resource for that

matter) does not mean it will be maximally exploited. The data presented here lead us to concur with Alfred Kroeber's assessment that while Nasca was a coastal culture, the Nasca were not a littoral people (1944:24). There is no evidence of an "economic symbiosis" between littoral and inland communities. The Nasca were an agrarian based coastal society in which marine resources played a relatively minor role in the dietary economy, despite apparent opportunity, subsistence logic, and iconographic indicators. We conclude Nasca use of the littoral was seasonal, transitory, and opportunistic. But the sea was a powerful force in Nasca mythology, the abode of fierce divinities and, as Mama Cocha – mother of all water, the most basic necessity of life in a desert.

Why do we find variation over time in the intensity of littoral exploitation along the South Coast, and what does this tell us about the cultures of this region? We cannot assume the factors which favored large EH settlements like El Chucho and Karwa in the Bahía de la Independencia were necessarily the same for LIP La Yerba at the Ica River mouth. From one perspective all three sites have immediate access to sand beaches, and the shell middens appear to be composed primarily of corresponding clam species (however, shell type ratios should be verified through formal malacological analysis from strata cuts). However, settlement patterns actually vary significantly, from bayside (El Chucho), to peninsula (Karwa) and estuary (La Yerba) locations, and from a series of linear mounds (El Chucho and Karwa) to a few roughly circular mounds (La Yerba). Does mound shape and number reflect the organization of social groups in these societies? Why are clusters of linear mounds not found south of the Bahía? What is the relationship of permanent littoral villages to inland communities?

Large oceanfront sites represent significant energy expenditure in long-term, if not year-round habitation. Clearly, the littoral zone provided significant return on labour investment

during the EH and LIP periods. Therefore the littoral zone can yield abundant food resources when environmental and cultural conditions are favorable.

During the EIP and MH the shores were not abandoned. Traffic continued, but there was less of it and for shorter times than in earlier and later periods. We predict solid evidence of EIP and MH littoral activities will be found (the Ica River estuary is a strong candidate). Our point here is not the simple presence/absence of a few sherds or radiocarbon dates, but the extent to which people of these periods exploited maritime resources. Construction of sedentary villages with materials and artifacts lugged across the desert is a strong measure of economic importance, or social imperative if valley politics were adverse. Absence of such investments turns our attention back to the social and economic arrangements of inland dwellers.

Use of the littoral zone is a reflection of interior conditions. We cannot fully illuminate the prehistory of the South Coast without taking both regions into account. The immediate mystery – why was the littoral more important in some periods than in others – is answerable. An agenda for future research should explore both natural and cultural environments. We need deep excavation at the Bahía and Ica estuary sites. Quantitative malacological analysis may reveal changes in species composition indicative of fluctuating sea temperatures, bottom structure, or economic specialization. Natural marine deposits in the immediate area of these sites should be cored for baseline standards. Archaeobotanical and faunal analyses will be essential in isolating the marine component, and documenting shore/inland interactions.

It will be instructive to compare archaeological remains at the estuaries of the Río Ica and Río Grande de Nazca, and in their nearby oases (both named Monte Grande). La Yerba should be compared with other LIP sites along the coast, especially the large village-mound of Laguna Grande at the north end of the Bahía (Engel 1981:71; Carmichael 1998:29, 72). For the Bahía

EH sites, we need to know if El Chucho and Karwa are sequential, overlapping, or contemporary occupations, and a formal comparison of site layouts, construction methods, midden constituents, and artifacts would be a major contribution. Garcia and Pinilla (1995) made an excellent start in setting Bahía archaeology in its regional context, and we eagerly await the results of new studies currently underway by the Paracas Archaeological Project (Dulanto et al 2013).

Answers to the foregoing questions and approaches will shed light on the factors – both environmental and cultural – which favored permanent villages in the littoral zone. Conversely, such findings will highlight elements not present during periods when littoral use was transitory. This is a form of indirect evidence reflecting on inland settlements during the EIP, which narrows the parameters of our inquiries into Nasca society.

From the first Native colonists to the Inca, Andean peoples continue to surprise us with their diversity of lifeways and organizational patterns. In the current example, relatively slight use of marine resources during the EIP along the Ica-Nazca littoral contrasts with earlier and later subsistence systems in the same area, and with earlier, contemporary, and later patterns elsewhere along the Peruvian coast. All forms represent adaptations to the physical and social environments of a given place and time. These conditions are dynamic on micro as well as macro levels. Static models applied diachronically and synchronically mask the highly nuanced tapestries of Andean cultures.

Acknowledgements

The authors express their gratitude to the Field Museum of Natural History, Chicago, and the Social Sciences and Humanities Research Council of Canada, for providing grants in support of

this research. Carmichael and Kennedy are especially grateful to their co-author, Lauren Cadwallader, for re-invigorating the isotope analysis.

¹ *“Near the mouth of Ica river, five minutes’ walk from the sea, there are two splendid sambaquis or shellmounds, each about 180 m. long from east to west, 100 m. wide, 50 m. high. These appear to be the first discovered on the Peruvian coast, and to resemble those of the southern Brazilian coast. Their bases may be low natural elevations, but probably 40 m. or more of their depth is almost pure shell kitchen-midden, only slightly mixed with sand. About a dozen varieties of marine mollusks are distinguishable. On these mounds I found erect house posts, insignificant adobe wall remains, fragments of textiles and wooden slings, a stone ax fragment, chips of stones, llama bones, whale vertebrae, and seaweeds, all well preserved on the surface. Also there were strewn about broken human leg bones, suggesting that they had been cracked for marrow. Potsherds indicated the civilization of the surface as the last pre-Inca one. On account of the expense of digging mounds as large as these, dynamiting to their interior was considered but given up as unfeasible”* (Uhle 1924:123 [Appendix A – Notes on Ica Valley, Field Reports 1899-1903]). Proulx (1970:31) dates Uhle’s observations to February 1901, citing a letter written at Ocucaje in the same month of that year, in which Uhle also relates he only abandoned the dynamite idea because some acquaintances convinced him the soil was too soft for good results.

Uhle made a small collection of artifacts at the mouth of the Ica River, which is now maintained at the Phoebe A. Hearst Museum of Anthropology, University of California, Berkeley. In a footnote to Uhle’s report above, Kroeber and Strong (1924:123) describe a sherd sample from the shell mounds and a nearby cemetery, stating, *“There are about a dozen sherds in the Museum (4-4671, 4673). Some of these are Late Ica; some almost certainly Middle Ica;*

and one or two suggest *Proto-Nazca* influence. One is incised.” Carmichael’s examination of Uhle’s handwritten Field Catalogue III, p.48 [28], (Phoebe A. Hearst Museum of Anthropology, University of California, Berkeley, Microfilm 65-11043), adds the following from the shell mounds at the river mouth:

4673a: broken leg bones

4673b: kinds of shell and crayfish represented there

4673c: sea-weed etc.

4673e,f,g,h,i: five woolen slings, probably used for catching sea-birds

4673k: ball of woolen yarn [letter j not used in catalogue]

4673l: fragments of pottery, proving that the remains found on the surface of the Sambaquis belonged to the later Chincha period.

4673 [m]: fragment of stone axe

² There is a large Preceramic site called Morro La Gringa on the north side of the Ica River mouth, across from La Yerba. It was excavated by Engel (1981:19-20) who defined three separate areas and a series of occupations. At the south end Engel encountered dwellings encircled by posts held in place by split rocks and cobbles, with walls formed of mats. This is the same area where he discovered part of a log rafted yielding a radiocarbon date corrected to circa A.D. 1,000, and Carmichael (1998:86-87) found LIP sherds on the surface. Morro La Gringa appears to have been a village seasonally reoccupied over a long period (Preceramic remains are definitely present), with some reuse roughly contemporary with La Yerba on the south bank.

³ This data is no longer available and therefore age and sex comparisons of the results are not possible.

⁴ This assumes the data to be parametric; however, the original data is not available, so this assumption cannot be tested nor can non-parametric statistical tests.

⁵ This finding in turn has implications for another “sacred cow” of Nasca archaeology, the notion that warfare was on the rise in Late Nasca because depictions of warriors and trophy heads increased (Roark 1965:56). Trophy heads, for example, certainly do increase as a percentage of primary motifs in Middle and Late Nasca (Table 1), but disagreement arises over what they represent – trophies of war, ritual sacrifices, or ceremonial combat (compare Proulx 1989 with Carmichael 1994; also see DeLeonardis 2000; Conlee 2007; Knudson et al. 2009). As shown here, marine and plant motif frequencies in Roark’s Referential Motif Category (assumed to reflect daily reality) are not congruent with archaeological evidence. If ceramic motifs are removed from the warfare assumption, what evidence remains? There are no fortifications, sites are not defensively located, and there are no examples of burned or destroyed sites. Some forms of conflict may have been more prevalent in Late Nasca times, but cultural reconstructions that begin with the assumption of inter-village or inter-valley warfare as a “fact” based on iconography alone are on shaky ground. Other, independent lines of evidence, such as trauma incidence in skeletal populations should be pursued (e.g. Kellner 2002; Tomasto 2009).

References Cited

Allen, Catherine J.

- 1981 The Nasca creatures: Some problems of iconography. *Anthropology* 5:43-70.
Department of Anthropology, State University of New York at Stony Brook.

Ambrose, Stanley H.

- 1991 Effects of diet, climate and physiology on nitrogen isotope abundances in terrestrial foodwebs. *Journal of Archaeological Science* 18(3):293-317.
- 1993 Isotopic analysis of paleodiets: Methodological and interpretive considerations. In *Investigations of Ancient Human Tissue: Chemical Analyses in Anthropology*, edited by M. K. Sandford, pp. 59-130. Gordon and Breach, Philadelphia.
- 2000 Controlled diet and climate experiments on nitrogen isotope ratios of rats. In *Biogeochemical Approaches to Paleodietary Analysis*, edited by S. H. Ambrose and M. A. Katzenberg, pp. 243-259. vol. Advances in Archaeological and Museum Science. Kluwer Academic/ Plenum Publishers, New York.

Aufderheide, Arthur C., Marc A. Kelley, Mario Rivera, Luz Gray, Larry L. Tieszen, Elysha Iversen, H. Roy Krouse, and Alvaro Carevic

- 1994 Contributions of Chemical Dietary Reconstruction to the Assessment of Adaptation by Ancient Highland Immigrants (Alto Ramirez) to Coastal Conditions at Pisagua, North Chile. *Journal of Archaeological Science* 21(4):515-524.

Babraj, John A., Daniel J. R. Cuthbertson, Kenneth Smith, Henning Langberg, Benjamin Miller, Michael R. Krogsgaard, Michael Kjaer, and Michael J. Rennie

- 2005 Collagen synthesis in human musculoskeletal tissues and skin. *American Journal of Physiology - Endocrinology and Metabolism* 289:864-869.

Beresford-Jones, David

- 2011 *The Lost Woodlands of Ancient Nasca*. The British Academy, Oxford University Press.

Beresford-Jones, David, Helen Lewis, and Steve Boreham

- 2009 Linking cultural and environmental change in Peruvian prehistory: Geomorphological survey of the Samaca Basin, Lower Ica Valley, Peru. *Catena* 78: 243-249.

Beresford-Jones, David G., Oliver Q. Whaley, Carmela Alarcón, and Lauren Cadwallader

- 2011 Two millennia of changes in human ecology: Archaeobotanical and invertebrate records from the lower Ica Valley, south coast Peru. *Vegetation History and Archaeobotany* 20(4):273-292.

Bird, Junius B. and John Hyslop

- 1985 The Preceramic Excavations at the Huaca Prieta Chicama Valley, Peru. *Anthropological Papers of The American Museum of Natural History*, Volume 62 Part 1:1-294.

Blasco, Concepción Bosqued and Luis J. Ramos Gómez

-
- 1980 *Cerámica Nazca*. Valladolid, Spain: Seminario Americanista de la Universidad de Valladolid.
- Bocherens, Hervé and Dorothée Drucker
2003 Trophic level isotopic enrichment of carbon and nitrogen in bone collagen: Case studies from recent and ancient terrestrial ecosystems. *International Journal of Osteoarchaeology* 13(1-2):46-53.
- Browne, David M. and José Pablo Baraybar
1988 An archaeological reconnaissance in the province of Palpa, Department of Ica, Peru. In *Recent Studies in Pre-Columbian Archaeology*, edited by N.J. Saunders and O. de Montmollin, 299-325. BAR International Series 421.
- Burger, Richard L. and Nikolaas J. van der Merwe
1990 Maize and the origin of highland Chavín civilization: An isotopic perspective. In *American Anthropologist* 92:85-95.
- Cadwallader, Lauren, David G. Beresford-Jones, Oliver Q. Whaley, and Tamsin C. O'Connell
2012 The signs of maize? A reconsideration of what $\delta^{13}\text{C}$ values say about palaeodiet in the Andean region. *Human Ecology* 40(4):487-509.
- Cadwallader, Lauren
2013 *Investigating 1500 Years of Dietary Change in the Lower Ica Valley, Peru, Using an Isotopic Approach*. Ph.D. dissertation, Department of Archaeology and Anthropology, University of Cambridge.
- Carmichael, Patrick H.
1988 *Nasca Mortuary Customs: Death and Ancient Society on the South Coast of Peru*. Ph.D. dissertation, Department of Archaeology, University of Calgary. University Microfilms International, No.8918465, Ann Arbor, Michigan.
1992 Interpreting Nasca iconography. In *Ancient Images, Ancient Thought: The Archaeology of Ideology*, edited by A. Shawn Goldsmith, Sandra Garvie, David Selin, and Jeanette Smith, pp.187-197. Proceedings of the 23rd Annual Chacmool Conference, Department of Archaeology, University of Calgary.
1994 The life from death continuum in Nasca imagery. *Andean Past* 4:81-90.
1995 Nasca burial patterns: Social structure and mortuary ideology. In *Tombs for the Living: Andean Mortuary Practices*, edited by Tom D. Dillehay, pp. 161-188. Dumbarton Oaks, Washington.
1998 Prehistory of the Ica-Nazca Littoral, Peru. (unpublished monograph).
2013 Regionalism in Nasca Style History. *Andean Past* 11: 213-227.
- Carmichael, Patrick H., Brenda V. Kennedy and Janice L. Lacapra
n.d. *Unbraiding the Past: A Forensic Examination of Nasca Hair Artifacts*. Unpublished manuscript.
- Chiou, Katherine L., Anita G. Cook, and Christine A. Hastorf

-
- 2013 Flotation versus dry sieving archaeological remains: A case history from the Middle Horizon southern coast of Peru. *Journal of Field Archaeology* 38 (1): 38-53.
- Cobo, Bernabé
1990 *Inca Religion and Customs. (original 1653)*. Translated and edited by Roland Hamilton, University of Texas Press, Austin.
- Conlee, Christina
2007 Decapitation and Rebirth: A Headless Burial from Nasca, Peru. *Current Anthropology* 48 (3):438-445.
- Conlee, Christina A., Michele R. Buzon, Aldo Noriega Gutierrez, Antonio Simonetti, and Robert A. Creaser
2009 Identifying foreigners versus locals in a burial population from Nasca, Peru: an investigation using strontium isotope analysis. *Journal of Archaeological Science*, 36: 2755-2764.
- Cook, Anita
1994 *Investigaciones de Reconocimiento Arqueológico en la Parte Baja del Valle de Ica: Informe Final 1988- 1990*. Vols. I-II. Report to the Instituto Nacional de Cultura, Lima, Peru.
- Cook, Anita G., and Nancy Parrish
2005 Gardens in the desert: Archaeobotanical analysis from the Lower Ica valley, Peru. *Andean Past* 7: 135-156.
- Cordy-Collins, Alana
1998 Survey Report for the Archaeological Zone of Carhua, Peru. *Prehistory of the Ica-Nazca Littoral, Peru*, pp. 33-37. Patrick H. Carmichael (unpublished monograph).
- DeLeonardis, Lisa
2000 The Body Context: Interpreting Early Nasca Decapitated Burials. *Latin American Antiquity*, 11 (4):363-386.
- DeNiro, Michael J.
1985 Postmortem preservation and alteration of *in vivo* bone collagen isotope ratios in relation to palaeodietary reconstruction. *Nature* 317:806-809.
- DeNiro, Michael J. and Christine A. Hastorf
1985 Alteration of $^{15}\text{N}/^{14}\text{N}$ and $^{13}\text{C}/^{12}\text{C}$ ratios of plant matter during the initial stages of diagenesis: Studies utilizing archaeological specimens from Peru. *Geochimica et Cosmochimica Acta* 49(1):97-115.
- DeNiro, Michael J. and Samuel Epstein

-
- 1978 Influence of diet on the distribution of carbon isotopes in animals. *Geochimica et Cosmochimica Acta* 42(5):495-506.
- 1981 Influence of diet on the distribution of nitrogen isotopes in animals. *Geochimica et Cosmochimica Acta* 45(3):341-351.
- Dulanto, Jahl, Carla Marquez, and Alejandro Rey de Castro
- 2013 Political and Economic Dynamics of Maritime Communities of the South Coast of Péru During the First Millenium BC: The Excavations of the Paracas Archaeological Project at Disco Verde and Puerto Nuevo. Paper presented at 32nd Annual Meeting of the Northeast Conference on Andean Archaeology and Ethnohistory, November 16-17, 2013, Yale University, New Haven, Connecticut.
- Eickmeier, William G. and Margaret M. Bender
- 1976 Carbon isotope ratios of Crassulacean Acid Metabolism species in relation to climate and phytosociology. *Oecologia* 25(4):341-347.
- El-Harake, Wassim A., Mikhail Furman, A., Brian Cook, K. Sreekumaran Nair, Jayme Kukowski, and Irwin G. Brodsky
- 1998 Measurement of dermal collagen synthesis rate *in vivo* in humans. *American Journal of Physiology - Endocrinology and Metabolism* 37(4):586-591.
- Engel, Frédéric-André
- 1966 *Paracas: Cien Siglos de Cultura Peruana*. Libreria-Editorial Juan Mejia Baca. Lima.
- 1981 *Prehistoric Andean Ecology, Vol. II: The Deep South*. Humanities Press and the Dept. of Anthropology, Hunter College, City University of New York.
- 1991 *Un desierto en tiempos prehispánicos. Río Pisco, Paracas, Río Ica*. Centro de Investigación de Zonas Aridas. Universidad Nacional Agraria. La Molina, Lima.
- Ericson, Jonathon E, Michael West, Charles H. Sullivan, and Harold W. Krueger
- 1989 The development of maize agriculture in the Viru valley, Peru. In *The Chemistry of Prehistoric Human Bone*, edited by T. Douglas Price, pp. 68-104.
- Erlandson, Jon M.
- 1988 The role of shellfish in prehistoric economies: A protein perspective. *American Antiquity* 53(1):102-109.
- Feldman, Robert A.
- 2009 Talking Dogs and New Clothes, or the Maritime Foundations Hypothesis Revisited. In *Andean Civilization*, edited by Joyce Marcus and Patrick Ryan Williams, pp. 89-98, Monograph 63, Cotsen Institute of Archaeology, University of California, Los Angeles.
- García Soto, Rubén, and José Pinilla Blenke
- 1995 Aproximación a una secuencia de fases con cerámicas tempranas de la región de Paracas. *Journal of the Steward Anthropological Society*, 23 (1-2): 43-81.

-
- González-Martín, Inmaculada, Claudio González Pérez, Jesús Hernández Méndez, and Carlos Sánchez González
2001 Differentiation of dietary regimene of Iberian swine by means of isotopic analysis of carbon and sulphur hepatic tissue. *Meat Science* 58:25-30.
- Hatch, Marshall D. and Charles R. Slack
1966 Photosynthesis by sugar-cane leaves: A new carboxylation reaction and the pathway of sugar formation. *Biochemical Journal* 101(1):103-111.
- Hedges, Robert E. M., John G. Clement, C. David L. Thomas, and Tamsin C. O'Connell
2007 Collagen turnover in the adult femoral mid-shaft: Modeled from anthropogenic radiocarbon tracer measurements. *American Journal of Physical Anthropology* 133:808-816.
- Hoefs, Jochen
2004 *Stable Isotope Geochemistry*. 5th ed. Springer, Berlin.
- Horn, Peter, Stefan Hölzl, Susanne Rummel, Göran Åberg, Solveig Schiegl, Daniela Biermann, Ulrich Struck, and Andreas Rossmann
2009 Humans and camelids in river oases of the Ica-Palpa-Nasca region in pre-Hispanic times: Insights from H-C-N-O-S-Sr isotope signatures. In *New Technologies for Archaeology: Multidisciplinary Investigations in Palpa and Nasca, Peru*, edited by M. Reindel and G. A. Wagner, pp. 173-192. Springer-Verlag, Berlin.
- Kanter, John and Kevin J. Vaughn
2012 Pilgrimage as costly signal: Religiously motivated cooperation in Chaco and Nasca. *Journal of Anthropological Archaeology* 31:66-82.
- Kelley, Marc A., Dianne R. Levesque, and Eric Weidl
1989 Contrasting patterns of dental disease in five early northern Chilean groups. In *Advances in Dental Anthropology*, edited by M. A. Kelley and C.S. Larsen, New York.
- Kellner, Corina
2002 *Coping with Environmental and Social Changes in Prehistoric Peru: Bioarchaeological Analysis of Nasca Populations*. Ph.D. dissertation, University of California, Santa Barbara.
- Kellner, Corina M., and Margaret J. Schoeninger
2008 Wari's imperial influence on local Nasca diet: The staple isotope evidence. *Journal of Anthropological Archaeology*, 27:226-243.
- Kennedy, Brenda V. and H. Roy Krouse
1989 Isotope fractionation by plants and animals: Implications for nutritional research. *Canadian Journal of Physiological Pharmacology* 68:960-972.

-
- Knudson, Kelly J., Sloan R. Williams, Rebecca Osborn, Kathleen Forgey, and Patrick Ryan Williams
2009 The geographic origins of Nasca trophy heads using strontium, oxygen, and carbon isotope data. *Journal of Anthropological Archaeology*, 28: 244-257.
- Kroeber, Alfred L.
1944 *Peruvian Archaeology in 1942*. Viking Fund Publications in Anthropology vol.4, Wenner-Gren Foundation for Anthropological Research, New York.
- Kroeber, Alfred L. and William Duncan Strong
1924 *The Uhle Pottery Collections From Ica*.
University of California Publications in American Archaeology and Ethnology, Vol. 21, No. 3, pp. 95-133.
- Kroeber, Alfred L. and Donald Collier
1998 *The Archaeology and Pottery of Nazca*, Peru, edited by Patrick H. Carmichael, Altamira Press, Walnut Creek, California.
- Lapiner, Alan
1976 *Pre-Columbian Art of South America*. New York: Abrams.
- Long, Eric, Richard Sweitzer, Duane Diefenbach, and Merav Ben-David
2005 Controlling for anthropogenically induced atmospheric variation in stable carbon isotope studies. *Oecologia* 146(1):148-156.
- Lumbreras, Luis G.
1983 *The Peoples and Cultures of Ancient Peru*. Smithsonian Institution Press, Washington. (sixth edition, original 1974).
- Lyon, T. D. B. and M. S. Baxter
1978 Stable carbon isotopes in human tissues. *Nature* 273(5665):750-751.
- Mächtle, B. and B. Eitel
2013 Fragile landscapes, fragile civilizations – How climate determined societies in the Pre-Columbian south Peruvian Andes. *Catena* 103:62-73.
- Macko, Stephen A., Michael H. Engel, Vladimir Andrushevich, Gert Lubec, Tamsin C. O'Connell, Robert E. M. Hedges, R. P. Ambler, B. Sykes, J. B. Griffiths, J. Bada, and G. Eglinton
1999 Documenting the diet in ancient human populations through stable isotope analysis of hair [and discussion]. *Philosophical Transactions: Biological Sciences* 354(1379):65-76.
- Marcus, Joyce

-
- 1987 *Late Intermediate Occupation at Cerro Azul, Peru*. University of Michigan Museum of Anthropology, Technical Report 20. Ann Arbor, Michigan.
- Massey, Sarah A.
1986 *Sociopolitical Change in the Upper Ica Valley, B.C. 400 to A.D. 400: Regional States on the South Coast of Peru*. Ph.D. dissertation, Dept. of Anthropology, University of California at Los Angeles, University Microfilm International, No.8621115, Ann Arbor, Michigan.
- Moseley, Michael E.
1975 *The Maritime Foundations of Andean Civilization*. Cummings Publishing Co., California.
- Nieves, Ana
2010 Tipología y cronología del arte ruprestre del Valle de Nasca y la Cuenca del Río Grande de Nasca, Departamento de Ica, Perú. *Quelca Rumi*, 1 (1): 17-31.
- O'Connell, Tamsin C. and Robert E. M. Hedges
1999 Investigations into the effect of diet on modern human hair isotopic values. *American Journal of Physical Anthropology* 108(4):409-425.
- O'Connell, Tamsin C., Robert E. M. Hedges, M. A. Healey, and H. R. W. Simpson
2001 Isotopic comparison of hair, nail and bone: Modern analyses. *Journal of Archaeological Science* 28:1247-1255.
- O'Connell, Tamsin C., Catherine J. Kneale, Nataša Tasevska, and Gunter G. C. Kuhnle
2012 The diet-body offset in human nitrogen isotopic values: A controlled dietary study. *American Journal of Physical Anthropology* 149(3):426-434.
- Odland, George F.
1991 Structure of the skin. In *Physiology, Biochemistry, and Molecular Biology of the Skin*, edited by L. A. Goldsmith, pp. 3-62. vol. 1. Oxford University Press, Oxford.
- O'Leary, Marion H.
1988 Carbon isotopes in photosynthesis. *BioScience* 38(5):328-336.
- ONERN
1971 Inventario, Evaluación y Uso Racional de los Recursos Naturales de la Costa: Cuenca del Río Grande. Vols. 1 & 2. Oficina Nacional de Evaluación de Recursos Naturales, Presidencia de la República, República del Perú, Lima.
- Orefici, Giuseppe
2012 *Cahuachi: Capital Teocrática Nasca*. Universidad de San Martín de Porres.
- Piacenza, Luigi

-
- 2005 Evidencias botánicas en asentamientos Nasca. *Boletín Museo de Arqueología y Antropología* 5(1):3-13.
- Paul, Anne
- 1991 Paracas: An ancient cultural tradition on the south coast of Peru. In *Paracas Art and Architecture*, edited by Anne Paul, 1-34. University of Iowa Press, Iowa City.
- Pozorski, Thomas and Sheila Pozorski
- 1987 *Early Settlement and Subsistence in the Casma Valley, Peru*. University of Iowa Press, Iowa City.
- 2012 Preceramic and Initial Period Monumentality within the Casma Valley of Peru. In *Early New World Monumentality*, Richard L. Burger and Robert M. Rosenswig (eds.), pp. 364-398, University Press of Florida.
- Privat, Karen L., Tamsin C. O'Connell, and Robert E. M. Hedges
- 2007 The distinction between freshwater- and terrestrial-based diets: methodological concerns and archaeological applications of sulphur stable isotope analysis. *Journal of Archaeological Science* 34(8):1197-1204.
- Proulx, Donald
- 1968 Local differences and time differences in Nasca pottery. *University of California Publications in Anthropology*, 5. University of California Press, Berkeley.
- 1970 Nasca Gravelots in the Uhle Collection from the Ica Valley, Peru. *Research Reports Number 5, Department of Anthropology, University of Massachusetts, Amherst*.
- 1983 The Nazca style. In *Art of the Andes: Pre-Columbian Sculptured and Painted Ceramics from the Arthur M. Sackler Collections*, ed. L. Katz, 87-105. Washington, D.C.: Arthur M. Sackler Foundation.
- 1989 Nasca Trophy Heads: Victims of Warfare or Ritual Sacrifice? In *Cultures in Conflict: Current Archaeological Perspectives*, edited by Diana Claire Tkaczuk and Briolan C. Vivian, pp.73-85. Proceedings of the 20th Annual Chacmool Conference, Department of Archaeology, University of Calgary.
- 2006 *A Sourcebook of Nasca Ceramic Iconography*. University of Iowa Press.
- 2007 Settlement Patterns and Society in South Coastal Peru: Report on a Survey of the Lower Rio Nasca and Rio Grande, 1998. Unpublished monograph, pdf version.
- Quilter, Jeffrey and Terry Stocker
- 1983 Subsistence Economies and the Origins of Andean Complex Societies. *American Anthropologist* 85:545-562.
- Quilter, Jeffrey, Bernardino Ojeda, Deborah Pearsall, Daniel Sandweiss, John Jones, and Elizabeth Wing
- 1991 The Subsistence Economy of El Paraíso, Peru. *Science* 251:277-283.
- Reindel, Markus
- 2009 Life at the edge of the desert – archaeological reconstruction of the settlement history in the valleys of Palpa, Peru. In *New Technologies for Archaeology: Multidisciplinary*

Investigations in Palpa and Nasca, Peru, edited by M. Reindel and G.A. Wagner, pp.439-461. Springer.

Reinhard, Johan

1988 *The Nasca Lines: A New Perspective on their Origin and Meaning* (4th edition). Editorial Los Pinos, Lima.

Richards, Mike P., Benjamin T. Fuller, Matt Sponheimer, Todd Robinson, and Linda Ayliffe

2003 Sulphur isotopes in palaeodietary studies: a review and results from a controlled feeding experiment. *International Journal of Osteoarchaeology* 13(1-2):37-45.

Ritz-Timme, Stefanie, I. Laumeier, and Matthew J. Collins

2003 Aspartic acid racemization: evidence for marked longevity of elastin in human skin. *British Journal of Dermatology* 149(5):951-959.

Roark, Richard P.

1965 From Monumental to Proliferous in Nasca pottery. *Ñawpa Pacha* 3:1-92.

Rodríguez de Sandweiss, María del Carman

1993 Malacological analysis. In *Cahuachi in the Ancient Nasca World*, by Helaine Silverman, pp. 294-299. University of Iowa Press, Iowa City.

Rostworowski de Diez Canseco, María

1981 *Recursos Naturales Renovables Y Pesca, Siglos XVI Y XVII*. Instituto de Estudios Peruanos, Lima.

Rowe, John H. and Dorothy Menzel (editors)

1973 *Peruvian Archaeology: Selected Readings* (5th edition: original 1967). Peek Publications, Palo Alto.

Rummel, Susanne, Stefan Hölzl and Peter Horn

2007 Isotopensignaturen von Bio- und Geo-Elementen in der Forensik. In *Biologische Spurenkunde, Bd. 1 Kriminalbiologie*, edited by B. Herrmann and K.-S. Saternus, pp. 381-407. Springer, Berlin.

Saitoh, Masaji, Makoto Uzuka, and Masao Sakamoto

1969 Rate of hair growth. In *Advances in Biology of Skin*, edited by W. Montagna and R. L. Dobson, pp. 183-201. vol. IX. Pergamon Press, Oxford.

Sandweiss, Daniel H.

2009 Early Fishing and Inland Monuments: Challenging the Maritime Foundations of Andean Civilization? In *Andean Civilization*, Joyce Marcus and Patrick Ryan Williams (eds.), pp. 39-54, Monograph 63, Cotsen Institute of Archaeology, University of California, Los Angeles.

-
- Sandweiss, Daniel H., James B. Richardson III, Elizabeth J. Reitz, Jeffrey T. Hsu, and Robert A. Feldman
1989 Early maritime adaptations in the Andes: Preliminary studies at the Ring Site, Perú. In *Ecology, Settlement and History in the Osmore Drainage Peru*, edited by D. Rice, C. Stanish, and P. R. Scarr, 35-84, BAR International Series 545, Part I.
- Sealey, Judith C. and Nikolaas J. van der Merwe
1986 Isotope assessment and the seasonal mobility hypothesis in the southwestern Cape of South Africa. *Current Anthropology* 27(2): 135-150.
- Schoeninger, Margaret J. and Michael J. DeNiro
1984 Nitrogen and carbon isotopic composition of bone collagen from marine and terrestrial animals. *Geochimica et Cosmochimica Acta* 48(4):625-639.
- Schreiber, Katherine J.
1998 Afterword: Nasca research since 1926 (1997). In *The Archaeology and Pottery of Nazca, Peru*. Alfred L. Kroeber and Donald Collier, edited by Patrick H. Carmichael. pp. 261-270. Altamira Press, Walnut Creek, California.
1999 Regional approaches to the study of prehistoric empires. In *Settlement Patterns in the Americas: Fifty Years since Viru*. Brian Billman and Gary Feinman (eds.), pp. 160-171. Smithsonian Institution Press, Washington.
- Schreiber, Katharina and Josué Lancho Rojas
2003 *Irrigation and Society in the Peruvian Desert: The Puquios of Nasca*. Lexington Books, New York.
- Schwarcz, Henry P.
1991 Some theoretical aspects of isotope paleodiet studies. *Journal of Archaeological Science* 18(3):261-275.
- Sealy, Judith C. and Nikolaas J. Van Der Merwe
1986 Isotope assessment and the seasonal mobility hypothesis in the southwestern Cape of South Africa. *Current Anthropology* 27(2):135-150.
- Sharp, Zachary
2007 *Principles of Stable Isotope Geochemistry*. Pearson Education, Inc., Upper Saddle River.
- Silverman, Helaine
1986 *Cahuachi: An Andean Ceremonial Center*. Ph.D. dissertation. Dept. of Anthropology, University of Texas at Austin. University Microfilms International, No.8706104, Ann Arbor.
1993 *Cahuachi in the Ancient Nasca World*. University of Iowa Press, Iowa City.
2002 *Ancient Nasca Settlement and Society*. University of Iowa Press, Iowa City.
- Thornton, Erin K., Susan D. Defrance, John Krigbaum, and Patrick R. Williams

-
- 2011 Isotopic evidence for Middle Horizon to 16th century camelid herding in the Osmore Valley, Peru. *International Journal of Osteoarchaeology* 21(5):544-567.
- Tieszen, Larry L., Thomas W. Boutton, K. G. Tesdahl, and N. A. Slade
1983 Fractionation and turnover of stable carbon isotopes in animal tissues: Implications for $\delta^{13}\text{C}$ analysis of diet. *Oecologia* 57:32-37.
- Tieszen, Larry L.
1991 Natural variations in the carbon isotope values of plants: Implications for archaeology, ecology, and paleoecology. *Journal of Archaeological Science* 18(3):227-248.
- Tieszen, Larry L. and Thomas W. Boutton
1989 Stable carbon isotopes in terrestrial ecosystem research. In *Stable Isotopes in Ecological Research*, edited by P. W. Rundel, J. R. Ehleringer and K. A. Nagy, pp. 167-195. Springer-Verlag, New York.
- Tieszen, Larry L. and Michael Chapman
1992 Carbon and nitrogen isotopic status of the major marine and terrestrial resources in the Atacama Desert of northern Chile. *Proceedings of the First World Congress on Mummy Studies*, pp. 409-426. Museo Arqueológico y Etnográfico de Tenerife, Tenerife.
- Tobin, Desmond J.
2005 The biogenesis and growth of human hair. In *Hair in Toxicology: An Important Bio-Monitor*, edited by D. J. Tobin, pp. 3-33. RSC Publishing, Cambridge.
- Tomast Cagigao, Elsa
2009 Talking Bones: Bioarchaeological analysis of individuals from Palpa. In *New Technologies for Archaeology: Multidisciplinary Investigations in Palpa and Nasca, Peru*, edited by M. Reindel and G.A. Wagner, pp.141-158. Springer.
- Townsend, Richard F.
1985 Deciphering the Nazca world: Ceramic images from ancient Peru. *Museum Studies* 2(2):116-139. The Art Institute of Chicago.
- Uhle, Max
1924 Appendix A: Notes on Ica Valley. In *The Uhle Pottery Collections From Ica*, pp. 121-123. Alfred L. Kroeber and William Duncan Strong, University of California Publications in American Archaeology and Ethnology, Vol. 21, No. 3, pp. 95-133.
- Urton, Gary
1982 *Report of Fieldwork in Nazca, Peru*. manuscript.
- Valdez, Lidio M.
1988 Patrones de Subsistencia Nasca: Una Perspectiva desde Kawachi y Tambo Viejo. Tesis presentado para optar la Licenciatura en Arqueología, Universidad de Huamanga,

-
- Ayacucho, Perú.
1994 Cahuachi: New evidence for an Early Nasca ceremonial role. *Current Anthropology* 35(5):675-679.
- Van Gijsegem, Hendrik
2004 Migration, Agency, and Social Change on a Prehistoric Frontier: The Paracas-Nasca Transition in the Southern Nasca Drainage, Peru. Ph.D. dissertation, University of California, Santa Barbara.
- Vaughn, Kevin J. and Moisés Linares Grados
2006 3,000 Years of Occupation in Upper Valley Nasca: Excavations at Upanca. *Latin American Antiquity* 17(4):595-612.
- Vaughn, Kevin J.
2009 *The Ancient Andean Village: Marcaya in Prehispanic Nasca*. University of Arizona Press, Tucson.
- Webb, Emily, Christine White, and Fred Longstaffe
2013 Dietary shifting in the Nasca region as inferred from the carbon- and nitrogen-isotope compositions of archaeological hair and bone. *Journal of Archaeological Science* 40(1):129-139.
- Wenstrup, Richard J., Saood Murad and Sheldon R. Pinnell
1991 Collagen. In *Physiology, Biochemistry, and Molecular Biology of the Skin*, edited by L. A. Goldsmith, pp. 481-508. Oxford University Press, Oxford.
- White, Christine D. and Henry P. Schwarcz
1994 Temporal trends in stable isotopes for Nubian mummy tissues. *American Journal of Physical Anthropology* 93(2):165-187.
- Wilson, Andrew S., Timothy Taylor, Maria C. Ceruti, Jose A. Chavez, Johan Reinhard, Vaughan Grimes, Wolfram Meier-Augenstein, Larry Cartmell, Ben Stern, Michael P. Richards, Michael Worobey, Ian Barnes, and M. Thomas P. Gilbert
2007 Stable isotope and DNA evidence for ritual sequences in Inca child sacrifice. *Proceedings of the National Academy of Sciences of the United States of America* 104(42):16456-16461.
- Zhao, F. J., J. S. Knight, Z. Y. Hu, and S. P. McGrath
2003 Stable sulfur isotope ratio indicates long-term changes in sulfur deposition in the Broadbalk Experiment since 1845. *Journal of Environmental Quality* 32:33-39.




Figure 1:
 South Coast of Peru with Carmichael's (1998) survey boundaries, archaeological sites, towns, and river valleys mentioned in the text.

PERIODS AND EPOCHS		TIME
Colonial		
Late Horizon		AD 1476 - 1534
8	Late Intermediate Period	AD 1000 - 1476
7		
6		
5		
4		
3		
2		
1		
4	Middle Horizon	AD 700 - 1000
3		
2		
1		
8	Early Intermediate Period	AD 1 - 700
7		
6		
5		
4		
3		
2		
1		
10	Early Horizon	700 BC - AD 1
9		
8		
7		
6		
5		
4		
3		
2		
1		
	Initial Period	1800 - 700 BC
	Preceramic	Prior to 1800 BC

Figure 2:

Traditional South Coast chronology (Pisco – Acari) adapted from Rowe and Menzel 1973. Numbers denote epochs (units of time) and corresponding style phases. Absolute dates are those used for Carmichael’s survey of the Ica-Grande littoral (Carmichael 1998:7).

SOUTHERN NASCA REGION (SNR) NAZCA - LAS TRANCAS VALLEYS			
EPOCHS Units of Time & Cultural Patterns	ABSOLUTE TIME	CERAMIC PHASES Units of Style	PERIODS
LORO	AD 600		MIDDLE HORIZON
	LATE NASCA		AD 500
MIDDLE NASCA	AD 400		
EARLY NASCA	AD 300		
	AD 200		
	AD 100		
PROTO-NASCA	AD 1		1
	AD -100	L A T E P A R A C A S	

Figure 3:

Current chronology for the Southern Nasca Region (Nasca – Las Trancas Valleys). For discussion see *Appendix: Revised Chronology* in Carmichael (2013).




Figure 4:

Location of the cemeteries from which Carmichael (1992) drew his pottery sample for motif analysis (Table 1), and also Kroeber's Nasca sites from which specimens were obtained for isotope analysis (Kroeber and Collier 1998).




Figure 5:

Mean carbon and nitrogen isotope value of bone collagen (black diamond) from the Kroeber collection (n=14; error bars are to 1σ) plotted against background isotope values for south coast plants and animals. Data for plant values is based on Cadwallader et al. (2012), DeNiro and Hastorf (1985), Thornton et al. (2011) and Tieszen and Chapman (1992). Animal data is based on Cadwallader (2013), Horn et al. (2009) and Tieszen and Chapman (1992). Modern data has been corrected for the Suess effect using Long et al. (2005). The dashed boxes on top of the C₃ and C₄ plant ranges represent the elevated nitrogen values expected for water stressed plants. Modified from Cadwallader (2013).




Figure 6:

Sulphur and carbon isotope values from hair samples compared against marine and terrestrial resources. Plant and animal data is taken from Horn et al. (2009) and Macko et al. (1999). Comparative human hair samples are taken from Aufderheide et al. (1994) - Pisagua mummies (northern Chile); Horn et al. (2009) - EIP mummies (Palpa, Peru); Macko et al. (1999) - Morro mummies (northern Chile); and Wilson et al. (2007) - Llullaillaco mummies (highland Argentina).




Figure 7:

Comparison of carbon and nitrogen bone collagen isotope data with other EIP south coast sites.

Table 1. Carmichael Nazca Basin Sample – Motif Frequencies for EIP

	EARLY (EIP 2–4)	LATE (EIP 5–8)
REFERENTIAL		
Marine	20 = 10.6%	2 = 0.7%
Birds	32 = 16.9%	2 = 0.7%
Plants	34 = 18.0%	5 = 1.9%
Artifacts	2 = 1.0%	5 = 1.9%
Trophy Heads	6 = 3.2%	22 = 8.5%
Animals	5 = 2.6%	12 = 4.6%
Human	7 = 3.7%	16 = 6.2%
Warriors	0 = 0.0%	4 = 1.5%
CONCEPTUAL	32 = 17.0%	111 = 43.0%
ABSTRACT	51 = 27.0%	79 = 31.0%
TOTALS	189	258

Sample consists of 61 Early gravelots with 189 vessels and 77 Late gravelots with 258 vessels from the Nazca River Valley = 138 gravelots with 447 vessels. Carmichael 1988:461-464, see Figure 4.

Table 2. Nasca Diet and Subsistence Project: Human Sample

SITE	TIME PERIOD	SPECIMENS		
		Bone	Hair	Soft Tissue
Kroeber Collection				
Aja	EIP	1		
Agua Santa	EIP	1		
Cahuachi	EIP	7	12	8

Cantayo	EIP	3	7	2
	MH	2		
	LIP	1		
Majoro Chico	EIP	9	1	
	LIP	1	1	2
Ocongalla	EIP	3	2	1
Soisongo	EIP	3	2	1
	MH	1	1	1
Total		27	26	15
Other				
Monte Grande	EIP	13		

Table 3. Results of Stable Isotope Analysis			
Tissue	¹³ C (N)	¹⁵ N (N)	³⁴ S (N)
Bone	-13.3 ± 0.8 (14)	+9.8 ± 1.3 (14)	
Hair	-15.0 ± 1.5 (17)	+7.9 ± 1.2 (16)	+2.1 ± 1.8 (10)
Soft Tissue	-15.2 ± 1.6 (5)	+10.9 ± 0.9 (5)	+1.3 ± 1.8 (10)

Table 4. Temporal Variation in Stable Isotope Values of Human Bone Collagen and Hair

Time Period	¹³C (N)	¹⁵N (N)	³⁴S (N)
<i>Bone Collagen</i>			
EIP (1-4)	-13.8 ± 0.7 (6)	+9.7 ± 1.8 (6)	
EIP (5-8)	-12.8 ± 0.6 (8)	+9.9 ± 0.7 (8)	
EIP (ALL)	-13.3 ± 0.8 (14)	+9.4 ± 1.3 (14)	
MH	-13.5 ± 0.4 (3)	+9.4 ± 0.5 (3)	
LIP	-11.0 (1)	+10.6 (1)	
<i>Hair</i>			
EIP (1-4)	-14.9 ± 1.4 (14)	+7.9 ± 1.3 (13)	+2.4 ± 1.6 (7)
EIP (5-8)	-15.8 ± 1.4 (3)	+8.1 ± 0.9 (3)	+1.3 ± 1.9 (3)
EIP (ALL)	-15.0 ± 1.5 (17)	+7.9 ± 1.2 (16)	+2.1 ± 1.8 (10)
MH	-14.4 (1)	+7.7 (1)	+1.1 (1)
LIP	-14.2 (1)	+8.4 (1)	