

1  
2  
3  
4 1  
5  
6  
7 2  
8  
9  
10 3  
11  
12 Effect of a two-step natural organic acid-treatment on  
13 microbial activity and lipid damage during blue whiting  
14  
15  
16  
17  
18 6  
19  
20  
21 7  
22 8  
23 9  
24 10  
25 11  
26 12  
27 13  
28 14  
29 14  
30 15  
31 16  
32 17  
33 18  
34 19  
35 20  
36 21  
37 22  
38 23  
39 24  
40 25  
41 26  
42 27  
43 28  
44 29  
45 30  
46 31  
47 32  
48 33  
49 34  
50 35  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Effect of a two-step natural organic acid-treatment on  
microbial activity and lipid damage during blue whiting  
(*Micromesistius poutassou*) chilling

Minia Sanjuás-Rey<sup>1</sup>, Bibiana García-Soto<sup>2</sup>, Jorge Barros-Velázquez<sup>1</sup>,  
José R. Fuertes-Gamundi<sup>2</sup> and Santiago P. Aubourg<sup>3,\*</sup>

<sup>1</sup> Department of Analytical Chemistry, Nutrition and Food Science, Escuela de Ciencias Veterinarias, Universidad de Santiago de Compostela, Lugo, Spain

<sup>2</sup> Cooperativa de Armadores de Pesca del Puerto de Vigo (ARVI), Vigo, Spain

<sup>3</sup> Department of Food Technology. Instituto de Investigaciones Marinas (CSIC), Vigo, Spain

\* Correspondent: [saubourg@iim.csic.es](mailto:saubourg@iim.csic.es)

## SUMMARY

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 A novel approach was carried out to inhibit quality loss in chilled blue whiting  
4 (*Micromesistius poutassou*). For it, a natural organic acid-mixture including ascorbic,  
5 citric and lactic acids was applied in a two-step processing. First, as an aqueous dipping  
6 medium previous to chilling storage; then, present in ice employed as chilling system.  
7 According to indices related to microbiological activity (namely, aerobe and  
8 psychrotroph counts) and lipid damage development, results obtained showed a  
9 profitable role of the organic acid-mixture; thus, a partial inhibition of both damage  
10 pathways was attained, this leading to an increased shelf life time of chilled blue  
11 whiting when compared to control fish. Lipid hydrolysis (free fatty acid formation)  
12 showed to be a more relevant event than lipid oxidation (peroxide and thiobarbituric  
13 acid reactive substance formation) in chilled blue whiting. Microbiological group counts  
14 remained as acceptable values for all kinds of samples throughout a 9-day chilling  
15 study.

16  
17  
18  
19 **Running Head:** Natural organic acid and blue whiting chilling

20 **Keywords:** *Micromesistius poutassou*, ascorbic, citric, lactic, dipping, icing, microbial  
21 activity, lipid damage

## 1. INTRODUCTION

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 Deterioration of marine species begins immediately upon capture or harvest, and the  
3 degree to which it continues depends directly on storage conditions. Flake ice has been  
4 the most employed method to cool and store fish products and partially inhibit  
5 detrimental effects on the commercial value. However, significant deterioration of  
6 sensory quality and nutritional value has been detected in chilled fish as a result of  
7 different damage pathways (Whittle et al., 1990; Olafsdóttir et al., 1997). To retard fish  
8 damage as long as possible, and accordingly extend shelf life, a wide number of  
9 preservative strategies to be combined to flake ice chilling have been tested  
10 satisfactorily such as previous chemical and physical treatments (Toledo-Flores & Zall,  
11 1992; Ashie et al., 1996) and employment of preservative packaging (Ozen & Floros,  
12 2001; Sivertsvik et al., 2002).

13 Among previous chemical treatments to chilling storage, natural organic acids have  
14 shown to represent a relevant choice because of their easy availability, low commercial  
15 cost and wide range of permitted concentrations for their use. Thus, ascorbic and citric  
16 acids (AA and CA, respectively) are widely known for their role as chelators, acidulants  
17 in biological systems and synergists of primary antioxidants, so that a profitable effect  
18 on minced fish (Hwang & Regenstein, 1988; Stodolnik et al., 1992), fish fillets (Badii &  
19 Howell, 2002; Kilinc et al., 2009; Pourashouri et al., 2009) and whole fish (Aubourg et  
20 al., 2004) has been observed. Further, lactic acid (LA) has been reported to be effective  
21 in suppressing Gram-negative bacteria, which are known to be the most important fish  
22 spoiler group; thus, LA pre-treatment has shown to be effective in preserving and  
23 extending shelf-life in fish fillets (Kim et al., 1995; Metin et al., 2001; Erkan, 2003) and  
24 coated fish (Gogus et al., 2006).

1 Blue whiting (*Micromesistius poutassou*) is a gadoid fish abundant in the northeast  
2 Atlantic. Previous research has shown its profitable nutritional support to human diet  
3 (Dagbjartsson, 1975; Martínez-Valverde et al., 2000), being most efforts focused to its  
4 employment in the manufacture of restructured products (Montero et al., 1998; Pérez-  
5 Mateos et al., 2002). However, since it is also directly consumed, blue whiting  
6 suitability to freezing (Huidobro & Tejada, 1995; Aubourg et al., 2007), cold-smoking  
7 (Gómez-Guillén et al., 2009) and canning (Oehlenschläger et al., 2008) processes has  
8 also been tested. Concerning its chilled storage, previous research accounts for the  
9 nucleotide degradation and volatile amine formation (Smith et al., 1980), lipid oxidation  
10 development (Aubourg et al., 1998) and trimethylamine oxide breakdown into  
11 formaldehyde and dimethylamine (Rey-Mansilla et al., 1999).

12 In the present research, a novel approach is carried out to inhibit quality loss in chilled  
13 blue whiting. For it, a natural organic acid-mixture including AA, CA and LA was  
14 applied in a two-step processing. First, as an aqueous dipping medium previously to  
15 chilling storage; then, present in the ice employed as chilling system. Microbial activity  
16 and lipid damage development were evaluated throughout the storage time and  
17 compared to control fish.

## 18 **2. MATERIALS AND METHODS**

### 19 **2.1. The natural organic acid-mixture and preliminary trials**

20 A commercial formula (BPS2) including a natural organic acid-mixture was supplied by  
21 *Atlantic one, S. L.* (Vigo, Spain) for employment in the present research. Such a product  
22 consists of a water-soluble viscous liquid including AA, CA and LA (1 meq acid/ 120  
23 mg product) in glycerol, being regarded as safe (GRAS) for use in foods according to  
24 European and American administrations (Madrid et al., 1994; Giese, 1996).  
25

1 Preliminary trials were carried out in order to assess the most convenient product  
2 concentration to be included both in the dipping medium as well as in the icing system.  
3 For it, the effect on sensory acceptance (appearance, texture, gills, odour and colour) of  
4 a wide concentration range (70-2000 ppm) of the organic acid-mixture was checked. As  
5 a result, 800 ppm concentration showed to provide the most convenient results when  
6 employed in both processing steps. Accordingly, this concentration was chosen for  
7 being employed in the present research.

## 8 9 **2.2. Raw fish, processing and sampling**

10 Fresh blue whiting (72 individuals) were caught near the Galician Atlantic coast  
11 (northwestern Spain) in Winter 2010 and transported on ice to the laboratory. The  
12 length and weight of the fish specimens were included in the following ranges: 22-25  
13 cm and 70-120 g, respectively.

14 Upon arrival in the laboratory, eight individual fishes were separated and considered as  
15 starting raw fish (day 0); for it, two different groups (4 individuals per group) were  
16 considered and analysed independently. The remaining fish were divided into two  
17 batches (32 individuals in each batch). The first batch (preserved fish; P batch) was  
18 dipped in a 800 ppm aqueous solution of the organic acid-mixture during 2 minutes (ca.  
19 290 g fish L<sup>-1</sup> solution); after that, the fish were placed in boxes and directly surrounded  
20 by ice prepared with a 800 ppm aqueous solution of the organic acid-mixture.

21 Fish corresponding to the second batch (control fish; C batch) was dipped in water  
22 during 2 minutes (ca. 290 g fish L<sup>-1</sup> solution) and then placed in boxes surrounded by  
23 traditional ice prepared with water.

24 In both batches, the fish individuals were surrounded by ice (with or without organic  
25 acid-mixture, respectively) at a 1:1 fish-to-ice ratio. Both batches were placed in a

1 refrigerated room (4 °C). Boxes employed allowed draining and ice was renewed when  
2 required. Fish samples from the two different batches were taken for analysis on days 2,  
3 5, 7 and 9. At each sampling point, eight individuals of each batch were taken for  
4 analysis, being considered into two groups (four individuals in each group) that were  
5 studied independently.

### 7 **2.3. Microbial activity measurement**

8 Samples of 10 g of fish muscle were dissected aseptically from chilled fish specimens,  
9 mixed with 90 ml of 0.1% peptone water (Merck, Darmstadt, Germany), and  
10 homogenised in a stomacher (AES, Combourg, France) as previously described (Ben-  
11 Gigirey et al., 1998, 1999).

12 In all cases, serial dilutions from the microbial extracts were prepared in 0.1% peptone  
13 water. Total aerobes were investigated by surface inoculation in plate count agar (PCA,  
14 Oxoid Ltd., London, UK), after incubation at 30 °C for 48 h. Psychrophiles were also  
15 investigated in PCA but incubation was carried out at 7-8 °C for 7 days.  
16 *Enterobacteriaceae* were investigated by pour plating on Violet Red Bile Agar  
17 (VRBA) (Merck, Darmstadt, Germany) after incubation at 37 °C for 24 h.  
18 Microorganisms exhibiting a proteolytic or lipolytic phenotype were investigated in  
19 casein-agar medium or tributyrin-agar, respectively, after incubation at 30 °C for 48 h,  
20 as previously described (Ben-Gigirey et al., 2000).

21 In all cases, bacterial counts were transformed into log CFU g<sup>-1</sup> muscle before  
22 undergoing statistical analysis.

#### **2.4. Proximate composition and pH assessment**

Moisture content was determined by the difference between the weight of fresh homogenised white muscle (1-2 g) and the weight recorded after 4 h at 105 °C. Results were calculated as g water kg<sup>-1</sup> muscle.

Lipids were extracted from the fish white muscle by the Bligh and Dyer (1959) method, by employing a single-phase solubilisation of the lipids using a chloroform-methanol (1:1) mixture. Quantification results were calculated as g lipid kg<sup>-1</sup> muscle.

The evolution of pH values in fish muscle along storage time was determined by means of a 6-mm diameter insertion electrode (Crison, Barcelona, Spain).

#### **2.5. Lipid damage assessment**

The peroxide value (PV) was determined spectrophotometrically (Beckman Coulter, DU 640, London, UK) in the lipid extract by previous peroxide reduction with ferric thiocyanate, according to the Chapman and McKay (1949) method. Results were expressed as meq active oxygen kg<sup>-1</sup> lipids.

The thiobarbituric acid (TBA) index was determined according to Vyncke (1970). This method is based on the reaction between a trichloroacetic acid extract of the fish muscle and TBA. Content on thiobarbituric acid reactive substances (TBARS) was spectrophotometrically measured at 532 nm and results were expressed as mg malondialdehyde kg<sup>-1</sup> muscle.

Formation of fluorescent compounds (Fluorimeter LS 45; Perkin Elmer España, Tres Cantos, Madrid, Spain) was determined by measurements at 393/463 nm and 327/415 nm as described by Aubourg et al. (1998). The relative fluorescence (RF) was calculated as follows:  $RF = F/F_{st}$ , where F is the fluorescence measured at each excitation/ emission maximum, and F<sub>st</sub> is the fluorescence intensity of a quinine

1 sulphate solution ( $1 \mu\text{g ml}^{-1}$  in  $0.05 \text{ M H}_2\text{SO}_4$ ) at the corresponding wavelength. The  
2 fluorescence ratio (FR) was calculated as the ratio between the two RF values:  $\text{FR} =$   
3  $\text{RF}_{393/463 \text{ nm}} / \text{RF}_{327/415 \text{ nm}}$ . The FR value was determined in the aqueous phase resulting  
4 from the lipid extraction of the fish muscle (Bligh & Dyer, 1959).  
5 Free fatty acid (FFA) content was determined in the lipid extract of the fish muscle by  
6 the Lowry and Tinsley (1976) method based on complex formation with cupric acetate-  
7 pyridine followed by spectrophotometric (715 nm) assessment. Results were expressed  
8 as g FFA  $\text{kg}^{-1}$  lipids.

## 10 **2.6. Sensory analysis**

11 Sensory analysis was conducted by a sensory panel consisting of five experienced  
12 judges, according to guidelines concerning fresh and refrigerated fish (DOCE, 1989).  
13 Panellists had been involved in sensory analysis of different kinds of fish foods during  
14 the last 10 years. Previously to the present experiment, a special training was carried out  
15 on chilled blue whiting.  
16 Four categories were ranked (Table 1): highest quality (E), good quality (A), fair quality  
17 (B) and unacceptable quality (C). Sensory assessment of the fish included the following  
18 parameters: skin and mucus development, external odour, gills and gill cavity, eyes and  
19 flesh odour (raw and cooked fish) and taste (cooked fish). At each sampling time, the  
20 fish muscle portions were presented to panellists in individual trays and were scored  
21 individually. The panel members shared samples tested.

## 23 **2.7. Statistical analysis**

24 Data obtained from the different microbial and chemical analyses were subjected to the  
25 ANOVA method ( $p < 0.05$ ) to explore differences by two different ways: processing



1 conditions effect and chilling time effect (Statsoft, Statistica, version 6.0, 2001);  
2 comparison of means was performed using a least-squares difference (LSD) method.  
3 Correlation analysis among parameters (chilling time, microbial indices, lipid damage  
4 indices and sensory acceptance) was also carried out; for sensory values, the Spearman  
5 test was employed. Linear fittings are expressed; otherwise, the kind of fitting  
6 (quadratic or logarithmic) is mentioned.

### 8 **3. RESULTS AND DISCUSSION**

#### 9 **3.1. Microbial activity assessment**

10 The comparative evolution of total aerobes for each batch during storage time can be  
11 seen in Figure 1. In general terms, the control batch exhibited higher microbial numbers  
12 than the counterpart batch stored in the natural-organic acid system. Thus, the most  
13 remarkable differences were observed on days 2, 5 and 9, the mean differences between  
14 batches being 0.57, 3.17 and 0.48 log CFU g<sup>-1</sup> muscle, respectively. Nevertheless, in the  
15 course of the nine days of chilled storage, significant differences (p<0.05) for aerobe  
16 assessment were only observed at day 5. Concerning its evolution during the chilling  
17 storage, aerobe counts did not provide a good correlation value with chilling time.

18 The comparative evolution of psychrophilic bacteria can be seen in Figure 2. The counts  
19 in the fish batch processed under the natural-organic acid condition were lower than in  
20 the control batch from day 5 until day 9. Thus, the average differences in the counts of  
21 psychrophiles at day 5 were 1.44 log CFU g<sup>-1</sup> muscle and for the following days  
22 differences of 0.46 and 0.91 log CFU g<sup>-1</sup> muscle were determined. However, and  
23 although the natural-organic acid system exhibited a better control of bacterial growth,  
24 significant differences (p<0.05) were only obtained at day 5, as for aerobe assessment.  
25 Concerning its evolution during the chilling storage, psychrophilic counts did not

1 provide a good correlation value with chilling time; compared to aerobe formation,  
2 good correlation was only obtained for the preserved batch ( $r^2 = 0.94$ ), being bad in the  
3 case of control fish ( $r^2 = 0.77$ ).

4 *Enterobacteriaceae* and proteolytic bacteria counts did not provide a clear tendency  
5 with chilling time (Table 2); thus, all values were included in the ranges 1.0-2.0 and 2.0-  
6 4.0 log CFU g<sup>-1</sup> muscle, respectively. With respect to proteolytic bacteria, the natural-  
7 organic acid batch exhibited counts that were 0.37, 0.20 and 0.63 log CFU g<sup>-1</sup> muscle  
8 lower than the control batch at days 2, 5 and 9, respectively. Likewise, the average  
9 counts for *Enterobacteriaceae* in the batch processed in the organic acid-mixture were  
10 0.24, 0.35 and 0.30 log CFU g<sup>-1</sup> muscle lower than the control batch at days 2, 5 and 9,  
11 respectively; however, no significant ( $p>0.05$ ) differences were obtained between both  
12 batches for both microbiological assessments.

13 Lipolytic bacteria provided a constant value (2 log CFU g<sup>-1</sup> muscle) for both batches  
14 during the 0-7-day period. At the end of the experiment, lipolytic counts were included  
15 in the 2.75-2.90 range, and no significant differences ( $p>0.05$ ) were found between both  
16 batches.

17 The microbial preservative effects observed in this study were quite in accordance with  
18 the results reported by Kilinc et al. (2009), who studied the microbiology of rainbow  
19 trout (*Oncorhynchus mykiss*) fillets dipped in solutions including organic salts (acetate,  
20 lactate and citrate). Thus, these authors reported that the immersion of fillets in such  
21 solutions slowed down the proliferation of total aerobic and psychrophilic bacteria.  
22 Likewise, Marshall et al. (1996) investigated the microbiology of catfish (*Silurus*  
23 *glanis*) fillets treated by dipping in acetic acid and lactic acid solutions, and found a  
24 partial inhibition of bacterial growth under these conditions. An inhibitory effect of  
25 microbial activity was also found in chilled catfish (*Silurus glanis*) fillets that were

1  
2  
3 1 previously dipped in a lactic acid solution (Kim et al., 1995); as a result, an extension of  
4  
5 2 the shelf life time was produced. Microbial activity was also partially inhibited by  
6  
7 3 previous lactic acid treatment in chilled chub mackerel (*Scomber japonicus*) (Metin et  
8  
9 4 al., 2001) and Mediterranean mackerel (*Scomber colias*) and mullet (*Mugil cephalus*)  
10  
11 5 (Erkan, 2003) fillets according to a lower microbial count and total volatile-base  
12  
13 6 nitrogen (TVB-N) content assessment throughout a 12-day and 9-day storage period,  
14  
15 7 respectively.

16  
17  
18  
19 8 In a previous experiment (Quitral et al., 2009), ice prepared from an aqueous solution  
20  
21 9 from rosemary and oregano extracts was applied to a fatty fish species (Chilean jack  
22  
23 10 mackerel; *Trachurus murphyi*). As in the present study, the employment of ice  
24  
25 11 including a preservative compound as chilling medium led to a partial inhibition of  
26  
27 12 microbial activity, according to the TVB-N content assessment.

28  
29  
30  
31 13 In the present study, values obtained in all samples for all kinds of microbial parameters  
32  
33 14 can be considered below the limit of acceptance. This conclusion agrees to previous  
34  
35 15 research on microbial activity development during blue whiting chilling, where  
36  
37 16 trimethylamine formation did not provide a great increase throughout a 8-day period  
38  
39 17 (Smith et al., 1980; Rey-Mansilla et al., 1999) and a strong content of TVB-N content  
40  
41 18 was only obtained after a 10-13-day chilled period (Aubourg et al., 2008).

42  
43  
44  
45  
46  
47  
48  
49

### 20 **3.2. Proximate analysis and pH assessment**

50  
51 21 Moisture and lipid contents were included in the ranges 797-831 and 3.8-4.4 (g kg<sup>-1</sup>  
52  
53 22 muscle), respectively. Such values agreed to previous results concerning this fish  
54  
55 23 species (Dagbjartsson, 1975; Aubourg et al., 1998; Aubourg et al., 2007). No  
56  
57 24 differences (p>0.05) in both constituents could be assessed as a result of the  
58  
59 25 preservative treatment. However, a slight increasing moisture content (p<0.05) could be  
60

1 observed with chilling time for both preserved and control fish batches that could be  
2 explained as a result of contact with ice during the chilled storage.

3 Concerning the pH analysis, values were included in the range 6.90-7.00 for both kinds  
4 of fish samples throughout the whole experiment (Table 2). No significant differences  
5 ( $p>0.05$ ) could be observed as a result of the preservative treatment, although lower  
6 mean values were observed in the 2-7-day period for the preserved batch, this in  
7 agreement to previously mentioned results on microbiological counts. Additionally, a  
8 slight increase in both batches could be observed in the 0-5-day period, so that a fair  
9 correlation value ( $r^2 = 0.86-0.89$ ; logarithmic fitting) with chilling time could be  
10 depicted for the complete experiment.

11 Results obtained in the present study concerning pH assessment agree to previous  
12 research. Thus, a pH decrease was also obtained in chub mackerel (*Scomber japonicus*)  
13 (Metin et al., 2001) and catfish (*Silurus glanis*) (Kim et al., 1995) chilled fillets as a  
14 result of a previous lactic acid treatment. Additionally, pH value was studied in Chilean  
15 jack mackerel (*Trachurus murphyi*) that was kept under ice prepared from a water  
16 solution of rosemary and oregano extracts (Quitral et al., 2009); as a result, a lower pH  
17 increase could be observed when compared to control mackerel that was attributed to  
18 the presence in ice of preservative compounds from both plant extracts.

### 19 20 **3.3. Lipid oxidation development**

21 Lipid oxidation was studied by means of different quality indices corresponding to  
22 different steps included in the fish lipid oxidation development.

23 Primary oxidation provided very low scores, that remained in all cases under the 4.50  
24 value (Table 3). The preservative treatment did not provide a clear effect ( $p>0.05$ ) on

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 peroxide formation; additionally, a clear tendency of peroxide formation could not be  
2 obtained as a result of the chilling time for both kinds of samples.

3 Concerning the secondary lipid oxidation development, again relatively low values were  
4 obtained, since lower scores than 0.60 were obtained in all cases (Table 3). When  
5 compared to control fish, the preservative treatment provided a lower ( $p<0.05$ ) TBARS  
6 formation in chilled fish in the 7-9-day period. For both kinds of samples, a progressive  
7 increase ( $p<0.05$ ) could be depicted during the chilled storage, so that good correlation  
8 values with chilling time ( $r^2 = 0.89-0.93$ ; quadratic fitting) were obtained.

9 The fluorescent compound analysis provided a higher formation in control fish, this  
10 being significant ( $p<0.05$ ) at days 2, 7 and 9 (Table 3); such results agree with the  
11 TBARS content, so that an inhibitory effect on lipid oxidation development was  
12 concluded for the preservative treatment. Starting fish showed a lower ( $p<0.05$ ) FR  
13 value than the remaining samples for both batches; then (2-9-day period), a significant  
14 increase with chilling time increase was not observed ( $p>0.05$ ) in any of the batches  
15 under study.

16 Lipid oxidation development has been recognised as a complex process where different  
17 kinds of molecules are produced, most of them unstable, susceptible to breakdown and  
18 originate lower weight compounds (Aubourg, 1993; Howell, 1995), or react with other  
19 molecules (nucleophilic-type, mostly) present in the fish muscle; consequently, the  
20 determination of each kind of compound cannot always provide an accurate method for  
21 the quality assessment. In the present research, no reliability for the primary lipid  
22 oxidation compound assessment (PV) was concluded, while a very short interaction  
23 compound formation (FR assessment) was produced. However, secondary (TBARS  
24 assessment) lipid oxidation compound measurement showed to be helpful in order to  
25 assess the lipid oxidation development throughout the whole chilled storage period.

1 The partial inhibition of oxidation development found in the present research for chilled  
2 blue whiting as a result of the organic acid-mixture treatment in two steps agrees to  
3 previous research where AA and CA have shown a profitable antioxidant effect when  
4 applied as a preliminary treatment to further storage. Thus, CA and its salts have shown  
5 to play a synergist role with primary antioxidants and oxygen scavengers during the  
6 refrigerated (Stodolnik et al., 1992; Khalil & Mansour, 1998) and frozen (Pourashouri et  
7 al., 2009; Aubourg et al., 2004) storage of marine species. In the same way, AA and its  
8 salts have been reported to act as antioxidants in minced fish (Hwang & Regenstein,  
9 1988), fish fillets (Badii & Howell, 2002) and whole fish (Aubourg et al., 2004) because  
10 of their oxygen scavenger and reducing roles.

11 Related to the second preservative step tested in the present research, ice prepared from  
12 a water solution from rosemary and oregano extracts was applied as chilling system to  
13 Chilean jack mackerel (*Trachurus murphyi*); as a result, a lower oxidation development  
14 (PV and TBARS assessments) in jack mackerel muscle could be observed during the  
15 chilled storage, being this inhibition attributed to polyphenolic compound presence in  
16 the icing system.

#### 17 18 **3.4. Lipid hydrolysis analysis**

19 Lipid hydrolysis development was measured by the FFA content evolution (Figure 3). A  
20 lower FFA formation ( $p < 0.05$ ) could be concluded for the P-batch in the 7-9-day period  
21 when compared to control fish, so that an inhibitory effect of the preservative treatment  
22 could be concluded. For both batches, a marked FFA formation increase ( $p < 0.05$ ) could  
23 be outlined during chilling storage according to previous research (Aubourg et al., 1998;  
24 Quitral et al., 2009), so that a good correlation value with chilling time ( $r^2 = 0.91$ ;  
25 logarithmic fitting) was obtained.

1 FFA formation during chilling storage has been reported to be produced as a result of  
2 endogenous enzyme activity and microbial activity (Whittle et al., 1990; Madrid et al.,  
3 1994). Before the end of the microbial lag phase (up to 5-8 days, depending on several  
4 factors), FFA formation has been reported to be produced mostly as a result of  
5 endogenous enzyme (namely, lipases and phospholipases) activity. Later on, microbial  
6 activity should gain importance, so that FFA formation is then mostly produced as a  
7 result of bacterial catabolic processes. According to this profile, present results on FFA  
8 formation in chilled blue whiting would show a first stage of FFA formation (0-5-day  
9 period) corresponding mostly to the endogenous enzymatic activity; in this period, no  
10 significant differences ( $p>0.05$ ) between batches are found. Then (7-9-day period),  
11 when microbial activity should gain importance, fish corresponding to the P-batch  
12 remains at a relatively constant FFA level, while its counterpart C-batch fish shows a  
13 marked ( $p<0.05$ ) FFA formation, so that a preservative effect would be accorded to the  
14 organic acid treatment.

15 To our knowledge, this inhibitory effect on FFA formation is described for the first time  
16 during the chilled storage of a lean fish species. As being molecules supporting  
17 acidulant properties, all three acids can be found responsible for this microbial  
18 inhibition (Whittle et al., 1990; Ashie et al., 1996). However, this effect has been  
19 reported to be specially important in the case of LA, as being recognised as effective in  
20 suppressing Gram-negative bacteria activity, which are known to be the most important  
21 fish spoiler group (Kim et al., 1995; Metin et al., 2001; Gogus et al., 2006).

22 Additionally, FFA formation was studied in Chilean jack mackerel (*Trachurus murphyi*)  
23 that was kept under ice prepared from a water solution of rosemary and oregano extracts  
24 (Quitral et al., 2009); as a result, a lower FFA content could be observed when

1 compared to control mackerel that was attributed to the presence of preservative  
2 compounds from both plants present in the icing system.

3 Relationship between lipid hydrolysis and oxidation has captivated a great attention.  
4 FFA are known to have detrimental effects on protein properties and oxidise faster than  
5 higher molecular weight lipid classes (namely, triglycerides and phospholipids) by  
6 providing a greater accessibility (lower steric hindrance) to oxygen and other pro-  
7 oxidant molecules. In the present research, the best correlation value between FFA  
8 formation and the different lipid oxidation indices was obtained in the case of the  
9 TBARS assessment ( $r^2 = 0.83-0.87$ ).

10

### 11 **3.5. Sensory acceptance**

12 A progressive acceptance decrease could be observed for both fish batches throughout  
13 the chilled storage, so that a good correlation value was obtained in all cases with  
14 chilling time ( $r^2 = 0.90-0.92$ ; quadratic fitting). Preserved fish maintained a good quality  
15 till day 5 and were still acceptable at the end of the experiment; however, control fish  
16 was only considered as good quality till day 2 and showed a shelf life time of 7 days.

17 Limiting attributes showed to be the external odour and gills appearance and odour.

18 Present research did not provide good correlation values for sensory acceptance and the  
19 different chemical lipid oxidation parameters, according to the fact that off-odours  
20 detected by the panel were putrid and not rancid. Thus, the best correlations were  
21 obtained with the TBA value ( $r^2 = 0.72-0.86$ ). Previous research has shown a strong  
22 incidence of FFA accumulation on fish quality loss, being their presence associated in  
23 some extent to the lack of sensory acceptability and strongly interrelated with off-odour  
24 development (Ingemansson et al., 1995; Refsgaard et al., 2000). In this sense, present



1  
2  
3 1 research provided a fair correlation value of sensory acceptance with FFA formation ( $r^2$   
4 = 0.81-0.90).  
5  
6 2

7  
8 3 Present scores concerning the shelf life increase in chilled blue whiting as a result of the  
9  
10 4 two-step organic acid treatment agree to previous research. Thus, CA and AA have  
11  
12 5 shown to provide shelf life increases when employed as a pre-treatment (namely,  
13  
14 6 aqueous solution dipping) to further frozen storage conditions (Pourashouri et al.,  
15  
16 7 2009). Concerning the chilling storage, pre-treatment by dipping on a lactic acid  
17  
18 8 solution also increased the shelf life in catfish (*Silurus glanis*) (Kim et al., 1995) and  
19  
20 9 chub mackerel (*Scomber japonicus*) (Metin et al., 2001) fillets.  
21  
22  
23  
24  
25  
26  
27  
28

#### 29 **4. FINAL REMARKS**

30 12 Assurance of both high quality and safety of chilled seafood is an important actual  
31  
32 13 challenge for fish traders and food technologists, so that commercial requirements are  
33  
34 14 always related to the search for valuable and practical technologies that may provide  
35  
36 15 enhancing possibilities. The present study, focused to blue whiting chilling, provides a  
37  
38 16 first approach to the employment of a novel two-step treatment including a natural  
39  
40 17 organic acid-mixture (AA, CA and LA), such acids providing complementary  
41  
42 18 preservative properties (acidulants, antioxidants and antimicrobians). In a first step, the  
43  
44 19 acid mixture was applied in an aqueous dipping medium previously to the chilling  
45  
46 20 storage; then, the acid mixture was included in the ice employed as chilling system.  
47  
48  
49

50 21 According to quality indices related to microbiological activity (microbiological group  
51  
52 22 counts) and lipid damage (hydrolysis and oxidation) development in chilled blue  
53  
54 23 whiting, results obtained showed a profitable role of the organic acid-mixture treatment;  
55  
56 24 thus, a partial inhibition of both damage pathways was attained, this leading to an  
57  
58 25 increased shelf life time when compared to control fish.  
59  
60

1 According to the results obtained in the present research, lipid hydrolysis showed to be  
2 a more relevant event than lipid oxidation in chilled blue whiting. Thus, FFA formation  
3 provided a great increase throughout the storage period, while lipid oxidation scores  
4 (peroxides and TBARS) remained relatively low. Meantime, counts obtained for the  
5 different microbiological groups remained as acceptable values throughout the whole  
6 experiment for individual fishes corresponding to both batches.

7 Further studies focused to the positive role of this two-step strategy are envisaged. A  
8 great effort ought to be addressed towards knowledge of the organic acid range where  
9 optimal effects could be attained, this including knowledge of the endogenous  
10 antioxidant composition corresponding to the fish species to be tested (synergism  
11 possibility analysis) and towards research checking the diffusion rate of organic acids  
12 from both the dipping and the icing media to the fish muscle.

13  
14

### 15 **Acknowledgements**

16 The authors thank the owner and crew of the CACHACHO and CHANS ships for their  
17 collaboration in the present study, and the Vigo's harbour fishing fleet in general for  
18 kindly providing the blue whiting fish. The authors also thank Mr. Marcos Trigo for his  
19 excellent technical assistance and ATLANTIC ONE, S. L. (Vigo, Spain) for providing  
20 the commercial organic acid-mixture BPS2. This work was supported by the Secretaría  
21 Xeral de I+D from the Xunta de Galicia (Galicia, Spain) through the Research Project  
22 PGIDIT 08 TAL 038E.

23  
24  
25

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## REFERENCES

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- 1 Ashie, I., Smith, J. & Simpson, B. (1996). Spoilage and shelf-life extension of fresh fish and shellfish. *Critical Reviews in Food Science and Nutrition*, **36**, 87-121.
- 2  
3  
4  
5 Aubourg, S. (1993). Review: Interaction of malondialdehyde with biological molecules: New trends about reactivity and significance. *International Journal of Food Science and Technology*, **28**, 323-335.
- 6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- 8 Aubourg, S., Lago, H., Sayar, N. & González, R. (2007). Lipid damage during frozen storage of gadiform species captured in different seasons. *European Journal of Lipid Science and Technology*, **109**, 608-616.
- 9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- 11 Aubourg, S., Medina, I. & Gallardo, J. (1998). Quality assessment of blue whiting (*Micromesistius poutassou*) during chilled storage by monitoring lipid damages. *Journal of Agricultural and Food Chemistry*, **46**, 3662-3666.
- 12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- 14 Aubourg, S., Pérez-Alonso, F. & Gallardo, J. (2004). Studies on rancidity inhibition in frozen horse mackerel (*Trachurus trachurus*) by citric and ascorbic acids. *European Journal of Lipid Science and Technology*, **106**, 232-240.
- 15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- 17 Badii, F. & Howell, N. (2002). Effect of antioxidants, citrate, and cryoprotectants on protein denaturation and texture of frozen cod (*Gadus morhua*). *Journal of Agricultural and Food Chemistry*, **50**, 2053-2061.
- 18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- 20 Ben-Gigirey, B., Vieites Baptista de Sousa, J., Villa, T. & Barros-Velázquez, J. (1998). Changes in biogenic amines and microbiological analysis in albacore (*Thunnus alalunga*) muscle during frozen storage. *Journal of Food Protection*, **61**, 608-615.

- 1  
2  
3 1 Ben-Gigirey, B., Vieites Baptista de Sousa, J., Villa, T. & Barros-Velázquez, J. (1999).  
4  
5 2 Histamine and cadaverine production by bacteria isolated from fresh and frozen  
6  
7 3 albacore (*Thunnus alalunga*). *Journal of Food Protection*, **62**, 933–939.  
8  
9  
10 4 Ben-Gigirey, B., Vieites Baptista de Sousa, J., Villa, T. & Barros-Velázquez, J. (2000).  
11  
12 5 Characterization of biogenic amine-producing *Stenotrophomonas maltophilia*  
13  
14 6 strains isolated from white muscle of fresh and frozen albacore tuna.  
15  
16 7 *International Journal of Food Microbiology*, **57**, 19-31.  
17  
18 8 Bligh, E. & Dyer, W. (1959). A rapid method of total extraction and purification.  
19  
20 9 *Canadian Journal of Biochemistry and Physiology*, **37**, 911-917.  
21  
22  
23 10 Chapman, R. & McKay, J. (1949). The estimation of peroxides in fats and oils by the  
24  
25 11 ferric thiocyanate method. *Journal of the American Oil Chemists' Society*, **26**,  
26  
27 12 360-363.  
28  
29  
30 13 Dagbjartson, B. (1975). Utilization of blue whiting, *Micromesistius poutassou*, for  
31  
32 14 human consumption. *Journal of the Fisheries Research Board of Canada*, **32**,  
33  
34 15 747-751.  
35  
36  
37 16 DOCE (1989). Baremo de Clasificación de Frescura. In: *Diario Oficial de las*  
38  
39 17 *Comunidades Europeas* (L5/21, 07.01.1989). Pp. 5-6. Brussels, Belgium:  
40  
41 18 European Commission.  
42  
43  
44 19 Erkan, N. (2003). Treatment of fillets (*Scomber colias* and *Mugil cephalus*) with sodium  
45  
46 20 lactate and propyl gallate: influence on shelf life and quality. *Archives für*  
47  
48 21 *Lebensmittelhygiene*, **54**, 60-63.  
49  
50  
51 22 Giese, J. (1996). Antioxidants: Tools for preventing lipid oxidation. *Food Technology*,  
52  
53 23 **50**, 73-80.  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 1 Gogus, U., Bozoglu, F. & Yurdugul, S. (2006). Comparative effects of lactic acid, nisin,  
4  
5 2 coating combined and alone applications on some postmortem quality criteria of  
6  
7 3 refrigerated *Sardina pilchardus*. *Journal of Food Quality*, **29**, 658-671.  
8  
9  
10 4 Gómez-Guillén, C., Gómez-Estaca, J., Giménez, B. & Montero, P. (2009). Alternative  
11  
12 5 fish species for cold-smoking process. *International Journal of Food Science*  
13  
14 6 *and Technology*, **44**, 1525-1535.  
15  
16  
17 7 Howell, N. (1995). Interaction of proteins with small molecules. In: *Ingredient*  
18  
19 8 *Interactions- Effects on Food Quality* (edited by A. Gaonkar). Pp. 269-289. New  
20  
21 9 York, USA: Marcel Dekker.  
22  
23  
24 10 Huidobro, A. & Tejada, M. (1995). Alteration of the electrophoretic pattern of  
25  
26 11 myofibrillar proteins in fish mince during frozen storage. *Zeitschrift für*  
27  
28 12 *Lebensmittel- Untersuchung und -Forschung*, **200**, 247-251.  
29  
30  
31 13 Hwang, K. & Regenstein, J. (1988). Protection of menhaden mince lipids from rancidity  
32  
33 14 during frozen storage. *Journal of Food Science*, **45**, 1120-1124.  
34  
35  
36 15 Ingemansson, T., Kaufmann, P. & Ekstrand, B. (1995). Multivariate evaluation of lipid  
37  
38 16 hydrolysis and oxidation data from light and dark muscle of frozen stored  
39  
40 17 rainbow trout (*Oncorhynchus mykiss*). *Journal of Agricultural and Food*  
41  
42 18 *Chemistry*, **43**, 2046-2052.  
43  
44  
45 19 Khalil, A. & Mansour, E. (1998). Control of lipid oxidation in cooked and uncooked  
46  
47 20 refrigerated carp fillets by antioxidant and packaging combinations. *Journal of*  
48  
49 21 *Agricultural and Food Chemistry*, **46**, 1158-1162.  
50  
51  
52 22 Kilinc, B., Cakli, S., Dincer, T. & Tolasa, S. (2009). Microbiological, chemical,  
53  
54 23 sensory, color, and textural changes of rainbow trout fillets treated with sodium  
55  
56 24 acetate, sodium lactate, sodium citrate, and stored at 4 °C. *Journal of Aquatic*  
57  
58 25 *Food Product Technology*, **18**, 3-17.  
59  
60

- 1  
2  
3 1 Kim, C., Hearnberger, J. & Eun, J. (1995). Gram-negative bacteria in refrigerated  
4  
5 2 catfish fillets treated with lactic culture and lactic acid. *Journal of Food*  
6  
7 3 *Protection*, **58**, 639-643.  
8  
9  
10 4 Lowry, R. & Tinsley, I. (1976). Rapid colorimetric determination of free fatty acids.  
11  
12 5 *Journal of the American Oil Chemists' Society*, **53**, 470-472.  
13  
14  
15 6 Madrid, A., Madrid, J. & Madrid, R. (1994). Chilling, freezing and ultra-freezing of fish  
16  
17 7 and derivatives. In: *Technology of fish and its derivatives*. Pp. 45-103. Madrid,  
18  
19 8 Spain: AMV Ediciones Mundi-Prensa.  
20  
21  
22 9 Marshall, D. & Kim, C. (1996). Microbiological and sensory analyses of refrigerated  
23  
24 10 catfish fillets treated with acetic and lactic acids. *Journal of Food Quality*, **19**,  
25  
26 11 317-329.  
27  
28  
29 12 Martínez-Valverde, I., Periago, M., Santaella, M. & Ros, G. (2000). The content and  
30  
31 13 nutritional significance of minerals on fish flesh in the presence and absence of  
32  
33 14 bone. *Food Chemistry*, **71**, 503-509.  
34  
35  
36 15 Metin, S., Erkan, N., Varlik, C. & Aran, N. (2001). Extension of shelf life of chub  
37  
38 16 mackerel (*Scomber japonicus* Houttuyn 1780) treated with lactic acid. *European*  
39  
40 17 *Food Research and Technology*, **213**, 174-177.  
41  
42  
43 18 Montero, P., Pérez-Mateos, M. & Borderías, J. (1998). Chilled storage of high pressure  
44  
45 19 and heat-treated gels of blue whiting (*Micromesistius poutassou*) muscle.  
46  
47 20 *Zeitschrift für Lebensmittel- Untersuchung und -Forschung*, **207**, 146-153.  
48  
49  
50 21 Oehlenschläger, J., Karl, H., Mitchell, M., Fagan, J. & Gormley, R. (2008). Evaluation  
51  
52 22 of quality parameters of canned and frozen blue whiting (*Micromesistius*  
53  
54 23 *poutassou*). *Journal of Fisheries Sciences*, **2**, 722-732.  
55  
56  
57 24 Olafsdóttir, G., Martinsdóttir, E., Oehlenschläger, J., Dalgaard, P., Jensen, B.,  
58  
59 25 Undeland, I., Mackie, I., Henehan, G., Nielsen, J. & Nilsen, H. (1997). Methods

- 1  
2  
3 1 to evaluate fish freshness in research and industry. *Trends in Food Science and*  
4  
5 2 *Technology*, **8**, 258-265.  
6  
7  
8 3 Ozen, B. & Floros, J. (2001). Effects of emerging food processing techniques on the  
9  
10 4 packaging materials. *Trends in Food Science and Technology*, **12**, 60-67.  
11  
12 5 Pérez-Mateos, M., Solas, T. & Montero, P. (2002). Carrageenans and alginate effects on  
13  
14 6 properties of combined pressure and temperature in fish mince gels. *Food*  
15  
16 7 *Hydrocolloids*, **16**, 225-233.  
17  
18 8 Pourashouri, P., Shabanpour, B., Aubourg, S., Daghigh Rohi, J. & Shabani, A. (2009).  
19  
20 9 An investigation of rancidity inhibition during storage of Wels catfish (*Silurus*  
21  
22 10 *glanis*) fillets by previous ascorbic and citric acid treatment. *International*  
23  
24 11 *Journal of Food Science and Technology*, **44**, 1503-1509.  
25  
26 12 Quitral, V., Donoso, M<sup>a</sup>L., Ortiz, J., Herrera, M<sup>a</sup>V., Araya, H. & Aubourg, S. (2009).  
27  
28 13 Chemical changes during the chilled storage of Chilean jack mackerel  
29  
30 14 (*Trachurus murphyi*): Effect of a plant extract-icing system. *Food Science and*  
31  
32 15 *Technology*, **42**, 1450-1454.  
33  
34 16 Refsgaard, H., Brockhoff, P. & Jensen, B. (2000). Free polyunsaturated fatty acids  
35  
36 17 cause taste deterioration of salmon during frozen storage. *Journal of the*  
37  
38 18 *Agricultural and Food Chemistry*, **48**, 3280-3285.  
39  
40 19 Rey-Mansilla, M., Sotelo, C. & Gallardo, J. (1999). Decomposition of trimethylamine  
41  
42 20 oxide during iced storage of blue whiting (*Micromesistius poutassou*). *Zeitschrift*  
43  
44 21 *für Lebensmittel- Untersuchung und -Forschung*, **208**, 267-269.  
45  
46 22 Sivertsvik, M., Jeksrud, W. & Rosnes, T. (2002). A review of modified atmosphere  
47  
48 23 packaging of fish and fishery products- significance of microbial growth,  
49  
50 24 activities and safety. *International Journal of Food Science and Technology*, **37**,  
51  
52 25 107-127.  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 1 Smith, J., Hardy, R., Thomson, A., Young, K. & Parsons, E. (1980). Some observations  
4  
5 2 on the ambient and chill storage of blue whiting (*Micromesistius poutassou*). In:  
6  
7 3 *Advances in Fish Science and Technology* (edited by J. Connell). Pp. 299-303.  
8  
9 4 Farnham, Surrey, England, UK: Fishing News Books.
- 10  
11  
12 5 Stodolnik, L., Blasiak, E. & Broszedzka, H. (1992). Effect of Tween 80, citric acid and  
13  
14 6 acetylsalicylic acids on changes in muscle tissue lipids of Baltic herrings during  
15  
16 7 storage. *Chłodnictwo*, **27**, 29-35.
- 17  
18  
19 8 Toledo-Flores, L. & Zall, R. (1992). Methods for extending the storage life of fresh  
20  
21 9 tropical fish. In: *Advances in Seafood Biochemistry* (edited by G. Flick & R.  
22  
23 10 Martin). Pp. 233-243. Lancaster, PA, USA: Technomic Publishing.
- 24  
25  
26 11 Vyncke, W. (1970). Direct determination of the thiobarbituric acid value in  
27  
28 12 trichloroacetic acid extracts of fish as a measure of oxidative rancidity. *Fette*  
29  
30 13 *Seifen Anstrichmittel*, **72**, 1084-1087.
- 31  
32  
33 14 Whittle, K., Hardy, R. & Hobbs, G. (1990). Chilled fish and fishery products. In:  
34  
35 15 *Chilled foods: The state of the art* (edited by T. Gormley). Pp. 87-116. New  
36  
37 16 York, USA: Elsevier Applied Science.
- 38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



## LEGENDS TO FIGURES

**Figure 1:** Comparative aerobe count (log CFU g<sup>-1</sup> muscle) assessment\* in chilled blue whiting processed under different conditions\*\*

\* Mean values; standard deviations are denoted by bars.

\*\* Processing conditions: C (control fish; fish dipped in water and then kept in traditional ice) and P (preserved fish; fish dipped in aqueous solution including 800 ppm of the organic acid-mixture and then kept in ice prepared from an aqueous 800 ppm solution of the acid mixture).

**Figure 2:** Comparative psychrotroph count (log CFU g<sup>-1</sup> muscle) assessment\* in chilled blue whiting processed under different conditions\*\*

\* Mean values are expressed; standard deviations are denoted by bars.

\*\* Processing conditions as expressed in Figure 1.

**Figure 3:** Free fatty acid (g kg<sup>-1</sup> lipids) assessment\* in chilled blue whiting processed under different conditions\*\*

\* Mean values are expressed; standard deviations are denoted by bars.

\*\* Processing conditions as expressed in Figure 1.

**TABLE 1****Scale employed for evaluating the sensory quality of chilled blue whiting**

<b>Attribute</b>	<b>Highest quality (E)</b>	<b>Good quality (A)</b>	<b>Fair quality (B)</b>	<b>Unacceptable (C)</b>
<b>Skin and mucus development</b>	Very intense pigmentation; transparent mucus	Milky mucus; insignificant pigmentation losses	Slightly greyish mucus; pigmentation without shine	Widely opaque mucus; important pigmentation losses
<b>External odour</b>	Sharply seaweed and shellfish smell	Weakly seaweed and shellfish smell	Incipiently putrid or ammonia odour	Putrid or ammonia odour
<b>Gills and gill cavity</b>	Brightly red; lamina perfectly separated; without odour	Rose coloured; lamina adhered in groups; without odour	Slightly pale; lamina adhered in groups; incipient fishy odour	Grey-yellowish colour; lamina totally adhered; intense ammonia odour
<b>Eyes</b>	Convex; transparent cornea; bright and black pupil	Convex and slightly sunken; slightly opalescent cornea; black and cloudy pupil	Flat; opalescent cornea; opaque pupil	Concave and milky cornea; Internal organs blurred
<b>Flesh odour and taste</b>	Sharply seaweedy; shellfish smell and taste	Weakly seaweedy; shellfish smell and taste	Incipiently putrid or ammonia odour and taste	Putrid or ammonia odour and taste

**TABLE 2**

**Evolution\* of *Enterobacteriaceae* and proteolytic counts (log CFU g<sup>-1</sup> muscle) and pH values in chilled blue whiting processed under different conditions\*\***

Chilling Time (days)	<i>Enterobacteriaceae</i>		Proteolytics		pH	
	C	P	C	P	C	P
0	1.70 (0.01)		3.00 (0.06)		6.80 (0.12)	
2	1.24 (0.35)	1.00 (0.01)	3.22 (1.04)	2.85 (0.21)	6.92 (0.01)	6.85 (0.13)
5	1.50 (0.28)	1.15 (0.21)	2.50 (0.71)	2.30 (0.43)	7.00 (0.08)	6.97 (0.02)
7	0.99 (0.01)	1.73 (1.06)	1.99 (0.01)	2.80 (1.14)	6.98 (0.07)	6.96 (0.01)
9	1.93 (0.11)	1.63 (0.21)	3.58 (0.71)	2.95 (0.35)	6.96 (0.01)	6.96 (0.01)

\* Mean values; standard deviations are indicated in brackets. For each parameter and for each chilling time, no significant ( $p > 0.05$ ) differences were obtained as a result of processing.

\*\* Processing conditions: C (control fish; fish dipped in water and then kept in traditional ice) and P (preserved fish; fish dipped in aqueous solution including 800 ppm of the organic acid-mixture and then kept in ice prepared from an aqueous 800 ppm solution of the acid mixture).

**TABLE 3**

**Evolution of lipid oxidation\* in chilled blue whiting processed under different conditions\*\***

Chilling Time (days)	PV (meq active oxygen kg <sup>-1</sup> lipids)		TBA value (mg malondialdehyde kg <sup>-1</sup> muscle)		FR	
	C	P	C	P	C	P
0	1.53 (0.51)		0.06 (0.03)		0.73 (0.06)	
2	0.71 a (0.28)	1.58 b (0.43)	0.02 (0.01)	0.03 (0.01)	1.23 b (0.14)	0.83 a (0.02)
5	3.39 (1.11)	1.99 (1.18)	0.27 (0.15)	0.24 (0.03)	1.38 (0.26)	1.04 (0.26)
7	0.57 (0.44)	1.25 (0.46)	0.38 b (0.04)	0.21 a (0.05)	1.09 b (0.02)	0.86 a (0.15)
9	4.27 (0.44)	3.89 (1.38)	0.57 b (0.04)	0.39 a (0.11)	1.32 b (0.12)	0.97 a (0.17)

\* Mean values; standard deviations are indicated in brackets. For each quality index and for each chilling time, mean values followed by different letters (a, b) denote significant differences as a result of processing. No letters are indicated when significant differences are not found ( $p > 0.05$ ).

\*\* Processing conditions as expressed in Table 2.

**TABLE 4**

**Evolution of sensory quality\* in chilled blue whiting processed under different conditions\*\***

Processing condition	Chilling Storage Time (days)				
	0	2	5	7	9
C	E	A	B	B	C
P		A	A	B	B

\* Quality categories: E (highest), A (good), B (fair) and C (unacceptable).

\*\* Processing conditions as expressed in Table 2.




Figure 1  
119x77mm (300 x 300 DPI)

30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60





Figure 2  
119x77mm (300 x 300 DPI)

Review



30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

figure 3  
119x77mm (300 x 300 DPI)