

## Manuscript

**ECOLOGY OF *SEPIA OFFICINALIS*****ANGEL GUERRA***ECOBIOIMAR. Instituto de Investigaciones Marinas (CSIC). Eduardo Cabello 6. 36208 Vigo. Spain.**Corresponding author: brc1@iim.csic.es*

**ABSTRACT-** This article comprises an up-dated review of the processes influencing the distribution and abundance of common cuttlefish *Sepia officinalis*, the interactions between the species and the main parameters of the environment in which it lives and its trophic, demographic and behavioural ecology.

Key words: CUTTLEFISH, *SEPIA OFFICINALIS*, CEPHALOPODA, ECOLOGY

**RÉSUMÉ.-** Le présent article est une révision actualisée des procès qui influencent la distribution et l'abondance de la seiche *Sepia officinalis*, des interactions entre cette espèce et les principaux paramètres de l'environnement où elle habite, comme aussi sur son écologie trophique, démographique et éthologique.

Mots clés: SEICHE, *SEPIA OFFICINALIS*, CÉPHALOPODES, ECOLOGIE

The geographical distribution of the common cuttlefish, *Sepia officinalis* L. 1758 covers all the Mediterranean Sea and the waters of the Eastern Atlantic from Southern Norway and Northern England to the northwestern coast of Africa. The species also lives in Madeira and in the Canary Islands (Khromov et al., 1998). The geographical distribution of *S. officinalis* and *Sepia hierredda* Rang, 1837 in the Eastern Central Atlantic shows that these species are sympatric. The Southern boundary of *S. officinalis* coincides approximately with the border between Mauritania and Senegal (16° N) and the Northern limit of *S. hierredda* is at Cape Blanc (21°N) (Guerra et al., 2001).

***Factors influencing the distribution and abundance of Sepia officinalis***

*S. officinalis* is a nekto-benthic species occurring predominantly on sandy and muddy bottoms from the coastline (2-3 m depth) to approximately 200 m depth, with the greatest abundance in the upper 100 m. Life in inshore water exposes this species to hydrologically unstable conditions, and because of this *S. officinalis* is relatively tolerant to variations in salinity. Animals have been observed in coastal lagoons at a salinity of 27 PSU in the Mediterranean (Mangold-Wirz, 1963). Observations from the Western Mediterranean and the NW Atlantic have shown that juveniles and adults can survive for some time at salinities  $18 \pm 2$  PSU if slowly acclimatised (Boletzky, 1983; Guerra and Castro, 1988). In culture tanks both embryonic development and growth of young cuttlefish usually occurs between  $37 \pm 3$  PSU (Domingues et al., 2001). However, Paulij et al. (1990) observed that some embryos of *S. officinalis* from eggs collected in the SW Netherlands hatched at a salinity of 26.5 PSU, but no hatching occurred below 23.9 PSU, and below 22.4 PSU embryos with morphological malformations were found.

Experiments with *S. officinalis* showed that shells of large animals implode between 150 and 200 m, whereas advanced embryonic specimens and newly hatched implode between 50 and 100 m. The larger individuals are occasionally caught at depths greater than the implosion depth of the juvenile shell parts. They apparently avoid implosion of the early shell portions by refilling these first-formed chambers with cameral liquid later in life (Ward and Boletzky, 1984).

The temperature limits of the species range from 10°C to 30°C. At temperatures below 10°C the individuals do not feed, stay inactive and die in a couple of days (Richard, 1971; Bettencourt, 2000). Hatchlings and young *S. officinalis* were successfully cultured in tanks with an open sea water system in which temperature reached 30°C (Domingues et al., 2001), and indeed the species lives in the lagoon system of the Ria Formosa (South Portugal), where temperature reaches  $27 \pm 3$ °C in summer (Domingues et al., 2002). Oxygen affinity expressed by  $P_{50}$  (partial pressure of gas at which the blood remains 50% saturated) as a function of temperature for *S. officinalis* showed that it increased from 12 mm Hg at 5° C to near 38 at 17°C, the slope of the linear regression being relatively low. This is an indication that the species does not have the ability to accommodate large or small temperature ranges in its natural habitat (Brix et al. 1994). Recent findings by Melzner et al. (2004) supported the

hypothesis of an oxygen limitation due to thermal tolerance (upper limit at about 26° C) caused by limited capacity and a loss in coordination of the components of the oxygen delivery system.

Johansen et al., (1982) studied the O<sub>2</sub> uptake in relation to body weight and concluded that the common cuttlefish is not very tolerant to low oxygen concentrations. Low oxygen concentrations can account for the absence or low abundance of *S. officinalis*. Thus, a comparison between the cuttlefish fisheries in the upwelling areas off the NW African coast and the Northern Benguela current undertaken by Guerra and Sánchez (1985) suggested that continuous eutrophic scenarios in shallow waters in the cores of the southern upwelling (25-28°S) This is where low oxygen concentrations are common, and appear to be the most important limiting factor for the development of cuttlefish populations.

The physiological processes for buoyancy in the cuttlebone of *S. officinalis* determine their role as major bioenergetic consumers. As indicated by Webber et al., (2000), buoyancy and activity of *S. officinalis* are typically higher at night (VO<sub>2</sub> from 93 to 120 mg O<sub>2</sub> kg<sup>-1</sup> h<sup>-1</sup>) than during the day (VO<sub>2</sub> between 77 and 93). Within the depth limits imposed by shell implosion, the cuttlefish is less energetically expensive with depth than any fish. Thus, it can competitively occupy top niches in the trophic web of shallow waters that involve daily vertical migrations.

Landing per unit effort of cuttlefish from a time series of 18 years in SW Spain indicated that the abundance of this species did not show a correlation with rainfall rates, river discharges and sea surface temperature (Sobrino et al., 2002). This reinforces the view that the common cuttlefish exhibit high physiological flexibility, which allows a great ability to endure changing environments, not only during its adult phase but also in the early juvenile stages (Sobrino et al., 2002). Jorge and Sobral (2004) indicated that strong precipitation had a negative influence on the cuttlefish abundance in the Ría de Aveiro (Central Portugal) and that, on the contrary, the cumulative effect of high values of solar radiation, temperature of the air, water transparency and salinity near the bottom seemed to positively influence the catches of this species.

Toxic effects of heavy metals can negatively influence the distribution and abundance of *S. officinalis*. As the species is a primary food source for many predators and also for human consumption, it is a potential threat for higher trophic levels (Bustamante et al., 2002). Concentration

and distribution of heavy metals in tissues of *S. officinalis* have observed high and selective bioaccumulation (Miramand and Bentley, 1992). Ecotoxicological studies using bioassays from isolated digestive gland cells demonstrated that some heavy metals (Cu, Zn and Ag) induced high disturbance of enzymatic systems (Le Bihan et al., 2004). The impact of these metals on survival and growth of eggs and juvenile cuttlefish can be very negative (Koueta pers. comm.). Culture experiments at different stages of the life cycle of *S. officinalis* using Zn and Cd tracers with sea water, sediments and food as uptake pathways showed that food is the likely primary track for bioaccumulation, and that the digestive gland plays a major role in the subsequent storage and presumed detoxification of these elements regardless of the uptake pathway (Bustamante et al., 2002). Malformed common cuttlefish caught in the Bay of Arcachon could be a product of the teratogenic effects of the antifouling compound TBT (Schipper and Boletzky, 1998).

*S. officinalis* does not form shoals neither in the wild nor in the laboratory, but in culture they tolerate one another except under extreme food deprivation. This tolerance is higher in young animals than in subadult and adult ones (Hanlon and Messenger, 1996). A feeding hierarchy first appearing after 4 months, which stabilizes after 5 months, has been found in this species (Warnke, 1994). Captive-rearing experiments indicated that behaviour of *S. officinalis* was strongly affected by housing conditions and suggested that this species is probably semi-solitary under natural conditions (Boal et al., 1999). The results obtained when studying the effects of crowding in cuttlefish cultured at different densities are somewhat contradictory. This could be due to the difficulty of comparing experiments undertaken under different conditions. Experiments completed by Domingues et al. (2003) showed that cuttlefish cultured in isolation had higher growth and survival rates than the ones maintained at relatively high densities, even when stressed. The authors observed agonistic behaviour related with competition for space with higher densities in tanks, indicating that density, or lack of space, appear to be more limiting than isolation.

To date, very few interspecific associations (excluding parasitism) have been reported for this species. Bacterial populations associated with *S. officinalis* have been localized mainly in the accessory nidamental glands, the renal appendages and the shell epithelium. The accessory nidamental glands show an intense orange-red coloration in mature females, and this colour is due to

carotenoid pigments, which occur in symbiotic bacteria (Van Den Braden et al., 1980). Five symbiotic bacterial taxa (*Agrobacterium*, *Roseobacter*, *Rhodobium-Xanthobacter*, *Sporochtya* and *Clostridium*) were identified in the tubules of the accessory nidamental glands, and three taxa of Pseudomonaceae were located in the renal appendages and the shell epithelium. All these bacteria, except Gram-positive ones, were also present in embryos, suggesting vertical transmission, i.e. maternal transmission at egg stage (Grigioni and Boucher-Rodoni, 2002).

The copepod *Metaxymolgus longicaudata* has also been found to be associated with this cuttlefish, but its role was not elucidated (Ho, 1983). Small specimens (mantle length, ML<65 mm) of *S. officinalis* and adult *S. elegans* (ML 45-65 mm) exhibited diets with similar prey types, although in different proportions. This may suggest a trophic competition between the two species at that size range (Castro and Guerra, 1990).

Apparent replacement of finfish by *Octopus vulgaris* and *S. officinalis* in the Sahara Bank (21°N-26°N) since the 1960s was attributed to a change in the ecosystem due to overexploitation of finfish. Balguerias et al. (2000) re-evaluated the history of these fisheries and suggested that the changes in the faunistic composition of the communities were caused by a combination of factors, including economic initiatives as well as oceanographic variations and competition for food. This ultimately favoured benthic cephalopod populations at the cost of most finfish populations.

The present-day geographical distribution pattern of the Sepiidae may have been generated by a complex mosaic of factors involving palaeo-oceanographical changes (Neige, 2003). Some attempts were made to clarify the taxonomic status of the genus *Sepia* L., 1758, which comprises approximately 100 species. Khromov et al., (1998) proposed a subdivision of the genus into six species complexes, which are not to be viewed as phylogenetic entities. Khromov (1998) suggested that five main stages could be seen in Sepiid radiation. One of the scenarios proposed by Khromov suggested that the *Sepia* sensu stricto forms (to which *S. officinalis* belongs) emerged in the Palaeogene (70 to 40 MYA). Khromov's scenario also suggested that these forms underwent a relatively recent radiation starting from the warm waters of the Tethyan Sea, and that the Western Mediterranean and Southern Atlantic European forms colonized the west coast of Africa. Recent studies on the biogeography of Sepiidae showed two radiation patterns: one to the Southern African

coasts, and the other to the 'East Indian' area. The Southern African pattern is characterized by high disparity for very different species richness values. This pattern may be caused by the coexistence of two independent phylogenetic clusters of species, one from the Atlantic and the other from the Indian Ocean. This has to be viewed in the paleogeographical context of the Eocene (60-40 MYA), where the Tethyan Sea was still open at its eastern end providing a connection between Europe, on the one hand, and the Indian Ocean and east African coast, on the other. At the end of the Eocene, this eastern corridor between the Mediterranean and Western India disappeared, involving a huge transformation in possible routes for cuttlefish migration. This could have produced two clusters of species, one in Europe and along the west African coast, the other in the Indian Ocean and along the east African coast. Mixing of these two clusters in Southern Africa may have produced the present pattern (Neige, 2003).

Allozyme (Pérez-Losada et al., 1999) and microsatellite markers (Pérez-Losada et al., 2002) display a highly significant subpopulation structuring of *S. officinalis* around the Iberian Peninsula, consistent with an isolation-by-distance model of low levels of gene flow. Distinct and significant clinal changes in allele frequencies between the Atlantic and the Mediterranean samples indicated, however, that these results might also be consistent with an alternative model of secondary contact and introgression between previously isolated and divergent populations. A pronounced 'step' change between SW Mediterranean samples associated with the Almería-Oran front suggest that this oceanographic feature may represent a contemporary barrier to gene flow.

Seasonal migrations between shallow and deeper waters are a well-known ecological feature of *S. officinalis*. In the Western Mediterranean populations a general tendency for the animals to migrate inshore in spring and summer for reproduction and move offshore in autumn was observed, although not all the animals migrate at the same time, size and age (Mangold, 1966). These migrations are of different distances, from a few dozens to several hundred nautical miles, and represent an important displacement of biomass, which has also been observed in other regions (Richard, 1971; Najaï, 1983; Boucaud-Camou and Boismery, 1991; Coelho and Martins, 1991; Le Goff and Daguzan, 1991b; Guerra and Castro, 1988; Jorge and Sobral, 2004). As pointed out by Boucaud-Camou and Boismery (1991), autumn *S. officinalis* offshore migration in winter at the English Channel is mainly influenced

not only by cooling of the littoral waters, but also by day-length reduction and decreased light intensity, which are other factors influencing maturation and spawning (Boletzky, 1983; Boucaud-Camou et al., 1991). Thus, the relatively deep milder waters at the central axis of the Channel seems to constitute the common hibernation area to all cuttlefishes in the Channel, which they leave at the end of the winter. Spring inshore displacements are mainly due to an increase of the temperature in littoral waters. These displacements were shown by tagging experiments (Boucaud-Camou and Boismery, 1991), but this spatial and temporal pattern is also supported by the analysis of geo-referenced data measured at both sides of the English Channel (Dunn, 1999; Denis and Robin, 2001; Royer, 2002; Wang et al., 2003). The role of strength of the Atlantic currents into the west part of the English Channel and the south part of the Celtic Sea was found to be the dominant influence on the timing of cuttlefish migration to these areas. Thus, the local abundance was positively correlated to sea surface temperature, with cuttlefish expanding their distribution further north in the spawning seasons in warm years and shifting in cool waters. The centre of high abundance in offshore deep water shifts north in warm winters and south in cool winters (Wang et al., 2003).

### ***Trophic ecology***

The diet of *S. officinalis* includes crustaceans, bony fishes, molluscs, polychaetes and nemertean worms (Nixon, 1987; Castro and Guerra, 1990; Pinczon du Sel, et al., 2000). Species composition within these prey groups depends upon the respective species composition and availability in each ecosystem. Main crustaceans prey items are mysids, shrimps, prawns, and crabs, but *S. officinalis* also feeds upon amphipods, isopods, and ostracods. The most important bony fishes found in the diet of the species were gobies, sand eels, whiting and wrasses, but cuttlefish can also prey upon some flatfishes. Among the cephalopods main food items include various sepiolids and sepiids species. Large cuttlefish are also cannibals, capturing and eating smaller individuals. Other small prey found in the stomach of this species, like bryozoans, foraminifera, bivalve molluscs and insects should be regarded with caution, because they can be the prey of prey, or accidentally ingested prey (Castro and Guerra, 1990). *S. officinalis* shows a wide range of diets and should therefore be considered as a trophic opportunistic animal. The species feeds exclusively on living animals, but in the laboratory it

has been fed with different kind of surimi and pelleted diets, and non- living food (Castro et al., 1993; Koueta and Boucaud-Camou, 1999; Perrin, 2004). Significant ontogenetic changes in the diet of the species with the progressive replacement of crustaceans by fishes have been found (Castro and Guerra, 1990). Ontogenetic changes in the prey size of this species are also well documented (Blanc et al., 1999; Blanc and Daguzan 2000). There were no, however, differences in feeding habits of male and female *S. officinalis* at any size, the feeding intensity of females increasing with sexual maturity, and no seasonal changes in diet were found (Castro and Guerra, 1990).

An attempt to establish the trophic position of *S. officinalis* in an estuarine community (a *Zostera* meadow in San Simón inlet, Ría de Vigo) was undertaken by Filgueira and Castro (2002) based on the analysis of the stable isotopes  $C^{13}$  and  $N^{15}$  in its muscle composition and from sympatric organisms. Significant decreases in  $\delta^{13}C$  and  $\delta^{15}N$  were found related with cuttlefish size (15-195 mm ML) when these values were converted to trophic level. These results disagree with an expected increase in values corresponding to trophic level with predator size, and are in contradiction with previous knowledge of the common cuttlefish feeding ecology. These authors proposed a working hypothesis based on spawning migration. As cuttlefish approach maturity, they migrate to shallow waters, such those of San Simón inlet, for spawning. Then, the smallest mature animals used in this study (60 mm ML for males and 80 mm ML for females) would probably not have left San Simón inlet yet, their isotopic composition representing the local food web. The largest animals present in San Simón were probably coming back from deeper waters, having an isotopic composition that does not depend initially on the local food web. Moreover, as the metabolic rate of large animals is lower than that of smaller ones, they would keep for longer time the isotopic signals from deeper waters before its body composition is in equilibrium with that of the shallow area. If the food web of San Simón shows higher delta ( $\delta$ ) isotopic values than that from outside, a predator, such as cuttlefish, growing in that habitat, should show a heavier isotopic composition than a predator outside this area. Therefore, C and N isotopic composition of cuttlefish from San Simón would be inadequate for estimating its trophic level, and for testing the hypothesis that the trophic level of a predator increases with body size, because large and small animals could belong to different trophic webs.



Analyses using carbon- and oxygen-isotope composition [ $\delta^{13}\text{C}$  ( $\text{CO}_3^{2-}$ ) and  $\delta^{18}\text{O}$  ( $\text{CO}_3^{2-}$ ), respectively] in the cuttlebone aragonite of wild and cultivated specimens of *S. officinalis* from NW Spain showed that seasonal changes in isotopic temperature revealed by these analyses agreed with changes in surrounding sea water temperature:  $\text{CaCO}_3$  was deposited in the cuttlebone all year round, a maximum life span of 2 years, a yearly spawning season, and the existence of variable growth rates among and within individuals can be inferred from isotopic temperatures (Bettencourt and Guerra, 1999).

The tentacles of *S. officinalis*, when ejected, reach the prey in less than 15 milliseconds at 25° C. Prey is dealt with summarily. Thus, prawns are paralysed and bitten within six seconds of capture and crabs are paralysed in about ten seconds. The immobilisation of prey is provoked by neurotoxins secreted by the posterior salivary glands (Hanlon and Messenger, 1996). If exists, external digestion of prey seems to be very weak and many pieces of exoskeleton are ingested (Guerra et al., 1988).

Despite the small size of mouth, cuttlefish can seize relatively large prey with their prehensile arms and tentacles. This, together with voracity, versatile feeding habits, and a highly evolved sensory system, allows them to occupy a broad trophic niche. Furthermore, migrations enable *S. officinalis* populations to exploit the temporal and spatial variability of productive systems and fluctuating populations of prey (Rodhouse and Nigmatullin, 1996). Visual detection of prey involves movement, contrast, size, shape and orientation. The visual attack in this ambush predator when facing a prawn exhibits three phases: attention, positioning, and seizure (Hanlon and Messenger, 1996). Data collected in two 24 h sampling operations carried out in August and February in the Ría de Vigo (NW Spain) suggested a 24 h feeding pattern for this species where most of the feeding occurred during darkness (Castro and Guerra, 1989). Such a feeding pattern has been also described in South Brittany (Pinczon du Sel et al., 2000) and in the Ría Formosa lagoon (Quintela and Andrade, 2002). These results suggest that *S. officinalis*, apart from visual detection, may also detect some prey by light emitted from their light organs, and that chemo- and mechanoreception (via statocysts and/or the lateral line analogue) cannot be ruled out. Predation of non-luminous prey can be also facilitated by dinoflagellate luminescence (Fleisher and Case, 1995).

Common cuttlefish have high absorption efficiency, which explains high growth rates and relatively low production of faeces. Forsythe et al. (1994) estimated a conversion efficiency of 59 % in animals cultivated at 24 ° C and fed with shrimps, which showed a growth rate of 6.5 and a feeding rate of 11.0 (both rates in % body mass per day).

With very few exceptions, there are no fishes that are specialist cephalopod predators. Among the elasmobranches, lower beaks of *S. officinalis* were found in the stomach content of *Prionace glauca* (Clarke and Steven, 1974). *S. officinalis* also occurred in the stomachs of *Scyliorhinus canicula*, *Mustelus mustelus* (Morte et al., 1997) and *Galeus melastomus* (Velasco et al., 2001). Among teleostei, it occurred in the stomach contents of *Merluccius merluccius* (Larrañeta, 1970; Velasco et al., 2001). Hatchling and juvenile common cuttlefish are preyed upon by *Serranus cabrilla* in *Posidonia* grass areas of the Mediterranean (Hanlon and Messenger, 1988). The species *Pollachius pollachius* exerts great predatory pressure on young cuttlefish in the French waters of north Brittany (Le Mao, 1985). In the Bay of Biscay, Velasco et al. (2001) found *S. officinalis* in the stomach contents of *Pagelus acarne*, *Aspitrigla cuculus*, *A. obscura*, *Lophius piscatorius*, *L. budegassa*, *Trisopterus luscus*, *Lepidorhombus whiffiagonis* and *L. boscii*. Young cuttlefish were observed in the stomach contents of *Dicentrarchus labrax*, *Labrus bergylta*, *Spondylisoma cantharus* and *Conger conger* in Morbihan Bay (Blanc and Daguzan, 1999).

A total of 12 specimens of *S. officinalis* were found in a Risso's dolphin (*Grampus griseus*) (Clarke and Pascoe, 1985). To date, the species has not been clearly identified in the stomach contents of other marine mammals, except in monk seals (*Monachus monachus*) from the Aegean Sea (Salman et al., 2001). However, some remains identified as *Sepia* sp, *Sepia* spp or simply Sepiidae were observed in the harbour porpoise (*Phocoena phocoena*), and in the dolphins *Tursiops truncatus*, *Delphinus delphis* and *Stenella coeruleoalba* (Santos, 1998).

It has been suggested that the ecological niche of a cephalopod species is more important in determining its risk of parasite infection than its phylogeny, and that *S. officinalis* should be included in one ecological coastal group (González et al., 2003). The virus-like particles found in the stomach epithelium of wild *S. officinalis* have a structure similar to vertebrate 'Retrovirus' (Hanlon and

Forsythe, 1990). Cultured in the laboratory, this species showed susceptibility to a highly virulent systemic infection by bacteria (*Pseudomonas* and *Vibrio*), which does not appear to be related to external injury (Hanlon and Forsythe, 1990). Diseases may be caused by other protists and metazoans such as fungi, coccidians, microsporidians, ciliates, dicyemids, diagenans, cestodes, nematodes, brachyurans, copepods and isopods (Hochberg, 1990). Many of these parasites are transmitted through the food web. Sexual stages of the coccidian *Aggregata eberthi* occur in the digestive tract of *S. officinalis*, and asexual stages infect the digestive tract of crustaceans. The complete life cycle of *A. eberthi* in NE Atlantic was only achieved when experimental infections showed that the prawns *Palaemon elegans* and *P. adpersus* are the intermediate hosts for this parasite (Gestal et al., 2002a).

### **Demographic Ecology**

Considering its reproductive traits, *S. officinalis* has been included in the group termed 'Intermittent terminal spawning' (Rocha et al., 2001). This reproductive pattern is characterized by the fact that all species included spawn once, ovulation is group-synchronous, spawning is monocyclic, egg lying occurs in separate batches, and somatic growth does not generally take place between spawning events. In such a species, spawning period tends to be relatively long. The main spawning season of *S. officinalis* in the Western Mediterranean and the Gulf of Tunis covers spring and summer, but winter spawning has also been observed (Mangold-Wirz, 1963; Najai, 1983). The spawning period extends from early spring and late summer in south and central Portugal and both the Atlantic and Mediterranean coast of South Spain, with a spawning peak in June and July (Villa, 1998; Tirado et al., 2003; Jorge and Sobral, 2004). The spawning season of this species within the downed estuarine valleys in NW of the Iberian Peninsula extends from early spring to late summer, but winter spawning has also been recorded (Guerra and Castro, 1988). The spawning season in the Bay of Biscay and the Gulf of Morbihan lasts for six months, from mid-March to late June (Le Goff and Daguzan, 1991a). Along both the north and the south coast of the English Channel the spawning season of *S. officinalis* extends from February to July (Dunn, 1999; Royer, 2002; Wang et al., 2003). Environmental factors (much milder winter conditions in some areas than in others) probably account

for most of the variations observed in *S. officinalis* spawning times (Boletzky, 1983). It also has been observed that a restriction of food intake in early life may delay maturation and extend life span in this species (Boletzky, 1979).

Studies on fecundity carried out by Laptikhovsky et al., (2003) showed that the potential fecundity (PF) of advanced maturing and mature pre-spawning *S. officinalis* in the Aegean Sea varies from 3,700 to 8,000 (mean 5,871) oocytes, whereas the number of large yolk oocytes increase with ML from 130 to 839. These authors also observed that spawning females have a PF of some 1,000-3,000 eggs below that of pre-spawning females. This provides evidence that intermittent spawning, which occurs in captivity (Boletzky, 1987), is a normal process in natural habitats, suggesting that common cuttlefish females release a number of eggs equivalent to about 50 % of PF during spawning, although many individual variants are possible under wild conditions.

*S. officinalis* generally lays eggs at depths rarely greater than 30 or 40 m. The eggs are attached in clusters to various plants, sessile animals such as tube worms, or dead structures such as drowned trees, cables or nets. No parental care has been reported in this species, but no major predation pressure on the eggs has been observed. The length of embryonic development is temperature dependent (Boletzky, 1983). Hatchling of this species has a mantle length that may vary from 6 to 9 mm, and it is strikingly similar to adult both in morphology and basic behaviour. Hatching generally occurs at a stage sufficiently advanced to enhance active feeding within hours after hatching. Young cuttlefishes can adapt to very low food intake and maintain growth rates much lower than normal. This provides a margin of safety allowing animals to survive under unfavourable conditions (Boletzky, 1983).

Common cuttlefish live for approximately two years, although some male individuals may attain a greater age. Females die shortly after spawning, although this event can extend over several weeks or even months in the laboratory. Mass mortality, after the spawning season, has been observed in the French and Spanish Atlantic coasts (Richard, 1971 and pers. obs.), but nothing of comparable intensity is known in the Mediterranean (Boletzky, 1983).

There can be several causes of death among the common cuttlefish in a population: removal by fishing, predation, diseases, accident, etc., each with its own rate. It is a usual practice in population

dynamic studies to consider a division into only two types: fishing, and natural mortality, which includes everything else. Each kind of mortality has its own instantaneous rate (Ricker, 1975). In an ideal scenario, the natural mortality could be differentiated by different causes. In practice, this is, however, very difficult. Preliminary results of a management exercise of *S. officinalis* gillnet fishery in San Simón inlet for the period 1997-2001 demonstrated that monthly instantaneous rate of natural mortality (M) over a six-month period (from November to April) ranged from 2.27 to 3.38, the mean being 2.70 (Outeiral, 2002; Rocha and Guerra, unpublished). These values were estimated by different methods exclusively based on the biological parameters obtained by Guerra and Castro (1988) and Bettencourt (2000) from the *S. officinalis* populations within the Galician Rías, and are similar to those calculated by Emam (1994) in the *Sepia prashadi* exploited population from the Gulf of Suez. The concept of an “instantaneous” rate can be troublesome for readers not familiarized with population dynamics. There is, however, an excellent explanation of this concept in Ricker (1975). The mean value of M estimated for the *S. officinalis* of San Simón inlet corresponds to an annual mortality rate (A) of approximately 93% of the total number of individuals of a given population, which is very high. That mortality rate suggests a catastrophic post-spawning mortality, which has been corroborated in the species by both field and laboratory observations. However, when M is used in different stock assessment methods (Pierce and Guerra, 1994) its value ranges from 0.1 to 0.6. The remaining mortality is due to fishing (F).

*S. officinalis* constitutes part of the diet of many marine predators at different stages of their life cycle, but natural mortality rates caused by predation have not yet been evaluated. Although various parasites are known in juvenile and adult *S. officinalis*, most of them do not appear to be important as a natural mortality factor at pre-reproductive stages (Hochberg, 1990). Nevertheless, a detrimental effect on gastrointestinal function by high digestive tract infections with *Aggregata eberthi* might result in a decrease or malfunction of absorption enzymes (Gestal et al., 2002b).

How many units of population conform *S. officinalis* within its area of distribution is something still unknown. However, from an exploitation point of view, cuttlefish concentrations within the English Channel are considered as a management unit. This is mainly due to the fact that

catch per unit effort is lower in the adjacent waters of the Bay of Biscay and the Celtic Sea than in the English Channel (Denis and Robin, 2001).

Common cuttlefish can exhibit variations in its life cycle along its geographic range. Around the Iberian Peninsula and in the Mediterranean Sea, spawning sizes range from 90 to 320 mm ML. This suggests the presence of two-year classes of breeders in the population, and the population structure of *S. officinalis* may superficially look simple, with seldom more than two annual cohorts or a cycle of alternating shorter and longer generations, at least as far as the female individuals are concerned (Boletzky, 1983). However, as cuttlefish may attain sexual maturity a very different sizes, spawning occurs over a long time period when compared to life span, the duration of egg development is dependent on temperature, and its growth is very depending on environmental factors (Bouchaud and Daguzan, 1990; Forsythe et al., 1994; Clarke et al., 1989; Dunn, 1999; Bettencourt, 2000; Domingues et al., 2002; Koueta and Boucaud-Camou, 2003). Recruitment of successive broods reveals subgroups, cohorts or 'micro-cohorts', whose age at recruitment varied significantly between seasons and cohorts, demonstrating different growth rates among them, and there are large interannual variations in recruitment (Challier et al., 2002; Challier, 2005), so a more complex demographic pattern is underlined. Factors affecting recruitment are, therefore, of key importance in understanding the population dynamics of this species (Challier, 2005).

A consistent biannual life cycle has been described in the English Channel (Dunn, 1999; Royer, 2002). Hatchlings born from July to September grow rapidly, and the juveniles migrate from the inshore nursery grounds in late autumn to overwintering grounds in deeper waters. The following spring they return inshore, and begin to exhibit the first signs of sexual maturation. Males start to mature at a mean ML of about 100 mm, and most are mature by September (about 13 months old). Female maturation begins slightly later, and takes longer, with the final stages of female maturation occurring during the following winter. After a second offshore migration to overwintering grounds the adult cuttlefish (approximately 18 months old) return inshore in the spring to spawn and then die. In south Brittany, some of the juveniles born from mid-March to late June begin their sexual development as early as November for males and late December for females. These precocious individuals require only one year to complete their life cycle, and constitute small size breeders (80-

100 mm ML). However, most individuals reproduce after a second offshore-inshore migration, which constitute a second group of breeders (130-350 mm ML). These two-year classes of breeding cuttlefish are not reproductively separated (Gauvrit et al. 1998).

### ***Behavioural ecology***

The seasonal migrations between shallow and deeper waters bring *S. officinalis* into contact with various types of soft and rocky bottoms. The ability of small juveniles to attach themselves to a hard substrate may be very important because it allows them to withstand strong water movement without being carried away. These animals are able to bury themselves in soft bottoms, and the behavioural pattern of this sand covering is well established at hatching (Boletzky, 1983).

The entire morphology of this species reflects adaptation to life near or on the bottom in a very complex environment. Moreover, *S. officinalis* has a considerable repertoire of defensive strategies involving a large number of chromatic, textural and postural components (Hanlon and Messenger, 1996). Detailed studies on defence sequences of young and adult *S. officinalis* show complicated and different behavioural patterns. However, these tactics were mainly observed under laboratory conditions and, therefore, caution must be invoked when extrapolating to wild conditions (Hanlon and Messenger, 1996). Dickel et al (2000) provided evidence that the environment during the 2<sup>nd</sup> and/or 3<sup>rd</sup> months of life was crucial to the ontogenetic development of memory in *S. officinalis*.

The reproductive behaviour in this species is well known (Hanlon and Messenger, 1996). A single pair can mate several times in succession, sometimes intermixed with egg lying. Under culture conditions temporary mate guarding by the male has been observed. However, when guarding relaxes, other mature males can copulate with the female. Therefore, there is evidence of promiscuity, at least in the laboratory, where some results obtained using microsatellite DNA (Guerra, unpublished data) and from behavioural studies (Hanlon et al., 1999) provide evidence that sperm competition may be a major feature of the mating behaviour in this species.

***Some general ecological remarks***

*S. officinalis* has life-cycle characteristics known in other coleoid cephalopods: early sexual maturation, extended spawning season, breeding once, catholic predator habits, rapid growth with variable growth rates depending on environmental factors, short life-span, little overlap of generations, and complex recruitment. However, in contrast to other species, mainly those that are pelagic throughout their life cycle (e.g. ommastrephid squids) or those with planktonic paralarval stages (e.g. some octopods), *S. officinalis* is not so vulnerable to predation and environmental variables. This is mainly due to the fact that the new hatchlings of this species are almost miniature adults, both in morphology and behaviour, and also because their physiological layout is highly flexible (Nixon and Mangold, 1998). In consequence, this species does not show as much unpredictability of distribution and density as other cephalopod species (Boyle and Boletzky, 1996).

As all cephalopods, *S. officinalis* shows relatively low levels of genetic variation (Sanjuan et al., 1996; Shaw and Pérez-Losada, 2000), and its population dynamics appears to be influenced principally by phenotypic plasticity in response to environmental variability, and the maintenance of this diversity balances the risks of mortality factors combining at any one time to cause periodic extinctions (Boyle and Boletzky, 1996). The change in the faunistic composition of the communities observed in the Sahara Bank showed that major ecological perturbations, such environmental shifts or imposed effects such as commercial fishing, have an important impact on *S. officinalis* and other cephalopod populations (Boyle and Boletzky, 1996).

Whether or not *S. officinalis* fits into the generalizable *r*-or *K*-strategies, and in view of the doubts expressed by Stearns (1992) on the value of these strategies to the interpretation of life histories, a theoretical framework for cephalopod life cycle is perhaps premature (Boyle and Rodhouse, 2005). It is best to consider that this species shows a complex set of covarying traits which constitute both strategies throughout its life cycle, as indicated by Calow (1987).

**ACKNOWLEDGEMENTS-** The author would like to thank his colleagues Drs. Angel F. González, Francisco Rocha, Santiago Pascual and Noussithé Koueta for the helpful discussions maintained on many of the aspects presented in this paper. I am also very grateful



to María T. Fernández for her constant and useful technical assistance. Finally, I wish to express my gratitude to Drs. Sigurd von Boletzky and José Pedro Andrade for their valuable comments and suggestions that largely improved the manuscript.

## REFERENCES

- Balguerías E, Quintero, ME, Hernández-González CL 2000. The origin of the Sahara Bank cephalopod fishery. *ICES J Mar Sci* 57: 15-23.
- Bettencourt V 2000. Idade e crescimento do choco, *Sepia officinalis* L. Ph. D. Thesis. Universidade do Alentejo. Portugal: 196 p.
- Bettencourt V, Guerra A 1999. Carbon -and oxygen- isotope composition of the cuttlebone of *Sepia officinalis*: a tool for predicting ecological information? *Mar Biol* 133(4): 651-657.
- Blanc A, Daguzan J 1999. Young cuttlefish *Sepia officinalis* (Mollusca: Sepiidae) in the Morbihan Bay (South Brittany, France): accessory prey of predators. *J mar biol Ass U K* 79:1133-1134.
- Blanc A, Daguzan J 2000. Size selectivity in the diet of the young cuttlefish *Sepia officinalis* (Mollusca: Sepiidae). *J mar biol Ass UK* 80:1137-1138.
- Blanc A, Pinczon du Sel G, Daguzan J 1999. Relationship between the length of the prey/predator for the most important prey of the cuttlefish *Sepia officinalis* L. (Mollusca, Cephalopoda). *Malacologia* 41: 139-145.
- Boal, JG, Hylton, RA, González, SA, and Hanlon, RT 1999. Effects of crowding on the social behaviour of cuttlefish (*Sepia officinalis*). *Contemp Top Lab Anim* 38(1):49-55.
- Boletzky Sv. 1979. Growth and life-span of *Sepia officinalis* under artificial conditions (Mollusca, Cephalopoda). *Rapp Comm Int Mer Medit*, 25/26: 159-168.
- Boletzky Sv 1983. *Sepia officinalis*. In Boyle, PR ed, Cephalopod Life Cycles. Vol I. Species account. Academic Press, London: 31-52.
- Boletzky Sv 1987. Fecundity variation in relation to intermittent or chronic spawning in the cuttlefish, *Sepia officinalis* L. (Mollusca, Cephalopoda). *Bull Mar Sci* 40 (2): 382-397.
- Boucaud-Camou E, Boismery, J 1991. The migrations of the cuttlefish (*Sepia officinalis* L.) in the English Channel. In Boucaud-Camou, E ed, The cuttlefish. Actes du 1<sup>er</sup> Symposium international

- sur la seiche. Centre de Publications de l'Université de Caen:179-189.
- Boucaud-Camou R, Koueta K, Boismery J Medhioub A 1991. Sexual cycle of *Sepia officinalis* L. from the Bay of Seine. In Boucaud-Camou, E ed, The cuttlefish. Actes du 1<sup>er</sup> Symposium international sur la seiche. Centre de Publications de l'Université de Caen:141-151.
- Bouchaud O, Daguzan J 1990. Étude expérimentale de l'influence de la température sur le déroulement embryonnaire de la seiche *Sepia officinalis* L. (Céphalopode, Sepiidae). *Cah Biol Mar* 31: 131-145.
- Boyle PR, Boletzky Sv 1996. Cephalopods populations: definition and dynamics. In Clarke MR ed, The role of cephalopods in the world's oceans. *Phil Trans Roy Soc London, Ser. B.* 351 (1343): 885-1002.
- Boyle PR, Rodhouse PG 2005. Cephalopods. Ecology and Fisheries. Blackwell Publishing, 464 p.
- Brix O, Colosimo A, Giardina B, 1994. Temperature dependence of oxygen binding to cephalopod haemocyanins: Ecological implications. In Pörtner O, O'Dor RK & Macmillan DL eds, Physiology of Cephalopods Molluscs. Gordon and Breach Publishers: 149-162.
- Bustamante P, Teyssié J-L, Fowler SW, Cotret O, Danis B, Miramand P, Warnau M 2002. Biokinetics of zinc and cadmium accumulation and depuration at different stages in the life cycle of the cuttlefish *Sepia officinalis*. *Mar Ecol Prog Ser* 231: 167-177.
- Calow, P 1987. Fact and theory-an overview. In: Boyle PR ed, Cephalopods Life Cycles, Vol. II: Comparative Reviews, Academic Press, London: 351-365.
- Castro BG, Guerra A 1989. Feeding pattern of *Sepia officinalis* (Cephalopoda, Sepioidea) in the Ría de Vigo (NW Spain). *J mar biol Ass UK* 69: 545-553.
- Castro BG, Guerra A 1990. The diet of *Sepia officinalis* and *Sepia elegans* (Cephalopoda, Sepioidea) from the Ría de Vigo. *Sci Mar* 54(4): 375-388.
- Castro BG, Di Marco, FP, DeRusha RH, Lee PG. 1993. The effects of surimi and pelleted diets on the laboratory survival, growth, and feeding rate of the cuttlefish *Sepia officinalis* L. *J Exp Mar Biol Ecol* 170: 241-252.

- Challier L. 2005. Variabilité de la croissance des céphalopodes juvéniles (*Sepia officinalis*, *Loligo forbesi*) et relation avec les fluctuations du recrutement, en Manche. Thèse Doctoral. Université de Caen: 181 p.
- Challier L, Royer J, Robin JP. 2002. Variability in age-at-recruitment and early growth in English Channel *Sepia officinalis* described with statolith analysis. *Aquat Living Resour* 15: 303-311.
- Clarke, MR Stevens JD. 1974. Cephalopods, blue sharks and migrations. *J mar biol Ass UK* 54: 949-957.
- Clarke MR, Pascoe PL 1985. The stomach contents of a Risso's dolphin (*Grampus griseus*) stranded at Thurlestone South Devon. *J mar biol Ass UK* 65: 663-665.
- Clarke A, Rodhouse PG, Holmes LJ, Pascoe PL 1989. Growth rate and nucleic acid ratio in cultured cuttlefish *Sepia officinalis* (Mollusca: Cephalopoda). *J Exp Mar Biol Ecol* 133: 229-240.
- Coelho L, Martins C 1991. Preliminary observations on the biology of *Sepia officinalis* in the Ria Formosa, Portugal. In Boucaud-Camou E ed, The cuttlefish. Actes du 1<sup>er</sup> Symposium international sur la seiche. Centre de Publications de l'Université de Caen: 169-184.
- Denis V, Robin J-P 2001. Present status of the French Atlantic fishery for cuttlefish (*Sepia officinalis*). *Fish Res* 52: 11-22.
- Dickel L, Bola, JG, Budelmann BU. 2000. The effect of early experience on learning and memory in cuttlefish. *Devel Psychol* 36: 101-110.
- Domingues PM, Kingston T, Sykes A, Andrade JP 2001. Growth of young cuttlefish, *Sepia officinalis* (L., 1758) at the upper end of the biological distribution temperature range. *Aquac Res* 32(1): 923-930.
- Domingues PM, Sykes A, Andrade JP 2002. The effects of temperature in the life cycle of two consecutive generations of the cuttlefish *Sepia officinalis* L. cultured in the Algarve (South Portugal). *Aquac Int* 10: 207-220.
- Domingues PM, Poidier R, Dickel L, Almansa E, Sykes A, Andrade JP. 2003. Effects of culture densities and live prey on growth and survival of juvenile cuttlefish, *Sepia officinalis*. *Aquac Int* 11: 225-242.
- Dunn MR 1999. Aspects of the stock dynamics and exploitation of cuttlefish, *Sepia officinalis*

- (Linnaeus, 1758) in the English Channel. *Fish Res* 40: 277-293.
- Emam WM 1994. Stock assessment of the cuttlefish *Sepia prashadi* (Mollusca: Cephalopoda) in the Gulf of Suez. *Indian J Mar Sci* 23: 35-38.
- Filgueira R, Castro BG 2002. Relationship between trophic position and size of cuttlefish *Sepia officinalis* L.: do stable isotopes give the right answer? In Rolán E & Troncoso J eds, II Congr Inter Soc Malac Europ, Vigo: 93.
- Fleisher KJ, Case, JF 1995. Cephalopod predation facilitated by dinoflagellate luminescence. *Biol Bull* 189:263-271.
- Forsythe JW, de Rusha RH, Hanlon RT 1994. Growth, reproduction and life-span of *Sepia officinalis* (Cephalopoda: Mollusca) cultures through seven consecutive generations. *J Zool Lond* 233: 175-192.
- Gauvrit, E, Pinczon du Sel, G, Blanc, A, Daguzan, J 1998. Detection of cross-mating between two generations of the cuttlefish *Sepia officinalis* by sperm reservoir analysis. *J Moll Stud* 64: 1-9.
- Gestal C, Guerra A, Pascual S, Azevedo C 2002a. On the life cycle of *Aggregata eberthi* and *A. octopiana* (Aplicomplexa, Aggregatidae) from Galicia (NE Atlantic). *Eur J Protistol* 37: 427-435.
- Gestal C, Páez de la Cadena M, Pascual, S 2002b. Malabsorption syndrome observed in the common octopus *Octopus vulgaris* infected with *Aggregata octopiana* (Protista: Apicomplexa). *Dis Aquat Org* 51: 61-65.
- González AF, Pascual S, Gestal C, Abollo E, Guerra, A 2003. What makes a cephalopod a suitable host for parasite? The case of Galician waters. *Fish Res* 60: 177-183.
- Grigioni S, Boucher-Rodoni R 2002. Symbiotic associations in *Sepia officinalis*. *Bull Mar Sci* 71(2):1124.
- Guerra A, Sánchez P 1985. Comparación de las pesquerías de cefalópodos de dos áreas de afloramiento de la costa occidental africana. *Simp Intern Aflor Inst Inves Pesq Barcelona*, vol. II: 749-760.
- Guerra A, Castro BG 1988. On the life cycle of *Sepia officinalis* (Cephalopoda, Sepioidea) in the Ría de Vigo (NW Spain). *Cah Bio Mar* 29: 395-405.

- Guerra A, Nixon M, Castro BG 1988. Initial stages of food ingestion by *Sepia officinalis*. *J Zool Lond* 214: 189-197.
- Guerra A, Pérez-Losada M, Rocha F, Sanjuán A 2001. Species differentiation of *Sepia officinalis* and *Sepia hierredda* (Cephalopoda: Sepiidae) based on morphological and allozyme analyses. *J mar biol. Ass UK* 81: 271-281.
- Hanlon RT, Messenger JB 1988. Adaptive coloration in young cuttlefish (*Sepia officinalis* L.): the morphology and development of body patterns and their relation to behaviour. *Phil Trans Roy Soc London B*, 320: 437-487.
- Hanlon RT, Forsythe JW 1990. Diseases caused by microorganisms. In Kinne O ed, Diseases of Marine Animals, Vol. III. Diseases of Mollusca: Cephalopoda. Biologische Anstalt Helgoland, Hamburg: 23-47.
- Hanlon RT, Messenger JB 1996. Cephalopod Behaviour. Cambridge University Press, 232 p.
- Hanlon RY, Ament SA, Gabr H 1999. Behavioral aspects of sperm competition in cuttlefish, *Sepia officinalis* (Sepioidea: Cephalopoda). *Mar Biol* 134: 719-728.
- Ho J-S 1983. *Metaxymolgus longicaudata* (Claus), a copepod associated with the cuttlefish, *Sepia officinalis* L. *J mar biol Ass UK* 63 :199-203.
- Hochberg FG 1990. Diseases caused by protists and metazoans. In Kinne O ed, Diseases of Marine Animals. Vol III. Diseases of Mollusca: Cephalopoda. Biol Anst Helgoland, Hamburg: 47-202.
- Jorge I, Sobral MP. 2004. Alguns aspectos da biología e ecología da população de choco, *Sepia officinalis* da região de Aveiro. *Relat Cient Téc IPIMAR, série digital*, nº 15: 29 p.
- Johansen K, Kornerup S, Lykkeboe G 1982. Factors affecting O<sub>2</sub>-uptake in the cuttlefish, *Sepia officinalis*. *J mar biol Ass UK* 62: 187-191.
- Khromov DN 1998. Distribution patterns of Sepiidae. In Voss NA, Vecchione M, Toll RB & Sweeney MJ. Systematics and Biogeography of Cephalopods. *Smith Cont Zool* 586:191-206.
- Khromov DN, Lu CC, Guerra A, Dong Zh, Boletzky Sv 1998. A synopsis of Sepiidae outside Australian waters (Cephalopoda: Sepioidea). In Voss, NA, Vecchione M, Toll RB & Sweeney MJ. Systematics and Biogeography of Cephalopods. *Smith Cont Zool* 586: 77- 130.

- Koueta N, Boucaud-Camou E. 1999. Food intake and growth in reared early juvenile cuttlefish *Sepia officinalis* L. (Mollusca, Cephalopoda). *J Exp Mar Biol Ecol* 240: 93-109.
- Koueta N, Boucaud-Camou E. 2003. Combined effects of photoperiod and feeding frequency on survival and growth of juvenile cuttlefish *Sepia officinalis* L. in experimental rearing. *J Exp Mar Biol Ecol* 296: 215-226.
- Larrañeta MG 1970. Sobre la alimentación, la madurez sexual y la talla de primera captura de *Merluccius merluccius* (L.). *Inv Pesq* 34: 267-280.
- Laptikhovsky, V., Salman, A., Önsoy, B., Katagan, T. 2003. Fecundity of the common cuttlefish, *Sepia officinalis* L. (Cephalopoda, Sepiidae): a new look at an old problem. *Sci Mar* 67: 279-284.
- Le Bihan E, Perrin A, Koueta N 2004. Development of a bioassay from isolated digestive glands cells of the cuttlefish *Sepia officinalis* L. (Mollusca Cephalopoda): effect of Cu, Zn and Ag on enzyme activities and cell viability. *J Exp Mar Biol Ecol* 309: 47-66.
- Le Goff R, Daguzan J 1991a. Growth and life cycles of the cuttlefish *Sepia officinalis* L. (Mollusca: Cephalopoda) in South Brittany (France). *Bull Mar Sci* 49:341-348.
- Le Goff R, Daguzan, J 1991b. Étude des déplacements de la seiche commune *Sepia officinalis* L. dans le Golfe du Morbihan au cours de la période de la reproduction. Premiers résultats. In Boucaud-Camou E ed, The cuttlefish. Centre de Publications de l' Université de Caen:167-177.
- Le Mao P. 1985. Place de la seiche *Sepia officinalis* (Mollusque, Céphalopode) dans les chaînes alimentaires du Golfe Normano-Breton. *Cah Biol Mar* 26: 331-340.
- Mangold-Wirz K 1963. Biologie des céphalopodes benthiques et nectoniques de la Mer Catalane. *Vie Milieu* 13 (suppl.), 285 p.
- Mangold K. 1966. *Sepia officinalis* de la Mer Calalane. *Vie Milieu* A 17: 961-1012.
- Melzner F, Bock C, Pörtner H.-O 2004. Coordination of ventilation and circulation in the cuttlefish *Sepia officinalis* in the light of an oxygen limitation of thermal tolerance. *ICES CM 2004 / CC*: 14:280-281.
- Miramand P, Bentley D 1992. Concentration and distribution of heavy metals in tissues of two cephalopods, *Eledone cirrhosa* and *Sepia officinalis*, from the French coast of the English Channel. *Mar Biol* 114: 407-414.

- Morte S, Redon MJ, Sanz-Brau A 1997. Feeding habits of juvenile *Mustelus mustelus* (Carchariformes, Triakidae) in the western Mediterranean. *Cah Biol Mar* 38: 103-107.
- Najai S 1983. Contribution à l'étude des céphalopodes de Tunisie. Application à l'espèce *Sepia officinalis* Linné, 1758. Thèse de 3<sup>ème</sup> cycle. Université de Tunis: 229 p.
- Neige P. 2003. Spatial pattern of disparity and diversity of the recent cuttlefishes (Cephalopoda) across the Old World. *J Biogeogr* 30: 1125-1137.
- Nixon M 1987. Cephalopod diets. In Boyle PR ed, Cephalopod Life Cycles. Vol II. Comparative reviews. Academic Press, London: 201-219.
- Nixon M, Mangold K 1998. The early life of *Sepia officinalis*, and the contrast with that of *Octopus vulgaris* (Cephalopoda). *J Zool Lond* 245: 407-421.
- Outeiral R 2002. La pesquería de *Sepia officinalis* en Galicia: Aplicación de un modelo basado en entrevistas sobre estadísticas pesqueras. Tesis de Licenciatura. Universidad de Vigo: 90 p.
- Paulij WP, Bogaards RH, Denucé JM 1990. Influence of salinity on embryonic development and the distribution of *Sepia officinalis* in the Delta Area (South Western part of The Netherlands). *Mar Biol* 107:17-23.
- Pérez-Losada M., Guerra A, Sanjuán A 1999. Allozyme differentiation in the cuttlefish *Sepia officinalis* from the NE Atlantic and the Mediterranean. *Heredity* 83: 280-289.
- Pérez-Losada M, Guerra A, Carvalho GR, Sanjuán A, Shaw PW 2002. Extensive population subdivision of the cuttlefish *Sepia officinalis* (Mollusca: Cephalopoda) around the Iberian Peninsula indicated by microsatellite DNA variation. *Heredity* 89: 417-424.
- Perrin MA. 2004. Étude expérimentale des capacités digestives chez la seiche, *Sepia officinalis* L. (Mollusque, Céphalopode): Impact de l'alimentation, indice de condition nutritionnelle et formulation d'un aliment artificiel. Thèse Doctoral. Université de Caen: 153 p.
- Pierce GJ, Guerra, A. 1994. Stock assessment methods used for cephalopod fisheries. *Fis Res* 21: 255-285.
- Pinczon du Sel G, Blanc A, Daguzan J. 2000. The diet of the cuttlefish *Sepia officinalis* L. (Mollusca: Cephalopoda) during its life cycle in the Northern Bay of Biscay (France). *Aquat Sci* 61:167-178.
- Quintela J, Andrade JP 2002. Diel feeding rhythms, daily ration and gastric evacuation rates of *Sepia*

- officinalis* in the Ria Formosa Lagoon (South Portugal). *Bull Mar Sci* 71(2):665-680.
- Richard A 1971. Contribution à l'étude expérimentale de la croissance et la maturation sexuelle de *Sepia officinalis*. Thèse de 3<sup>ème</sup> cycle. Université de Lille: 264 p.
- Ricker WE 1975. Computation and interpretation of biological statistics of fish populations. *Bull Fish Res Board Canada* 191, 382 p.
- Rocha F, Guerra A, González AF 2001. A review of the reproductive strategies in cephalopods. *Biol Rev* 76:291-304.
- Rodhouse PG, Nigmatullin ChM 1996. Role as consumers. In Clarke, M.R. (Ed.). The role of cephalopods in the world's oceans. *Phil Trans Roy Soc London Ser B* 351 (1343):1003-1022.
- Royer J 2002. Modélisation des stocks de céphalopodes de Manche. Thèse de 3<sup>ème</sup> cycle. Université de Caen: 242 p.
- Salman A Bilecenoglu M Güçlüsoy H. 2001. Stomach contents of two Mediterranean monk seals (*Monachus monachus*) from the Aegean Sea, Turkey. *J mar biol Ass UK* 81, 719-720.
- Sanjuan A, Pérez-Losada M Guerra A. 1996. Intra and interespecific genetic differentiation in three species of *Sepia* (Mollusca: Cephalopoda) from Galician waters (NW Iberian Peninsula). *Mar Biol* 126: 253-259.
- Santos MB 1998. Feeding ecology of harbour porpoises, common and bottlenose dolphins and sperm whales in the Northeast Atlantic. Ph.D. Thesis. University of Aberdeen: 284 p.
- Schipp R, Boletzky Sv. 1998. Congenital malformation of the systemic heart of *Sepia officinalis* L.: morphological, phylogenetic and ecotoxicological aspects. In Payne, AI et al., eds, Cephalopod Biodiversity, Ecology and Evolution. *S Afr J mar Sci* 20: 25-27.
- Shaw PW, Pérez-Losada M. 2000. Polimorphic microsatellites in the common cuttlefish *Sepia officinalis* (Cephalopoda). *Mol Ecol* 9:237-238.
- Sobrino I, Silva L, Bellido JM, Ramos F 2002. Rainfall, river discharges and sea temperature as factors affecting abundance of two coastal benthic cephalopod species in the Gulf of Cádiz (SW Spain). *Bull Mar Sci* 71:851-865.
- Stearns, SC 1992. The Evolution of Life Histories. New York, Oxford University Press.
- Tirado C, Rodríguez de la Rúa A, Bruzón MA, López JI, Salas C, Márquez I. 2003. La reproducción



- del pulpo (*Octopus vulgaris*) y el choco (*Sepia officinalis*) en la costa andaluza. *Junta de Andalucía. Consejería de Agricultura y Pesca*, 159 p.
- Van Den Branden C, Gillis M, Richard A 1980. Carotenoid producing bacteria in the accessory nidamental glands of *Sepia officinalis*. *Comp Biochem Physiol* 66B: 331-334.
- Velasco F, Olaso I, Sánchez F 2001. The role of cephalopods as forage for the demersal fish community in the Southern Bay of Biscay. *Fish Res* 52: 65-77.
- Villa H 1998. Estudio de alguns aspectos da biologia reproductiva de la espécie *Sepia officinalis* (Linnaeus, 1758). Msc Thesis. Univ Algarve: 82 p.
- Ward PD, Boletzky Sv 1984. Shell implosion depth and implosion morphologies in three species of *Sepia* (Cephalopoda) from the Mediterranean Sea. *J mar biol Ass UK* 64 :955-966.
- Wang J, Pierce GJ, Boyle PR, Denis V, Robin JP, Bellido JM 2003. Spatial and temporal patterns of cuttlefish (*Sepia officinalis*) abundance and environmental influences - a case study using trawl fishery data in French Atlantic coastal, English Channel, and adjacent waters. *ICES J Mar Sci* 60:1149-1158.
- Warnke K 1994. Some aspects of social interaction during feeding in *Sepia officinalis* (Mollusca: Cephalopoda) hatched and reared in the laboratory. *Vie Milieu* 44 (2):125-131.
- Webber DM, Aitken JP, O'Dor RK 2000. Costs of locomotion and vertic dynamics of cephalopods and fish. *Phys Bioch Zool* 73(6): 651-662.