

## Phytochemical fingerprinting of vegetable *Brassica oleracea* and *Brassica napus* by simultaneous identification of glucosinolates and phenolics

Journal:	<i>Phytochemical Analysis</i>
Manuscript ID:	Draft
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	
Complete List of Authors:	Velasco, Pablo; Misión Biológica de Galicia (CSIC), Plant Genetics Francisco, Marta Moreno, Diego Ferrerres, Federico Garcia-Viguera, Cristina Cartea, Elena; Misión Biológica de Galicia (CSIC), Plant Genetics
Keywords:	<i>Brassica oleracea</i> , <i>Brassica napus</i> , glucosinolates, phenolic compounds, LC/UV-PAD/ESI-MSn



1  
2  
3 1 **Phytochemical fingerprinting of vegetable *Brassica oleracea* and *Brassica napus* by**  
4  
5 2 **simultaneous identification of glucosinolates and phenolics**  
6  
7  
8 3

9  
10 4 Pablo Velasco<sup>a\*</sup>, Marta Francisco<sup>a</sup>, Diego A. Moreno<sup>b</sup>, Federico Ferreres<sup>b</sup>, Cristina García-  
11  
12 5 Viguera<sup>b</sup>, María Elena Cartea<sup>a</sup>  
13  
14 6

15  
16  
17 7 a. Misión Biológica de Galicia (CSIC), PO Box 28, E-36080 Pontevedra, Spain

18 8 b. Department of Food Science and Technology, CEBAS-CSIC, Campus Universitario de  
19  
20 9 Espinardo-Edificio 25, PO Box 164, Espinardo, E-30100 Murcia, Spain  
21  
22

23  
24 10 \* Corresponding author:

25  
26 11 Pablo Velasco

27  
28 12 Address: Misión Biológica de Galicia (CSIC), PO Box 28, E-36080 Pontevedra, Spain

29  
30 13 Tel: 0034-986854800

31  
32 14 Fax: 0034-986841362

33  
34 15 E-mail: [pvelasco@mbg.cesga.es](mailto:pvelasco@mbg.cesga.es)  
35  
36  
37  
38  
39 16

40  
41 17 **Abstract**

42  
43 18 For the first time, intact glucosinolates and phenolic compounds were simultaneously  
44  
45 19 identified in kale, cabbage and leaf rape by LC-UV (PAD)-(ESI) MSn. This study led to the  
46  
47 20 identification of 12 glucosinolates and 32 phenolic compounds which included quinic acid  
48  
49 21 esters of hydroxycinnamic acids, hydroxycinnamic acyl gentobiosides and flavonoids. This  
50  
51 22 study provided a deeper and complete identification of health-promoting compounds in kale  
52  
53 23 and cabbage than previously reported and the new identification of the major phenolic  
54  
55 24 compounds in leaf rape.  
56  
57  
58  
59  
60

1  
2  
3 25 **KEYWORDS:** *Brassica oleracea*, *Brassica napus*, glucosinolates, phenolic compounds,  
4  
5 26 flavonol glycosides, hydroxycinnamic acids, LC/UV-PAD/ESI-MSn  
6  
7  
8 27  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

## 28 INTRODUCTION

29 Brassica vegetables have been related with the prevention of degenerative diseases and  
30 different types of cancer as well as cardiovascular health-promotion (Cartea and Velasco,  
31 2008; Traka and Mithen, 2009). Compounds that appear to contribute to these health-related  
32 properties of brassicas and other food plants include isothiocyanates and their cognate  
33 glucosinolates, phenolics, including flavonoids, and other non-nutrients (Jahangir et al., 2009;  
34 Jeffery and Araya, 2009) inducing a variety of physiological functions as direct or indirect  
35 antioxidants, regulation of enzyme proteins and activities and controlling apoptosis and cell  
36 cycle (Duthie et al., 2000).

37 Glucosinolates are a large group of sulphur-containing secondary plant metabolites, which  
38 occur in all Brassica crops. A wide variety of glucosinolates exists but all share a common  
39 structure comprises a  $\beta$ -thioglucoside N-hydroxysulfates, a  $\beta$ -D glucopyranosyl moiety and a  
40 variable side-chain derived from methionine, tryptophan or phenylalanine. Upon cellular  
41 disruption, glucosinolates are hydrolyzed to various bioactive breakdown products by the  
42 endogenous enzyme myrosinase. Isothiocyanates and indole glucosinolate metabolites (in  
43 particular indol-3-carbinol) are two major groups of autolytic breakdown products of  
44 glucosinolates. Both of them exhibit protective activities against many types of cancer. *In*  
45 *vitro* and *in vivo* studies have reported that these compounds may affect many stages of  
46 cancer development, including the induction of detoxification enzymes (Phase II enzymes)  
47 and the inhibition of activation enzymes (Phase I enzymes) (Mithen et al., 2003) but also  
48 antiproliferative mechanisms like cell cycle arrest or apoptosis (Clarke et al., 2008).

49 Phenolic compounds are a large group of phytochemicals widespread in plant kingdom.  
50 Depending on their structure they can be classified into simple phenols, phenolic acids,  
51 hydroxycinnamic acid derivatives and flavonoids. The most common are the flavonoids  
52 which are built upon C<sub>6</sub>-C<sub>3</sub>-C<sub>6</sub> flavone skeleton. Flavonoids and hydroxycinnamic acid

1  
2  
3 53 derivatives are widely distributed in plants and are important biologically active constituents  
4  
5 54 of the human diet. In *Brassica* foods the flavonoids are complex, with up to five sugar  
6  
7  
8 55 residues present, and these may be further substituted with hydroxycinnamic residues (Vallejo  
9  
10 56 et al., 2004). The bioavailability and activity of different glycosides depends on their  
11  
12 57 substituents (Cermak et al., 2003). For this reason, it is important to characterize and quantify  
13  
14 58 the different derivatives of phenolic compounds. These compounds have direct antioxidant  
15  
16 59 and free radical-scavenging activities but can also induce expression of various genes  
17  
18 60 encoding metabolic enzymes thought to decrease the risk of various diseases and disorders  
19  
20 61 (Bennett et al., 2006).  
21  
22 62 The Brassicaceae family has been widely investigated for glucosinolate (Kushad et al., 1999;  
23  
24 63 Padilla et al., 2007c; Cartea et al., 2008a; 2008b) and for phenolic composition (Llorach et al.,  
25  
26 64 2003a; Vallejo et al., 2004; Ferreres et al., 2005; 2006; Romani et al., 2006; Sousa et al.,  
27  
28 65 2008). Nowadays, the profile of different *Brassica* species is well established. The analysis of  
29  
30 66 these compounds by different methods is laborious and time consuming. Both phenolic  
31  
32 67 compounds and glucosinolates have beneficial properties on human health and synergic  
33  
34 68 effects could appear between both classes of metabolites. For this reason, a method to extract  
35  
36 69 and analyze these compounds at the same time would be very useful. Bennet et al (2003;  
37  
38 70 2006) used a method for analyzing both kinds of compounds on different species. Later, this  
39  
40 71 method was applied to a set of turnip green and turnip top local populations (*B. rapa* var.  
41  
42 72 *rapa*) (Francisco et al., 2009) but, as far as we know, the method has not been used for other  
43  
44 73 *Brassica* crops, like *B. oleracea acephala* and *capitata* groups, and *B. napus*.  
45  
46 74 In Galicia (northwestern Spain), different *Brassica* species are used as leaf vegetable products  
47  
48 75 for human and also for animal consumption. Kales (*Brassica oleracea acephala* group),  
49  
50 76 cabbages (*B. oleracea capitata* group), leaf rape (*B. napus pabularia* group), and turnip tops  
51  
52 77 and turnip greens (*B. rapa rapa* group) are the most important Brassica crops in this region.  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 78 At the Misión Biológica de Galicia (CSIC, Spain), a collection of local varieties of *B.*  
4  
5 79 *oleracea* and *B. napus* is kept as part of the *Brassica* genus germplasm bank. In previous  
6  
7  
8 80 reports, this collection was characterized based on morphological and agronomical traits  
9  
10 81 (Picoaga et al., 2003; Rodriguez et al., 2005; Padilla et al., 2007a; Soengas et al., 2008; Vilar  
11  
12 82 et al., 2008) and the profile of desulphoglucosinolates in leaves was studied (Cartea et al.,  
13  
14  
15 83 2008a; 2008b; Velasco et al., 2008). To date, no information is available on content of intact  
16  
17 84 glucosinolates and phenolic compounds in these species. Therefore, for a more  
18  
19 85 comprehensive assessment that allows the nutritional study, the objective of this work was the  
20  
21 86 identification of glucosinolates, flavonoids and hydroxycinnamic acids in a representative  
22  
23  
24 87 variety of kale, cabbage and leaf rape. Identification was carried out by LC-UV photodiode  
25  
26  
27 88 array detection (PAD)-electrospray ionization (ESI) MSn.  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## 89 **EXPERIMENTAL**

90 **Plant material.** One local variety of each crop was evaluated in this study: a kale variety  
91 named as 'MBG-BRS0468' (*B. oleracea acephala*), a white cabbage variety named as 'MBG-  
92 BRS0057' (*B. oleracea capitata*), and a leaf rape variety named as 'MBG-BRS0063' (*B.*  
93 *napus pabularia*). These varieties are in the germplasm collection at the Misión Biológica de  
94 Galicia (CSIC) and were selected based on previous agronomic and nutritional evaluations  
95 (Rodríguez et al., 2005; Cartea et al., 2008a; 2008b; Soengas et al., 2008; Vilar et al., 2008).  
96 The populations were planted in multipot-trays and seedlings were transplanted into the field  
97 at the five or six leaves stage to collect leaves in well-developed plants. After harvesting on  
98 dry ice, the material was immediately transferred to the laboratory and frozen at -80°C, prior  
99 to their lyophilisation. The dried material was powdered using an IKA-A10 (IKA-Werke  
100 GmbH & Co.KG) mill and the powder was used for analysis.

101  
102 **Sample preparation.** Extraction and the LC gradient for glucosinolate and phenolic analyses  
103 is a multi-purpose chromatographic method that simultaneously separates glucosinolates and  
104 phenolics (Bennett et al., 2003; 2006). Fifty mg of each sample were extracted in 1.5 mL 70%  
105 MeOH at 70°C for 30 min with vortex mixing every 5min to facilitate the extraction. The  
106 samples were centrifuged (13000g, 15min, 4°C). The supernatants were collected and  
107 methanol was completely removed using a rotary evaporator under vacuum at 37°C. The dry  
108 material obtained was redissolved in 1mL of ultrapure water and filtered through a 0.20 µm  
109 syringe PTFE filters (Anotop<sup>TM</sup>, Whatman International Ltd, UK).

110  
111 **Alkaline hydrolysis.** For the study of acyl flavonoid derivatives, an alkaline hydrolysis was  
112 carried out to eliminate acid moieties like p-coumaroyl (*m/z* 146) and caffeoyl (*m/z* 162),  
113 which coincide with those of rhamnosyl and hexosyl residues respectively and, therefore, a

1  
2  
3 114 miss-assignment can occur in MS analysis. Saponification was performed as follows: 1 mL of  
4  
5 115 the extract plus 1 mL of 2M NaOH (up to pH 9–10) for 12 h at room temperature in a  
6  
7  
8 116 stoppered test tube under N<sub>2</sub> atmosphere. The alkaline hydrolysis products were acidified with  
9  
10 117 concentrated HCl (up to pH 1–2) and directly analysed by LC/UV- PAD/ESI-MSn.  
11  
12

13 118  
14  
15 119 **LC/UV- PAD/ESI-MSn analyses.** Chromatographic analyses were carried out on a Luna  
16  
17 120 C18 column (250mm × 4.6mm, 5µm particle size; Phenomenex, Macclesfield, UK). The  
18  
19 121 mobile phase was a mixture of (A) Trifluoro acetic acid (TFA) 0.1% and (B) acetonitrile/TFA  
20  
21 122 (99.9:0.1). The flow rate was 1mL min<sup>-1</sup> in a linear gradient starting with 0% B at 0–5 min,  
22  
23 123 reaching 17% B at 15–17 min, 25% B at 22min, 35% B at 30min, 50% B at 35 min, 99% B at  
24  
25 124 50min and at 55-65min 0% B. The flow rate was 1mL min<sup>-1</sup>, and the injection volume 20µL.  
26  
27 125 Chromatograms were recorded at 330 nm for flavonoid glycosides and acylated derivatives  
28  
29 126 and 227 nm for glucosinolates. The LC/UV- PAD/ESI-MSn analyses were carried out in an  
30  
31 127 Agilent HPLC 1100 series equipped with a photodiode array detector and mass detector in  
32  
33 128 series (Agilent Technologies, Waldbronn, Germany). The HPLC consisted of a binary pump  
34  
35 129 (model G1312A), an autosampler (model G1313A), a degasser (model G1322A), and a  
36  
37 130 photodiode array detector (model G1315B). The HPLC system was controlled by a  
38  
39 131 ChemStation software (Agilent, v. 08.03). The mass detector was an ion trap spectrometer  
40  
41 132 (model G2445A) equipped with an electrospray ionisation interface and was controlled by  
42  
43 133 LCMSD software (Agilent, v. 4.1). The ionisation conditions were adjusted at 350°C and 4kV  
44  
45 134 for capillary temperature and voltage, respectively. The nebulizer pressure and flow rate of  
46  
47 135 nitrogen were 65.0 psi and 11 L min<sup>-1</sup>, respectively. The full scan mass covered the range  
48  
49 136 from *m/z* 100 up to *m/z* 1500. Collision-induced fragmentation experiments were performed  
50  
51 137 in the ion trap using helium as the collision gas, with voltage ramping cycles from 0.3 up to 2  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 138 V. Mass spectrometry data were acquired in the negative ionisation mode. MS<sub>n</sub> is carried out  
4  
5  
6 139 in the automatic mode on the more abundant fragment ion in MS<sub>(n-1)</sub>.  
7  
8 140  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

## 141 RESULTS AND DISCUSSION

### 142 Glucosinolate identification.

143 The MS product ions obtained from glucosinolates were dominated by intense  $[M-H]^-$  ions  
144 and other fragments of the composition of the side chain. Glucosinolates fragmentation  
145 revealed two groups of typical fragments, one associated with the common moiety of the  
146 glucosinolates (aglycone) and the other providing useful diagnostic ions for the identification  
147 of the variable side chain. As it has been described by other authors (Fabre et al., 2007;  
148 Rochfort et al., 2008), the MS<sup>2</sup> fragmentation of the aglycone side chain produces specific  
149 ions at  $m/z$  195, 241, 259 and 275. The fragmentation of the ion  $m/z$  259 (MS<sup>3</sup>  $[(M-H) \rightarrow 259]^-$   
150 ) gave rise to the fragments at  $m/z$  139, 97 (corresponding to the sulphate group ( $[SO_4H]^-$ )),  
151 and  $m/z$  81 ( $[SO_3H]^-$ ). The  $m/z$  97 fragment ion is formed with high abundance in negative  
152 ion ESI method (Mellon et al., 2002). Therefore fragments at  $m/z$  259 and  $m/z$  97 were used as  
153 diagnostic ions of glucosinolates. These fragment ions were a very useful preliminary  
154 screening method for determining the presence of glucosinolates in plant extracts.

155 The other group of fragments are compound-specific products ions that allowed structure  
156 elucidation. All glucosinolates provided of constant neutral loss under the fragmentation  
157 conditions. The most intense and consistent was the combined loss of sulphur trioxide and  
158 anhydroglucose  $[M-H-242]^-$ . Other MS fragmentation pathways were to losses of glucose  
159 radical  $[M-H-162]^-$  and/or thioglucose moiety  $[M-H-196]^-$  after the H-rearrangement of the  
160 side chain to the sulphur atom in thioglucose moiety (Kokkonen et al., 1991). The indolic  
161 glucosinolates were characterized also by the examination of characteristic product ions from  
162 their specific R. This group contain two nitrogen atoms and the  $m/z$  values of their  
163 deprotonated molecules were thus at odd mass numbers. Neoglucobrassicin and 4-  
164 methoxyglucobrassicin exhibit identical molecular masses and fragmentation ions, they were

1  
2  
3 165 differentiated by comparison with reported elution sequence during reversed phase HPLC  
4  
5  
6 166 (Kushad et al., 1999).  
7  
8 167 Therefore the molecular ion  $[M-H]^-$  of glucosinolates, their fragment ion pattern and the  
9  
10 168 retention times allowed the identification of eight glucosinolates in kale and cabbage which  
11  
12 169 exhibited the same glucosinolate profile and eleven glucosinolates in *B. napus* (Figure 1). In  
13  
14  
15 170 total were identified the following intact glucosinolates:  $m/z$  422 glucoiberin,  $m/z$  388  
16  
17 171 progoitrin,  $m/z$  358 sinigrin,  $m/z$  438 glucoraphanin,  $m/z$  402 gluconapoleiferin,  $m/z$  372  
18  
19  
20 172 gluconapin,  $m/z$  463 4-hydroxyglucobrassicin,  $m/z$  386 glucobrassicinapin,  $m/z$  447  
21  
22 173 glucobrassicin,  $m/z$  422 gluconasturtiin,  $m/z$  477 4-methoxyglucobrassicin and  $m/z$  477  
23  
24 174 neoglucobrassicin. The mass spectral information of the glucosinolates identified is  
25  
26  
27 175 summarized in Table 1. The GS profile found in these three crops was similar to those  
28  
29 176 reported by other authors in *B. oleracea* and *B. napus* leaves (Cartea et al., 2008a; 2008b).  
30  
31 177 Cartea *et al.* (Cartea et al., 2008a; 2008b) studied the glucosinolate content of the collection of  
32  
33  
34 178 kale, cabbage and leaf rape kept at the MBG by HPLC-DAD and some differences were  
35  
36 179 found regarding the current work. Eight glucosinolates were identified in the *B. napus*  
37  
38 180 collection by HPLC-DAD. In the present work, 4-methoxyglucobrassicin was not detected  
39  
40  
41 181 and four other glucosinolates (4-hydroxyglucobrassicin, glucoiberin, sinigrin and  
42  
43 182 glucoraphanin) were also identified although all of them in trace quantities (Table 1, Fig.1).  
44  
45  
46 183 With regard to kales and cabbages, the same authors found and quantified 10 and 15  
47  
48 184 glucosinolates, respectively (Cartea et al., 2008a; 2008b). In this work, the most abundant  
49  
50 185 glucosinolates (i.e. glucoiberin, sinigrin, or glucobrassicin) were identified in both crops but  
51  
52  
53 186 other minor glucosinolates like progoitrin, glucoiberin, glucoalyssin, and  
54  
55 187 glucobrassicinapin, were not found in kale and cabbage (Table 1, Fig.1).  
56  
57  
58 188  
59  
60 189

### 189 **Flavonoids identification.**

1  
2  
3 190 The HPLC-DAD chromatogram of Brassica vegetable extracts revealed the existence of  
4  
5 191 glycosylated derivatives of three flavonoids with substitution in position 3, that is, kaempferol  
6  
7  
8 192 (267, 300sh and 349 nm), quercetin (255, 267sh, and 355 nm) and isorhamnetin (255, 268sh,  
9  
10 193 294sh, and 354 nm). In addition, acylated flavonoids were detected in the extract, and their  
11  
12 194 UV spectra, characterized with a maximum with a high absorption at 330 nm and a little  
13  
14 195 maximum between 255 and 268 nm, suggested that the flavonoid-glycoside molecules were  
15  
16 196 linked with hydroxycinnamic acid derivatives, in which sinapic, ferulic, caffeic and *p*-  
17  
18 197 coumaric acids were the most abundant. The results are summarized in Table 2. The  
19  
20 198 chromatographic profiles, recorded at 330 nm, of the naturally occurring phenolic compounds  
21  
22 199 in leaf rape, kale and cabbage extracts and the deacylated phenolic compounds, obtained after  
23  
24 200 alkaline hydrolysis, are shown in (Fig. 2).

25  
26  
27 201 Alkaline hydrolysis was performed to reduce the complexity of the naturally occurring  
28  
29 202 compounds present in the plant extracts due to the release of the hydroxycinnamic acids by  
30  
31 203 cleavage of the ester linkage between the acids and the glycosides (Martens, 2002). After  
32  
33 204 alkaline hydrolysis, the chromatogram showed, apart from several hydroxycinnamic acids,  
34  
35 205 various flavonoid glycosides and disappearance of the acylated derivatives (Fig. 2).

#### 26 206 *Deacylated flavonoids.*

27  
28 207 The MS ion trap analysis of the saponified extracts of the leaves showed mainly the presence  
29  
30 208 of several kaempferol derivatives but quercetin and isorhamnetin were also determined. The  
31  
32 209 presence of the ion at  $m/z$  285/284 [Agl-H/2H]<sup>-</sup> as base peak for compounds **2**, **3**, **4**, **7**, **8** and **9**  
33  
34 210 show them as kaempferol derivatives (3,5,7,4'-tetrahydroxyflavone), while on compounds **1**  
35  
36 211 and **6** this ion was  $m/z$  301/300 indicative of quercetin derivatives ((3,5,7,3',4'-  
37  
38 212 pentahydroxyflavone pentahydroxyflavone) and  $m/z$  315 for **5** isorhamnetin derivatives  
39  
40 213 (3,5,7,4'-tetrahydroxy-3'-methoxyflavone) (Table 2). The MS<sup>2</sup>[M-H]<sup>-</sup> fragmentation analysis  
41  
42 214 of compounds **1-5** showed ions [M-H-162]<sup>-</sup> as base peak, indicated a loss of glycosidic

1  
2  
3 215 residue at 7 position (Ferrerres et al., 2004). According to previous studies (Llorach et al.,  
4  
5 216 2003b; Ferrerres et al., 2004; Vallejo et al., 2004) the fragmentation pattern and the relative  
6  
7 217 abundance of the obtained ions indicated that compounds **1-3** are glycosylated with a  
8  
9 218 hexoside in position 7 and a di- or trihexoside in the 3 position. The first fragmentation of the  
10  
11 219 deprotonated molecular ion  $[M - H]^-$  in this compounds is expected to always be due to the  
12  
13 220 breakdown of the O-glycosidic bond at the 7-position (Ferrerres et al., 2004). The remaining  
14  
15 221 glycosyl moieties of the flavonoid molecule are expected to be linked to the hydroxyl at the 3-  
16  
17 222 position on the flavonol aglycone. The fragmentation  $MS^3[(M-H) \rightarrow (M-H-162)]^-$  of **1-3**  
18  
19 223 showed losses to come from interglycosidic fragmentations at position 3 of the ring which in  
20  
21 224 agree with previous Brassica works, mentioned above, suggest the (1→2) interglycosidic  
22  
23 225 linkage between the disaccharide moieties of the flavonoids (mainly sophorosides). So, these  
24  
25 226 compounds were tentatively characterized as (**1**) quercetin-3-*O*-sophoroside-7-*O*-glucoside;  
26  
27 227 (**2**) kaempferol-3-*O*-sophorotrioside-7-*O*-glucoside and (**3**) kaempferol-3-*O*-sophoroside-7-*O*-  
28  
29 228 glucoside. Compounds **4** and **5** were characterized as flavonoids with two sugar moieties  
30  
31 229 linked to different phenolic hydroxyl (di-*O*-glycosidics). According to Ferrerres et al. (2004),  
32  
33 230 in these 3,7-di-*O*-glucosides, a base peak ion at  $[M-H-162]^-$  in the  $MS^2[M-H]^-$  mode is  
34  
35 231 always observed (Table 2). On the other hand, for compounds **6, 7** and **8** it was observed the  
36  
37 232 fragment ion  $[M-H-180]^-$  and the appearance of  $[Agl-H]^-$  as base peak in the  $MS^2[M-H]^-$ ,  
38  
39 233 together with the observed fragmentations, indicating them as flavonol-*O*-diglycosidics. The  
40  
41 234 UV spectra and MS fragmentation for compounds **7** and **8** show that they are kaempferol-3-  
42  
43 235 *O*-dihexosides isomers. The  $[M-H-180]^-$  ion was not observed in the fragmentation of **7**;  
44  
45 236 while this ion is very important for the compound **8**, indicating a interglycosidic linkage  
46  
47 237 (1→2) for this compound. Thus, they were identified as (**4**) kaempferol-3,7-di-*O*-glucoside;  
48  
49 238 (**5**) isorhamnetin-3,7-di-*O*-glucoside; (**6**) quercetin-3-*O*-sophoroside; (**7**) kaempferol-3-*O*-  
50  
51 239 diglucoside; and (**8**) kaempferol-3-*O*-sophoroside. Compound **9** was identified as a  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 240 monoglycoside kaempferol derivative with hydroxyl at 3 position free: (9) kaempferol-7-O-  
4  
5 241 glucoside.

6  
7  
8 242 In native extracts of kale, cabbage and leaf rape we observed compounds previously described  
9  
10 243 by other authors in *Brassica* spp.(Llorach et al., 2003a; 2003b; Ferreres et al., 2004; Vallejo et  
11  
12 244 al., 2004; Ferreres et al., 2005; 2006). The compound 5 was found only in *B. napus* extracts in  
13  
14  
15 245 trace quantities.

16  
17 246 *Acylated flavonoids.*

18  
19  
20 247 Several of the flavonoids in the Brassica sample extracts had UV spectra with a broad  
21  
22 248 maximum absorbance around 330-340 nm (Table 2), suggesting they were acylated with  
23  
24 249 hydroxycinnamic acids (Fig. 2). The comparison of the HPLC-DAD chromatogram of the  
25  
26  
27 250 extracts with that of the saponified extract (Fig.2) indicated the existence of acylated  
28  
29 251 compounds in high amounts. The MS study of these compounds allowed the detection of a  
30  
31 252 total of 13 acylated flavonol glycosides (compounds 10-22). These compounds were acyl  
32  
33  
34 253 derivatives from compounds 1 (10, 13), 3 (11, 12, 14, 15, 16, 17), 6 (21) and 8 (18, 19, 20,  
35  
36 254 22). The fragmentation of some of these acylated derivatives showed in MS<sup>2</sup> a base peak  
37  
38  
39 255 resultant from the loss of the sugar in the 7 position [(M-H)-162]<sup>-</sup>. This fragmentation is  
40  
41 256 typical from flavonid-3-O-(acyl)glycoside-7-O-hexoside and has been widely described in  
42  
43 257 different *Brassicac*s (Llorach et al., 2003a; Ferreres et al., 2005; 2006; 2008). Other important  
44  
45  
46 258 ion also detected was due to the loss of the acyl radical and/or the sugar and acids from the  
47  
48 259 [M-H]<sup>-</sup>. The resulting fragmentation after the loss of the sugar residues at 7 position (-  
49  
50  
51 260 MS<sup>3</sup>[(M-H)→(M-H-162)]<sup>-</sup>), showed that the acid loss is easily detected and that the acylation  
52  
53 261 is always present on the sugars at 3 position in these compounds. Losses of 162, 206, 176,  
54  
55 262 146 and 192 *m/z* has been identified as caffeic acid, sinapic acid, ferulic acid, *p*-coumaric  
56  
57  
58 263 acid and methoxycaffeic, respectively. The ion corresponding to the flavonoid 3-O-glycoside  
59  
60 264 was always the base peak in MS<sup>3</sup> of these compounds. Thus, they had been characterized as

1  
2  
3 265 acylated derivatives of quercetin-3-*O*-sophoroside-7-*O*-glucoside with caffeoyl (**10**) and  
4  
5 266 sinapoyl (**13**), and kaempferol-3-*O*-sophoroside-7-*O*-glucoside with methoxycaffeoyl (**11**),  
6  
7 267 caffeoyl (**12** and **17**), sinapoyl (**14**), feruloyl (**15**) and *p*-coumaroyl (**16**). Compounds **18-22**  
8  
9 268 presented a fragmentation MS<sup>2</sup> [M-H]<sup>-</sup> similar to the MS<sup>3</sup> of previous compounds (Table 2),  
10  
11 269 what is expected of flavonoids with glycosilation on a single phenolic hydroxyl. In addition  
12  
13 270 an ion resulting from loss of the acyl radical and fragment *m/z* 180 (162 +18) [(M-H)-acyl-  
14  
15 271 180]<sup>-</sup> that comes from the interglycosidic breakdown was observed in the fragmentation of  
16  
17 272 **19-22**, confirming the structure of flavonoid-*O*-diglycosides. These compounds have been  
18  
19 273 characterized as acyl derivatives of kaempferol-3-*O*-sophoroside with *p*-coumaroyl (**18**),  
20  
21 274 methoxycaffeoyl (**19**), caffeoyl (**20**), sinapoyl (**22**) and quercetin-3-*O*-(feruloyl) sophoroside  
22  
23 275 (**21**).  
24  
25  
26  
27  
28  
29  
30  
31

### 32 277 *Hydroxycinnamic acids identification.*

33  
34 278 Ten hydroxycinnamic acids and derivatives (**3CQAc**, **3pCoQAc**, **SG**, **4CQAc**, **4FQAc**, **SA**,  
35  
36 279 **A**, **B**, **C**, and **D**) were detected in leaves of leaf rape, kale and cabbage (Table 3, Fig. 2). The  
37  
38 280 most abundant in the three crops were 3-caffeoyl quinic acid (**3CQAc**) (*Rt* 17.3 min; UV  
39  
40 281 295sh, 325 nm; MS: 353, MS<sub>2</sub>(353): 191 (100%), 179 (62%)), 3-*p*-coumaroyl quinic acid  
41  
42 282 (**3pCoQAc**) (*Rt* 19.1 min; UV 311 nm; MS: 337, MS<sub>2</sub>(337): 191 (7%), 179 (100%)) [30], and  
43  
44 283 sinapoylglucoside (**SG**) (*Rt* 20.5 min; UV 329 nm; MS: 285, MS<sub>2</sub>(285): 291 (100%), 223  
45  
46 284 (85%)). 4-caffeoyl quinic (**4CQAc**) (*Rt* 19.6 min; UV 295sh, 326 nm; MS: 353, MS<sub>2</sub>(353):  
47  
48 285 191 (16%), 179 (53), 173 (100%), 191 (16), 135 (12%)) and 4-feruloyl quinic (**4FQAc**) (*Rt*  
49  
50 286 22.9 min; UV 325 nm; MS: 367, MS<sub>2</sub>(367): 191 (5%), 173 (3%), 163 (100%), 119 (5%))  
51  
52 287 were also identified in kale and cabbage but not in leaf rape. Derivatives formed from the  
53  
54 288 interaction of hydroxycinnamic acids with quinic acid and glucose were previously reported  
55  
56 289 in kale, pak choi, Chinese leaf mustard, turnip greens and turnip tops (Rochfort et al., 2006;  
57  
58  
59  
60



1  
2  
3 290 Ayaz et al., 2008; Ferreres et al., 2008; Francisco et al., 2009; Olsen et al., 2009). Sinapic acid  
4  
5 291 (SA) (*Rt* 27.3 min; UV 329 nm; MS: 223, MS<sup>2</sup>(223): 208 (35%), 179(30%), 164(100%)) was  
6  
7 292 a compound present in high quantities in *B. napus* and detected in trace amounts in the two *B.*  
8  
9 293 *oleracea* crops.  
10  
11 294 Other identified hydroxycinnamic acid derivatives were sinapic and ferulic acids esterified  
12  
13 295 carrying more than one hexose moiety (compounds **A**, **B**, **C** and **D**). It could be observed in  
14  
15 296 all cases, the loss of 224 *m/z* from the deprotonated molecular ion, corresponding to sinapic  
16  
17 297 acid (Table 3). Compound **A** and **D** also presented ferulic acid and displayed the loss of this  
18  
19 298 acid (194 *m/z*). By comparison with data reported earlier in other *Brassicac*s (Llorach et al.,  
20  
21 299 2003b; Ferreres et al., 2006) these compounds were tentatively identified as: 1,2-  
22  
23 300 disinapoylgentiobioside (**A**), 1-sinapoyl-2-feruloylgentiobioside (**B**), 1, 2, 2'-  
24  
25 301 trisinapoylgentiobioside (**C**), and 1,2'-disinapoyl-2-feruloylgentiobioside (**D**). These results  
26  
27 302 are in accordance with compounds detected in other *Brassicac* species, like turnip tops  
28  
29 303 (Romani et al., 2006; Francisco et al., 2009), tronchuda cabbage (Ferreres et al., 2006)  
30  
31 304 broccoli (Vallejo et al., 2004) and now for first time in kale and leaf rape.  
32  
33  
34  
35  
36  
37  
38  
39  
40

## 41 306 CONCLUSIONS

42  
43 307 A method to simultaneously extract and identify glucosinolates and phenolic compounds  
44  
45 308 (Bennett et al., 2003; 2006) is used for first time in *B. oleracea acephala* and *capitata* groups  
46  
47 309 and *B. napus* crops. Twelve intact glucosinolates and 32 phenolic compounds were  
48  
49 310 simultaneously characterized by LC/UV-PAD/ESI-MS<sup>n</sup>. The major flavonoid kaempferol  
50  
51 311 was found as mono-, di-, and triglycosides acylated with different hydroxycinnamic acids.  
52  
53 312 The major hydroxycinnamic acid derivatives were present as esters of sinapic acid, ferulic  
54  
55 313 acid, and glycosides. The highest difference between species is the number of sugar residues  
56  
57  
58  
59  
60



1  
2  
3 314 present on the flavonol core. It may also be expected that the different glycosides possess  
4  
5 315 different biological activities for human health.  
6  
7  
8 316 This study shows that kale, cabbage and leaf rape are a good source of phenolic antioxidants.  
9  
10 317 The main naturally occurring phenolic compounds identified were flavonols and  
11  
12 318 hydroxycinnamic acids. The majority of the flavonoids found in these varieties are  
13  
14  
15 319 kaempferol glycosylated and acylated with different hydroxycinnamic acids. Quercetin and  
16  
17 320 isorhamnetin derivatives were also found. Kaempferol and quercetin are the most prevalent  
18  
19 321 flavonoids in the *Brassicaceae* family. Kaempferol is known to be a strong antioxidant and  
20  
21 322 quercetin also a potent free radical scavenger and is considered to be a protective against  
22  
23 323 cardiovascular disease. Cabbage is a well established crop at world level as cauliflower or  
24  
25 324 broccoli. Kale and leaf rape are minority crops in many parts of the world but they are a good  
26  
27 325 source of nutritive compounds and due to their rusticity (Rodriguez et al., 2005; Padilla et al.,  
28  
29 326 2007b) could be a good substitute of different *Brassica* species under hard climatic  
30  
31 327 conditions.  
32  
33  
34  
35  
36  
37  
38  
39 328

### 329 **Acknowledgements**

40  
41 330 Research supported by the National Plan for Research and Development (AGL2009-09922)  
42  
43 331 and the Excm. Diputación Provincial de Pontevedra. Marta Francisco acknowledges a I3P  
44  
45 332 Fellowship from the CSIC. The authors thank Rosaura Abilleira and Susana Calvo for all the  
46  
47 333 invaluable help in the laboratory work.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

336 **References**

- 337 Ayaz FA, S Hayirlioglu-Ayaz, S Alpay-Karaoglu, J Gruz, K Valentova, J Ulrichova, and M  
338 Strnad. 2008. Phenolic acid contents of kale (*Brassica oleraceae* L. var. *acephala*  
339 DC.) extracts and their antioxidant and antibacterial activities. *Food Chem.* 107:19-25.
- 340 Bennett RN, EAS Rosa, FA Mellon, and PA Kroon. 2006. Ontogenic profiling of  
341 glucosinolates, flavonoids, and other secondary metabolites in *Eruca sativa* (salad  
342 rocket), *Diplotaxis eruroides* (wall rocket), *Diplotaxis tenuifolia* (wild rocket), and  
343 *Bunias orientalis* (Turkish rocket). *J. Agric. Food Chem.* 54:4005-4015.
- 344 Bennett RN, FA Mellon, N Foidl, JH Pratt, MS Dupont, L Perkins, and PA Kroon. 2003.  
345 Profiling glucosinolates and phenolics in vegetative and reproductive tissues of the  
346 multi-purpose trees *Moringa oleifera* L. (horseradish tree) and *Moringa stenopetala* L.  
347 *J. Agric. Food Chem.* 51:3546-3553.
- 348 Cartea ME, and P Velasco. 2008. Glucosinolates in Brassica foods: Bioavailability in food  
349 and significance for human health. *Phytochem. Rev.* 7:213-229.
- 350 Cartea ME, VM Rodriguez, A de Haro, P Velasco, and A Ordas. 2008a. Variation of  
351 glucosinolates and nutritional value in nabicol (*Brassica napus pabularia* group).  
352 *Euphytica* 159:111-122.
- 353 Cartea ME, P Velasco, S Obregon, G Padilla, and A de Haro. 2008b. Seasonal variation in  
354 glucosinolate content in *Brassica oleracea* crops grown in northwestern Spain.  
355 *Phytochemistry* 69:403-10.
- 356 Cermak R, S Landgraf, and S Wolfram. 2003. The bioavailability of quercetin in pigs  
357 depends on the glycoside moiety and on dietary factors. *J. Nutr.* 133:2802-2807.
- 358 Clarke JD, RH Dashwood, and E Ho. 2008. Multi-targeted prevention of cancer by  
359 sulforaphane. *Cancer Lett.* 269:291-304.
- 360 Duthie GG, SJ Duthie, and JAM Kyle. 2000. Plant polyphenols in cancer and heart disease:  
361 implications as nutritional antioxidants. *Nutrition Research Reviews* 13:79-106.
- 362 Fabre N, V Poinot, L Debrauwer, C Vigor, J Tulliez, I Fouraste, and C Moulis. 2007.  
363 Characterisation of glucosinolates using electrospray ion trap and electrospray  
364 quadrupole time-of-flight mass spectrometry. *Phytochem. Anal.* 18:306-319.
- 365 Ferreres F, R Llorach, and A Gil-Izquierdo. 2004. Characterization of the interglycosidic  
366 linkage in di-, tri-, tetra- and pentaglycosylated flavonoids and differentiation of  
367 positional isomers by liquid chromatography/electrospray ionization tandem mass  
368 spectrometry. *J. Mass Spectrom.* 39:312-321.
- 369 Ferreres F, C Sousa, V Vrchovska, P Valentao, JA Pereira, RM Seabra, and PB Andrade.  
370 2006. Chemical composition and antioxidant activity of tronchuda cabbage internal  
371 leaves. *Eur. Food Res. Technol.* 222:88-98.
- 372 Ferreres F, P Valentao, JA Pereira, A Bento, A Noites, RM Seabra, and PB Andrade. 2008.  
373 HPLC-DAD-MS/MS-ESI screening of phenolic compounds in *Pieris brassicae* L.  
374 reared on *Brassica rapa* var. *rapa* L. *J. Agric. Food Chem.* 56:844-853.
- 375 Ferreres F, P Valentao, R Llorach, C Pinheiro, U Cardoso, JA Pereira, C Sousa, RM Seabra,  
376 and PB Andrade. 2005. Phenolic compounds in external leaves of tronchuda cabbage  
377 (*Brassica oleracea* L. var. *costata* DC). *J. Agric. Food Chem.* 53:2901-2907.
- 378 Francisco M, DA Moreno, ME Cartea, F Ferreres, C Garcia-Viguera, and P Velasco. 2009.  
379 Simultaneous identification of glucosinolates and phenolic compounds in a  
380 representative collection of vegetable *Brassica rapa*. *J Chromatogr A* 1216:6611-9.
- 381 Jahangir M, HK Kim, YH Choi, and R Verpoorte. 2009. Health-Affecting Compounds in  
382 Brassicaceae. *Comprehensive Reviews in Food Science and Food Safety* 8:31-43.
- 383 Jeffery EH, and M Araya. 2009. Physiological effects of broccoli consumption. *Phytochem.*  
384 *Rev.* 8:283-298.

- 1  
2  
3 385 Kokkonen PS, J Vandergreef, WMA Niessen, UR Tjaden, GJ Tenhove, and G Vandewerken.  
4 386 1991. Identification of intact glucosinolates using direct coupling of high-performance  
5 387 Liquid-Chromatography with continuous-flow frit fast-atom-bombardment tandem  
6 388 mass-spectrometry. *Biological Mass Spectrometry* 20:259-267.
- 7 389 Kushad MM, AF Brown, AC Kurilich, JA Juvik, BP Klein, MA Wallig, and EH Jeffery.  
8 390 1999. Variation of glucosinolates in vegetable crops of *Brassica oleracea*. *J. Agric.*  
9 391 *Food Chem.* 47:1541-1548.
- 10 392 Llorach R, JC Espin, FA Tomas-Barberan, and F Ferreres. 2003a. Valorization of cauliflower  
11 393 (*Brassica oleracea* L. var. *botrytis*) by-products as a source of antioxidant phenolics.  
12 394 *J. Agric. Food Chem.* 51:2181-2187.
- 13 395 Llorach R, A Gil-Izquierdo, F Ferreres, and FA Tomas-Barberan. 2003b. HPLC-DAD-  
14 396 MS/MS ESI characterization of unusual highly glycosylated acylated flavonoids from  
15 397 cauliflower (*Brassica oleracea* L. var. *botrytis*) agroindustrial byproducts. *J. Agric.*  
16 398 *Food Chem.* 51:3895-3899.
- 17 399 Martens DA. 2002. Identification of phenolic acid composition of alkali-extracted plants and  
18 400 soils. *Soil Sci. Soc. Am. J.* 66:1240-1248.
- 19 401 Mellon FA, RN Bennett, B Holst, and G Williamson. 2002. Intact glucosinolate analysis in  
20 402 plant extracts by programmed cone voltage electrospray LC/MS: Performance and  
21 403 comparison with LC/MS/MS methods. *Anal. Biochem.* 306:83-91.
- 22 404 Mithen R, K Faulkner, R Magrath, P Rose, G Williamson, and J Marquez. 2003.  
23 405 Development of isothiocyanate-enriched broccoli, and its enhanced ability to induce  
24 406 phase 2 detoxification enzymes in mammalian cells. *Theor. Appl. Genet.* 106:727-734.
- 25 407 Olsen H, K Aaby, and GIA Borge. 2009. Characterization and Quantification of Flavonoids  
26 408 and Hydroxycinnamic Acids in Curly Kale (*Brassica oleracea* L. Convar. *acephala*  
27 409 Var. *sabellica*) by HPLC-DAD-ESI-MSn. *J. Agric. Food Chem.* 57:2816-2825.
- 28 410 Padilla G, ME Cartea, and A Ordas. 2007a. Comparison of several clustering methods in  
29 411 grouping kale landraces. *J. Am. Soc. Hort. Sci.* 132:387-395.
- 30 412 Padilla G, ME Cartea, P Soengas, and A Ordas. 2007b. Characterization of fall and spring  
31 413 plantings of Galician cabbage germplasm for agronomic, nutritional, and sensory  
32 414 traits. *Euphytica* 154:63-74.
- 33 415 Padilla G, ME Cartea, P Velasco, A de Haro, and A Ordas. 2007c. Variation of glucosinolates  
34 416 in vegetable crops of *Brassica rapa*. *Phytochemistry* 68:536-545.
- 35 417 Picoaga A, ME Cartea, P Soengas, L Monetti, and A Ordas. 2003. Resistance of kale  
36 418 populations to lepidopterous pests in northwestern Spain. *J. Econ. Entomol.* 96:143-  
37 419 147.
- 38 420 Rochfort SJ, M Imsic, R Jones, VC Trenerry, and B Tomkins. 2006. Characterization of  
39 421 flavonol conjugates in immature leaves of pak choi [*Brassica rapa* L. Ssp *chinensis* L.  
40 422 (Hanelt.)] by HPLC-DAD and LC-MS/MS. *J. Agric. Food Chem.* 54:4855-4860.
- 41 423 Rochfort SJ, VC Trenerry, M Imsic, J Panozzo, and R Jones. 2008. Class targeted  
42 424 metabolomics: ESI ion trap screening methods for glucosinolates based on MSn  
43 425 fragmentation. *Phytochemistry* 69:1671-1679.
- 44 426 Rodriguez VM, ME Cartea, G Padilla, P Velasco, and A Ordas. 2005. The nabicol: A  
45 427 horticultural crop in northwestern Spain. *Euphytica* 142:237-246.
- 46 428 Romani A, P Vignolini, L Isolani, F Ieri, and D Heimler. 2006. HPLC-DAD/MS  
47 429 characterization of flavonoids and hydroxycinnamic derivatives in turnip tops  
48 430 (*Brassica rapa* L. subsp *sylvestris* L.). *J. Agric. Food Chem.* 54:1342-1346.
- 49 431 Soengas P, ME Cartea, P Velasco, G Padilla, and A Ordas. 2008. Morphologic and agronomic  
50 432 diversity of *Brassica napus* crops. *J. Am. Soc. Hort. Sci.* 133:48-54.
- 51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 433 Sousa C, M Taveira, P Valentao, F Fernandes, JA Pereira, L Estevinho, A Bento, F Ferreres,  
4 434 RM Seabra, and PB Andrade. 2008. Inflorescences of Brassicacea species as source of  
5 435 bioactive compounds: A comparative study. *Food Chem.* 110:953-961.  
6 436 Traka M, and R Mithen. 2009. Glucosinolates, isothiocyanates and human health. *Phytochem.*  
7 437 *Rev.* 8:269-282.  
8 438 Vallejo F, FA Tomas-Barberan, and F Ferreres. 2004. Characterisation of flavonols in  
9 439 broccoli (*Brassica oleracea* L. var. *italica*) by liquid chromatography-UV diode-array  
10 440 detection-electrospray ionisation mass spectrometry. *J. Chromatogr.* 1054:181-193.  
11 441 Velasco P, P Soengas, M Vilar, ME Cartea, and M del Rio. 2008. Comparison of  
12 442 glucosinolate profiles in leaf and seed tissues of different Brassica napus crops. *J. Am.*  
13 443 *Soc. Hort. Sci.* 133:551-558.  
14 444 Vilar M, ME Cartea, G Padilla, P Soengas, and P Velasco. 2008. The potential of kales as a  
15 445 promising vegetable crop. *Euphytica* 159:153-165.  
16 446  
17 447  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

Table 1. List of identified glucosinolates with the corresponding retention times and MS data in extracts of leaf rape (*B. napus pabularia* group), cabbage (*B. oleracea capitata* group), and kale crops (*B. oleracea acephala* group)..

Code	Compound	Rt (min)	m/z [M-H] <sup>-</sup>	MS <sup>2</sup> [M-H] <sup>-</sup> m/z (%)	Leaf rape	Kale	Cabbage
<b>GIB</b>	Glucoiberin	4.8	422	358(100), 342(1), 291(2), 275(3), 259(5), 195(2), 180(6)	X	X	X
<b>PRO</b>	Progoitrin	5.9	388	332(29), 308(12), 301(17), 275(40), 259(100), 240(18) 210(78), 195(14), 192(24), 154(16), 136(42), 130(30)	X	-	-
<b>SIN</b>	Sinigrin	6.6	358	278(2), 275(7), 259(100), 241(28), 195(4), 180(3), 162(8), 135(2), 116(5)	X	X	X
<b>GRA</b>	Glucoraphanin	7.4	436	420(6), 372(100), 356(1), 291(2), 275(1), 259(3), 194(2), 162(4)	X	X	X
<b>GNL</b>	Gluconapoleiferin	12.3	402	306(53), 275(20), 259(100), 240(7), 225(10), 215(24), 195(10), 163(13), 160(9), 145(18), 140(23), 120(9)	X	-	-
<b>GNA</b>	Gluconapin	12.5	472	292(4), 275(30), 259(100), 241(37), 227 (9), 195(25), 176(8), 139 (11), 130(10)	X	-	-
<b>4-OHGBS</b>	4-Hydroxigluco brassicin	13.1	463	403(2), 383(10), 365(6), 300(6), 285(73), 267(100), 259(17), 240(25), 220(17), 169(30), 160(23), 132(5)	X	X	X
<b>GBN</b>	Glucobrassicinapin	15.7	386	306(4), 275(21), 259(100), 241(33), 208(12), 195(6), 163(4), 144(18) 139(8)	X	-	-
<b>GBS</b>	Glucobrassicin	16.8	447	367(22), 291(6), 275(34), 259(100), 251(17), 241(11), 224(3), 205(28), 195(11), 144(9)	X	X	X
<b>GST</b>	Gluconasturtiin	19.5	422	342(9), 275(24), 259(100), 244(7), 229(8), 195(13), 180(24), 169(1), 163(5), 145(4), 140(6), 119(4)	X	X	X
<b>4-MGBS</b>	4-Methoxyglucobrassicin	20.4	477	397(25), 299(14), 291(69), 275(98), 259(62), 241(80), 235(100), 198(80), 195 (16), 144(70)	-	X	X
<b>NGBS</b>	Neoglucobrassicin	23.5	477	447(68), 446(100), 429(16), 385(5), 273(12), 259(16), 241(26), 224(4), 205(5)	X	X	X


Table 2. Data of glycosilated flavonoids from saponificated extract (compounds 1-9) and acylated flavonoids from native extracts of leaf rape (*B. napus pabularia* group), cabbage (*B. oleracea capitata* group), and kale crops (*B. oleracea acephala* group). (compounds 10-22).

Peak	Compound	Rt (min)	UV (nm)	[M-H] <sup>-</sup> (m/z)	MS <sup>2</sup> [M-H] <sup>-</sup> m/z (%)	MS <sup>3</sup>	Leaf rape	Kale	Cabbage
1	Querc-3- <i>O</i> -shop-7- <i>O</i> -glc	16.9	255, 267sh, 351	787	625(100), 463(5), 301(3)	463(23), 445(21), 300(100)	-	X	X
2	Kaempf-3- <i>O</i> -triglc-7- <i>O</i> -glc	17.0	266, 316sh, 348	933	771(100)	489(15), 429(19), 285(100)	X	X	X
3	Kaempf-3- <i>O</i> -soph-7- <i>O</i> -glc	17.4	266, 317sh, 347	771	609(100)	489(25), 429(21), 285(100)	X	X	X
4	Kaempf-3,7-di- <i>O</i> -glc	19.6	265, 317sh, 347	609	488(8), 447(100), 285(34)	285(100)	X	X	X
5	Isorhmnt-3,7-di- <i>O</i> -glc	19.9	255, 267sh, 352	639	519(12), 477(100), 315(19)	314(100)	X	-	-
6	Querc-3- <i>O</i> -soph	20.5	-	625	463(10), 445(51), 300(100)	-	-	X	X
7	Kaempf-3- <i>O</i> -diglc	22.8	266, 297sh, 347	609	447(100), 285(80)	285(100)	-	X	X
8	Kaempf-3- <i>O</i> -soph	27.4	266, 297sh, 347	609	447(21), 429(65), 285(100)	-	-	X	X
9	Kaempf-7- <i>O</i> -glc	28.1	266, 318sh, 367	447	285(100)	-	X	X	X
10	Querc-3- <i>O</i> -(caffeoyl)-shop-7- <i>O</i> -glc	17.8	253, 269sh, 299sh, 337	949	787(100), 625(50)	625(100)	X	X	X
11	Kaempf-3- <i>O</i> -(methoxycaffeoyl)-soph-7- <i>O</i> -glc	18.4	269, 331	963	801(100), 609(7)	609(100), 285(12)	X	X	X
12	Kaempf-3- <i>O</i> -(caffeoyl)-soph-7- <i>O</i> -glc	18.6	269, 331	933	771(100), 609(7)	609(100), 285(10)	X	X	X
13	Querc-3- <i>O</i> -(sinapyl)-shop-7- <i>O</i> -glc	18.8	270, 337	993	831(100), 787(60), 625(65)	625(100)	X	X	X
14	Kaempf-3- <i>O</i> -(sinapoyl)-soph-7- <i>O</i> -glc	19.6	269, 331	977	815(100), 609(3)	609(100), 285(10)	X	X	X
15	Kaempf-3- <i>O</i> -(feruloyl)-soph-7- <i>O</i> -glc	20.0	Coeluting with 16	947	785(100), 609(6)	609(5)	X	X	X
16	Kaempf-3- <i>O</i> -(p-coumaroyl)-soph-7- <i>O</i> -glc	20.1	Coeluting with 15	917	755(100), 609(5),	-	X	X	X
17	Kaempf-3- <i>O</i> -(caffeoyl)-soph-7- <i>O</i> -glc (isomer)	20.8	269, 331	933	771(100), 609(9)	609(100)	X	-	-
18	Kaempf-3- <i>O</i> -(p-coumaroyl)-soph	23.4	269, 317	755	609(100), 429(1), 284(15)	429(35), 285(100)	-	X	X
19	Kaempf-3- <i>O</i> -(methoxycaffeoyl)-soph	24.3	268, 330	801	609(100), 429(5), 285(10)	429(25), 285(100)	X	X	X
20	Kaempf-3- <i>O</i> -(caffeoyl)-soph	24.7	269, 330	771	609(100), 429(4), 285(9)	428(34), 285(100)	X	X	X
21	Querc-3- <i>O</i> -(feruloyl)-soph	25.1	-	801	625(100), 445(5), 301(4)	445(15), 301(100)	X	X	X
22	Kaempf-3- <i>O</i> -(sinapoyl)-soph	25.6	269, 330	815	609(100), 429(3), 285(9)	429(30), 285(100)	X	X	X

Table 3. Data of hydroxycinnamic acids from extracts of leaf rape (*B. napus pabularia* group), cabbage (*B. oleracea capitata* group), and kale crops (*B. oleracea acephala* group).

Peak	Compound	Rt (min)	UV (nm)	[M-H] <sup>-</sup> (m/z)	MS <sup>2</sup> [M-H] <sup>-</sup> m/z (%)	MS <sup>3</sup> [(M-H)→(M-H-224)] <sup>-</sup>	Leaf rape	Kale	Cabbage
<b>3CQAc</b>	3-Caffeoyl quinic acid	17.3	295sh, 325	353	191 (100), 179 (62)	-	X	X	X
<b>3pCoQAc</b>	3- <i>p</i> -Coumaroyl quinic acid	19.1	311	337	191 (7), 179 (100)	-	X	X	X
<b>4CQAc</b>	4-Caffeoyl quinic acid	19.6	295sh, 326	353	191 (16), 179 (53), 173 (100), 135 (12)	-	-	X	X
<b>SG</b>	Sinapylglucoside	20.5	329	285	291(100%), 223 (85%)	-	X	X	X
<b>4FQAc</b>	4-Feruloyl quinic acid	22.9	325	367	191 (5), 173 (3), 163 (100), 119 (5)	-	-	X	X
<b>SA</b>	Sinapic acid	27.3	329	223	208 (35), 179 (30), 164 (100)	-	X	<i>In traces</i>	<i>In traces</i>
<b>A</b>	1,2-Disinapoylgentiobioside	28.1	330	753	529 (100), 223 (1)	223 (100), 205 (59)	X	X	X
<b>B</b>	1-Sinapoyl-2-feruloylgentiobioside	28.7	330	723	529 (11), 499 (100), 259 (4)	259 (28), 246 (20), 217 (53)	X	X	X
<b>C</b>	1,2,2'-Trisinapoylgentiobioside	30.5	330	959	735 (100)	717 (15), 529 (100)	X	X	X
<b>D</b>	1,2'-Disinapoyl-2-feruloylgentiobioside	31.2	330	929	735 (4), 705 (100)	499 (100)	X	X	X




Figure 1. Glucosinolate profile of leaf rape (*B. napus pabularia* group), cabbage (*B. oleracea capitata* group), and kale crops (*B. oleracea acephala* group).



1  
2  
3  
4 **GIB**= Glucoiberin. **PRO**= Progoitrin. **SIN**= Sinigrin. **GRA**=Glucoraphanin. **GNL**=Gluconapoleiferin. **GNA**=Gluconapin. **4-OHGBS**=4-Hydroxigluco brassicin.  
5  
6 **GBN**=Glucobrassicinapin. **GBS**=Glucobrassicin. **GST**=Gluconasturtiin. **4-MGBS**=4-Methoxygluco brassicin. **NGBS**=Neogluco brassicin.  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

For Peer Review

Figure 2. Phenolic profiles of leaf phenolics in rape (*B. napus pabularia* group), cabbage (*B. oleracea capitata* group), and kale (*B. oleracea acephala* group).

1  
2  
3  
4 **1**=Querc-3-*O*-shop-7-*O*-glc. **2**=Kaempf-3-*O*-triglc-7-*O*-glc. **3**=Kaempf-3-*O*-soph-7-*O*-glc. **4**=Kaempf-3,7-di-*O*-glc. **5**=Isorhmnt-3,7-di-*O*-glc. **6**=Querc-3-*O*-soph.  
5 **7**=Kaempf-3-*O*-diglc. **8**=Kaempf-3-*O*-soph. **9**=Kaempf-7-*O*-glc. **10**=Querc-3-*O*-(caffeoyl)-shop-7-*O*-glc. **11**=Kaempf-3-*O*-(methoxycaffeoyl)-soph-7-*O*-glc.  
6 **12**=Kaempf-3-*O*-(caffeoyl)-soph-7-*O*-glc. **13**=Querc-3-*O*-(sinapyl)-shop-7-*O*-glc. **14**=Kaempf-3-*O*-(sinapoyl)-soph-7-*O*-glc. **15**=Kaempf-3-*O*-(feruloyl)-soph-7-*O*-glc.  
7 **16**=Kaempf-3-*O*-(*p*-coumaroyl)-soph-7-*O*-glc. **17**=Kaempf-3-*O*-(caffeoyl)-soph-7-*O*-glc (isomer). **18**=Kaempf-3-*O*-(*p*-coumaroyl)-soph. **19**=Kaempf-3-*O*-  
8 (methoxycaffeoyl)-soph. **20**=Kaempf-3-*O*-(caffeoyl)-soph. **21**=Querc-3-*O*-(feruloyl)-soph. **22**=Kaempf-3-*O*-(sinapoyl)-soph. **3CQAc**=3-Caffeoyl quinic acid.  
9 **3pCoQAc**=3-*p*-Coumaroyl quinic acid. **4CQAc**=4-Caffeoyl quinic acid. **SG**=Sinapylglucoside. **4FQAc**=4-Feruloyl quinic acid. **SA**=Sinapic acid. **A**=1,2-  
10 Disinapoylgentiobioside. **B**=1-Sinapoyl-2-feruloylgentiobioside. **C**=1,2,2'-Trisinapoylgentiobioside. **D**=1,2'-Disinapoyl-2-feruloylgentiobioside  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

For Peer Review