

1 **Effect of different chemical compounds as coadjutants of 4-hexylresorcinol on**
2 **appearance of deepwater pink shrimp (*Parapenaeus longirostris*) during chilled**
3 **storage**

4 Oscar Martínez-Alvarez*, Carmen Gómez-Guillén and Pilar Montero

5 Instituto del Frío (C.S.I.C.), José Antonio Novais 10, 28040 Madrid (Spain)

6 Tel: +345492300; Fax: +345493627

7 Author to whom correspondence should be addressed [e-mail
8 oscar_martine50@hotmail.com]

9 **ABBREVIATED RUNNING TITLE: Appearance of pink shrimp immersed in 4-**
10 **hexylresorcinol-based formulas**

11
12 **ABSTRACT**

13 Different chemical compounds (kojic acid, cumic acid phytic acid, sodium metabisulphite,
14 magnesium carbonate, sorbic acid, and different protease inhibitors) were used as
15 coadjutants in 4-hexylresorcinol-based melanosis-inhibiting formulas tested by immersion
16 on pink shrimp (*Parapenaeus longirostris*). Increasing concentrations of 4-hexylresorcinol
17 delayed the occurrence of melanosis during storage. However, they could not prevent the
18 appearance of yellow-greenish colouration in the cephalothorax, which diminished the
19 consumer acceptability of shrimps. The incorporation of protease inhibitors (EDTA,
20 disodium dihydrogen pyrophosphate, iodoacetic acid, egg white and PMSF) into the 4-
21 hexylresorcinol-based blends improved acceptability through storage, suggesting protease
22 activity post mortem contributes to the final acceptability of crustaceans.

23 **Keywords:** melanosis, pink shrimp, *Parapenaeus longirostris*, polyphenol oxidase, 4-
24 hexylresorcinol.

25 INTRODUCTION

26 Melanosis is a surface colouration that initiates as soon as crustaceans are removed from
27 water and come into contact with the oxygen in the atmosphere. The reaction involves the
28 oxidation of phenols to quinones by polyphenol oxidase (PPO), providing the precursors of
29 insoluble polymeric pigments (Otwell & Marshall, 1986). Currently, prevention of black
30 spots involves application of sulphite derivatives, although the authorised residual content
31 in flesh is not very effective to delay the appearance of melanosis in some species such as
32 pink shrimp (*Parapenaeus longirostris*). The use of high doses of sulphites is not a good
33 alternative given that it is related to allergic reactions, as well as other health risks
34 associated with its use (Taylor & Bush, 1987). It emphasizes the need for a safe effective
35 sulphite alternative. 4-hexylresorcinol (4-HR), is for example a competitive inhibitor of
36 PPO “generally recognized as safe” (GRAS) for use in the prevention of crustacean
37 melanosis (Frankos et al., 1991). The use of 4-HR is permitted in the US, Canada,
38 Australia, and some Latin American countries (Montero, Martínez-Alvarez & Gómez-
39 Guillén, 2004). In 2003, the Scientific Committee on Food of the European Commission
40 considered 4-HR to be toxicologically acceptable for the prevention of melanosis in
41 shrimps under specific conditions of use (provided residues in crustacean meat do not
42 exceed 2 mg/kg). In 2004, a proposal of amending the EU Directive 95/2/EC on food
43 additives other than colorants and sweeteners was adopted by the European Commission
44 and presented to the European Parliament in order to authorize 4-HR as an alternative to
45 sulphites for preventing the browning of crustaceans. Its effectiveness as melanosis-
46 inhibiting compound has been demonstrated both in laboratory tests and on board
47 (McEvily, Iyengar & Otwell, 1991; Otwell, Iyengar & McEvily, 1992; Guandalini, Ioppolo,
48 Mantovani, Stacchini & Giovannini, 1998; Montero, López-Caballero & Pérez-Mateos,

49 2001a; Montero et al., 2004). However, the effective doses are different depending on
50 several factors such as species, physiological states, and application method. In addition,
51 the experimental conditions cannot be held in the same account, as those used in
52 commercial boats, where fishermen manipulate a considerable amount of shrimp and
53 accurate weight measurement is difficult. That results in amount of melanosis-inhibiting
54 formulations used on board not corresponding to those used in the laboratory.

55 A commercial formulation composed of 4-HR and NaCl (Everfresh®, Opta Food
56 Ingredients) is used in different countries with positive results in delaying melanosis.
57 However, for deepwater pink shrimp (*Parapenaeus longirostris*) this product has proved to
58 be quite ineffective using the concentrations and application times recommended by the
59 manufacturer (unpublished data). That encourages the search for more potentially effective
60 4-HR-based formulations, mainly for species very susceptible to melanosis such as pink
61 shrimp. Montero et al. (2004) reported the application of citric acid, ascorbic acid and
62 acetic acid in association with 4-HR to enhance the appearance of shrimp, accentuating
63 their natural pink colouration. Despite the fact that this formulation inhibited melanosis
64 itself, the viscera turned yellowish during storage, which is attributed to primary post
65 mortem spoilage. The incorporation of coadjuvants in 4-HR based formulations could result
66 in the prevention of the prevalence of yellow shades in the head, thereby improving
67 appearance. These chemicals could be different food-grade compounds for which anti-
68 browning action has been demonstrated in vegetables, fruits and crustaceans. Phytic acid is
69 a natural antioxidant (Graf, Empson & Eaton, 1987) and chelating agent which
70 demonstrates melanosis-inhibiting activity, as been described in vegetables (Hicks et al.,
71 2004), but rarely in crustaceans (Guang-Li, Yuan-hong, Shu-qing & Xue-lin, 1996). Kojic
72 acid is an antibiotic substance produced by several species of *Aspergillus* and *Penicillium*,

73 and is widely used in Japan as a food additive for preventing enzymatic browning. Kojic
74 acid acts by interfering with the uptake of O₂ required for the enzymatic reaction and/or
75 reduction of quinone compounds to diphenols to prevent melanin formation (Chen, Wei, &
76 Marshall, 1991a). Cumic acid, abundant in seeds of *Cuminum cyminum* (cumin), acts by
77 binding to the coupled binuclear copper active site with the carboxylic group and can be
78 classified as a non-competitive inhibitor (Kubo & Kinst-Hori, 1998). Another chemical,
79 magnesium carbonate, is decomposed into carbon dioxide, which interferes with the
80 utilization of oxygen by PPO (Gutierrez Alsina, 1976). Furthermore, several food
81 preservatives, such as potassium sorbate and sulphite derivatives, could also be used to
82 delay the appearance of melanosis mediated by some spoilage bacteria, according to
83 Chinivasagam, Bremner & Reeves, (1998). Other substances show a double role, such as
84 ethylenediaminetetraacetic acid (EDTA), and disodium dihydrogen pyrophosphate (PPi)
85 (Iyengar & McEvily, 1992; Montero et al., 2004). They can chelate the copper prosthetic
86 group at the PPO active site, slowing down the enzymatic reaction, and can also inhibit
87 digestive metalloproteases responsible for PPO activation. Moreover, the inhibition of
88 digestive proteases should diminish free tyrosine concentration in flesh, which is a natural
89 substrate for PPO. Egg white can also be used as protease inhibitor, as it contains serine and
90 cysteine proteases. Other specific non-food grade inhibitors such as phenylmethylsulphonyl
91 fluoride (PMSF) and Iodoacetic acid (IAA) could also be used to study the role of different
92 proteases in the decomposition of viscera during storage. PMSF is an inhibitor of serine
93 proteases, while Iodoacetic acid (IAA) is an inhibitor of cysteine proteases.

94

95 The main objective of this work was to assess the effect of several chemicals used as
96 coadjuvants in 4-HR-based formulations on the occurrence of black spots and yellow-green

97 shades in the head of the deepwater pink shrimp (*Parapenaeus longirostris*) during chilled
98 storage. All formulations were used by dipping on board, simulating conditions habitually
99 used by fishermen. Preservatives, natural melanosis-inhibiting chemicals and protease
100 inhibitors infrequently studied in crustaceans were studied. The effect of these formulations
101 on acceptability was also evaluated through storage.

102 **MATERIALS AND METHODS.**

103 Deepwater pink shrimp (*Parapenaeus longirostris*) were caught off the South coast of
104 Spain (Cádiz) by trawl in March and May. Mean shrimp weights were 8.03 ± 1.54 g. On
105 board they were separated from the by-catch, washed with seawater and separated into
106 groups. Different 4-HR based formulations (Table 1) were then applied by immersion for 1
107 hour (shrimp:seawater relation of 1:2), one group immersed in seawater being considered
108 as control. All melanosis-inhibiting blends (Table 1) included 0.1% or 0.25 % of 4-
109 hexylresorcinol (Sigma-Chemical, St Louis, MO, USA), and in almost all blends it was
110 used in combination with 0.5 % L-ascorbic acid (Sigma Chemical, St Louis, USA), 0.5 %
111 citric acid (Panreac Chemical, Barcelona, Spain), and 0.3% acetic acid (Panreac Chemical,
112 Barcelona, Spain). For simplicity, these formulations were designated as ACRA 0. 1%
113 (with 0.1% 4-HR) and ACRA 0.25% (with 0.25% 4-HR). Resorcinol alone or these basic
114 mixtures were accompanied by different chemical compounds to determine their capacity
115 to improve the appearance of shrimps: magnesium carbonate, di sodium di hydrogen
116 pyrophosphate (PPi), and potassium sorbate (Panreac Chemical, Barcelona, Spain);
117 iodoacetic acid (IAA, Sigma Chemical, St Louis, MO, USA), phenylmethylsulphonyl
118 fluoride (PMSF, Sigma Chemical), kojic acid (5-hydroxy-2 hydroxy-methyl- γ -pyrone,
119 Sigma Chemical), phytic acid, and ethylenediaminetetraacetic acid (EDTA) (Sigma
120 Chemical); cumic acid (4-isopropilbenzoic acid, Aldrich Chemical, Milwaukee, WI, USA),

121 sodium metabisulphite (Sigma Aldrich Chemie, Germany); and dried egg white powder
122 (Degussa, Barcelona, Spain). Each formulation was assayed in two lots (1.5 kg of shrimps
123 per lot). Once the treatment time was finished, the shrimps were removed, placed in
124 perforated polystyrene boxes with a capacity of 2 kg, and covered with flaked ice. On
125 arrival of the trawler at the port, the boxes were taken by refrigerated truck to the Instituto
126 del Frío in Madrid, where they were stored in ice at 2 °C.

127 **Sensorial analyses**

128 A taste panel composed of eleven trained panellists visually inspected 14 shrimp per
129 treatment group every two days during the time of chilled storage. This conventional
130 sensory assessment was used to identify those significant sensory appearance attributes
131 which best define the quality loss of pink shrimp during storage. Those included general
132 outward appearance (acceptability), the condition of the cephalothorax-tail junction, the
133 presence of yellow-greenish colouration beneath the head cuticle and also that of black
134 spots on the shell. The outward appearance was assessed as total percentage of shrimps
135 considered as acceptable for being sold on the market. The presence of yellow-greenish
136 colourations was evaluated as percentage of individuals with visual presence of this shade
137 in the head. The condition of the cephalothorax-tail junction was evaluated as percentage of
138 individuals with cephalothorax and tail separated. Finally, the presence of black spots in the
139 shell was assessed according to a visual scale from 1 to 4 (Montero et al., 2004), in which
140 1 = complete absence of black spots; 2 = a few small spots on the carapace; 3 =
141 considerable spotting on the carapace; 4 = substantial spotting over the entire shrimp. The
142 cephalothorax-tail junction was not affected by the different formulas tested, and for that
143 reason this parameter was not further discussed.

144 **Statistical analyses**

145 The melanosis scores were regressed on time in storage using the SPSS computer software
146 program (SPSS Inc., Chicago, Ill., USA). Linear or polynomial regressions were plotted
147 through data dispersion, showing in all cases regression coefficients $R^2 \geq 0.90$. The data
148 from the different variables analysed were used as the data matrix in different principal
149 component analyses (PCA).

150 **RESULTS AND DISCUSSION.**

151 **Non treated samples (Control samples).** A noticeable increase in all parameters was
152 observed during iced storage, but especially in the case of melanosis, which showed the
153 earliest and more pronounced changes. Firstly, melanosis was observed mainly in the
154 cephalothorax and pleopods. Later, melanosis was also detected in abdomen cuticle and
155 telson, covering the whole surface after seven days of chilled storage (Fig 1). Regarding
156 acceptability, black spots decreased the perceived quality of the shrimp by the test panel,
157 and only approximately 25 % of shrimps were considered as acceptable two days after
158 capture.

159 **Inhibitory effect of 4-hexylresorcinol combined with acids.** The use of the melanosis-
160 inhibiting formula composed of 0.1% 4-HR, 0.5% L-ascorbic acid, 0.5 % citric acid, and
161 0.3% acetic acid led to an absence of black spots in shrimps for the first four days of chilled
162 storage (Fig. 1a). This formula with 0.25 % 4-HR proved to be more effective at inhibiting
163 melanosis (Fig 1b). The effectiveness of 4-HR as inhibitor of PPO activity from crustaceans
164 has been extensively reported (McEvily et al., 1991; Otwell et al., 1992; Guandalini et al.,
165 1998; Montero et al., 2004), although the effect depends on the application method, the
166 quantity applied, and mainly on the crustacean specie. The presence of organic acids in the
167 4-HR-based formula produces an acidic pH, that should negatively affect the PPO activity
168 (Gökoglu, 2004). Furthermore, ascorbic acid causes the chemical reduction of the pigment

169 precursors, and citric acid may have a dual inhibitory effect on PPO: lowering the pH and
170 chelating the copper at the active site of the enzyme (Iyengar & McEvily, 1992). Both
171 ascorbic and citric acid could synergistically interact in the inhibition process, according to
172 Lee-Kim, Hwang & Kim (1997).

173 Despite the finding that ACRA formula inhibited the PPO activity, the heads (beneath
174 cuticle) turned greenish-yellow during storage, affecting acceptability (Table 2). The
175 factor(s) which promote(s) the appearance of this colouration is/are unknown. However, it
176 coincides with the appearance of black spots on carapace, so it is possible that yellow-
177 greenish colouration is in some way related to melanosis process. Autolysis of shrimp
178 viscera, or spoilage bacteria capable of producing melanin (Chinivasagam et al., 1998)
179 could also be involved in the appearance of yellow-greenish shades in heads.

180 Kojic acid was incorporated in 4-HR based formulas as it would complement the inhibitory
181 effect of 4-HR (Chen et al., 1991a). The activity of kojic acid in the prevention of
182 melanosis involves two mechanisms: direct inhibition of PPO, and the chemical reduction
183 of the pigment or pigment precursors to colourless compounds (Iyidogan & Bayindirli,
184 2004). Direct inhibition of PPO is produced as kojic acid is a competitive, slow-binding
185 inhibitor, like 4-HR (Jiménez & García-Carmona, 1997). The melanosis-inhibiting effect of
186 kojic acid has been observed in several crustacean species, such as *Penaeus monodon*
187 (Chen, Wei, Rolle, Otwell, Balaban & Marshall, 1991b) *Penaeus japonicus* (Montero,
188 Avalos & Pérez-Mateos, 2001b), but the effect of this chemical together with 4-HR has not
189 been studied in crustaceans. In this context positive results of melanosis-inhibiting
190 formulations including 4-HR and kojic acid have been described on apples (Iyidogan &
191 Bayindirli, 2004). The effectiveness of kojic acid against melanosis could be higher when
192 mixed with ascorbic acid and citric acid, and those chemicals constitute a Japanese product

193 for inhibiting PPO in foods (Chen et al., 1991b). A mixture of ascorbic acid and kojic acid
194 has also been patented for use as an anti-browning agent in foods (Fukusawa, Wakabayashi
195 & Natori, 1982). Nevertheless, the use by immersion of formulations with 4-HR, organic
196 acids and 0.1% kojic acid had no significant effect on appearance of shrimps from two days
197 of chilled storage (Fig 1 and Table 1). It is worth noting that the bleaching mechanism of
198 quinones by kojic acid seems to be related to a redox reaction leading to an oxidised yellow
199 derivative of the inhibitor (Kahn, 1995), and it could enhance the intensity of yellow colour
200 in head during storage. The author suggested that kojic acid may not be desirable in certain
201 products, but that would depend on the concentration of the inhibitor and the type of food
202 in which it is used.

203 The incorporation of 0.1% cumic and/or phytic acid in the 4-HR-based formulations was
204 also tested (Fig. 1a and 1b). Cumic acid is a non-competitive inhibitor of PPO (Kubo &
205 Kinst-Hori, 1998). Furthermore, phytic acid is an antioxidant and chelating agent, so it
206 would inhibit not only PPO, but also digestive metalloproteases. Intensity of melanosis was
207 attenuated only when those chemicals were included in 0.1% 4-HR-based formulations, and
208 for 9 days of chilled storage. When cumic and/or phytic acid were incorporated in the
209 0.25% 4-HR based formula, the intensity of yellow shade in the head was lower after 7
210 days of chilled storage. However, this was not enough to improve the general appearance of
211 shrimps, according to the test panel.

212 **Inhibitory effect of 4-hexylrerorcinol combined with sulphites.** Sodium metabisulphite
213 was tested in association with 4-HR. This chemical is usually incorporated in commercial
214 melanosis-inhibiting formulations. The use of sulphite derivatives is limited by the
215 regulatory authorities of many countries, which have indicated a maximum concentration of
216 sulphites and derivatives in different foods. However, the appearance of melanosis occurs

217 more rapidly in several shrimp species such as pink shrimp, and for that reason dosage of
218 sulphites exceeds the usually permitted concentrations (Gómez-Guillén, Martínez-Alvarez,
219 Llamas Marcos & Montero, 2005). The use of 4-HR together with sulphite-derivatives in
220 commercial formulations could therefore be of high interest to reduce sulphite
221 concentrations in the flesh of shrimp. Figure 2 depicts the effect of 0.1% 4-HR combined
222 with variable concentrations of sodium metabisulphite routinely used on board (0.62% and
223 1.25%). In both cases, their effect on melanosis and also on the evidence of yellow-
224 greenish colouration beneath cuticle was similar to that achieved by dipping shrimps in a
225 solution of 4-HR alone (Table 3). Nonetheless, lower concentrations of 4-HR were more
226 effective to prevent melanosis than that used for sodium metabisulphite.

227 **Inhibitory effect of 4-HR combined with acids and magnesium carbonate.** Magnesium
228 carbonate exerts a positive effect on melanosis prevention when it is used together with
229 citric, ascorbic, and sodium metabisulphite (Gutierrez Alsina, 1976). Magnesium carbonate
230 is decomposed into carbon dioxide, which interferes in the utilization of oxygen by PPO,
231 preventing decomposition of the viscera. Therefore, the effect of both magnesium
232 carbonate and 4-HR on melanosis could be complementary. The incorporation of 0.5%
233 magnesium carbonate in ACRA-0.1% formulations slightly prevented melanosis, but only
234 after 9 and 11 days of storage (Fig 3). This effect was not noticeable when magnesium
235 carbonate was used together with 0.25 % 4-HR. Moreover, the intensity of yellow-greenish
236 colouration was not reduced (Table 3), registering figures of 100% after 4 days of storage.
237 After 7 days, this colouration was not observed by the panellists due to the intensity of
238 melanosis in the head. Higher concentrations of magnesium carbonate (7 %) decreased
239 final quality after 7 days of storage and did not have any effect on the other parameters
240 judged.

241 **Inhibitory effect of 4-hexylrerorcinol combined with acids and protease inhibitors.**

242 The use of protease-inhibitors (EDTA and PPI) together with 4-HR and organic acids
243 increased acceptability, as shrimps were preserved well during nine days of chilled storage.

244 It should also be noted that, although exhibiting a decrease in sensorial appraisal during
245 storage, melanosis did not reach the threshold rejection value (≤ 2) after 9 days of iced
246 storage, except when egg white was used (Fig. 4). Despite the presence of yellow shades
247 beneath cuticle registering figures higher than 56% after 4 days of storage, almost 100% of
248 shrimps were surprisingly considered acceptable by the panellists during the whole period.

249 That is because the use of protease inhibitors decreased the intensity of the yellow shade in
250 head (Table 4), maintaining the fresh pink appearance. This result is ascribed to the natures
251 of EDTA and PPI, which are metalloprotease inhibitors and consequently may, at least
252 partially, inhibit the formation of free tyrosine and phenylalanine, substrates for the action
253 of PPO. In addition, they can inhibit the PPO activity by chelating the copper prosthetic
254 group at the PPO-active site or reducing the level of copper available for incorporation into
255 the enzyme (McEvily, Iyengar & Otwell, 1992).

256 On the other hand, egg white was added as it contains several inhibitor compounds,
257 including ovomucoid and ovoinhibitor, which are serine-protease inhibitors, and cystatin,
258 which is a cysteine-protease inhibitor (Stevens, 1991). Nonetheless, egg white exerts a
259 detrimental effect on appearance (Table 5). It could be due to the fact that egg white could
260 not be completely dissolved into the dipping solution. Egg white also contains proteins
261 which can be hydrolyzed by endogenous and microbial proteases, producing free
262 aminoacids which could be good substrates for PPO.

263 PMSF and IAA were also incorporated in the 4-HR-based formulas. Both were used only to
264 gain scientific knowledge about their possible mechanism of action in the context of

265 enzymatic browning of crustacean, since they are non food grade chemicals. PMSF is a
266 characteristic inhibitor of serine proteases, while IAA is a thiol-blocking reagent used to
267 inhibit cysteine proteases in many studies (Hameed & Haard, 1985; Yamashita &
268 Konagaya, 1990; Gómez-Guillén, Hurtado & Montero, 2002). Shrimps treated with this
269 formulation showed the lowest melanosis index during storage, probably because the use of
270 protease inhibitors could at least partially inhibit the formation of free tyrosine and
271 phenylalanine, both substrates for the action of PPO. On the other hand, overall quality was
272 improved, but only after 9 days of chilled storage (Table 5), probably because the formula
273 with PMSF and IAA attenuated the yellow shade of head. The slight improvement in the
274 visual aspect could indicate that the appearance of greenish colouration and decomposition
275 of viscera are connected, since serine and cysteine proteases are the main enzymes implied
276 in this process. The use of cystein and serine protease inhibitors should also be important as
277 these proteases are implied in the mechanism of activation of pro-PPO post mortem
278 (Söderhall & Cerenius, 1992; Zotos & Taylor 1997).

279 **Effect of preservatives in 4-HR-based formulations.** Spoilage bacteria could cause
280 melanosis in stored prawns, according to Chinivasagam et al. (1998). To detect the effect of
281 microorganisms on appearance of melanosis or greenish colouration, the preservative
282 potassium sorbate was incorporated in formulations containing 4-HR plus acids. Melanosis
283 in samples treated with and without preservatives evolved similarly during storage (Fig 5),
284 which evidenced the negligible effect of microorganisms on appearance of melanosis.
285 Similar results were observed regarding greenish colouration (Table 5). Furthermore,
286 almost 100 % of shrimps were considered acceptable by the panellists after 7 days of
287 storage.

288 **Principal Component Analyses.** Different principal component analyses (PCA) were
289 performed in addition to sensorial analyses. The aim here was to identify significant
290 differences that might have been overlooked previously in simple sensorial analyses. These
291 further analyses were performed separately for samples treated with formulations including
292 the same chemical compound. The corresponding principal component analyses (PCA) data
293 matrix are shown in Table 7. In all cases, the global data matrix was reduced to 2 principal
294 components (PC), which together account for 84-92 % of the explained variance.
295 The analyses of PC1 showed in all cases that yellow-greenish colouration correlated
296 positively with the time of storage and negatively with the acceptability. That means none
297 of the chemicals used prevented the appearance of yellow shades beneath the head cuticle,
298 which led, together with the appearance of melanosis, to the progressive loss of quality.
299 Even though 4-HR could not stop the appearance of the yellow-greenish colouration, the
300 acceptability and melanosis inhibition were increased with higher concentrations of this
301 chemical, as it is shown by the second component (PC 2, 36 % variance, Table 7a). The
302 incorporation of magnesium carbonate, phytic acid, kojic acid, cumic acid or sulphites in
303 the 4-HR-based formulas did not improve the general acceptability of shrimps (Table 7b-f).
304 On the other hand, the presence of protease inhibitors in the melanosis-inhibiting formulas
305 was related to better general acceptability (Table 7g, PC 2, 26% variance). However, PC 1
306 confirm that parameter acceptability is greatly influenced by both dark and yellow spots.
307 As yellow-greenish colouration was judged only as presence or absence, it is possible that
308 the protease inhibitors improved general appearance by reducing the intensity of this tone
309 beneath the head cuticle.

310 **CONCLUSIONS**

311 The use of increasing concentrations of 4-hexylresorcinol is related to better consumer
312 acceptability of pink shrimp (*Parapenaeus longirostris*) during chilled storage, also
313 inhibiting the evidence of melanosis. Although 4-hexylresorcinol can lessen the appearance
314 of black spots in the carapace, it can not prevent the previous manifestation of yellow-
315 greenish colourations underneath head cuticle. The incorporation of protease inhibitors as
316 coadjuvants in 4-hexylresorcinol-based formulations improved general appearance, but this
317 effect was not statistically significant. This suggests that the appearance of greenish
318 colourations could be related to decomposition of the viscera mediated by proteases, which
319 could facilitate liberation of substrates for PPO such as tyrosine. On the other hand, the
320 incorporation of another natural chemicals as kojic, cumic and phytic acid in 4-
321 hexylresorcinol based formulations did not improve the overall acceptability of shrimps
322 during storage. Finally, the incorporation of preservatives did not have any effect on
323 appearance, suggesting micro-organisms do not participate in the appearance of anomalous
324 colourations in the cephalothorax.

325

326 **ACKNOWLEDGEMENTS**

327 This research was financed under project between Consejería de Agricultura y Pesca (Junta
328 de Andalucía) and Consejo Superior de Investigaciones Científicas (C.S.I.C.).

329

330 **REFERENCES.**

331

332 - Chen, J.S., Wei, C., & Marshall, M.R. (1991a). Inhibition Mechanism of Kojic Acid on
333 Polyphenol Oxidase? *Journal of Agricultural and Food Chemistry*, 39 (11): 1897-1901.

- 334 - Chen, J.S., Wei, C., Rolle, R.S., Otwell, W.S., Balaban, M.O., & Marshall, M.R.
335 (1991b). Inhibitory Effect of Kojic Acid on Some Plant and Crustacean Polyphenol
336 Oxidases? *Journal of Agricultural and Food Chemistry*, 39, 1396-1401.
- 337 - Chinivasagam, H.N., Bremner, H.A., & Reeves, R. (1998). Can spoilage bacteria cause
338 blackspot (melanosis) in stored prawns? *Letters in Applied Microbiology*, 27 (1), 5-8.
- 339 - Frankos, V.H., Schmitt, D.F., Haws, L.C., McEvily, A.J., Iyengar, R., Iller, S.A.,
340 Munro, I.C., Clydesdale, F.M., Forbes, A.L., & Sauer, R.M. (1991). *Regulatory*
341 *Toxicology and Pharmacology*, 14, 202-212.
- 342 - Fukusawa, R., Wakabayashi, H., & Natori, T. (1982). Japanese patent 57 40875.
- 343 - Gökoglu, N. (2004). The effect of organic acid treatments on the melanosis inhibition
344 and shelf-life in shrimp (*Penaeus japonicus*). *Acta Alimentaria*, 33 (2): 191-199.
- 345 - Gomez-Guillén, M.C., Hurtado, J.L., & Montero, P. (2002). Autolysis and protease
346 inhibition effects on dynamic viscoelastic properties during thermal gelation of squid
347 muscle. *Journal of Food Science*, 67 (7): 2491-2496.
- 348 - Gómez-Guillén, M.C., Martínez-Alvarez, O., Llamas Marcos, A., & Montero, P. (2005).
349 Melanosis inhibition and SO₂ residual levels in shrimps (*Parapenaeus longirostris*)
350 after different sulphite based treatments. *Journal of Science of Food and Agriculture*,
351 85, 1143-1148.
- 352 - Graf, E., Empson, K.L., & Eaton, J.W. (1987). Phytic acid. A natural antioxidant.
353 *Journal of Biological Chemistry*, 262 (24), 11647-11650.
- 354 - Guandalini, E., Ioppolo, A., Mantovani, A., Stacchini, P., & Giovannini, C. (1998).
355 *Food Additives and Contaminants*, 15, 171-180.

- 356 - Guang-li, Y., Yuan-hong, W., Shu-qing, L., & Xue-lin, T. (1996). Effects of low
357 molecular weight chitosan (LMC-1) on shrimp preservation. *Chinese Journal of*
358 *Oceanology and Limnology*, 14(2), 189–192.
- 359 - Gutiérrez Alsina, L. (1976). Preservation of crustacean. USA Patent nº 3982030.
- 360 - Hameed, K. S., & Haard, N.F. (1985). Isolation and characterization of cathepsin-c
361 from atlantic short finned squid illex-illecebrosus. *Comparative biochemistry and*
362 *physiology B-Biochemistry & molecular biology*, 82 (2), 241-246.
- 363 - Hicks, K. B., Haines, R.M., Tong, C.B.S., Sapers, G. M., El Atawy, Y., Irwin, P.L., &
364 Seib, P.A. (1996). Inhibition of enzymatic browning in fresh fruit and vegetable juices
365 by soluble and insoluble forms of beta-cyclodextrin alone or in combination with
366 phosphates. *Journal of Agricultural and Food Chemistry*, 44 (9): 2591-2594.
- 367 - Iyengar R., & McEvily A. J. (1992). Anti-browning agents: alternatives to the use of
368 sulfites in foods. *Trends in Food Science & Technology*, 31, 60-64.
- 369 - Iyidogan N. F., & Bayindirli A. 2004. Effect of L-cysteine, kojic acid and 4-
370 hexylresorcinol combination on inhibition of enzymatic browning in Amasya apple
371 juice. *Journal of Food Engineering*, 62(3), 299–304.
- 372 - Jiménez, M., & García-Carmona, F. (1997). 4-Substituted resorcinols (sulfite
373 alternatives) as slow-binding inhibitors of tyrosinase catecholase activity. *Journal of*
374 *Agricultural and Food Chemistry*, 45, 2061-2065.
- 375 - Kahn, V. (1995). Effect of kojic acid on the oxidation of DL-DOPA, norepinephrine,
376 and dopamine by mushroom tyrosinase. *Pigment Cell Research*, 8, 234-240.
- 377 - Kubo, I., & Kinst-Hori, I. (1998). Tyrosinase inhibitors from Cumin. *Journal of*
378 *Agricultural and Food Chemistry*, 46, 5338-5341.

- 379 - Lee-Kim, M.S., Hwang, E.S., & Kim, K.H (1997). Inhibition studies on burdock
380 polyphenyl oxidase (PPO) activity. *Journal of Food Processing and Preservation*, 21
381 (6), 485-494.
- 382 - McEvily, A. J., Iyengar, R., & Otwell, W. S. (1991). Sulfite alternative prevents
383 shrimps melanosis. *Food Technology*, 45(9), 80-86.
- 384 - McEvily, A.J., Iyengar, R., & Otwell, W.S. (1992). Inhibition of enzymatic browning in
385 foods and beverages. *CRC Critical Reviews in Food Science and Nutrition*, 32(3), 253-
386 273.
- 387 - Montero, P., López-Caballero, M. E., & Pérez-Mateos, M. (2001a). The effect of
388 inhibitors and high pressure treatment to prevent melanosis and microbial growth on
389 chilled prawns (*Penaeus japonicus*). *Journal of Food Science*, 66 (8), 1201-1206.
- 390 - Montero, P., Ávalos, A. & Pérez-Mateos, M. (2001b). Characterization of
391 polyphenoloxidase of prawns (*Penaeus japonicus*). Alternatives to inhibition: additives
392 and high-pressure treatment. *Food Chemistry*, 75, 317-324.
- 393 - Montero, P., Martínez-Alvarez, O., & Gómez-Guillén, M.C. (2004). Effectiveness of
394 onboard application of 4-hexylresorcinol in inhibiting melanosis in shrimp
395 (*Parapenaeus longirostris*). *Journal of Food Science*, 68 (8), 643-647.
- 396 - Otwell, W.S., Iyengar, R., & Mc Evily, A.J. (1992). Inhibition of shrimp melanosis by
397 4-hexylresorcinol. *Journal of Aquatic Food Product Technology*, 1(1), 53-65.
- 398 - Otwell, W. S., & Marshall, M. (1986). Screening alternatives to sulfiting agents to
399 control shrimp melanosis. Florida Cooperative Extension Service, Sea Grant Extension
400 Program. Technical Paper No. 26.
- 401 - Söderhäll, K. & Cerenius, L. (1992). Crustacean immunity. *Annual Review of Fish*
402 *Diseases*, 3-23.

- 403 - Stevens, L. (1991). Egg white protein. Mini-review. *Comparative Biochemistry and*
404 *Physiology*, 100B, 1-9.
- 405 - Taylor, S.L., & Bush, R. K. (1987). Sulphites as food Ingredients. *Food Technology in*
406 *Australia*, 39 (11), 47-51.
- 407 - Yamashita, M., & Konagaya, S. (1990). Participation of cathepsin 1 into extensive
408 softening of the muscle of chum salmon caught during spawning migration. *Nippon*
409 *Suisan Gakkaishi*, 56 (8): 1271-1277.
- 410 - Zotos, A., & Taylor, K. D. A. (1997). Studies on the influence of small molecule
411 factor(s) on protease activities in Norway lobster (*Nephrops norvegicus*). *Food*
412 *Chemistry*, 59 (1): 19-25.

413

414 **FIGURE CAPTIONS**

415 - Figure 1: Melanosis during the chilled storage of shrimp treated with 4-HR based
416 formulas including kojic acid (0.1%), cumic acid (0.1%) or phytic acid (0.1%). a) Formulas
417 with 0.1% 4-hexylresorcinol plus organic acids; b) Formulas with 0.25% 4-HR plus organic
418 acids. ACRA: 4-HR+ascorbic acid (0.5%)+acetic acid (0.05 N)+citric acid (0.5%); Koj:
419 kojic acid; Cum; cumic acid; Phy: phytic acid.

420 - Figure 2: Intensity of melanosis during the chilled storage of shrimp treated with 4-
421 hexylresorcinol (0.25 % or 0.1%) combined with citric acid (0.5%), ascorbic acid (0.5%),
422 acetic acid (0.05 N), and sodium metabisulphite (1% or 0.625%). ACRA: 4-
423 HR+ascorbic+acetic+citric acid; Sul: sodium metabisulphite.

424 - Figure 3: Appearance of melanosis during the chilled storage of shrimp treated with 4-
425 hexylresorcinol (0.25 % or 0.1%) combined with citric acid (0.5%), ascorbic acid (0.5%),

426 acetic acid (0.05 N), and magnesium carbonate (7% or 0.5%). ACRA: 4-
427 HR+ascorbic+acetic+citric acid; Mg Car: magnesium carbonate.

428 - Figure 4: Intensity of melanosis during the chilled storage of shrimps treated with 4-
429 hexylresorcinol (0.25 % or 0.1%) combined with organic acids, and the protease inhibitors
430 EDTA (225 ppm), PPI (1%), egg white (1% and 2%), Iodoacetic acid (210 ppm),and PMSF
431 (1 %)). ACRA 0.25%: 4-HR (0.25%)+ascorbic acid (0.5%)+acetic acid (0.05 N)+citric acid
432 (0.5%); EDTA: Ethylenediaminetetraacetic acid; PPI: Sodium pyrophosphate; Ew: Egg
433 white; IAA: Iodoacetic acid; PMSF: Phenylmethylsulphonyl fluoride.

434 - Figure 5: Melanosis during the chilled storage of shrimp treated with 4-hexylresorcinol
435 (0.25 % or 0.1%) combined with organic acids, the protease inhibitors EDTA (225 ppm)
436 and PPI (1%) and potassium sorbate. ACRA 0.25%: 4-HR (0.25%)+ascorbic acid
437 (0.5%)+acetic acid (0.05 N)+citric acid (0.5%). Sor: Potassium sorbate (2%).

438

439 **TABLES**

440 Table 1. Melanosis-inhibiting blends used by immersion after capture. ACRA: 0.5%
 441 ascorbic acid+0.5% citric acid+0.3% acetic acid+0.1% 4-HR (ACRA 0.1%) or 0.25 % 4-
 442 HR (ACRA 0.25%). R: 4-HR; Cit: Citric acid; Koj: Kojic acid; Cum: Cumic acid; Phy:
 443 Phytic acid; Mg Car: Magnesium carbonate; Sul: sodium metabisulphite; EDTA:
 444 Etilenediaminetetraacetic acid; Sor: Potassium sorbate; PPI: Sodium pyrophosphate; Ew:
 445 Egg white; IAA: Iodoacetic acid; PMSF: Phenylmethylsulphonyl fluoride.

446

MELANOSIS-INHIBITING BLEND	Koj (%)	Cum (%)	Phy (%)	Mg Car (%)	Sul (%)	Sor (%)	EDTA (ppm)	PPi (%)	EW (%)	IAA (%)	PMSF (%)
CONTROL											
ACRA 0.1%											
ACRA 0.1%+Koj	0.1										
ACRA 0.1%+Cum		0.1									
ACRA 0.1%+Phy			0.1								
ACRA 0.1%+Cum+Phy		0.1	0.1								
ACRA 0.1%+Mg Car				0.5							
ACRA 0.25%											
ACRA 0.25%+Koj	0.1										
ACRA 0.25%+Cum		0.1									
ACRA 0.25%+Phy			0.1								
ACRA 0.25%+Cum+Phy		0.1	0.1								
ACRA 0.25%+EDTA+PPi							225	1			
ACRA 0.25%+EDTA+PPi+Ew 1%							225	1	1		
ACRA 0.25%+EDTA+PPi+Ew 2%							225	1	2		
ACRA 0.25%+EDTA+PPi+IAA+PMSF							225	1		1	1
ACRA 0.25%+Sor											2
ACRA 0.25%+EDTA+PPi+Sor							2	225	1		
ACRA 0.25%+Mg Car				0.5							
ACRA 0.25%+Mg Car 7%				7							
R 0.1%											
R 0.1%+Sul 1.25%											1.25
R 0.1%+Sul 0.625%											0.625
Sul 1.25											1.25

447 Table 2. Prevalence of yellow-greenish colourations and acceptability (%) of shrimp treated
 448 with 4-HR based formulas including kojic acid (0.1%), cumic acid (0.1%) or phytic acid
 449 (0.1%). a) Formulas with 0.1% 4-hexylresorcinol plus organic acids; b) Formulas with
 450 0.25% 4-HR plus organic acids. ACRA: 4-HR+ascorbic acid (0.5%)+acetic acid (0.05
 451 N)+citric acid (0.5%); Koj: kojic acid; Cum; cumic acid; Phy: phytic acid.
 452 a)

Yellow-greenish shades (%)	DAYS OF STORAGE					
	0	2	4	7	9	11
Control	0	12.50	12.50	33.33		
ACRA 0.1%	0	16.67	42.86	70	75	100
ACRA 0.1%+Koj	0	16.67	100	100	100	100
ACRA 0.1%+Cum	0	41.67	50	80	75	100
ACRA 0.1%+Phy	0	33.33	57.14	80	100	100
ACRA 0.1%+Cum+Phy	0	33.33	57.14	80	100	100
Acceptability (%)	0	2	4	7	9	11
Control	100	27.58	0	0	0	0
ACRA 0.1%	100	100	100	80	0	0
ACRA 0.1%+Koj	100	100	100	0	0	0
ACRA 0.1%+Cum	100	83.3	100	50	8.33	0
ACRA 0.1%+Phy	100	83.3	100	50	0	0
ACRA 0.1%+Cum+Phy	100	100	100	50	0	0

453 b)

Yellow-greenish shades (%)	DAYS OF STORAGE					
	0	2	4	7	9	11
Control	0	12.5	12.5	33.33		
ACRA 0.25%	0	8.33	28.57	80	100	100
ACRA 0.25%+Koj	0	33.33	14.29	80	91.67	100
ACRA 0.25%+Cum	0	33.33	21.43	50	90	81.25
ACRA 0.25%+Phy	0	16.67	14.29	30	80	87.5
ACRA 0.25%+Cum+Phy	0	33.33	21.43	50	80	87.5
Acceptability (%)	0	2	4	7	9	11
Control	100	27.58	0	0	0	0
ACRA 0.25%	100	100	100	90	8.33	0
ACRA 0.25%+Koj	100	100	100	100	8.33	6.25
ACRA 0.25%+Cum	100	100	100	100	16.67	0
ACRA 0.25%+Phy	100	100	100	100	8.33	6.25
ACRA 0.25%+Cum+Phy	100	100	100	100	0	0

454

455 Table 3. Presence of yellow-greenish colouration beneath cuticle and acceptability (%) of
 456 shrimps dipped in melanosis-inhibiting solutions including 4-HR and/or sodium
 457 metabisulphite. R: 4-HR; Sul: sodium metabisulphite.

458

DAYS OF STORAGE

Yellow-greenish shades (%)	0	2	4	7	9
Control	0	12.50	12.50	33.33	
R 0.1%	0	14.29	59.38	98.61	88.89
Sul 1.25%	0	24.40	25	33.33	
R 0.1%-Sul 1.25%	0	9.52	75	77.78	100
R 0.1%-Sul 0.625%	0	15.48	62.50	84.26	100
Acceptability (%)	0	2	4	7	9
Control	100	27.58	0	0	0
R 0.1%	100	80.95	69.64	0	0
Sul 1.25%	100	42.86	2.08	0	0
R 0.1%-Sul 1.25%	100	79.76	29.17	27.78	0
R 0.1%-Sul 0.625%	100	75.60	45.83	16.67	0

459

460 Table 4. Presence of yellow-greenish colouration beneath head cuticle (%) and
 461 acceptability (%) of shrimps treated with different melanosis-inhibiting blends including
 462 magnesium carbonate. ACRA: 4-HR (0.1 or 0.25%)+ascorbic acid (0.5%)+acetic acid (0.05
 463 N)+citric acid (0.5%); Mg Car: Magnesium carbonate.

464

Yellow-greenish shades (%)	DAYS					
	0	2	4	7	9	11
Control	100	27.58	0	0	0	0
ACRA 0.1%	100	100	100	80	0	0
ACRA 0.25%	100	100	100	90	8.33	0
ACRA 0.1%+Mg Car 0.5%	100	100	100	20	0	0
ACRA 0.25%+Mg Car 0.5%	100	100	100	60	0	6.25
ACRA 0.25%+Mg Car 7%	100	100	100	0	0	0
Acceptability (%)	0	2	4	7	9	11
Control	100	27.58	0	0	0	0
ACRA 0.1%	100	100	100	80	0	0
ACRA 0.25%	100	100	100	90	8.33	0
ACRA 0.1%+Mg Car 0.5%	100	100	100	20	0	0
ACRA 0.25%+Mg Car 0.5%	100	100	100	60	0	6.25
ACRA 0.25%+Mg Car 7%	100	100	100	0	0	0

465

466 Table 5: Presence of yellow-greenish shades beneath cuticle (%) and acceptability (%) of
 467 shrimps treated with different melanosis-inhibiting blends. ACRA 0.25%: 4-HR
 468 (0.25%)+ascorbic acid (0.5%)+acetic acid (0.05 N)+citric acid (0.5%); EDTA:
 469 Etilenediaminetetraacetic acid; PPI: Sodium pirophosphate; Ew: Egg white; IAA:
 470 Iodoacetic acid; PMF: Phenylmethilsulphonil fluoride.
 471

Yellow-greenish shades (%)	DAYS					
	0	2	4	7	9	11
ACRA 0.25%	0	8.33	28.6	80	100	100
ACRA 0.25%+EDTA+PPI+Ew 1%	0	58.3	57.1	70		
ACRA 0.25%+EDTA+PPI+Ew 2%	0	41.7	71.4	40	40	
ACRA 0.25%+EDTA+PPI+IAA+PMSF	0	0	56.3	56.3	60	80
Acceptability (%)	0	2	4	7	9	11
ACRA 0.25%	100	100	100	90	8.33	0
ACRA 0.25%+EDTA+PPI+Ew 1%	100	100	100	0	0	0
ACRA 0.25%+EDTA+PPI+Ew 2%	100	100	83.3	0	0	0
ACRA 0.25%+EDTA+PPI+IAA+PMSF	100	100	100	83.3	83.33	67

472

473 Table 6: Development of yellow-greenish colouration in the cephalothorax and
 474 acceptability of shrimps dipped in different melanosis-inhibiting solutions. EDTA:
 475 Etilenediaminetetraacetic acid; PPI: Sodium pyrophosphate; Sor: Potassium sorbate.

476

	DAYS OF STORAGE					
Yellow-greenish shades (%)	0	2	4	7	9	11
ACRA 0.25%+Sor	0	41.7	42.9	75	100	100
ACRA 0.25%+EDTA+PPI+Sor	0	41.7	57.1	80	80	100
ACRA 0.25%	0	8.33	28.6	80	100	100
Acceptability (%)	0	2	4	7	9	11
ACRA 0.25%+Sor	100	100	85.7	100	0	0
ACRA 0.25%+EDTA+PPI+Sor	100	100	85.7	100	16.67	0
ACRA 0.25%	100	100	100	90	8.33	0

477

478 Table 7: Principal Component Analyses (data matrix) on the basis of the increasing
 479 concentrations of 4-HR in the melanosis-inhibiting formulas (a), or else of the presence of
 480 kojic acid (b), cumic acid (c), phytic acid (d), sulphites (e), magnesium carbonate (f), and
 481 protease inhibitors (EDTA, PPI, Ew, IAA, PMSF) (g) in the 4-HR-inhibiting formulas.

482

a)	PC 1 (49% variance)	PC 2 (36 % variance)
4-hexylresorcinol content	0.316	0.864
Storage period	0.971	-0.199
Yellow-greenish colouration	0.966	0.056
Acceptability	-0.593	0.689
Melanosis score	0.349	-0.745

483

b)	PC 1 (62 % variance)	PC 2 (27 % variance)
Kojic acid	0.008	0.985
Storage period	0.960	-0.047
Yellow-greenish colouration	0.908	0.109
Acceptability	-0.833	0.339
Melanosis score	0.960	0.095

484

c)	PC 1 (68 % variance)	PC 2 (22 % variance)
Cumic acid	0.035	0.990
Storage period	0.972	0.043
Yellow-greenish colouration	0.938	0.104
Acceptability	-0.886	0.251
Melanosis score	0.893	0.159

485

d)	PC 1 (68 % variance)	PC 2 (24 % variance)
Phytic acid	0.017	0.987
Storage period	0.967	-0.003
Yellow-greenish colouration	0.952	-0.069
Acceptability	-0.832	0.458
Melanosis score	0.927	0.181

486

487

488

489

e)	PC 1 (67 % variance)	PC 2 (22 % variance)
Sulphites	-0.002	0.968
Storage period	0.978	-0.070
Yellow-greenish colouration	0.886	-0.211
Acceptability	-0.972	-0.089
Melanosis score	0.824	0.322

490

f)	PC 1 (67 % variance)	PC 2 (23 % variance)
Magnesium carbonate	0.004	0.989
Storage period	0.965	-0.012
Yellow-greenish colouration	0.923	0.002
Acceptability	-0.888	0.341
Melanosis score	0.873	0.276

491

g)	PC 1 (58 % variance)	PC 2 (26 % variance)
Protease inhibitors	0.033	0.979
Storage period	0.944	-0.054
Yellow-greenish colouration	0.856	-0.038
Acceptability	-0.725	0.577
Melanosis score	0.867	-0.004

492

493

494

495

496

497