

## FUNCTIONAL BENEFITS OF PSYLLIUM FIBER SUPPLEMENTATION

Julia Wärnberg<sup>1,2</sup>, Ascensión Marcos<sup>3</sup>, Gloria Bueno<sup>4</sup> and Luís A. Moreno<sup>5</sup>

<sup>1</sup>Department of Preventive Nutrition and Public Health, Faculty of Medicine, University of Navarra, Pamplona, Spain; <sup>2</sup>Unit for Preventive Nutrition, Department of Biosciences and Nutrition, Karolinska Institutet, Stockholm; <sup>3</sup>Immunonutrition Group, Department of Metabolism and Nutrition, Instituto del Frio-ICTAN, Consejo Superior de Investigaciones Científicas (CSIC), Madrid, Spain; <sup>4</sup>Grupo GENUD (Growth, Exercise, Nutrition and Development), Departamento de Pediatría, University of Zaragoza, Spain; <sup>5</sup>Grupo GENUD (Growth, Exercise, Nutrition and Development), Escuela Universitaria de Ciencias de la Salud, University of Zaragoza, Spain

[Received December 23, 2008; Accepted April 15, 2009]

**ABSTRACT:** *Commonly called Psyllium (Ispaghula Psyllium, Plantago ispaghula and Plantago ovata), has a long history of use as a dietary fiber supplement, primarily as a gentle bulk-forming laxative to promote the regulation of large bowel function, but it has as well a potential role in the treatment and prevention of bowel diseases such as diverticulosis, irritable bowel syndrome and inflammatory bowel disease, and could even play a protective role in the prevention of colon cancer. During the last decade, dietary supplementation with Psyllium has been shown to lower blood cholesterol level (especially LDL cholesterol) and to maintain blood glucose homeostasis, which together are the most effective preventive measures against diabetes and cardiovascular diseases. Much interest is being shown by the general public, the scientific community, and federal US regulators in the medical application of foods with specific health benefits (functional foods), foods that favorably modify physiologic function. There is also considerable interest in health claims by the industry to produce functional foods and to ensure efficacy of action. The intent of this review is to summarize the functional benefits of Psyllium fiber consumption and to explore the potential application this fiber has for first-line dietary prevention of these diseases and disorders.*

**KEY WORDS:** Cholesterol lowering, Dietary supplementation, Intestinal regulation, Soluble fiber.

---

*Corresponding Author:* Dr. Luís A. Moreno, Grupo GENUD (Growth, Exercise, Nutrition and Development), Escuela Universitaria de Ciencias de la Salud, University of Zaragoza, Spain; E-mail: lmoreno@unizar.es

---

### INTRODUCTION

Commonly called Psyllium (Ispaghula Psyllium, *Plantago ispaghula* and *Plantago ovata*), has a long history of use as a dietary

fiber supplement, primarily as a gentle bulk-forming laxative to promote the regulation of large bowel function. Also, during the last decade, dietary supplementation with psyllium has been shown to lower blood cholesterol levels, and especially LDL cholesterol (Williams *et al.*, 1995; Brown *et al.*, 1999; Anderson *et al.*, 2000a; Anderson *et al.*, 2000b) as well as to maintain blood glucose homeostasis, which together are the most effective preventive measures against diabetes and cardiovascular diseases (CVD). Considering that cardiovascular diseases is one of the major causes of death in most Western countries, and that dietary intervention should be the first-line approach, increasing soluble dietary fiber has been recommended as safe and practical for cardiovascular disease prevention.

Much interest has been shown internationally by the general public, the scientific community, and federal US regulators in the medical application of foods with specific health benefits (functional foods), foods that favorably modify physiologic function. There is also considerable interest in health claims by the industry to produce functional foods and to ensure efficacy of action. The intent of this review is to summarize the documented functional benefits of Psyllium fiber consumption.

### WHAT IS PSYLLIUM ?

According to the European Pharmacopoeia, Psyllium seed (*Psyllii semen*) consists of the ripe, whole, dry seeds of *Plantago afra* L. (*Plantago psyllium* L.) or *Plantago indica* L. (*Plantago arenaria* Waldstein and Kitaibel). Husk and seeds from *Plantago ovata* (*Plantaginis ovatae*) is however commonly also referred to as Psyllium, although the correct definition of *Plantago ovata* Forssk. (*P. ispaghula* Roxb.) is Ispaghula husk and seeds. Ispaghula, or commonly psyllium husk, consists of the epispem and collapsed adjacent layers removed from the seeds of *Plantago ovata* Forssk and are obtained by milling the seed to remove the hulls. We are, in this review of Psyllium, referring to water-soluble fiber derived

## 2 *Psyllium fiber supplementation*

from all above mentioned plants.

The physiologically active component of psyllium husk is shown to be a highly branched, neutral arabinoxylan consisting of a xylose backbone and arabinose- and xylose-containing side chains. In contrast to arabinoxylans in cereal grains that are extensively fermented, psyllium husk possesses a structural feature, as yet unidentified, that hinders its fermentation by typical colonic microflora (Marlett and Fischer, 2003). Psyllium is classified as a mucilaginous fiber due to its powerful ability to form a gel in water and through animal and human feeding experiments, it has been shown that a gel-forming fraction, amounting to some 55–60% of the husk, is responsible for both the laxative and cholesterol-lowering activities (Marlett and Fischer, 2002). Other viscous, non-nutrient polysaccharides, such as  $\beta$ -glucans and pectins, lower blood cholesterol levels by the same mechanism as psyllium, but these substances have negligible effects on bowel function. They are rapidly and completely fermented in the gut, whereas psyllium husk largely survives, increasing stool output and imparting a gel-like consistency to the excreta (Marteau *et al.*, 1994; Marlett and Fischer, 2003; Fischer *et al.*, 2004).

### PSYLLIUM AND GASTROINTESTINAL REGULATION

#### Constipation and diarrhea

Most dietary fiber sources promote laxation by increasing colonic contents, which stimulate propulsion. Unfermented or incompletely fermented fiber and the accompanying moisture it holds contributes to increase stool mass. These fibers also provide substrate for microbial growth and induces therefore a greater bacterial mass (Stephen and Cummings, 1980; Chen *et al.*, 1998), another factor for increased colonic content. Another benefit is that psyllium containing stools contains an unfermented gel which functions as an emollient and lubricant leading to a greater ease of passage of the stools (Marlett *et al.*, 2000). Psyllium fiber is widely used as a fiber supplement for the treatment of constipation and has in clinical trials repeatedly reported significantly increased levels of stool moisture, as well as wet and dry stool weight in healthy subjects (Prynne and Southgate, 1979; Spiller *et al.*, 1979; Stevens *et al.*, 1988; Tomlin and Read, 1988; Gelissen *et al.*, 1994; Marteau *et al.*, 1994; Levitt *et al.*, 1996; Marlett *et al.*, 2000) and in patients with gastrointestinal disease (Eastwood *et al.*, 1978; Kumar *et al.*, 1987; Prior and Whorwell, 1987; Thorburn *et al.*, 1992; Gelissen *et al.*, 1994; Ashraf *et al.*, 1995; Cheskin *et al.*, 1995; McRorie *et al.*, 1998). It has been proposed that each gram of Psyllium fiber increases stool weight an average of 5.9–6.1g (Marteau *et al.*, 1994; Marlett *et al.*, 2000) compared with 4.9–5.4g and 3.4–4.5g for consumed wheat bran fiber or oat bran fiber, respectively (Cummings, 1993; Chen *et al.*, 1998).

On the opposite of the desired effect against constipation, and because of its great ability to retain water, Psyllium has also been shown to slow down the gastric emptying time and colon transit, this being of benefit for individuals with fecal incontinence from liquid stools or diarrhea (Washington *et al.*, 1998; Bliss *et al.*, 2001).

#### Bowel diseases

Psyllium supplementation may be helpful in the treatment of irritable bowel diseases, inflammatory bowel disease and ulcerative colitis. While the beneficial effects of Psyllium in treatment of irritable bowel syndrome is most probably associated with its anticonstipation activity, the beneficial effects on ulcerative colitis and inflammatory bowel disease seem to be due to anticonstipation activity together with increased levels of the short-chain fatty acid butyrate (Bijkerk *et al.*, 2004). Anaerobic fermentation of Psyllium fiber in the intestines results in a considerable production of the short-chain fatty acids acetate, propionate, and butyrate (Mortensen *et al.*, 1992; Marteau *et al.*, 1994; Nordgaard *et al.*, 1996; Pylkas *et al.*, 2005) which have anti-inflammatory and anti-oxidant properties (Tedelind *et al.*, 2007; Hamer *et al.*, 2008) as well as being an important source of energy-yielding substrates to the colonic mucosa (Mortensen and Clausen, 1996).

Psyllium supplementation is widely used also for hemorrhoids and diverticulitis, although evidence has been questioned (Tan and Seow-Choen, 2007). Its consumption has been shown to be beneficial for hemorrhoids with improvement in reduction of bleeding on contact and of congested hemorrhoidal cushions (Webster *et al.*, 1978; Perez-Miranda *et al.*, 1996; Alonso-Coello *et al.*, 2005). The treatment should, however, last more than a month as no amelioration was noted in another trial of 30 days. Less evidence is available for the effectiveness of fiber in the treatment of uncomplicated diverticulitis (Petruzzello *et al.*, 2006), yet widely used among patients to relieve overall symptoms.

#### Other conditions

##### *Colon cancer prevention*

Short chained fatty acids and specifically butyric acid nourishes the colonocytes, which is important in colon cancer prevention (Clausen *et al.*, 1991). *In vitro*, butyrate also exerts potent inhibition of inflammation and carcinogenesis, reinforcing various components of the colonic defence barrier and decreasing oxidative stress (Hamer *et al.*, 2008). As mentioned above, the colonic anaerobic fermentation of Psyllium yields a considerable production of this short-chain fatty acid (Mortensen *et al.*, 1992; Marteau *et al.*, 1994; Nordgaard *et al.*, 1996; Pylkas *et al.*, 2005) and therefore its consumption could have a preventive effect against cancer. Additionally, *in vitro* assays have proposed that an antineoplastic activity against cancer cells is provided by phytosterol beta-sitosterol that has been isolated from Psyllium (Nakamura *et al.*, 2004; Nakamura *et al.*, 2005). Together with the well known protective effect of fiber consumption for colon cancer prevention, strengthen, but doesn't explain, the possible benefits of Psyllium supplementation in colon cancer prevention.

##### *Prebiotic effect*

Prebiotics are food ingredients that stimulate selectively the growth and activity of bifidobacteria and lactobacilli in the gut and thereby benefit health (Cummings and Macfarlane, 2002). The defining property of prebiotics is their effect on the microflora of the large bowel (Kanauchi *et al.*, 2003; Rodriguez-Cabezas *et al.*, 2003). Poor microbiologic evidences are nevertheless available

concerning the ability of Psyllium derivatives to promote the growth of bifidobacteria in human gut and thus act as a prebiotic. A recent study suggests that Psyllium seed husk can be metabolized by bifidobacteria only after partial hydrolysis (Elli *et al.*, 2008). This study in women demonstrates that the bifidogenic potential can only be detected in healthy subjects with low levels of fecal bifidobacteria before Psyllium supplementation.

#### *Anti-inflammatory properties*

Recent evidence suggests that inflammation may be an important mediator in the association between consumption of dietary fiber and CVD. Cross-sectional research has demonstrated an association between dietary fiber and levels of C-reactive protein (CRP), a clinical indicator of inflammation (King *et al.*, 2003; Ma *et al.*, 2006). Less consistent evidence exists for fiber supplementation. In a randomized crossover intervention trial of a naturally high-fiber diet (30-g/d) or a Psyllium-supplemented diet (30 g/d), lean normotensive participants experienced a greater relative reduction in CRP measurements (40% vs. 10%) than the obese hypertensive persons (King *et al.*, 2007). In another trial, by the same authors, again, no significant differences were observed in several inflammatory serum markers including CRP after a Psyllium supplementation for 3 months (King *et al.*, 2008).

### **PSYLLIUM AND CARDIOVASCULAR DISEASE PROTECTION**

The role of dietary fiber in the prevention of cardiovascular disease has been the subject of considerable attention in the last decade. The functional benefits are now very well recognized and several national agencies concerned about cardiovascular health are acknowledging a role for fiber and especially soluble fiber, such as Psyllium, in cholesterol reduction. The FDA was one of the first national agencies to recognize a role for fiber in cardiovascular disease risk reduction (FDA, 1998). Products that contain 0.75 g  $\beta$ -glucan or 1.78 g psyllium/serving are permitted to carry a health claim stating that the product “will reduce the risk of coronary heart disease” (FDA, 1998). The FDA further determined that 4 servings of these foods are likely to provide the effective daily dose. In the European Union, a common regulation on nutrition and health claims made on foods was introduced in 2007 (EC Regulation, 2008). This Regulation provides opportunities for the use of health claims on foods in Europe, including reduction of disease risk claims. Although the Regulation will not be fully implemented until January 2010 some European countries have applied voluntary codes of practice on health claims for foods, awaiting the Regulation (Asp and Bryngelsson, 2008). Generally most national codes already claim that soluble fiber such as  $\beta$ -glucan or Psyllium reduces cholesterol, and prevents cardiovascular diseases.

Although not as documented as for cholesterol reduction, Psyllium supplementation is also widely used for weight control and to maintain glucose homeostasis that together with cholesterol reduction are the most effective preventive measures against cardiovascular diseases.

#### **Lipid lowering**

Already in 1965 it was observed that Psyllium administered as a hydrophilic mucilloid (i.e., the commercial bulk-forming laxative) reduced serum triglycerides by 9% after supplementation with 9.6 g psyllium/day for 5 wk (Garvin *et al.*, 1965). To lower serum cholesterol the first dietary fibers of interest were pectin and guar gum (Jenkins *et al.*, 1975); later attention focused on oat  $\beta$ -glucan and Psyllium (Brown *et al.*, 1999).

In a large number of studies, Psyllium has generally solidly proved to lower blood lipid levels, and specially LDL cholesterol (Anderson *et al.*, 1988; Bell *et al.*, 1990; Neal and Balm, 1990; Pastors *et al.*, 1991; Everson *et al.*, 1992; Sprecher *et al.*, 1993; Gupta *et al.*, 1994; Ganji and Kies, 1996; Olson *et al.*, 1997; Anderson *et al.*, 2000a; Anderson *et al.*, 2000b; Jenkins *et al.*, 2002; Cicero *et al.*, 2007; Sola *et al.*, 2007; Uehleke *et al.*, 2008). Together with LDL cholesterol reductions, results are also observed in triglycerides and apolipoprotein B with no concluding effect on HDL-cholesterol. The postprandial exposure to triglycerides seems to be lower if the meal is supplemented with Psyllium (Khossousi *et al.*, 2008).

Also in children has the effect of Psyllium been studied for the control of hypercholesterolemia (Glassman *et al.*, 1990; Dennison and Levine, 1993; Davidson *et al.*, 1996). The effect of Psyllium to lower LDL-cholesterol serum concentrations in the management of hypercholesterolemic children, ranged from 3 to -23%; the effect in HDL-cholesterol from -4 to 3%; and the effect on triglycerides from 9 to -20% (Moreno *et al.*, 2003).

The mechanism by which Psyllium lowers LDL cholesterol concentrations is not entirely clear but several mechanisms may play a synergetic effect (Turley and Dietschy, 1995). The primary mechanism of action is likely to be related to the ability to stimulate bile acid synthesis and loss (Kritchevsky and Story, 1974; Miettinen and Tarpila, 1989). Summing up are as well an increased fecal fat loss (Ganji and Kies, 1994), and presumably a lower fat intake (Turnbull and Thomas, 1995) with Psyllium supplementation.

#### **Diabetes Control**

The aim of the management of type 2 diabetes patients is the normalisation of blood glucose values and HbA1C (American Diabetes Association, 2005). The advantages of high-fiber diets for diabetic patients include the lowering of blood glucose and cholesterol, as well as weight reduction and maintenance. It is especially soluble fiber that has beneficial effects on carbohydrate metabolism. Several studies have assessed the effect of Psyllium in the control of type 2 diabetes (Jarjis *et al.*, 1984; Pastors *et al.*, 1991; Frati Munari *et al.*, 1998; Rodriguez-Moran *et al.*, 1998; Anderson *et al.*, 1999; Sierra *et al.*, 2001; Sierra *et al.*, 2002; Moreno *et al.*, 2003; Ziai *et al.*, 2005; Cicero *et al.*, 2007; Khossousi *et al.*, 2008). The percentage change in glucose levels after Psyllium supplementation ranged from -12 to -20%.

Psyllium may slow the absorption of carbohydrates in the small intestine, thereby blunting postprandial glucose peaks (Ellis *et al.*, 1991; Wolever *et al.*, 1991; Fairchild *et al.*, 1996).

##### **Hypertension**

Successful control of hypertension (both systolic and diastolic blood pressure) is also important in diabetes control. Psyllium, but not guar gum lowered blood pressure in a 6 months trial of hypertensive and obese subjects (Cicero *et al.*, 2007). Nevertheless, as weight reduction was observed and confounders were not controlled for, more studies need to confirm these results. Other studies have not shown any changes in blood pressure (Bell *et al.*, 1990).

##### **Weight control**

Psyllium may have a satiating effect on appetite, which may help people to reduce energy intake by earlier termination of meals (Delargy *et al.*, 1995; Turnbull and Thomas, 1995; Cicero *et al.*, 2007). Psyllium supplementation (20g pre-meal and another 20g postmeal) significantly increased feelings of fullness after meals, and significantly reduced *ad libitum* fat intake (Turnbull and Thomas, 1995).

##### **PSYLLIUM AS A FUNCTIONAL FOOD**

There are various commercial uses of Psyllium in food, pharmaceutical and other industries. Although obtaining dietary fiber from whole foods is preferable because it is accompanied by additional nutrients and phytochemicals, a fiber supplement should be recommended to close the fiber gap.

It has been a continuous effort to improve the physicochemical, functional, sensory, and biological properties of psyllium for promoting its food utilization and enhancing its safety. It is a great challenge to disperse psyllium in water or aqueous solutions even with vigorous agitation because of its extremely strong water-absorbing capacity. The readers are referred to an excellent review focused on the approaches to improve the functionality (Yu *et al.*, 2009). Psyllium is most frequently added to breakfast cereals, meal replacements, bread, biscuits and other bakery products to improve the fiber content of the food but is also added to juices, shakes, yogurt, soups and even ice creams. It can also be used as a thickener in drinks or frozen deserts. Although Psyllium fiber has positive physiologic benefits, its high viscosity could make it difficult to incorporate into food products and a task is to get it acceptable to consumers. Their clinical relevance has been questioned because of the extreme unpalatability of products tested with high content of viscose fiber (Ellis *et al.*, 1991). To incorporate lower amounts of fiber into different foods during the day is therefore important to attain the recommended amount for cholesterol lowering (FDA, 1998).

##### **POSSIBLE ADVERSE EFFECT OF PSYLLIUM SUPPLEMENTATION**

Psyllium may alter nutrient and drug absorption, decrease caloric availability by reducing food intake or suppressing appetite, increase bloating and flatulence, cause abdominal pain, and elicit anaphylactic symptoms. Many studies detected the possible adverse effects of Psyllium intake, although others demonstrated that Psyllium is generally safe for human consumption.

Psyllium is known to be the cause of occupational allergy (rhinitis, asthma) in health care and pharmaceutical workers. With the increased prevalence of consumption due to its cholesterol lowering effects has led to its recognition as an emerging food allergen (Lantner *et al.*, 1990). Clinical manifestations of allergy range from upper respiratory tract symptoms on inhalation to anaphylaxis on ingestion (Khalili *et al.*, 2003)

Fermentation of dietary fiber by anaerobic bacteria in the large intestine produces gas, which may be related to complaints of distention or flatulence as well as bloating and diarrhoea. When dietary fiber is increased, fluid intake should also be increased, and fiber intake should gradually be increased to allow the gastrointestinal tract time to adapt (Uehleke *et al.*, 2008). Normal laxation may be achieved with smaller amounts of dietary fiber, and the smallest dose that results in normal laxation should be accepted. Cholesterol reduction however, needs higher doses for optimal effects and this could compromise the adherence to treatment.

Potential negative effects of dietary fiber include reduced absorption of calcium, iron, trace metals, and certain vitamins (Heaney and Weaver, 1995; Asvarujanon *et al.*, 2004). It is unlikely that healthy adults who consume fiber in amounts within the recommended ranges will have problems with nutrient absorption; however, dietary fiber recommendations of 25 g/day may not be appropriate for children and the elderly (Slavin and Greenberg, 2003).

Because of the wide individual variability of responses to dietary fiber and the potential problems with large doses, the smallest dose of dietary fiber that produces the desired result should always be used.

##### **CONCLUSIONS**

We consider that strong scientific evidence exists that regular consumption of Psyllium provides modest reductions in blood levels of LDL cholesterol as well as total cholesterol. This reduction does however not appear to have effects on HDL cholesterol while more research is needed to clarify if Psyllium consumption reduces ApoA or triglyceride levels. There is good scientific evidence for the use of Psyllium supplementation for constipation as most studies observe an increased stool weight, an increase in bowel movements per day, and a decrease in total gut transit time. There is also good scientific evidence that Psyllium has an effective stool bulking effect, against diarrhea. Unclear scientific evidence exists to relieve or prevent the symptoms of hemorrhoids or inflammatory bowel disease or irritable bowel syndrome. Although Psyllium supplementation taken with meals could improve blood glucose homeostasis, better evidence is necessary to conclude its benefits for diabetes control. Obesity related disorders such as dyslipidemia and hyperglycemia could improve by Psyllium supplementation, but further studies are needed to elucidate mechanisms involved and how its consumption affects body weight. Finally, although promising studies attempt to show anti-carcinogenic effect of Psyllium, and specifically in colon cancer, more studies are needed to determine if Psyllium can help prevent cancer.

**CONFLICT-OF-INTEREST STATEMENT**

None of the authors have any conflict of interests to declare.

**REFERENCES**

5th edition of the European Pharmacopoeia. The Council of Europe (EDQM). Online version accessed from <http://online.pheur.org>.

Alonso-Coello P, Guyatt G, Heels-Ansdell D, Johanson JF, Lopez-Yarto M, Mills E and Zhou Q (2005) Laxatives for the treatment of hemorrhoids. *Cochrane database of systematic reviews*, CD004649.

American Diabetes Association (2005) Clinical Practice Recommendations 2005. *Diabetes Care* **28**, S1–S79.

Anderson JW, Allgood LD, Lawrence A, Altringer LA, Jerdack GR, Hengehold DA and Morel JG (2000a) Cholesterol-lowering effects of psyllium intake adjunctive to diet therapy in men and women with hypercholesterolemia: meta-analysis of 8 controlled trials. *American Journal of Clinical Nutrition* **71**, 472–479.

Anderson JW, Allgood LD, Turner J, Oeltgen PR and Daggy BP (1999) Effects of psyllium on glucose and serum lipid responses in men with type 2 diabetes and hypercholesterolemia. *American Journal of Clinical Nutrition* **70**, 466–473.

Anderson JW, Davidson MH, Blonde L, Brown WV, Howard WJ, Ginsberg H, Allgood LD and Weingand KW (2000b) Long-term cholesterol-lowering effects of psyllium as an adjunct to diet therapy in the treatment of hypercholesterolemia. *American Journal of Clinical Nutrition* **71**, 1433–1438.

Anderson JW, Zettwoch N, Feldman T, Tietyen-Clark J, Oeltgen P and Bishop CW (1988) Cholesterol-lowering effects of psyllium hydrophilic mucilloid for hypercholesterolemic men. *Archives of internal medicine* **148**, 292–296.

Ashraf W, Park F, Lof J and Quigley EM (1995) Effects of psyllium therapy on stool characteristics, colon transit and anorectal function in chronic idiopathic constipation. *Alimentary pharmacology and therapeutics* **9**, 639–647.

Asp N-G and Bryngelsson S (2008) Health Claims in Europe: New Legislation and PASSCLAIM for Substantiation. *Journal of Nutrition* **138**, 1210S–1215.

Asvarujanon P, Ishizuka S and Hara H (2004) Inhibitory effects of psyllium on rat mineral absorption were abolished by reduction of viscosity with partial hydrolysis. *Bioscience, biotechnology, and biochemistry* **68**, 1737–1742.

Bell LP, Hectorn KJ, Reynolds H and Hunninghake DB (1990) Cholesterol-lowering effects of soluble-fiber cereals as part of a prudent diet for patients with mild to moderate hypercholesterolemia.

*American Journal of Clinical Nutrition* **52**, 1020–1026.

Bijkerk CJ, Muris JW, Knottnerus JA, Hoes AW and de Wit NJ (2004) Systematic review: the role of different types of fibre in the treatment of irritable bowel syndrome. *Alimentary pharmacology and therapeutics* **19**, 245–251.

Bliss DZ, Jung HJ, Savik K, Lowry A, LeMoine M, Jensen L, Werner C and Schaffer K (2001) Supplementation with dietary fiber improves fecal incontinence. *Nursing research* **50**, 203–213.

Broader JH, Gunn IF and Alexander-Williams J (1974) Evaluation of a bulk-forming evacuant in the management of haemorrhoids. *The British journal of surgery* **61**, 142–144.

Brown L, Rosner B, Willett WW and Sacks FM (1999) Cholesterol-lowering effects of dietary fiber: a meta-analysis. *American Journal of Clinical Nutrition* **69**, 30–42.

Chen H-L, Haack VS, Janecky CW, Vollendorf NW and Marlett JA (1998) Mechanisms by which wheat bran and oat bran increase stool weight in humans. *American Journal of Clinical Nutrition*, 711–719.

Cheskin LJ, Kamal N, Crowell MD, Schuster MM and Whitehead WE (1995) Mechanisms of constipation in older persons and effects of fiber compared with placebo. *Journal of the American Geriatrics Society* **43**, 666–669.

Cicero AF, Derosa G, Manca M, Bove M, Borghi C and Gaddi AV (2007) Different effect of psyllium and guar dietary supplementation on blood pressure control in hypertensive overweight patients: a six-month, randomized clinical trial. *Clinical and experimental hypertension* **29**, 383–394.

Clausen MR, Bonnen H and Mortensen PB (1991) Colonic fermentation of dietary fibre to short chain fatty acids in patients with adenomatous polyps and colonic cancer. *Gut* **32**, 923–928.

Cummings JH (1993) The effect of dietary fibre on faecal weight and composition. In *Dietary fibre in human nutrition.*, pp. 263–350 [S GA, editor]. Boca Raton, FL: CRC Press.

Cummings JH and Macfarlane GT (2002) Gastrointestinal effects of prebiotics. *British Journal of Nutrition* **87 Suppl 2**, S145–151.

Davidson MH, Dugan LD, Burns JH, Sugimoto D, Story K and Drennan K (1996) A psyllium-enriched cereal for the treatment of hypercholesterolemia in children: a controlled, double-blind, crossover study. *American Journal of Clinical Nutrition* **63**, 96–102.

Delargy HJ, Burley VJ, O’Sullivan KR, Fletcher RJ and Blundell JE (1995) Effects of different soluble: insoluble fibre ratios at breakfast on 24-h pattern of dietary intake and satiety. *European Journal of Clinical Nutrition* **49**, 754–766.

## 6 *Psyllium fiber supplementation*

Dennison BA and Levine DM (1993) Randomized, double-blind, placebo-controlled, two-period crossover clinical trial of psyllium fiber in children with hypercholesterolemia. *Journal of Pediatrics* **123**, 24-29.

Eastwood MA, Smith AN, Brydon WG and Pritchard J (1978) Comparison of bran, ispaghula, and lactulose on colon function in diverticular disease. *Gut* **19**, 1144-1147.

EC Regulation (2008) Regulations (EC) No 109/2008 on nutrition and health claims made on foods. Official Journal of the European Union 13.2.2008, pp. 14-15. [THE EUROPEAN PARLIAMENT AND OF THE COUNCIL, editor.

Elli M, Cattivelli D, Soldi S, Bonatti M and Morelli L (2008) Evaluation of prebiotic potential of refined psyllium (*Plantago ovata*) fiber in healthy women. *Journal of clinical gastroenterology* **42 Suppl 3 Pt 2**, S174-176.

Ellis PR, Dawoud FM and Morris ER (1991) Blood glucose, plasma insulin and sensory responses to guar-containing wheat breads: effects of molecular weight and particle size of guar gum. *British Journal of Nutrition* **66**, 363-379.

Everson GT, Daggy BP, McKinley C and Story JA (1992) Effects of psyllium hydrophilic mucilloid on LDL-cholesterol and bile acid synthesis in hypercholesterolemic men. *Journal of lipid research* **33**, 1183-1192.

Fairchild RM, Ellis PR, Byrne AJ, Luzio SD and Mir MA (1996) A new breakfast cereal containing guar gum reduces postprandial plasma glucose and insulin concentrations in normal-weight human subjects. *British Journal of Nutrition* **76**, 63-73.

FDA (1998) Food labeling: health claims; soluble fiber from certain foods and coronary heart disease. In *Fed Regist* [US Food and Drug Administration, editor.

Fischer MH, Yu N, Gray GR, Ralph J, Anderson L and Marlett JA (2004) The gel-forming polysaccharide of psyllium husk (*Plantago ovata* Forsk). *Carbohydrate research* **339**, 2009-2017.

Fрати Munari AC, Benitez Pinto W, Raul Ariza Andraca C and Casarrubias M (1998) Lowering glycemic index of food by acarbose and *Plantago psyllium* mucilage. *Archives of medical research* **29**, 137-141.

Ganji V and Kies CV (1994) Psyllium husk fibre supplementation to soybean and coconut oil diets of humans: effect on fat digestibility and faecal fatty acid excretion. *European Journal of Clinical Nutrition* **48**, 595-597.

Ganji V and Kies CV (1996) Psyllium husk fiber supplementation to the diets rich in soybean or coconut oil: hypocholesterolemic effect in healthy humans. *International journal of food sciences and nutrition* **47**, 103-110.

Garvin JE, Forman DT, Eiseman WR and Phillips CR (1965) Lowering of human serum cholesterol by an oral hydrophilic colloid. *Proceedings of the Society for Experimental Biology and Medicine. Society for Experimental Biology and Medicine (New York, N.Y.)* **120**, 744-746.

Gelissen IC, Brodie B and Eastwood MA (1994) Effect of *Plantago ovata* (psyllium) husk and seeds on sterol metabolism: studies in normal and ileostomy subjects. *American Journal of Clinical Nutrition* **59**, 395-400.

Glassman M, Spark A, Berezin S, Schwarz S, Medow M and Newman LJ (1990) Treatment of type IIa hyperlipidemia in childhood by a simplified American Heart Association diet and fiber supplementation. *American journal of diseases of children (1960)* **144**, 973-976.

Gupta RR, Agrawal CG, Singh GP and Ghatak A (1994) Lipid-lowering efficacy of psyllium hydrophilic mucilloid in non insulin dependent diabetes mellitus with hyperlipidaemia. *The Indian journal of medical research* **100**, 237-241.

Hamer HM, Jonkers D, Venema K, Vanhoutvin S, Troost FJ and Brummer RJ (2008) Review article: the role of butyrate on colonic function. *Alimentary pharmacology and therapeutics* **27**, 104-119.

Heaney RP and Weaver CM (1995) Effect of psyllium on absorption of co-ingested calcium. *Journal of the American Geriatrics Society* **43**, 261-263.

Jarjis HA, Blackburn NA, Redfern JS and Read NW (1984) The effect of ispaghula (Fybogel and Metamucil) and guar gum on glucose tolerance in man. *British Journal of Nutrition* **51**, 371-378.

Jenkins DJ, Newton C, Leeds AR and Cummings JH (1975) Effect of pectin, guar gum, and wheat fibre on serum-cholesterol. *Lancet* **1**, 1116-1117.

Jenkins DJA, Kendall CWC, Vuksan V, Vidgen E, Parker T, Faulkner D, Mehling CC, Garsetti M, Testolin G, Cunnane SC, Ryan MA and Corey PN (2002) Soluble fiber intake at a dose approved by the US Food and Drug Administration for a claim of health benefits: serum lipid risk factors for cardiovascular disease assessed in a randomized controlled crossover trial. *American Journal of Clinical Nutrition* **75**, 834-839.

Kanauchi O, Mitsuyama K, Araki Y and Andoh A (2003) Modification of intestinal flora in the treatment of inflammatory bowel disease. *Current pharmaceutical design* **9**, 333-346.

Khalili B, Bardana EJ, Jr. and Yunginger JW (2003) Psyllium-associated anaphylaxis and death: a case report and review of the literature. *Annals of allergy, asthma and immunology : official publication of the American College of Allergy, Asthma, and Immunology* **91**, 579-584.

- Khossousi A, Binns CW, Dhaliwal SS and Pal S (2008) The acute effects of psyllium on postprandial lipaemia and thermogenesis in overweight and obese men. *British Journal of Nutrition* **99**, 1068-1075.
- King DE, Egan BM and Geesey ME (2003) Relation of dietary fat and fiber to elevation of C-reactive protein. *The American journal of cardiology* **92**, 1335-1339.
- King DE, Egan BM, Woolson RF, Mainous AG, 3rd, Al-Solaiman Y and Jesri A (2007) Effect of a high-fiber diet vs. a fiber-supplemented diet on C-reactive protein level. *Arch Intern Med* **167**, 502-506.
- King DE, Mainous AG, 3rd, Egan BM, Woolson RF and Geesey ME (2008) Effect of psyllium fiber supplementation on C-reactive protein: the trial to reduce inflammatory markers (TRIM). *Ann Fam Med* **6**, 100-106.
- Kritchevsky D and Story JA (1974) Binding of bile salts in vitro by nonnutritive fiber. *Journal of Nutrition* **104**, 458-462.
- Kumar A, Kumar N, Vij JC, Sarin SK and Anand BS (1987) Optimum dosage of ispaghula husk in patients with irritable bowel syndrome: correlation of symptom relief with whole gut transit time and stool weight. *Gut*, 28150-28155.
- Lantner RR, Espiritu BR, Zumerchik P and Tobin MC (1990) Anaphylaxis following ingestion of a psyllium-containing cereal. *JAMA : the journal of the American Medical Association* **264**, 2534-2536.
- Levitt MD, Furne J and Olsson S (1996) The relation of passage of gas an abdominal bloating to colonic gas production. *Annals of internal medicin* **124**, 422-424.
- Ma Y, Griffith JA, Chasan-Taber L, Olendzki BC, Jackson E, Stanek EJ, III, Li W, Pagoto SL, Hafner AR and Ockene IS (2006) Association between dietary fiber and serum C-reactive protein. *American Journal of Clinical Nutrition* **83**, 760-766.
- Marlett JA and Fischer MH (2002) A poorly fermented gel from psyllium seed husk increases excreta moisture and bile acid excretion in rats. *Journal of Nutrition* **132**, 2638-2643.
- Marlett JA and Fischer MH (2003) The active fraction of psyllium seed husk. *The Proceedings of the Nutrition Society* **62**, 207-209.
- Marlett JA, Kajs TM and Fischer MH (2000) An unfermented gel component of psyllium seed husk promotes laxation as a lubricant in humans. *American Journal of Clinical Nutrition* **72**, 784-789.
- Marteau P, Flourie B, Cherbut C, Correze JL, Pellier P, Seylaz J and Rambaud JC (1994) Digestibility and bulking effect of ispaghula husks in healthy humans. *Gut* **35**, 1747-1752.
- McRorie JW, Daggy BP, Morel JG, Diersing PS, Miner PB and Robinson M (1998) Psyllium is superior to docusate sodium for treatment of chronic constipation. *Alimentary pharmacology and therapeutics* **12**, 491-497.
- Miettinen TA and Tarpila S (1989) Serum lipids and cholesterol metabolism during guar gum, plantago ovata and high fibre treatments. *Clinica chimica acta; international journal of clinical chemistry* **183**, 253-262.
- Moreno LA, Tresaco B, Bueno G, Fleta J, Rodriguez G, Garagorri JM and Bueno M (2003) Psyllium fibre and the metabolic control of obese children and adolescents. *Journal of physiology and biochemistry* **59**, 235-242.
- Mortensen PB and Clausen MR (1996) Short-chain fatty acids in the human colon: relation to gastrointestinal health and disease. *Scandinavian journal of gastroenterology. Supplement* **216**, 132-148.
- Mortensen PB, Clausen MR, Bonnen H, Hove H and Holtug K (1992) Colonic fermentation of ispaghula, wheat bran, glucose, and albumin to short-chain fatty acids and ammonia evaluated in vitro in 50 subjects. *JPEN. Journal of parenteral and enteral nutrition* **16**, 433-439.
- Nakamura Y, Trosko JE, Chang CC and Upham BL (2004) Psyllium extracts decreased neoplastic phenotypes induced by the Ha-Ras oncogene transfected into a rat liver oval cell line. *Cancer Letters* **203**, 13-24.
- Nakamura Y, Yoshikawa N, Hiroki I, Sato K, Ohtsuki K, Chang CC, Upham BL and Trosko JE (2005) Beta-sitosterol from psyllium seed husk (*Plantago ovata* Forsk) restores gap junctional intercellular communication in Ha-ras transfected rat liver cells. *Nutrition and cancer* **51**, 218-225.
- Neal GW and Balm TK (1990) Synergistic effects of psyllium in the dietary treatment of hypercholesterolemia. *The Southern medical journal* **83**, 1131-1137.
- Nordgaard I, Hove H, Clausen MR and Mortensen PB (1996) Colonic production of butyrate in patients with previous colonic cancer during long-term treatment with dietary fibre (*Plantago ovata* seeds). *Scandinavian journal of gastroenterology* **31**, 1011-1020.
- Olson BH, Anderson SM, Becker MP, Anderson JW, Hunninghake DB, Jenkins DJ, LaRosa JC, Rippe JM, Roberts DC, Stoy DB, Summerbell CD, Truswell AS, Wolever TM, Morris DH and Fulgoni VL, 3rd (1997) Psyllium-enriched cereals lower blood total cholesterol and LDL cholesterol, but not HDL cholesterol, in hypercholesterolemic adults: results of a meta-analysis. *Journal of Nutrition* **127**, 1973-1980.
- Pastors JG, Blaisdell PW, Balm TK, Asplin CM and Pohl SL (1991) Psyllium fiber reduces rise in postprandial glucose and insulin

## 8 *Psyllium fiber supplementation*

concentrations in patients with non-insulin-dependent diabetes. *The American journal of clinical nutrition* **53**, 1431-1435.

Perez-Miranda M, Gomez-Cedenilla A, Leon-Colombo T, Pajares J and Mate-Jimenez J (1996) Effect of fiber supplements on internal bleeding hemorrhoids. *Hepatogastroenterology* **43**, 1504-1507.

Petruzzello L, Iacopini F, Bulajic M, Shah S and Costamagna G (2006) Review article: uncomplicated diverticular disease of the colon. *Alimentary pharmacology and therapeutics* **23**, 1379-1391.

Prior A and Whorwell PJ (1987) Double blind study of ispaghula in irritable bowel syndrome. *Gut* **28**, 1510-1513.

Prynne CJ and Southgate DAT (1979) The effects of a supplement of dietary fibre on faecal excretion by human subjects *The British journal of nutrition* **41**, 495-503.

Pylkas AM, Juneja LR and Slavin JL (2005) Comparison of different fibers for in vitro production of short chain fatty acids by intestinal microflora. *Journal of medicinal food* **8**, 113-116.

Rodriguez-Cabezas ME, Galvez J, Camuesco D, Lorente MD, Concha A, Martinez-Augustin O, Redondo L and Zarzuelo A (2003) Intestinal anti-inflammatory activity of dietary fiber (*Plantago ovata* seeds) in HLA-B27 transgenic rats. *Clinical nutrition (Edinburgh, Scotland)* **22**, 463-471.

Rodriguez-Moran M, Guerrero-Romero F and Lazcano-Burciaga G (1998) Lipid- and glucose-lowering efficacy of *Plantago Psyllium* in type II diabetes. *Journal of diabetes and its complications* **12**, 273-278.

Sierra M, Garcia JJ, Fernandez N, Diez MJ and Calle AP (2002) Therapeutic effects of psyllium in type 2 diabetic patients. *European Journal of Clinical Nutrition* **56**, 830-842.

Sierra M, Garcia JJ, Fernandez N, Diez MJ, Calle AP and Sahagun AM (2001) Effects of ispaghula husk and guar gum on postprandial glucose and insulin concentrations in healthy subjects. *European Journal of Clinical Nutrition* **55**, 235-243.

Slavin JL and Greenberg NA (2003) Partially hydrolyzed guar gum: clinical nutrition uses. *Nutrition* **19**, 549-552.

Sola R, Godas G, Ribalta J, Vallve JC, Girona J, Anguera A, Ostos M, Recalde D, Salazar J, Caslake M, Martin-Lujan F, Salas-Salvado J and Masana L (2007) Effects of soluble fiber (*Plantago ovata* husk) on plasma lipids, lipoproteins, and apolipoproteins in men with ischemic heart disease. *The American journal of clinical nutrition* **85**, 1157-1163.

Spiller GA, Shipley EA, Chernoff MC and Cooper WC (1979) Bulk laxative efficacy of a psyllium seed hydrocolloid and of a mixture of cellulose and pectin. *Journal of clinical pharmacology* **19**, 313-320.

Sprecher DL, Harris BV, Goldberg AC, Anderson EC, Bayuk LM, Russell BS, Crone DS, Quinn C, Bateman J, Kuzmak BR and Allgood LD (1993) Efficacy of psyllium in reducing serum cholesterol levels in hypercholesterolemic patients on high- or low-fat diets. *Annals of internal medicine* **119**, 545-554.

Stephen AM and Cummings JH (1980) Mechanism of action of dietary fibre in the human colon. *Nature* **284**, 283-284.

Stevens J, VanSoest PJ, Robertson JB and Levitsky DA (1988) Comparison of the effects of psyllium and wheat bran on gastrointestinal transit time and stool characteristics. *Journal of the American Dietetic Association* **88**, 323-326.

Tan KY and Seow-Choen F (2007) Fiber and colorectal diseases: separating fact from fiction. *World journal of gastroenterology: WJG* **13**, 4161-4167.

Tedelind S, Westberg F, Kjerrulf M and Vidal A (2007) Anti-inflammatory properties of the short-chain fatty acids acetate and propionate: a study with relevance to inflammatory bowel disease. *World journal of gastroenterology: WJG* **13**, 2826-2832.

Thorburn HA, Carter KB, Boldberg JA and Finlay IG (1992) Does ispaghula husk stimulate the entire colon in diverticular disease? *Gut* **33**, 352-356.

Tomlin J and Read NW (1988) The relation between bacterial degradation of viscous polysaccharides and stool output in human beings. *The British journal of nutrition* **60**, 467-475.

Turley SD and Dietschy JM (1995) Mechanisms of LDL-cholesterol lowering action of psyllium hydrophilic mucilloid in the hamster. *Biochimica et biophysica acta* **1255**, 177-184.

Turnbull WH and Thomas HG (1995) The effect of a *Plantago ovata* seed containing preparation on appetite variables, nutrient and energy intake. *International journal of obesity and related metabolic disorders: journal of the International Association for the Study of Obesity* **19**, 338-342.

Uehleke B, Ortiz M and Stange R (2008) Cholesterol reduction using psyllium husks - do gastrointestinal adverse effects limit compliance? Results of a specific observational study. *Phytomedicine* **15**, 153-159.

Washington N, Harris M, Mussellwhite A and Spiller RC (1998) Moderation of lactulose-induced diarrhoea by psyllium: effects on motility and fermentation. *The American journal of clinical nutrition* **67**, 317-321.

Webster DJ, Gough DC and Craven JL (1978) The use of bulk evacuant in patients with haemorrhoids. *The British journal of surgery* **65**, 291-292.



Williams CL, Bollella M, Spark A and Puder D (1995) Soluble fiber enhances the hypocholesterolemic effect of the step I diet in childhood. *Journal of the American College of Nutrition* **14**, 251-257.

Wolever TM, Vuksan V, Eshuis H, Spadafora P, Peterson RD, Chao ES, Storey ML and Jenkins DJ (1991) Effect of method of administration of psyllium on glycemic response and carbohydrate digestibility. *Journal of the American College of Nutrition* **10**, 364-371.

Yu LL, Lutterodt H and Cheng Z (2009) Beneficial health properties of psyllium and approaches to improve its functionalities. *Advances in food and nutrition research* **55**, 193-220.

Ziai SA, Larijani B, Akhoondzadeh S, Fakhrzadeh H, Dastpak A, Bandarian F, Rezai A, Badi HN and Emami T (2005) Psyllium decreased serum glucose and glycosylated hemoglobin significantly in diabetic outpatients. *Journal of ethnopharmacology* **102**, 202-207.

Karp, C.L., Flick, L.M., Park, K.W., Softic, S., Greer, T.M., Keledjian, R., Yang, R., Uddin, J., Guggino, W.B., Atabani, S.F., Belkaid, Y., Xu, Y., Whitsett, J.A., Accurso, F.J., Wills-Karp, M. and Petasis, N.A. (2004) Defective lipoxin-mediated anti-inflammatory activity in the cystic fibrosis airway. *Nature Immunology* **5**, 388-392.

Kent, G., Oliver, M., Foskett, J.K., Frndova, H., Durie, P.R., Forstner, J., Forstner, G.G., Riordan, J.R., Percy, D. and Buchwald, M. (1996) Phenotypic abnormalities in long term surviving cystic fibrosis mice. *Pediatric Research* **40**, 233-241.

