

Neutralising antibodies for West Nile virus in horses from Brazilian Pantanal

Alex Pauvolid-Corrêa^{1/+}, Maria Alejandra Morales², Silvana Levis², Luis Tadeu Moraes Figueiredo⁴, Dinair Couto-Lima¹, Zilca Campos³, Marcia Furlan Nogueira³, Edson Elias da Silva⁵, Rita Maria Ribeiro Nogueira¹, Hermann Gonçalves Schatzmayr^{1†}

¹Laboratório de Flavivírus ⁵Laboratório de Enterovírus, Instituto Oswaldo Cruz-Fiocruz, Ministério da Saúde, Avenida Brasil 4365, 21045-900 Rio de Janeiro, RJ, Brasil ²Laboratório de Arbovírus, Instituto Nacional de Enfermidades Virales Humanas Dr Julio I Maiztegui, Pergamino, BsAs, Argentina ³Embrapa Pantanal, Ministério da Agricultura Pecuária e Abastecimento, Corumbá, MS, Brasil ⁴Centro de Pesquisa em Virologia, Faculdade de Medicina de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, SP, Brasil

Despite evidence of West Nile virus (WNV) activity in Colombia, Venezuela and Argentina, this virus has not been reported in most South American countries. In February 2009, we commenced an investigation for WNV in mosquitoes, horses and caimans from the Pantanal, Central-West Brazil. The sera of 168 horses and 30 caimans were initially tested using a flaviviruses-specific epitope-blocking enzyme-linked immunosorbent assay (blocking ELISA) for the detection of flavivirus-reactive antibodies. The seropositive samples were further tested using a plaque-reduction neutralisation test (PRNT₅₀) for WNV and its most closely-related flaviviruses that circulate in Brazil to confirm the detection of specific virus-neutralising antibodies. Of the 93 (55.4%) blocking ELISA-seropositive horse serum samples, five (3%) were seropositive for WNV, nine (5.4%) were seropositive for St. Louis encephalitis virus, 18 (10.7%) were seropositive for Ilheus virus, three (1.8%) were seropositive for Cacipacore virus and none were seropositive for Rocio virus using PRNT₅₀, with a criteria of \geq four-fold antibody titre difference. All caimans were negative for flaviviruses-specific antibodies using the blocking ELISA. No virus genome was detected from caiman blood or mosquito samples. The present study is the first report of confirmed serological evidence of WNV activity in Brazil.

Key words: West Nile virus - Pantanal - neutralisation - equines - caimans - Brazil

In 1937, an unknown neurotropic virus called West Nile virus (WNV) was isolated from the blood of a low-grade febrile woman from northeastern Uganda (Smithburn et al. 1940). Afterwards, WNV was detected in Africa, the Middle East, Asia and Europe and was associated mainly with sporadic or size-limited outbreaks of WNV fever, which was usually a self-limited and uncomplicated febrile syndrome (Couissinier-Paris 2006). However, in the mid-1990s, a major WNV fever epidemic in Europe with a high rate of neurological infections occurred in the lower Danube valley and Bucharest areas in Romania (Tsai et al. 1998). Concomitantly, human and animal outbreaks of WNV fever with neurological disorders were reported in the old world, which warned of the change of WNV fever impact on animal and public health (Tber Abdelhaq 1996, Platonov et al. 2001). In 1999, WNV definitively reemerged as a worldwide major

public health and veterinarian concern when it was detected in the Western Hemisphere for the first time as the cause of human and equine encephalitis outbreaks concomitant with extensive mortality in birds in the north-eastern United States of America (USA) (CDC 1999). The virus rapidly spread through North America and towards Latin America (Komar & Clark 2006). There have been recent reports of serological evidence of WNV activity in Colombia (Mattar et al. 2005), Venezuela (Bosch et al. 2007) and especially Argentina, where anti-WNV antibodies have been detected in birds (Diaz et al. 2008) and the virus was isolated for the first time in South America from the brains of horses that had died of encephalitis (Morales et al. 2006). However, most South American countries, including the largest, Brazil, remain without reports of WNV infection. Alerted by the periodical outbreaks of WNV encephalitis in the USA (Lindsey et al. 2010) and the serological evidence of WNV activity in South America, we performed this study.

The Brazilian Pantanal was selected as a suitable place for the study of WNV circulation because of the following factors: (i) natural foci of WNV infections have been mostly described in wetland ecosystems, such as deltas and flood plains, as reported in France (Joubert et al. 1970), Venezuela (Bosch et al. 2007), Romania (Tsai et al. 1998) and Italy (Autorino et al. 2002), (ii) the local movement of migratory birds (Nunes & Tomas 2008), (iii) the unsuccessful efforts of the detection of WNV-specific antibodies in the free-ranging birds in Brazilian coastal areas (MS 2003, 2004), (iv) the recent detec-

Financial support: CNPq, FAPERJ, FIOCRUZ

† In memoriam

APC is a recipient of the scholarship of CNPq in the Doctor of Tropical Medicine Program at Fiocruz. This work is part of APC's PhD Thesis at the Instituto Oswaldo Cruz-Fiocruz, Ministério da Saúde, Rio de Janeiro, RJ, Brazil, presented in part at the XXI National Meeting of Virology, Gramado, RS, Brazil, October 17-20th, 2010.

+ Corresponding author: pauvolid@ioc.fiocruz.br

Received 16 December 2010

Accepted 11 May 2011


tion of arbovirus vectors blood-feeding on human and equine hosts in the area (Pauvolid-Corrêa et al. 2010a) and, finally, (v) the serological evidence of arbovirus activity in the area (Iversson et al. 1993), which includes St. Louis encephalitis virus (SLEV) (Pauvolid-Corrêa et al. 2010b), a species that is antigenically related to WNV (Smithburn 1942). In view of these data, the hypothesis that the Brazilian Pantanal is one of the most susceptible areas to WNV circulation in Brazil (Pauvolid-Corrêa & Varella 2008) seemed promising.

The Pantanal is a tropical, seasonal wetland of approximately 140,000 km² that is fed by tributaries of the upper Paraguay River in the centre of South America, which covers mainly Brazilian but also Paraguayan and Bolivian territories and is classified as one of the largest freshwater wetland ecosystems in the world (Alho 2005). The Brazilian Pantanal, which represents 85% of the total Pantanal area, is located within the states of Mato Grosso do Sul and Mato Grosso in Central-West Brazil, which are recognised as South and North Pantanal, respectively (Alho et al. 1988). This region is ecologically classified into sub-regions according to vegetation, flooding and physiography (Silva & Abdon 1988). The Nhecolândia sub-region that is situated in South Pantanal is one of the largest, comprising approximately one fifth of the total area, and is characterised by hundreds of shallow lakes that display varying degrees of salinity and coalescence with the system during floods (Adámoli 1982).

Recent studies have demonstrated that some crocodilian species may be infected by WNV (Steinman et al. 2003, Jacobson et al. 2005) and that *Alligator mississippiensis* may also have a viremic load that affords the infection of blood-feeding Culicidae vectors, which suggests that in areas with high population densities of these alligators, juvenile individuals could play an important role in WNV transmission (Klenk et al. 2004). Taking these data into account, the unsuccessful efforts to detect WNV circulation in the avian hosts in Brazil and the widespread, high prevalence of the Crocodylia species, *Caiman crocodilus yacare*, in the Brazilian Pantanal (Campos et al. 2005), our strategy was to investigate the WNV circulation in the potential vectors, dead-end hosts and natural secondary amplifying hosts through the collection and testing of mosquitoes and serum samples of horses and wild caimans from the Nhecolândia sub-region. No morbidity was observed during the sampling period and only apparently healthy horses and caimans were sampled in the present study.

MATERIALS AND METHODS

Study area - In February 2009, mosquitoes and blood samples from horses and wild caimans were collected in six different cattle ranches of the Nhecolândia sub-region (18°20' 19°40'S and 57°14' 55°00'W) during the rainy season. Considering the free circulation of mosquitoes, horses and wild caimans in large ranches, sample collections for the study were undertaken in a 700-square-kilometre (70,000 ha) area (Figure). The collections for this study were authorised by the Brazilian Institute of Environment and Natural Resources (license IBAMA 18363-1/2009).



Sampling locations of West Nile virus investigation conducted in the Pantanal, Brazil.

Samples collections - Adult mosquitoes were captured at sites that were randomly selected using CDC automatic light traps and manual aspirators while landing to blood-feed on horses and research team members as routinely reported. Living mosquito specimens were transported to a field laboratory where they were immobilised by chilling and the species were identified through direct observation of the morphological characteristics according to the taxonomic keys (Lane 1953, Consoli & Lourenço-de-Oliveira 1994). The mosquitoes were grouped into pools of up to 50 individuals that were sorted with respect to trap, site of capture and species and they were then stored in liquid nitrogen in the field.

Caimans were captured from sites where a high concentration of these animals was observed. The animals were seized by boat in lentic systems that were formed by intermittent rivers and taken to the banks. All animals were weighed, measured, gendered, numbered and bled before release. The whole procedure was performed only with physical restraint and each animal was worked and released within 15 min. Caiman blood samples were obtained by puncture of the internal jugular vein between the 1st and 2nd cervical vertebrae as described previously (Olson et al. 1975).

Blood samples were taken from horses of different ages from different ranches. Concomitantly, an interview was conducted with the rural workers of the ranches that were visited who were asked if they had seen horses with any abnormality, such as clinical signs that involved the central or peripheral nervous system. We also collected information about the sex, age and breed of the horses that were present. Additional information included whether the horse had recently received vaccine or medical treatment and whether any horse had been moved into or out of South Pantanal.

Epitope-blocking enzyme-linked immunosorbent assay (blocking ELISA) - Caiman and horse sera were first screened for flaviviruses-reactive antibodies using a blocking ELISA as described previously (Blitvich et al. 2003a, c). Briefly, the ability of the sera to block the binding of the monoclonal antibody 6B6C-1 to the WNV antigen was compared to the blocking ability of negative control horse serum. The monoclonal antibody 6B6C-1 is appropriate to screen for any flavivirus-reactive antibody. Data are expressed as relative percentages. Samples were considered seropositive when the inhibition values produced were $\geq 35\%$.

Plaque-reduction neutralisation test (PRNT) - The seropositive sera of the blocking ELISA was sequentially heat-inactivated and submitted to a PRNT₉₀ as described previously (Russell et al. 1967). Briefly, two-fold dilutions that ranged from 1:10-1:320 of each blocking ELISA seropositive sample were tested for their ability to neutralise plaque formation by the chimeras SLEV and WNV that were derived from highly attenuated Yellow fever virus that contain the premembrane-envelope (prM-E) and prM-E and envelope protein genes, respectively (Arroyo et al. 2001, Pugachev et al. 2004, Komar et al. 2009), and Cacipacore virus (CPCV), Ilheus virus (ILHV) and Rocio virus (ROCV), which are the Brazilian flaviviruses that are phylogenetically related to WNV (Kuno et al. 1998). Serum was considered seropositive to a virus species when it reduced at least 90% of the formation of plaques of this virus at $\geq 1:40$ dilutions and its neutralising antibody titre was \geq four-fold greater than what was observed for the other tested flaviviruses. The WNV seropositive samples were also submitted to PRNT₉₀ for Bussuquara virus (BSQV).

Virus isolation - The caiman sera and mosquito samples were assayed for virus isolation. The caiman serum samples at 1:5 dilutions were inoculated into VERO cell culture flasks (Stim 1969). The macerated mosquito pool suspensions were inoculated into C6/36 cell culture flasks (White 1987). The incubated cell flasks were observed during 14 days to look for any evidence of viral cytopathic effect (CPE).

Reverse-transcriptase polymerase chain reaction (RT-PCR) - The molecular approach was performed in all caiman serum samples and the mosquito samples in which the inoculated flasks showed CPE in virus isolation. These samples were submitted to RNA extraction using a commercial kit (QIAamp® Viral RNA Mini Kit QIAGEN, USA) in accordance with the manufacturer's instructions and was followed by a generic nested RT-PCR, which sequenced for the detection and identification of arbovirus members of the *Flavivirus* and *Alphavirus* genus, as described previously (Sánchez-Seco et al. 2001, 2005).

RESULTS

Serum samples of 168 horses from four of the six studied cattle ranches were sampled. Most of these equines were adult, tame, healthy thoroughbreds of the Cavalo Pantaneiro breed. All 168 horse sera were first tested using the blocking ELISA and 93 (55.4%) were se-

ropositive. When this subset of 93 seropositive samples was tested using PRNT₉₀, five (3%) were seropositive for WNV, nine (5.4%) were seropositive for SLEV, 18 (10.7%) were seropositive for ILHV, three (1.8%) were seropositive for CPCV, but none were seropositive for ROCV, using the criteria of a \geq four-fold antibody titre difference among the five flaviviruses that were assayed. All of the WNV-seropositive samples were $< 1:10$ for BSQV (Table). From the remaining 58 (34.5%) blocking ELISA seropositive samples, 56 (33.3%) samples were considered seropositive for an undetermined flavivirus using the < 4 -fold antibody titre difference and two (1.2%) were considered seronegative with titres $< 1:10$ for all of the flaviviruses that were assayed by PRNT₉₀.

From the five horses that were seropositive for WNV, two horses had dates of birth available and were four and five years old at the time of blood collection. All five seropositive animals had no history of journey outside South Pantanal or WNV vaccination. All WNV-seropositive samples were retested for confirmation and the antibody serum titres were 1:80 ($n = 1$), 1:160 ($n = 2$) and $\geq 1:320$ ($n = 2$). The sera of these two horses with antibody serum titres $\geq 1:320$ were retested with higher dilutions and an antibody serum titre of 1:320 was confirmed. These five WNV-seropositive horses came from three out of the four ranches where horses were sampled.

Concerning SLEV of the nine seropositive horses from the four ranches sampled, six had dates of birth available with ages that ranged from four-11 years old. All of the SLEV-seropositive horses had never left South Pantanal and their antibody serum titres ranged from 1:40- $\geq 1:320$.

Of 18 ILHV-seropositive horses, 12 had available dates of birth with ages that ranged from three-13 years old in animals from all of the ranches where horses were sampled. None of these animals had a history of travel and their antibody serum titres ranged from 1:40- $\geq 1:320$.

From the three horses that were seropositive for CPCV, two had available dates of birth and were 12 years old at the time of blood collection. These two animals were from the same ranch, had never left the region and had antibody serum titres for CPCV of 1:160. Most of the blocking ELISA-seropositive samples had antibody titers $< 1:10$ for ROCV. Of the samples with antibody titres $\geq 1:10$ for ROCV, only one had a titre $\geq 1:40$. Because the antibody titre difference for the other flaviviruses assayed was less than four-fold, it was considered seropositive for an undetermined flavivirus. Of the 93 blocking ELISA-seropositive horses, 56 (60.2%) with titre differences less than four-fold were considered seropositive for an undetermined flavivirus. The two blocking ELISA-seropositive samples that had serum titres $< 1:10$ for all five flaviviruses assayed were considered as seronegative in the present study.

Thirty caiman specimens up to 1.5 m in length were captured from two distinct sites of the same ranch. The sera from the 30 caimans displayed negative results to virus isolation, RT-PCR and blocking ELISA and were subsequently not submitted to PRNT₉₀ methods.

A total of 1,204 mosquitoes from 10 species were captured at 19 sites that were randomly selected on five ranches. Specimens of the following species were captured:

Ochlerotatus scapularis, *Psorophora albigena*, *Sabethes albiprivus*, *Culex declarator*, *Culex quinquefasciatus*, *Mansonia titillans*, *Anopheles albiparvus* s.l., *Psorophora ferox*, *Anopheles triannulatus* s.l. and *Psorophora ciliata*. The mosquitoes were grouped into 65 pools of up to 50 individuals that were sorted with respect to trap, site of capture and species. Of a total of the 65 mosquito pools

that were assayed, one (1.5%) pool with 22 specimens of *O. scapularis* exhibited CPE four days post-infection (dpi). The same CPE was observed on subsequent passages by inoculation into new C6/36 cell culture flasks four dpi. Nevertheless, despite the CPE, the pool and the supernatant from the sequential passages was negative to flaviviruses and alphaviruses as measured by RT-PCR.

TABLE
Plaque-reduction neutralisation test antibody titres for SLEV, ILHV, WNV, ROCV, CPCV and BSQV in epitope-blocking enzyme-linked immunosorbent assay-reactive horses which were seropositive for a determined species considering the four-fold titre difference criteria

ID	Ranch	Age	Sex	Tame	Travel	SLEV ^a	ILHV	WNV ^a	ROCV	CPCV	BSQV	Interpretation
136	B	12	F	T	N	1:20	1:80	< 1:10	1:20	1:20	-	ILHV
140	B	8	F	T	N	1:20	≥ 1:160	< 1:10	< 1:10	1:10	-	ILHV
145	B	11	F	T	N	1:40	≥ 1:320	1:10	1:10	1:80	-	ILHV
158	B	6	F	T	N	1:80	1:10	1:20	< 1:10	1:20	-	SLEV
160	B	12	F	T	N	< 1:10	> 1:40	< 1:10	< 1:10	< 1:10	-	ILHV
162	B	3	F	T	N	< 1:10	> 1:40	< 1:10	< 1:10	< 1:10	-	ILHV
174	B	4	F	U	N	1:40	< 1:20	< 1:10	< 1:10	1:10	-	SLEV
182	B	4	F	U	N	1:40	< 1:10	< 1:10	< 1:10	< 1:10	-	SLEV
183	B	4	F	U	N	1:10	1:40	< 1:10	< 1:10	1:10	-	ILHV
190	A	13	G	T	N	1:40	≥ 1:160	< 1:10	< 1:10	< 1:10	-	ILHV
194	A	12	G	T	N	1:20	< 1:40	1:20	1:20	1:160	-	CPCV
197	A	13	G	T	N	< 1:10	> 1:40	< 1:10	< 1:10	< 1:10	-	ILHV
199	A	12	G	T	N	< 1:10	> 1:40	< 1:10	< 1:10	< 1:10	-	ILHV
203	A	12	G	T	N	1:20	< 1:40	< 1:10	1:20	1:160	-	CPCV
207	A	13	G	T	N	< 1:10	1:40	< 1:10	< 1:10	< 1:10	-	ILHV
208	A	8	G	T	N	1:160	1:40	1:20	1:10	1:40	-	SLEV
211	A	11	G	T	N	≥ 1:320	< 1:40	1:10	< 1:10	1:10	-	SLEV
214	A	5	G	T	N	1:20	< 1:40	1:320	< 1:10	< 1:20	< 1:10	WNV
216	A	7	G	T	N	1:20	≥ 1:320	1:80	1:20	1:80	-	ILHV
220	A	5	G	T	N	1:160	< 1:40	1:10	< 1:10	< 1:10	-	SLEV
241	A	4	G	U	N	< 1:10	≥ 1:80	< 1:10	< 1:10	< 1:10	-	ILHV
250	A	4	G	U	N	< 1:10	< 1:40	1:160	< 1:10	< 1:20	< 1:10	WNV
253	F	UNK	G	T	N	< 1:10	> 1:40	< 1:10	< 1:10	< 1:10	-	ILHV
268	F	UNK	G	T	N	1:80	< 1:10	< 1:10	< 1:10	< 1:10	-	SLEV
269	F	UNK	G	T	N	1:80	< 1:40	1:20	< 1:10	1:20	-	SLEV
270	F	UNK	F	T	N	< 1:10	1:40	< 1:10	< 1:10	< 1:10	-	ILHV
271	F	UNK	S	U	N	1:10	≥ 1:160	1:20	< 1:10	1:40	-	ILHV
280	F	UNK	F	U	N	1:20	< 1:10	1:80	< 1:10	< 1:20	< 1:10	WNV
283	E	UNK	G	T	N	< 1:10	1:40	< 1:10	< 1:10	< 1:20	-	ILHV
285	E	UNK	G	T	N	< 1:10	< 1:40	1:160	< 1:10	1:20	< 1:10	WNV
290	E	UNK	G	T	N	< 1:10	1:80	< 1:10	< 1:10	< 1:10	-	ILHV
303	E	UNK	G	T	N	< 1:10	1:10	< 1:10	< 1:10	1:40	-	CPCV
305	E	UNK	G	T	N	≥ 1:320	< 1:40	1:20	1:10	< 1:10	-	SLEV
306	E	UNK	G	T	N	1:10	< 1:40	1:320	< 1:10	1:80	< 1:10	WNV
309	E	UNK	G	T	N	1:10	1:80	< 1:10	< 1:10	< 1:10	-	ILHV

a: chimera; BSQV: Bussuquara virus; CPCV: Cacipacore virus; F: female; G: gelding; ID: identification of animal; ILHV: Ilheus virus; N: no travel outside South Pantanal; ROCV: Rocio virus; S: stallion; SLEV: Saint Louis encephalitis virus; T: tamed; U: untamed; UNK: unknown; WNV: West Nile virus; -: non-submitted.

DISCUSSION

Flaviviruses serological test result interpretation is sometimes complex and requires careful evaluation (Komar 2001), especially where data from areas of intense flaviviruses co-circulation is expected, such as for South American countries. Considering the complex cross-reactivity of flaviviruses, several studies have adopted ELISA as a screening method followed by more virus-specific serological tests, such as PRNT methods (Blitvich et al. 2003b, Morales-Betoulle et al. 2006, Diaz et al. 2008). In the present study, the use of blocking ELISA as a screening method followed by PRNT₉₀ for the most related sylvatic Brazilian flaviviruses allowed for the detection of specific neutralising antibodies to WNV in 3% of the horses tested from the Pantanal region for the first time in Brazil.

Based on the results that were obtained with the blocking ELISA, 93 horse sera presented antibodies to flaviviruses. When the same sera were tested using PRNT₉₀ for SLEV, WNV, ILHV, ROCV and CPCV, except for two samples, none had a serum titre < 1:10 for all of the flaviviruses that were tested, which disclosed the high specificity of the blocking ELISA method. According to our data, in conjunction with the fact that blocking ELISA does not require the use of multiple capture antibodies, which designated this test as useful for virtually any vertebrate species, the present study corroborated the application of blocking ELISA as an efficient screening method for flaviviruses serosurveys.

We have searched for the circulation of WNV in a country with an intense circulation of other flaviviruses. Therefore, to avoid misinterpretation based on serum cross-reactions it was necessary to employ a maximum specificity PRNT and only consider seropositive samples that promote a $\geq 90\%$ reduction in the number of WNV plaques as has been described previously in South and Central American serosurveys (Quirin et al. 2004, Mattar et al. 2005). Besides SLEV, ROCV, ILHV and CPCV, other Brazilian sylvatic flaviviruses that are related to WNV include the BSQV and Iguape virus (IGUV) (Kuno et al. 1998). Despite the geographic dispersal of BSQV, which is mainly located in the Caribbean and Central America (Gubler et al. 2007), the WNV seropositive samples were also submitted to PRNT₉₀ for BSQV to minimise potential high-titered cross reaction antibodies. However, considering that IGUV was only isolated from sentinel mice in the Southeastern Brazil (Coimbra et al. 1993) and that serosurveys suggest infrequent occurrence of overt infection, in the present study IGUV was not included in the PRNT methods.

To minimise the cross-reaction results in this serosurvey to WNV, we considered the seropositivity for WNV and for each of the other flaviviruses assayed, only sera with a 1:40 monotypic reaction or with an antibody titre higher than 1:40 and a four-fold higher to a species than its other related flaviviruses that were assayed. However, considering these criteria, 56 (60.2%) of the 93 blocking ELISA-seropositive samples were seropositive for an undetermined flavivirus species. The high prevalence of these inconclusive results may be attributed to the cross-reacting antibodies that are usually observed among species of a Japanese encephalitis virus group (JEVG)

(Tarr & Hammon 1974) and to the original antigenic sin phenomenon, where some individuals that are sequentially infected by a heterologous flavivirus species can boost antibody levels against the original virus, which produces heterologous cross-reacting antibodies (Inouye et al. 1984). Considering the similar maintenance cycles of the flaviviruses that were assayed together with the hypothesis of its co-circulation in the South Pantanal, the original antigenic sin phenomenon may have had an effect on the difference of antibody titres among the flaviviruses that were assayed and some sequential infections by a heterologous flavivirus species may not have been detected. Of the 56 horses that were seropositive for an undetermined flavivirus species, 13 (23.2%) had antibody titres $\geq 1:40$ for WNV, nine (69.2%) of which had antibody titres $\geq 1:80$. Therefore, it is possible that the other 13 (7.7%) of the 168 horses that were tested had contact with WNV. It is noteworthy that neutralising antibody titres are frequently more than two-fold lower in PRNT with chimeras WNV and SLEV than with wild virus (Pugachev et al. 2004, Komar et al. 2009). Therefore, it is also quite possible that some low titre WNV and SLEV-seropositive samples were not detected.

The detection of specific neutralising antibodies to WNV in 3% of horses tested in the Pantanal region from Brazil was reported in this study. This prevalence is similar to the values previously reported for horses from other countries of Central and South America, such as Venezuela, where a prevalence of 4.3% was described (Bosch et al. 2007); Colombia, with a prevalence of 9% (Mattar et al. 2005), and El Salvador, with the highest seropositivity to WNV of 13.7% (Cruz et al. 2005). Despite the adoption of the same criteria for the detection of WNV-seropositive animals, it is important to note the difference of the flaviviruses that were used in the PRNT methods that were performed in different areas.

The CPCV, which is a Brazilian virus of JEVG, is as phylogenetically related to WNV as SLEV and is even more closely related to WNV than ILHV, BSQV or ROCV, which belong to other flavivirus serocomplexes (Kuno et al. 1998). Although CPCV is a non-pathogenic arbovirus to date, we detected many samples with neutralising antibodies for both CPCV and WNV, which suggest that in future serosurveys for WNV and other sylvatic flaviviruses to be conducted in Brazil, the investigation for CPCV should be included as a differential diagnosis.

The highest seropositivity for ILHV was expected because the circulation of ILHV in South Pantanal horses was reported in the 1990s (Iversson et al. 1993). Although ROCV, which is an indigenous Brazilian flavivirus species, was the causative agent of an extensive encephalitis epidemic in Southeastern Brazil in the 1970s (Tiriba et al. 1976), the detection of neutralizing antibodies for ROCV nowadays is unusual. Our findings corroborate this data because except for one sample, all blocking ELISA-seropositive horses had PRNT antibody titres < 1:40 for ROCV. One should be aware that, although phylogenetically related to ILHV (Medeiros et al. 2007) that presented the highest prevalence in our study, none of the horses tested were seropositive for ROCV, which demonstrates the high-quality specificity of the PRNT assay.

Based on our serological data, we reported indirect evidence that argues for the silent activity of WNV in the Brazilian Pantanal. Considering this hypothesis and taking into account the age available for the seropositive animals of four and five years old, we can infer that at least during the period from 2005-2009, WNV circulated in the area. The detection of SLEV-seropositive four-year-old horses suggested that some were infected by SLEV between 2005-2009, which corroborates the recent report of the detection of neutralising antibodies to SLEV in horses sampled in 2007 in the same area of the present study (Pauvolid-Corrêa et al. 2010b). For CPCV, the two seropositive horses with available dates of birth were 12 years old; therefore, the period of probable circulation was difficult to determine. The circulation of ILHV in South Pantanal was described in the 1990s (Iversson et al. 1993) and this putative circulation has remained. However, according to the age of our 18 ILHV-seropositive animals, we can only deduce that between 2006-2009, ILHV circulated in the area. Although a better estimation of recent circulation of these flaviviruses for future surveillance should be based on IgM assays, our data suggested the viral activity after 2005 of at least three flaviviruses, including WNV, in the Brazilian Pantanal.

We encourage further more encompassing surveys for WNV in the South Pantanal and other bordering areas that especially focus on virus isolation to definitively confirm the circulation of WNV in the region, and consequently, identify vectors and vertebrate hosts that are involved in the virus local maintenance cycle and transmission. Despite of the serological evidence of horse infections by WNV in the Pantanal region presented here, morbidity was not observed, which suggests that WNV could be circulating through silent enzootic cycles in an area with pristine conditions. In fact, WNV has not been related to horse disease in most of the South American countries with the exception of Argentina (Morales et al. 2006). However, WNV is an important cause of encephalitis in horses in the USA and should be considered in the future in the differential diagnosis of neurological infections in horses from South Pantanal.

Despite of the small number of caimans that were sampled, the negative results obtained using virus isolation, RT-PCR and blocking ELISA suggested that the participation of caimans in flaviviruses cycles in South Pantanal is improbable. Nevertheless, the high prevalence of this crocodilian species in the Pantanal and the high concentration that was observed in some remaining collections of water during the dry season associated to the present data of serological evidence of the activity of members of JEVG in the area justify the need of further arbovirus investigations in local caimans.

All mosquito species identified in the present study were previously reported in the Nhecolândia sub-region (Pauvolid-Corrêa et al. 2010a). Moreover, the non-detection of *Flavivirus* and *Alphavirus* RNA in the *O. scapularis* pool that caused CPE in C6/36 cells implied a virus identity that was not related to WNV, SLEV or alphaviruses, which would have been indicated by the RT-PCR. Further investigations of the other arbovirus groups are currently in progress.

Finally, we have reinforced that the Brazilian Pantanal is an interesting scenario where sylvatic flaviviruses cycles are probably maintained in silent progression in a relatively balanced ecosystem. The recently increased environmental degradation of this sensitive, complex biome could lead to changes in the intricate ecological relationships that disturb natural enzootic arbovirus cycles, which potentially favours local arbovirus outbreaks.

ACKNOWLEDGEMENTS

To the cordial people of the Pantanal, who have wisely managed to keep the Pantanal preserved through the centuries, especially the rural workers of the ranches sampled including Marcelo Santos, Francisco Rondon, Luis Fernando Santiago, Gilberto Silva, Marcelo Dias, José Mendes, Marcos Moraes, Luiz Graça, Luiz da Silva, Benis da Silva and Sebastião Gonzaga, to Henrique de Jesus, Augusto Silva, Cremilson Gonçalves, Murilo Maciel and Oziel da Silva, of Embrapa Pantanal to Victoria Luppó, Pablo Baroni, Lucas Bogdanic, Stella Fuster and Silvina Coenaga, of Instituto Nacional de Enfermidades Virales Humanas, to Vanessa Muller, of Faculdade de Medicina de Ribeirão Preto, and to Heitor Herrera, Fernando Tavares, Flávia dos Santos, Ana Bispo and Elba Lemos, of Fio-cruz, for their contributions.

REFERENCES

- Adâmolí I 1982. *O Pantanal e suas relações fitogeográficas com os cerrados. Discussão sobre o conceito "Complexo Pantanal"*, In Anais do 32º Congresso Nacional de Botânica, Teresina, p. 109-119.
- Alho CJR 2005. The Pantanal. In LH Fraser, PA Keddy, *The world's largest wetlands: ecology and conservation*, Cambridge University Press, New York, p. 203-271.
- Alho CJR, Lacher TE, Gonçalves HC 1988. *Environmental degradation in the Pantanal ecosystem. Bioscience* 38: 164-171.
- Arroyo J, Miller CA, Catalan J, Monath TP 2001. Yellow fever vector live-virus vaccines: West Nile virus vaccine development. *Trends Mol Med* 7: 350-354.
- Autorino GL, Battisti A, Deubel V, Ferrari G, Forletta R, Giovannini A, Lelli R, Murri S, Scicluna MT 2002. West Nile virus epidemic in horses, Tuscany Region, Italy. *Emerg Infect Dis* 8: 1372-1378.
- Blitvich BJ, Bowen RA, Marlenee NL, Hall RA, Bunning ML, Beaty BJ 2003a. Epitope-blocking enzyme-linked immunosorbent assays for detection of West Nile virus antibodies in domestic mammals. *J Clin Microbiol* 41: 2676-2679.
- Blitvich BJ, Fernandez-Salas I, Contreras-Cordero JF, Marlenee NL, Gonzalez-Rojas JI, Komar N, Gubler DJ, Calisher CH, Beaty BJ 2003b. Serologic evidence of West Nile virus infection in horses, Coahuila state, Mexico. *Emerg Infect Dis* 9: 853-856.
- Blitvich BJ, Marlenee NL, Hall RA, Calisher CH, Bowen RA, Roehrig JT, Komar N, Langevin SA, Beaty BJ 2003c. Epitope-blocking enzyme-linked immunosorbent assays for the detection of serum antibodies to West Nile virus in multiple avian species. *J Clin Microbiol* 41: 1041-1047.
- Bosch I, Herrera F, Navarro J, Lentino M, Dupuis A, Maffei J, Jones M, Fernández E, Pérez N, Pérez-Emán J, Guimarães AE, Barrera R, Valero N, Ruiz J, Velásquez G, Martinez J, Comach G, Komar N, Spielman A, Kramer L 2007. West Nile virus, Venezuela. *Emerg Infect Dis* 13: 651-653.
- Campos ZMS, Coutinho ME, Mourão G 2005. *Metodologia para contagem e avaliação de ninhos de jacaré-do-pantanal usando o ultraleve*, Embrapa Pantanal, Circular Técnica 58, Corumbá, 3 pp.

- CDC - Centers for Disease Control and Prevention 1999. Outbreak of West Nile-like viral encephalitis-New York, 1999. *Morb Mortal Wkly Rep* 48: 845-849.
- Coimbra TLM, Nassar ES, Nagamori AH, Ferreira IB, Pereira LE, Rocco IM, Ueidato M, Romano NS 1993. **Iguape: a newly recognized flavivirus from São Paulo state, Brazil.** *Intervirology* 36: 144-152.
- Consoli RAGB, Lourenço-de-Oliveira R 1994. *Principais mosquitos de importância sanitária no Brasil*, Editora Fiocruz, Rio de Janeiro, 228 pp.
- Couissinier-Paris P 2006. West Nile virus in Europe and Africa: still minor pathogen, or potential threat to public health? *Bull Soc Pathol Exot* 99: 348-354.
- Cruz L, Cardenas VM, Abarca M, Rodriguez T, Reyna RF, Serpas MV, Fontaine RE, Beasley DWC, Travassos Da Rosa APA, Weaver SC, Tesh RB, Powers AM, Suarez-Rangel G 2005. Short report: serological evidence of West Nile virus activity in El Salvador. *Am J Trop Med Hyg* 72: 612-615.
- Diaz LA, Komar N, Visintin A, Juri MJD, Stein M, Allende RL, Spinsanti L, Konigheim B, Aguilar J, Laurito M, Almirón W, Contigiani M 2008. West Nile virus in birds, Argentina. *Emerg Infect Dis* 14: 689-691.
- Gubler DJ, Kuno G, Markoff L 2007. Flaviviruses. In DM Knipe, PM Howley, DE Griffin, RA Lamb, SE Straus, MA Martin, B Roizman, *Field's Virology*, 5th ed., Lippincott Williams & Wilkins, Philadelphia, p. 1153-1252.
- Inouye S, Matsuno S, Tsurukubo Y 1984. "Original antigenic sin" phenomenon in experimental flavivirus infections of guinea pigs: studies by enzyme-linked immunosorbent assay. *Microbiol Immunol* 28: 569-574.
- Iversson LB, Silva RAMS, Travassos da Rosa APA, Barros VLRS 1993. Circulation of Eastern equine encephalitis, ilheus, maguari and tacaigua viruses in equines of the Brazilian Pantanal, South America. *Rev Inst Med Trop Sao Paulo* 35: 355-359.
- Jacobson ER, Ginn PE, Troutman JM, Farina L, Stark L, Klenk K, Burkhalter KL, Komar NJ 2005. West Nile virus infection in farmed American alligators (*Alligator mississippiensis*) in Florida. *Wildl Dis* 41: 96-106.
- Joubert L, Oudar J, Hannoun C, Beytout D, Corniou B, Guillon JC, Panthier R 1970. Epidemiology of the West Nile virus: study of a focus in Camargue. IV. Meningo-encephalomyelitis of the horse. *Ann Inst Pasteur (Paris)* 118: 239-247.
- Klenk K, Snow J, Morgan K, Bowen R, Stephens M, Foster F, Gordy P, Beckett S, Komar N, Gubler D, Bunning M 2004. Alligators as West Nile virus amplifiers. *Emerg Infect Dis* 10: 2150-2155.
- Komar N 2001. West Nile virus surveillance using sentinel birds. *Ann NY Acad Sci* 951: 58-73.
- Komar N, Clark GG 2006. West Nile virus activity in Latin America and Caribbean. *Rev Panam Salud Publica* 19: 112-117.
- Komar N, Langevin S, Monath TP 2009. Use of a surrogate chimeric virus to detect West Nile virus-neutralizing antibodies in avian and equine sera. *Clin Vaccine Immunol* 16: 134-135.
- Kuno G, Chang GJ, Tsuchiya KR, Karabatsos N, Cropp B 1998. Phylogeny of the genus flavivirus. *J Virol* 72: 73-83.
- Lane J 1953. *Neotropical Culicidae*, Editora Universidade de São Paulo, São Paulo, 1112 pp.
- Lindsey NP, Staples JE, Lehman JA, Fischer M 2010. Surveillance for human West Nile virus disease - United States, 1999-2008. *MMWR Surveill Summ* 59: 1-17.
- Mattar S, Edwards E, Laguado J, González M, Alvarez J, Komar N 2005. West Nile virus antibodies in Colombian horses. *Emerg Infect Dis* 9: 1497-1498.
- Medeiros DB, Nunes MR, Vasconcelos PF, Chang GJ, Kuno G 2007. Complete genome characterization of Rocio virus (Flavivirus: Flaviviridae), a Brazilian flavivirus isolated from a fatal case of encephalitis during an epidemic in São Paulo state. *J Gen Virol* 88: 2237-2246.
- Morales MA, Barrandeguy M, Fabbri C, Garcia JB, Vissani A, Trono K, Gutierrez G, Pigretti S, Menchaca H, Garrido N, Taylor N, Fernandez F, Levis S, Enría D 2006. West Nile virus isolation from equines in Argentina, 2006. *Emerg Infect Dis* 12: 1559-1561.
- Morales-Betoulle ME, Morales H, Blitvich BJ, Powers AM, Davis EA, Klein R, Córdón-Rosales C 2006. West Nile virus in horses, Guatemala. *Emerg Infect Dis* 6: 1038-1039.
- MS - Ministério da Saúde, Brasil 2003. Secretaria de Vigilância em Saúde. Primeiro inquérito sorológico em aves migratórias e nativas do Parque Nacional da Lagoa do Peixe/RS para detecção do vírus do Nilo ocidental. *Boletim eletrônico epidemiológico* 3: 3-12.
- MS - Ministério da Saúde, Brasil 2004. Secretaria de Vigilância em Saúde. Inquérito sorológico em aves migratórias e residentes de Galinhos/RN para detecção do vírus da febre do Nilo ocidental e outros vírus. *Boletim eletrônico epidemiológico* 4: 1-12.
- Nunes AP, Tomas WM 2008. *Aves migratórias e nômades ocorrentes no Pantanal*, Embrapa Pantanal, Corumbá, 123 pp.
- Olson GA, Hessler JR, Faith RE 1975. Technics for blood collection and intravascular infusion of reptiles. *Lab Anim Sci* 25: 783-786.
- Pauvolid-Corrêa A, Tavares FN, Alencar J, Santos SJ, Murta M, Serra-Freire NM, Pellegrin AO, Gil-Santana H, Guimarães AE, Silva EE 2010a. Preliminary investigation of Culicidae species in South Pantanal, Brazil and their potential importance in arbovirus transmission. *Rev Inst Med Trop Sao Paulo* 52: 17-24.
- Pauvolid-Corrêa A, Tavares FN, Costa EV, Burlandy FM, Murta M, Pellegrin AO, Nogueira MF, Da Silva EE 2010b. **Serologic evidence of the recent circulation of Saint Louis encephalitis virus and high prevalence of equine encephalitis viruses in horses in the Nhecolândia sub-region in South Pantanal, Central-West Brazil.** *Mem Inst Oswaldo Cruz* 105: 829-833.
- Pauvolid-Corrêa A, Varella RB 2008. Epidemiological aspects of West Nile fever. *Rev Bras Epidemiol* 11: 463-472.
- Platonov AE, Shipulin GA, Shipulina AY, Tyutyunnik EN, Frolochkina TI, Lanciotti RS, Yazyshina S, Platonova OV, Obukhov IL, Zhukov AN, Vengerov YY, Pokrovskii VI 2001. Outbreak of West Nile virus infection, Volgograd region, Russia, 1999. *Emerg Infect Dis* 7: 128-132.
- Pugachev KV, Guirakhoo F, Mitchell F, Ocran SW, Parsons M, Johnson BW, Kosoy OL, Lanciotti RS, Roehrig JT, Trent DW, Monath TP 2004. Construction of yellow fever/St. Louis encephalitis chimeric virus and the use of chimeras as a diagnostic tool. *Am J Trop Med Hyg* 71: 639-645.
- Quirin R, Salas M, Zientara S, Zeller H, Labie J, Murri S, Lefrançois T, Petitclerc M, Martinez D 2004. West Nile virus, Guadeloupe. *Emerg Infect Dis* 10: 706-708.
- Russell PK, Nisalak A, Sukhachana P, Vivona S 1967. A plaque reduction test for dengue virus neutralizing antibodies. *J Immunol* 99: 285-290.
- Sánchez-Seco MP, Rosario D, Domingo C, Hernández L, Valdés K, Guzmán MG, Tenorio A 2005. **Generic RT-nested-PCR for detection of flaviviruses using degenerated primers and internal control followed by sequencing for specific identification.** *J Virol Methods* 126: 101-109.
- Sánchez-Seco MP, Rosario D, Quiroz E, Guzmán G, Tenorio A 2001. A generic nested-RT-PCR followed by sequencing for detection and identification of members of the *Alphavirus* genus. *J Virol Methods* 95: 153-161.

- Silva JV, Abdon MM 1988. Delimitação do Pantanal brasileiro e suas sub-regiões. *Pesq Agrop Bras* 33: 1703-1711.
- Smithburn KC 1942. Differentiation of the West Nile virus from the viruses of St. Louis and Japanese b encephalitis. *J Immunol* 44: 25-31.
- Smithburn JS, Hughes TP, Burke AW, Paul JH 1940. A neurotropic virus isolated from the blood of a native of Uganda. *Am J Trop Med Hyg* 20: 471-492.
- Steinman A, Banet-Noach C, Tal S, Levi O, Simanov L, Perk S, Malkinson M, Shpigel N 2003. West Nile virus infection in crocodiles. *Emerg Infect Dis* 9: 887-888.
- Stim TB 1969. Arbovirus plaquing in two simian kidney cell lines. *J Gen Virol* 5: 329-338.
- Tarr GC, Hammon WM 1974. Cross-protection between group B arboviruses: resistance in mice to Japanese B encephalitis and St. Louis encephalitis viruses induced by dengue virus immunization. *Infect Immun* 9: 909-915.
- Tber Abdelhaq A 1996. West Nile fever in horses in Morocco. *Bull Off Int Epizoot* 11: 867-869.
- Tiriba AC, Miziara AM, Lourenço R, Costa CRB, Costa CS, Pinto GH 1976. Encefalite humana primária epidêmica por arbovírus observada no litoral sul de São Paulo. *Rev Assoc Med Bras* 22: 415-420.
- Tsai TF, Popovici F, Cernescu C, Campbell GL, Nedelcu NI 1998. West Nile encephalitis epidemic in southeastern Romania. *Lancet* 352: 767-771.
- White LA 1987. Susceptibility of *Aedes albopictus* C6/36 cells to viral infection. *J Clin Microbiol* 25: 1221-1224.