

Article

Received 10 Nov 2010
Accepted 16 Feb 2011
Available online 23 Sep 2011

Keywords:

alkaloids
antiproliferative activity
diterpenes
medicinal plants

ISSN 0102-695X
<http://dx.doi.org/10.1590/S0102-695X2011005000174>

Constituents and antiproliferative activity of extracts from leaves of *Croton macrobothrys*

Lucimar B. Motta,¹ Cláudia M. Furlan,¹ Deborah Y. A. C. Santos,¹ Maria L. F. Salatino,¹ Joaquim M. Duarte-Almeida,² Giuseppina Negri,² João E. de Carvalho,³ Ana Lúcia T. G. Ruiz,³ Inês Cordeiro,⁴ Antonio Salatino*¹

¹Instituto de Biociências, Universidade de São Paulo, Brazil,

²Departamento de Psicobiologia, Universidade Federal de São Paulo, Brazil,

³Divisão de Farmacologia e Toxicologia, Universidade Estadual de Campinas, Brazil,

⁴Seção de Curadoria do Herbário, Instituto de Botânica, Brazil.

Abstract: *Croton macrobothrys* Baill, Euphorbiaceae, is a tree from the Atlantic Forest in Southern Brazil, used in traditional medicine and popularly known as "dragon's blood" and "pau-sangue". Leaf *n*-hexane, dichloromethane and methanol extracts were analyzed by GC/MS and evaluated for their *in vitro* antiproliferative activity on cell lines 786-0 (kidney), HT-29 (colon), K562 (leukemia), NCI-ADR/RES (drug resistant ovary), NCI-H460 (lung), MCF-7 (mammary), PC-3 (prostate), OVCAR-3 (ovary), U251 (glioma) and UACC-62 (melanoma). The dichloromethane extract exhibited activity against all cell lines at the concentration 25 µg/mL, in particular on cell lines NCI-H460 (GI50 0.33 µg/mL) and K562 (GI50 0.91 µg/mL). Relevant constituents in dichloromethane extract are the alkaloids corydine and salutaridine, as well as the diterpenes geranylgeraniol and crotonin-derived clerodanes.

Introduction

Over 85000 plant species have been documented as used worldwide in traditional medicine. The World Health Organization (WHO) estimates that almost 75% of the world's population employs plant-based traditional remedies (Liu et al., 2008). Over 60% of the currently used anticancer chemotherapeutics are derived in one way or another from plants (Cragg & Newman, 2009; Tan et al., 2006). The major categories of plant-derived compounds with anticancer properties are alkaloids and terpenoids (Gupta et al., 2005); examples are vincristine and vinblastine from *Catharanthus roseus* (L.) G. Don (Johnson et al., 1963; Carvalhaes et al., 2002) and taxol and docetaxel from *Taxus brevifolia* Nutt. (Wani et al., 1971).

Croton is a genus of Euphorbiaceae comprising around 1300 species, widespread in tropical regions of the New and Old Worlds. Several species have a long role in traditional medicine in Africa, Asia and South America (Salatino et al., 2007; Perazzo et al., 2007). Red latex of *C. draconoides* Mull. Arg., *C. lechleri* Müll. Arg., *C. palanostigma* Klotzsch and *C. urucurana* Baill., popularly known as "sangre de drago" (dragon's blood), are used medicinally by indigenous and rural communities (Riina, et al., 2009). In upper Amazonia, *Croton* latex is used to treat

tuberculosis and cancer, sometimes combined with other medicinal plants (e.g., uña-de-gato, or *Uncaria tomentosa* (Willd. ex Roem. & Schult.) DC. (Maxwell, 1990). *C. palanostigma* has been used by indigenous people of the region of Pucallpa (Peru) to treat tumors (Jones 2003). The red latex of *C. lechleri* has been shown to have anti-tumor activity (Rossi et al., 2003). Shoots of *C. hieronymi* Griseb. have been shown to have strong activity against lung A-549 carcinoma cells, mouse lymphoma and human colon carcinoma (Catalán et al., 2003). The dichloromethane extract of leaves of *C. zambesicus* Müll. Arg. showed *in vitro* cytotoxicity against human cervix carcinoma cells (Block et al., 2002). *Croton* species are abundant sources of active substances against cancer, such as diterpenoids (clerodane, furoclerodane and acyclic diterpenes) and alkaloids (e.g. taspine) (Salatino et al., 2007).

Croton macrobothrys Baill. is a tree from the Atlantic Forest in Eastern Brazil, popularly known as "dragon's blood", "pau-sangue" and "sangre-de-drago-de-folha-miúda" and used in the treatment of several diseases (Caruzo, 2005 and Gouveia et al., 2007). The purpose of this study was to identify relevant constituents of leaf extracts of *C. macrobothrys* and evaluate their *in vitro* antiproliferative activity against several malignant tumor cell lines.

Materials and Methods

Plant material

Leaves of *Croton macrobothrys* Baill, Euphorbiaceae were collected in April 2009, in the reserve of the Atlantic Forest of Paranapiacaba, municipality of Santo André-SP (southeast Brazil). A voucher specimen (Lima 999) was deposited in the Herbarium Maria Eneyda P. K. Fidalgo (SP) of the Institute of Botany (SEMA, São Paulo state).

Preparation of plant extracts

Material (30 g) from dried powdered leaves was extracted in Soxhlet sequentially with *n*-hexane, dichloromethane and methanol, for 6 h with each solvent. The extracts were concentrated under reduced pressure and evaporated to dryness under nitrogen flow.

GC/MS analyses

Identification of the extract constituents of was performed with 1 μ L of 2 mg/mL extracts by gas chromatography/mass spectrometry, using a system Hewlett-Packard 6890/5975B, DB-5ht fused silica capillary column (30 m x 0.32 mm x 0.10 μ m) and helium as carrier gas at 1 mL/min. The temperature program of the column started at 150 $^{\circ}$ C (1 min), rising 6 $^{\circ}$ C/min to 320 $^{\circ}$ C; injector and detector temperatures were 250 $^{\circ}$ C. Mass spectra were obtained at 70 eV and scans ranged from 40 to 450 daltons at 2 scans/s (modified from Tansakul et al., 1998). Identification of the compounds followed comparisons of mass spectra with NIST library and literature data (Ichimura et al., 1986; Zdero et al., 1992; Pizzoferrato et al. 1993; Pereira et al. 1999; Quintana et al., 2003; Souza et al., 2006).

Antiproliferative activity assays

Cell cultures

Cancer cell lines used were human 786-0 (kidney), HT-29 (colon), K562 (leukemia), NCI-ADR/RES (drug resistant ovary), NCI-H460 (lung), MCF-7 (mammary), NCI-PC-3 (prostate), OVCAR-3 (ovary), U251 (glioma), UACC-62 (melanoma) and a normal cell line VERO (kidney epithelial cells of African green monkey). Stock cell cultures were grown in medium containing RPMI 1640, supplemented with 5% of fetal bovine serum. Experimental cultures were supplemented also with peniciline:streptomcicine (10 μ g/mL:10 UI/mL).

Antiproliferative assay

Cells (100 μ L cells/well, inoculation density from $3-6 \times 10^4$ cell/mL) in 96-well plates were exposed to various sample concentrations (0.25 to 250 μ g/mL, 100 μ L/well) in DMSO/RPMI 1640 at 37 $^{\circ}$ C, 5% of CO₂ in air for 48 h. Final DMSO concentration did not affect cell viability. Cells were then fixed with 50% trichloroacetic acid and cell proliferation was determined by spectrophotometric quantification (540 nm) of cellular protein content, using the sulforhodamine B assay. Doxorubicin (DOX; 0.025-25 μ g/mL) was used as positive control. Three measurements were obtained at the beginning of incubation (time zero, T_0) and 48 h post-incubation for compound-free (C) and tested (T) cells. Cell proliferation was determined according to the equation $100 \times [(T-T_0)/C-T_0]$, for $T_0 < T \leq C$, and $100 \times [(T-T_0)/T_0]$, for $T \leq T_0$ and a concentration-response curve for each cell line was plotted using software Origin 7.5 (OriginLab Corporation) (Monks et al, 1991).

Data analysis

Using the concentration-response curve for each cell line, GI50 (concentration causing 50% growth inhibition; Shoemaker, 2006) was determined by means of non-linear regression analysis, using software Origin 7.5 (OriginLab Corporation). The average activity (mean of log GI50) of the extracts tested was also determined using MS Excel software (Fouche et al, 2008).

Results and Discussion

Retention times and mass spectra data of constituents in the extracts analyzed are given in Table 1. Only one compound (corydine, an alkaloid) was detected in the methanol extract. Major constituents of the *n*-hexane extract are the steroid β -sitosterol and the triterpenoid β -amyrin (Table 1); campesterol and squalene are other relevant constituents, while corydine, geranylgeraniol and a clerodane derivative are minor compounds of the hexane extract. The dichloromethane extract exhibited the widest diversity of constituents. Among the major constituents the acyclic diterpenoid geranylgeraniol was detected and two clerodanes were tentatively identified as crotonin derivatives (Table 1). Corydine and salutaridine (both alkaloids) are important compounds in the dichloromethane extract, while stigmasterol and β -sitosterol are minor constituents.

All compounds detected in the present work have been reported for *Croton* species. Acyclic diterpenes have been found in *C. kerrii* Airy Shaw, *C. stellatopilosus* H. Ogba and *C. sublyratus* Kurz (Sato et al., 1980; Salatino et al., 2007). Corydine and other isoquinoline alkaloids have been reported as major constituents of

C. hemiargyreus Müll. Arg. and *C. echinocarpus* Baill. (Pereira et al., 1999). Salutaridine has been reported as occurring in *C. balsamifer* Jacq. (Chambers et al., 1966) and *C. salutaris* Casar. (Barnes and Soeiro, 1981). β -Amyrin, β -sitosterol, stigmasterol and campesterol were identified in *C. betulaster* Müll. Arg., *C. hieronymi*, *C. draco* Schltdl. & Cham., *C. cajucara* Benth. and *C. urucurana* Baill. Derivatives of squalene are constituents of *C. hieronymi* Griseb. (Salatino et al., 2007).

Coherent with a wider diversity of compounds, the dichloromethane extract was shown to have higher

antiproliferative activity than the other two extracts. Toward most cell lines, GI50 values of the dichloromethane extract are substantially lower than those of the *n*-hexane and methanol extracts (Table 2). Activities were more pronounced against NCI-H460 and K562 cancer cell lines (GI50<1 μ g/mL, Table 2). Moderate activity (GI50<10 μ g/mL) of the dichloromethane extract was noted against cell lines U251, 786-0, OVCAR-3 and VERO (Table 2).

Extracts of plants from other taxa have shown to exert activity against the same cell lines used in the present work. For example, dichloromethane and

Table 1. Relevant constituents of leaf extracts of *Croton macrobothrys* and respective data of GC-MS analyses.

RT (min)	MW	Fragment ion (intensity, %)	Compound	Relative amount (%)		
				Hex	DCM	MeOH
8.9	290	290 (1), 272 (7), 257 (5), 203 (5), 187 (9), 161 (13), 147 (11), 133 (18), 119 (36), 107 (29), 93 (53), 81 (55), 69 (100)	geranylgeraniol	0.7	20.7	-
16.9	341	341 (70), 340 (28), 326 (51), 324 (31), 310 (100), 295 (9), 281 (12), 155 (21), 42 (41),	corydine	0.1	8.5	2.1
18.0	316	316 (2), 222 (27), 107 (28), 95 (63), 93 (34), 81 (98), 69 (100), 55 (29), 43 (48), 41 (43)	clerodane crotonin derivative	-	18.2	-
18.5	327	327 (100), 312 (66), 299 (36), 284 (77), 242 (24), 226 (24), 87(31), 73(49)	salutaridine	-	2.0	-
19.6	374	374 (5), 356 (12), 341 (27), 324 (62), 309 (7), 261 (28), 178 (50), 96 (100), 95 (77), 81 (46)	clerodane crotonin derivative	0.4	29.4	-
20.6	400	400 (2), 382 (47), 145 (58), 107 (54), 95 (50), 91 (53), 81 (56), 55 (59), 41 (43), 43 (100)	campesterol	2.2	-	-
21.0	412	412 (2), 354 (18), 351 (12), 300 (14), 271 (22), 255 (26), 159 (44), 105 (46), 91 (47), 83 (70), 81 (72), 79 (43), 43 (100)	stigmasterol	2.9	0.2	-
21.6	414	414 (35), 396 (52), 145 (64), 105 (64), 107 (59), 95 (56), 91 (55), 81 (63), 55 (62), 57 (55), 43 (100)	β -sitosterol	15.4	0.9	-
22.1	426	393 (1), 359 (1), 279 (2), 218 (100), 203 (34), 189 (20), 95 (34), 81 (31), 44 (30), 93 (29), 69 (29)	β -amyrin	11.0	-	-
22.9	410	395 (6), 269 (16), 229 (21), 207 (50), 189 (45), 175 (35), 107 (61), 95 (81), 81 (75), 69 (80), 55 (91), 43 (100)	squalene	2.0	-	-

RT: retention time; MW: molecular weight; DCM: dichloromethane; Hex: hexane; MeOH: methanol; -: not detected or trace amounts.

Table 2. Antiproliferative activity (GI50, μ g/mL) of leaf extracts *Croton macrobothrys* on culture cell linesa.

Material tested	Cell lines											Mean log GI50 ^c
	U251	UACC-62	MCF-7	NCI-ADR/RES	786-0	NCI-H460	PC-3	OVCAR-3	HT-29	K562	VERO	
Doxorubicin ^b	0.025	0.028	0.14	0.093	0.034	<0.025	0.052	0.12	0.033	0.054	0.66	-1.24 P
Hexane	36.53	83.32	28.09	70.21	159.60	30.71	65.21	51.41	140.08	67.87	>250	1.86 I
Dichloromethane	8.90	25.38	46.41	25.59	7.47	0.33	13.34	7.54	18.66	0.91	5.06	0.89 M
Methanol	27.75	29.04	11.66	26.93	28.28	6.08	28.26	24.47	16.31	7.45	120.05	1.33 W

^aU251: glioma; UACC-62: melanoma; MCF-7: mammary; NCI-ADR/RES: drug resistant ovary; 786-0: kidney; NCI-H460: lung; NCI-PC-3: prostate; OVCAR-3: ovary; HT-29 colon; K562: leukemia; VERO: Kidney epithelial cells of African green monkey. ^bPositive control. ^cNCI's criteria (Foucher et al, 2008): I: Mean log GI50>1.5= inactive; W: weak activity: Mean log GI50>1.10-1.5; M, moderate activity: Mean log GI50>0-1.1; P, potent activity: Mean log GI50<0.

methanol crude extracts of dried leaves of *Aspidosperma tomentosum* Mart., Apocynaceae, displayed antiproliferative activity in a concentration-dependent way against some cell lines used in the present work. The dichloromethane extract presented higher inhibition toward the lung cells (NCI460) (Kohn et al., 2006). The crude dichloromethane extract of *Virola sebifera* Aubl., Myristicaceae, was shown to be highly active, with selectivity toward NCI460 (Denny et al., 2007). The cytotoxicity of the dichloromethane crude extract, obtained from the aerial parts of *Pothomorphe umbellata* (L.) Miq., Piperaceae, was evaluated against nine human cancer cell lines and presented antiproliferative activity against all cell lines, including leukemia (K-652) (Sacoman et al., 2008).

In vitro antiproliferative activity of the latex of *Croton lechleri* was determined against leukemia K562 cells line (Rossi et al., 2003). In our experiment, dichloromethane extract of *C. macrobothrys* exhibited high selectivity against the same cell line (GI50 0.91 µg/mL, Table 2). The activity may be accounted for geranylgeraniol, the major component in the extract (Table 1), and corydine, an aporphine alkaloid. Acyclic diterpenoids, such as plaunotol from *C. sublyratus*, have been recently reported as inhibitor of angiogenesis (Kawai et al., 2005; Yoshikawa et al., 2009). Corydine has shown inhibitory activity against several mouse tumor cell lines such as leukemia P388 and L1210, melanoma B16, bladder cancer MBC2, and colon cancer Colon 26 (Kondo et al., 1990; Bruneton, 1999). Other alkaloids from *Croton* species have been shown to be active against cancer; examples are taspine (from *C. lechleri*) and julocrotol, isojulocrotol and julocrotone, from *C. cuneatus* Klotzsch (Salatino et al., 2007). Also clerodanes may be involved in the observed activity: the furoclerodane croblongifolin from *C. oblongifolius* Sieber ex Spreng. was shown to be cytotoxic (Roengsumran et al., 2002) and trans-dehydrocrotonin from *C. cajucara* exhibited anti-tumor effect (Melo et al., 2004). The next step in our investigation is to test antiproliferative activity of fractions and isolated compounds from dichloromethane extracts, obtained from column chromatography.

The use of material from *C. macrobothrys* in popular medicine or phytotherapy requires evaluation of possible undesirable side effects. For example, 25 cases of hepatotoxicity were documented among people from Amazonia, which were ascribed to consumption of sacaca (*Croton cajucara*) along 36 months (Soares, 2006). Twenty-one cases corresponded to acute, three to chronic and one to fulminating hepatitis. This issue requires further investigation, because a recent study did not detect significant alterations on hepatic transferases in animals treated with *C. cajucara* (Rodrigues et al., 2010). Nothing is known about toxic effects of *C.*

macrobothrys.

The results of the present work, suggesting that leaves of *C. macrobothrys* may contain antiproliferative active compounds, are in agreement with previous evidences, which have shown that *Croton* species are likely sources of substances useful for the development of new drugs (Salatino et al. 2007).

Acknowledgments

The authors are grateful to Frederick Cancer Research & Development Center of the National Cancer Institute (Frederick, MA, USA) for the kind provision of cell lines, and to Fundação de Amparo à Pesquisa do Estado de São Paulo (08/10595-3 auxílio) (08/09942-0 bolsa), CAPES and CNPq for financial support.

References

- Barnes RA, Soeiro OM 1981. The alkaloids of *Croton salutaris*. *Phytochemistry* 20: 543-544.
- Block S, Stevigny C, De Pauw-Gillet MC, De Hoffman E, Llabrés G, Adjakidjé V, Quertin-Leclercq J 2002. ent-Trachyloban-3β-ol, a new cytotoxic diterpene from *Croton zambesicus*. *Planta Med* 68: 647-648.
- Bruneton J 1999. *Pharmacognosie: Phytochimie Plantes médicinales*, 2ème Edition, Lavoisier Tec & Doc: Paris.
- Chambers C, Haynes LJ, Stuart KL 1966. Norsinoacutine and salutaridine isolated from *Croton balsamifera* Jacq. *Chem Commun* 14: 449.
- Caruzo MBR 2005. *Estudo taxonômico e biogeográfico do gênero Croton L. (Euphorbiaceae) no Estado de São Paulo, Brasil*. Dissertação de Mestrado, Instituto de Biociências USP, São Paulo.
- Catalán CAN, Heluani CS, Kotowicz C, Gedris TE, Herz W 2003. A linear sesterterpene, two squalene derivatives and two peptide derivatives from *Croton hieronymi*. *Phytochemistry* 64: 625-629.
- Cragg GM, Newman DJ 2009. Nature: a vital source of leads for anticancer drug development. *Phytochem Rev* 8: 313-331.
- Carvalhoes SF, Costa DL, Mazzei JL, Taddei LEM, d'Avila LA 2002. Alternative extraction of alkaloid anticarcinogens from Brazilian "vinca rosea" using ion exchange chromatography. *Rev Bras Farmacogn* 12: 83-84.
- Denny C, Zacharias ME, Kohn LK, Foglio MA, Carvalho JE 2007. Atividade antiproliferativa dos extratos e da fração orgânica obtidos das folhas de *Virola sebifera* Aubl. (Myristicaceae). *Rev Bras Farmacogn* 17: 598-603.
- Fouche G, Cragg GM, Pillay P, Kolesnikova N, Maharaj VJ, Senabe J 2008. *In vitro* anticancer screening of South African plants. *J Ethnopharmacol* 119: 455-461.
- Gouveia TC, Florsheim SMB, Pastore OT, Lima IL 2007.

- Morfologia, anatomia do lenho e densidade básica de *Croton floribundus* Spreng e *Croton macrobothrys* Baillon. *IF Série Registros* 31: 45-49.
- Gupta R, Gabrielsen B, Ferguson SM 2005. Nature's medicine: Traditional knowledge and intellectual property management. Case studies from the National Institutes of Health (NIH), USA. *Curr Drug Discov Technol* 2: 203-219.
- Ichimura K, Yamanakab H, Chibab K, Shinozucac T, Shikid Y, Saitob K, Kusanod, S, Ameniya S, Oyamac K, Nozakic Y, Katoc K 1986. Simultaneous quantitative measurement of fourteen adrenal steroids by capillary column gas chromatography-mass spectrometry, and its clinical application. *J Chromatogr* 374: 5-16.
- Johnson IS, Armstrong JG, Gorman M 1963. The vinca alkaloids: a new class of oncolytic agents. *Cancer Res* 23: 1390-1397.
- Jones K 2003. Review of Sangre de Drago (*Croton lechleri*) - A South American tree sap in the treatment of diarrhea, inflammation, insect bites, viral infections, and wounds: traditional uses to clinical research. *J Altern Complem Med* 9: 877-896.
- Kawai K, Tsuno NH, Kitayama J, Okaji Y, Yazawa K, Asakage M, Yamashita H, Watanabe T, Takahashi K, Nagawa H 2005. Anti-angiogenic properties of plaunotol. *Anticancer Drugs* 16: 401-407.
- Kohn LK, Pizão PE, Foglio MA, Antônio MA, Amaral MCE, Bittric V, Carvalho JE 2006. Antiproliferative activity of crude extract and fractions obtained from *Aspidosperma tomentosum* Mart. *Rev Bras Pl Med* 8: 110-115.
- Kondo Y, Imai Y, Hojo H, Endo T, Nozoe S 1990. Suppression of tumor cell growth and mitogen response by aporphine alkaloids, dicentrine, glaucine, corydine and apomorphine. *J Pharmacobio-dynam* 13: 426-431.
- Liu Y, Wang MW 2008. Botanical drugs: Challenges and opportunities. Contribution to Linnaeus Memorial Symposium 2007. *Life Sci* 82: 445-449.
- Maxwell N 1990. *Witch Doctor's Apprentice: Hunting for Medicinal Plants in the Amazon*, 3rd Edition, Citadel Press: New York, NY.
- Melo PS, Justo G Z, Durán N, Haun M 2004. Natural killer cell activity and anti-tumour effects of dehydrocrotonin and its synthetic derivatives. *Eur J Pharmacol* 487: 47-54.
- Monks A, Scudeiro D, Skehan P, Shoemaker R, Paull K, Vistica D, Hose C, Langley J, Cronise P, Vaigro-Wolff A, Gray-Goodrich M, Campbell H, Mayo J, Boyd M 1991. Feasibility of a high-flux anticancer drug screen using a diverse panel of cultured human tumor cell lines. *J Natl Cancer I* 83: 757-766.
- Pereira AS, Amaral ACF, Barnes RA, Cardoso FR, Aquino Neto FR 1999. Identification of Isoquinoline alkaloids in crude extracts by high temperature gas chromatography-mass spectrometry. *Phytochem Analysis* 10: 254-258.
- Pizzoferrato L, Nicoli S, Lintas C 1993. GC-MS characterization and quantification of sterols and cholesterol oxidation products. *Chromatographia* 3: 269-274.
- Perazzo F, Carvalho JCT, Rodrigues M, Morais EKL, Maciel MAM 2007. Comparative anti-inflammatory and antinociceptive effects of terpenoids and an aqueous extract obtained from *Croton cajucara* Benth. *Rev Bras Farmacogn* 17: 521-528.
- Quintana A, Reinhard J, Faure R, Uva P, Bagneres AE, Massiot G, Clément J 2003. Interspecific variation in terpenoid composition of defensive secretions of european Reticulitermes Termites. *J Chem Ecol* 29: 639-652.
- Riina R, Berry PE, Van E BW 2009. Molecular phylogenetics of the dragon's blood *Croton* section *Cyclostigma* (Euphorbiaceae), a polyphyletic assemblage unraveled. *Syst Bot* 34: 360-374.
- Rodrigues G, Marcolin E, Bona S, Porawski M, Lehmann M, Marroni NP 2010. Effects of *Croton cajucara* Benth (sacaca) in diabetic rats. *Arq Gastroenterol* 47: 301-305.
- Roengsumran S, Musikul K, Petsom A, Vilaivan T, Sangvanich P, Pompakakul S, Puthong S, Chaichantipyuth C, Jaiboon N, Chaichit N 2002. Croblongifolin, a new anticancer clerodane from *Croton oblongifolius*. *Planta Med* 68: 274-277.
- Rossi D, Bruni R, Bianchi N, Chiarabelli C, Gambari R, Medici A, Lista A, Paganetto G 2003. Evaluation of the mutagenic, antimutagenic and antiproliferative. *Phytomedicine* 10: 139-144.
- Sacoman JL, Monteiro KM, Possenti A, Figueira GM, Foglio MA, Carvalho JE 2008. Cytotoxicity and antitumoral activity of dichloromethane extract and its fractions from *Pothomorphe umbellata*. *Braz J Med Biol Res* 41: 411-415.
- Salatino A, Salatino MLF, Negri G 2007. Traditional uses, Chemistry and Pharmacology of *Croton* species (Euphorbiaceae). *J Brazil Chem Soc* 18: 11-33.
- Sato A, Ogiso A and Kuwano H 1980. Acyclic diterpenes from *Croton kerrii*. *Phytochemistry* 19: 2207-2209.
- Shoemaker RH 2006. The NCI60 human tumour cell line anticancer drug screen. *Nat Rev Cancer* 6: 813-826.
- Soares MC 2004. Would sacaca, *Croton cajucara* Benth (Euphorbiaceae) be an hepatotoxic plant like Germander, *Teucrium chamaedrys* L. (Labiatae) *Rev Soc Bras Med Trop* 37: 96-97.
- Souza MAA, Souza SR, Veiga Junior VF, Cortez JKPC, Leal RS, Dantas TNC, Maciel MA 2006. Composição química do óleo fixo de *Croton cajucara* e determinação das suas propriedades fungicidas. *Rev Bras Farmacogn* 16: 599-610.
- Tansakul T, De-Eknamkul W 1998. Geranylgeranyol-18-hidroxyase: the last enzyme on the plaunotol biosynthetic pathway in *Croton sublyratus*. *Phytochemistry* 47: 1241-1247.
- Tan G, Gyllenhaal C, Sorjarto DD 2006. Biodiversity as a source of anticancer drugs. *Curr Drug Targets* 7: 265-277.
- Wani MC, Taylor HL, Wall ME 1971. Plant antitumor agents.

The isolation and structure of taxol, a novel leukemic and antitumor agent from *Taxus brevifolia*. *J Am Chem Soc* 93: 2325-2327.

Yoshikawa N, Yamada J, Tsuno NH, Okaji Y, Kawai K, Tsuchiya T, Satomi Yoneyama S, Tanaka J, Shuno Y, Nishikawa T, Nagawa H, Oshima N, Takahashi K 2009. Plaunotol and Geranylgeraniol Induce Caspase-Mediated Apoptosis in Colon Cancer. *J Surg Res* 153: 246-253.

Zdero C, Bohlmann F, King RM1992. Clerodane derivatives from *Diplostegium*. *Phytochemistry* 31: 213-216.

***Correspondence**

Antonio Salatino
Instituto de Biociências, Universidade de São Paulo, São Paulo
Rua do Matão 277, Cidade Universitária, 05508-090
asalatin@ib.usp.br
Tel.: +55 11 3091 7546; 3091 7416