



Universidade de São Paulo

Biblioteca Digital da Produção Intelectual - BDPI

Departamento de Alimentos e Nutrição Experimental - FCF/FBA Artigos e Materiais de Revistas Científicas - FCF/FBA

2012

INFLUENZA VIRUSES IN ADULT DOGS RAISED IN RURAL AND URBAN AREAS IN THE STATE OF SAO PAULO, BRAZIL

REVISTA DO INSTITUTO DE MEDICINA TROPICAL DE SAO PAULO, SAO PAULO, v. 54, n. 6, p.
311-314, NOV-DEC, 2012

<http://www.producao.usp.br/handle/BDPI/41473>

Downloaded from: Biblioteca Digital da Produção Intelectual - BDPI, Universidade de São Paulo

INFLUENZA VIRUSES IN ADULT DOGS RAISED IN RURAL AND URBAN AREAS IN THE STATE OF SÃO PAULO, BRAZIL

Dalva Assunção Portari MANCINI(1), Rita Maria Zucattelli MENDONÇA(1), Aparecida Santo Pietro PEREIRA(1), Adélia Hiroko Nagamori KAWAMOTO(1), Camila Infantsi VANNUCCHI(2), José Ricardo PINTO(1), Enio MORI(3) & Jorge MANCINI FILHO(4)

SUMMARY

In 1970, searching for the interspecies transmission of influenza viruses led to the first study on influenza viruses in domestic animals. Birds and mammals, including human beings, are their natural hosts; however, other animals may also play a role in the virus epidemiology. The objective was to investigate the incidence of influenza viruses in adult dogs raised in rural (9, 19.56%) and urban (37, 80.43%) areas in the state of São Paulo, Brazil. Dog serum samples were examined for antibodies to influenza viruses by the hemagglutination inhibition (HI) test using the corresponding antigens from the circulating viruses in Brazil. Dogs from rural areas presented antibodies to influenza A H₃N₂, and influenza A H₇N₇ and H₃N₈. In rural areas, dog sera displayed mean titers as 94.37, 227.88, 168.14, 189.62 HIU/25 µL for subtypes H₁N₁, H₃N₂, H₇N₇, H₃N₈, respectively. About 84% and 92% of dogs from urban areas exhibited antibodies to human influenza A H₁N₁ and H₃N₂, respectively, with statistical difference at $p < 0.05$ between the mean titers of antibodies to H₁N₁ and H₃N₂. About 92% and 100% were positive for H₇N₇ and H₃N₈, respectively. In dogs from urban areas, the mean titers of antibodies against influenza A H₁N₁, H₃N₂, H₇N₇ and H₃N₈, were 213.96, 179.42, 231.76, 231.35 HIU/25 µL respectively. The difference among them was not statistically significant at $p > 0.05$. In conclusion, these dogs were positive for both human and equine influenza viruses. The present study suggests the first evidence that influenza viruses circulate among dogs in Brazil.

KEYWORDS: Influenza A; Dogs; Serology; Epidemiology.

INTRODUCTION

Influenza is an acute and highly contagious viral disease that afflicts the respiratory tract and is transmitted through contact with nasopharyngeal secretions. It is known to be caused by RNA viruses of the *Orthomyxoviridae* family. In their lipoprotein envelope are two glycoproteins, hemagglutinin (H) and neuraminidase (N), whose structures differ among virus strains. Sixteen hemagglutinin (H1 - H16) and nine neuraminidase (N1 - N9) subtypes have been identified in influenza A viruses. Influenza A viruses can infect both humans and animals, but influenza B viruses only infect humans^{1,17}.

Hemagglutinin (H) and neuraminidase (N) undergo considerable antigenic variation in influenza A and B viruses, whereas these same proteins display minor alterations in types B and C. Such changes lead to the emergence of new strains. Influenza incidence peaks during winter and can be very recurrent. The main symptoms are: high temperature (± 38 °C), sneezing- and coughing, which can evolve into pneumonia in both humans and animals. Influenza types A and B are responsible for seasonal flu epidemics. Type C also causes the disease; however, the symptoms are milder¹⁰.

Influenza virus receptors on host cells differ among mammalian species, viruses bind to receptors on tracheal epithelial cells, while duck viruses attach to the cells of the intestine. Avian (H₁, H₃, H₉, H₁₁) and equine (H₃ and H₇) strains have receptor-binding specificity for acetylneuraminic acid with an alpha 2,3-N-linkage to galactose (α -2,3 NeuAcGal). Human strains (H₁ and H₃), on the other hand, have specificity for the alpha 2,6-linkage. Pigs present both receptors, which makes them susceptible to human and avian influenza viruses. They are considered to be intermediate hosts for the adaptation of avian influenza viruses to humans and "mixing vessels" for novel combinations between avian and human influenza viruses, which can result in the emergence of new pandemic strains. The resulting new viruses may then be able to infect horses and humans¹³. The proteolytic cleavage of the viral receptor proteins is also a very important activation mechanism in the pathogenesis of influenza viruses^{18,20}.

The host range of influenza viruses has been widely studied in order to characterize the role of some animals in the transmission chain - once considered unlikely to infect others. The susceptibility of domestic animals to an influenza type A virus was first reported in 1972¹². Recently, influenza viruses were isolated from dogs with respiratory

(1) Laboratório de Virologia, Divisão de Desenvolvimento Científico, Instituto Butantan, São Paulo, SP, Brazil.

(2) Clínica Médica Veterinária, Departamento de Reprodução Animal, Faculdade de Medicina Veterinária e Zootecnia, USP, São Paulo, SP, Brazil.

(3) Departamento de Medicina Veterinária Preventiva e Saúde Animal – Laboratório de Biologia Molecular Aplicada e Sorologia, Faculdade de Medicina Veterinária e Zootecnia, USP, São Paulo, SP, Brazil.

(4) Laboratório de Lípidos, Departamento de Bioquímica do Alimento, Faculdade de Farmácia, USP, São Paulo, SP, Brazil.

Correspondence to: Dalva Assunção Portari Mancini, Laboratório de Virologia, Divisão de Desenvolvimento Científico, Instituto Butantan, Av. Vital Brasil 1500, 05503-900 São Paulo, SP, Brasil. Tel.: 55 11 3726-7222 Ext. 2152. E-mail: dapmancini@butantan.gov.br

infections. Among the isolated viruses, a specific strain, named Influenza A/canine/Florida/43/2004, was observed to be very closely related to the equine influenza A subtype H₃N₈. This monophyletic origin of US canine influenza viruses of subtype H₃N₈, which have circulated among dogs since the early 1990s, very likely indicates a single trans-species transmission event directly from horse to dog, possibly facilitated by direct contact in a racing environment^{3,15}.

Dog-to-dog contact transmission of avian-origin influenza A viruses subtypes H₃N₂ and H₃N₈ has been observed in a recent experimental study of Beagle dogs. The dogs showed clinical signs after 14 days post-inoculation, confirming the interspecies transmission of the influenza virus¹⁶. Domestic animals (cats and dogs) may also enable the adaptation of avian influenza viruses to mammals, especially carnivores that feed on infected birds⁵.

Recently, HEINEMANN *et al.*⁶ reported serologic evidence of the equine influenza virus A subtype H₇N₇ circulation in some regions of north Brazil.

Reptilian and crocodylian species, as well as domestic animals (cats and dogs), can be potential hosts of influenza viruses. Recently, influenza type A viruses were detected in crocodylians from two locations in Florida (USA)².

In one of our previous studies, poisonous snakes (heterothermic animals) showed seropositivity for human influenza A viruses subtypes H₁N₁ and H₃N₂⁹.

To date, no studies on influenza viruses in dogs have been carried out in Brazil.

OBJECTIVE

This work was fueled by concerns about the dog-to-human contact transmission of influenza viruses. Therefore this work aimed to study the role of adult dogs from rural and urban areas in the state of São Paulo, Brazil, in the epidemiology of the influenza virus infections. Considering the originality of the present study, the hope is that it could contribute towards the research and the knowledge of this study field.

MATERIALS AND METHODS

Animals: Forty-six healthy dogs, assisted at the clinic of the Faculty

of Veterinary Medicine at the University of São Paulo - Brazil, were grouped according to whether they were raised in nine rural or 37 urban areas in the state of São Paulo. The protocol for animal studies, CEEA number 2230/2011, was approved by the Animal Experimentation Ethics Committee of the School of Veterinary and Animal Sciences of the University of São Paulo.

Serum: Blood serum samples were taken, inactivated by heat treatment at 56 °C for 30 min and then treated with Kaolin (20%) and erythrocytes (50%) according to MANCINI *et al.*⁸, DESHPANDE *et al.*⁴ and OIEN *et al.*¹¹.

Serology: The hemagglutination inhibition (HI) test was performed according to MANCINI *et al.*⁸. Four HA units in 25 µL of viruses were used. Titers were expressed as hemagglutination inhibition units per 25 µL of sera (HIU/25 µL).

Antigens: Antigens of the influenza viruses as A/SP/2/95 = A/Beijing/353/89 (H₃N₂), A/SP/1/91= A/Singapore/6/86 (H₁N₁), A/Eq1/SP/56 (A Eq1 H₇N₇) and A/Eq2/1/SP/85 (A Eq2 H₃N₈) were used. Influenza virus samples were grown in MDCK (Madin Darby canine kidney) cell cultures according to MANCINI *et al.*⁸.

RESULTS

The serological analysis was obtained through the hemagglutination inhibition (HI) test which showed that 15.27, 6.52 and 6.52% of the dogs exhibited low antibody titers (≤ 40 HIU/25 µL) to influenza A viruses subtypes H₁N₁, H₃N₂ and H₇N₇, suggesting no protective response against these subtypes. The other animals tested presented higher antibody titers, ranging from 40 to ≥ 320 HIU/25 µL (Table 1).

Many serologic methods such as the neutralization, Elisa, could be used to search the antibodies to influenza virus. But the hemagglutination inhibition test is usually the choice for investigating the influenza virus^{13,14}.

As seen in Figure 1, the average number of protective antibody titers (≥ 40 HIU/25 µL) to human H₁N₁ and H₃N₂ and equine influenza H₇N₇, H₃N₈, in rural areas were 94.37, 227.88, 168.14, 189.62 HIU/25 µL, respectively. In this case, the difference (*p* < 0.05) among mean titers for human influenza A subtypes H₁N₁ and H₃N₂ was statistically significant.

Figure 2 shows the average of protective antibody titers (≥ 40HIU/25 µL) against human and equine influenza viruses in dogs

Table 1
Analysis of antibody titers for influenza A virus subtypes in dog sera from urban and rural areas. Values expressed as HIU/25µL

Virus Subtypes	HIU/25 µL					Total
	< 40	40/80	> 80/160	> 160/320	> 320	
H ₁ N ₁	07(15.217%)	07(15.271%)	16(34.782%)	09(19.565%)	07(15.217%)	(100%)
H ₃ N ₂	03(6.521%)	13(28.260%)	08(17.391%)	15(32.608%)	07(15.217%)	(100%)
H ₃ N ₈	-	08(17.391%)	14(30.434%)	17(36.956%)	07(15.217%)	(100%)
H ₇ N ₇	03(6.521%)	08(17.391%)	10(21.739%)	22(47.826%)	03(6.521%)	(100%)

(HIU = Hemagglutination inhibition units).





Fig. 1 - Analysis of mean titers of protective antibodies (≥ 40 HIU/25 μ L) for human and equine influenza A virus subtypes in dogs from rural areas.



Statistical difference is not significant $P > 0.05$

Fig. 2 - Analysis of mean titers of protective antibodies (≥ 40 HIU/25 μ L) for human and equine influenza A virus subtypes in dogs from urban areas.

from urban areas. Mean titers of 213.96, 179.42, 231.76 and 231.35 HIU/25 μ L were observed for human influenza A subtypes H₁N₁ and H₃N₂ and equine influenza A subtypes H₇N₇ and H₃N₈ viruses, respectively. No statistically significant difference among mean titers was observed ($p > 0.05$). When dogs from urban and rural areas were compared for the presence of protective antibody titers (≥ 40 HIU/25 μ L) against human and equine influenza A subtypes H₁N₁, H₃N₂, H₇N₇ and H₃N₈, no statistically significant difference ($p > 0.05$) was observed (Fig 3).




Fig. 3 - Comparative analysis of percentages of dogs from urban versus rural areas exhibiting protective antibody titers (≥ 40 HIU/25 μ L) for human and equine influenza A virus subtypes.

DISCUSSION

Nine (19.56%) and 37 (80.43%) dogs from rural and urban areas, respectively, exhibited high mean titers of antibodies (≥ 40 HIU/25 μ L) against influenza A viruses subtypes H₁N₁, H₃N₂, H₇N₇ and H₃N₈, suggesting that these dogs had been in contact with human and equine influenza viruses. Our results are in agreement with a recent serological study performed via the hemagglutination inhibition test on pet dogs participating in a flyball tournament in Pennsylvania (PA, USA), which showed positive results for antibodies against an influenza A virus subtype H₃N₈¹⁴.

Healthy dogs, kept in close contact with equine influenza virus (EIV)-infected horses, were reported as developing antibodies against the virus, which suggests that interspecies transmission of EIV to dogs could have occurred as result of contact between these animal species¹⁹.

Similar observations were also reported for dogs in Japan. Their serum samples responded positively to human and equine influenza A viruses subtype H₃ in the hemagglutination inhibition test (HI)¹³.

A careful analysis of acute and convalescent dog serum samples, history and clinical signs are important in diagnosing canine influenza. In a study on the etiology of the disease, its causative agent, a type A virus, was found to be closely related to the equine influenza A virus subtype H₃N₈⁷.

Results from this study and the literature suggest evidence that dogs can harbor the influenza viruses and may play a role in the interspecies transmission and epidemiology of the virus. Furthermore, recent studies on the distribution of receptor molecules in the canine respiratory tract detected that SA α 2,3Gal was the dominant receptor, although SA α 2,6Gal could be detected in the trachea, bronchi and bronchioles. These reports are consistent with the hypothesis that the sero-positive dogs found in this study were infected with H₃N₂ influenza A virus originating from human sources¹³.

The high percentage (100%) of the protective antibodies to influenza virus detected in all evaluated dogs from rural and urban areas suggests that animals have had several contacts with different virus subtypes, during their lives. The subtype H₇N₇, considered the original avian strain revealed to be present in the dogs population. Also, HEINEMANN *et al.* (2009) reported that the influenza A, subtype H₇N₇, was present in equine populations raised in north from Brazil⁶.

It's also possible to conclude that influenza viruses could be circulating among dogs in many parts of the world, including Brazil.

RESUMO

Incidência do vírus influenza em cães adultos criados em áreas rural e urbana do estado de São Paulo, Brasil

A transmissão interespecífica do vírus influenza é relatada em estudo sobre influenza com animais domésticos desde 1970. Pássaros e mamíferos, incluindo o homem, são seus hospedeiros naturais, porém outros animais podem participar da sua epidemiologia. Foi investigada a incidência do vírus influenza em cães adultos criados em zonas rural

(9, 19,56%) e urbana (37, 80,43%), do Estado de São Paulo. Os soros dos cães foram examinados pelo teste de inibição da hemaglutinação (IH), usando antígeno dos vírus influenza circulantes no Brasil. Nos cães rurais foram detectados títulos médios de 94,37, 227,88, 168,14 e 189,62 UIH/25 µL (unidades inibidoras de hemaglutinação/25 µL) para os subtipos H₁N₁, H₃N₂, H₇N₇, H₃N₈ de vírus influenza A, respectivamente, com diferenças estatisticamente significativas (p<0,05) entre as médias de títulos de anticorpos contra H₁N₁ e H₃N₂. Cerca de 84% e 92% dos cães urbanos responderam aos vírus influenza A humano H₁N₁ e H₃N₂, respectivamente e destes 92% e 100% foram positivos para os vírus eqüinos H₇N₇ e H₃N₈, respectivamente. Para esses cães as médias de títulos de anticorpos para os vírus influenza A H₁N₁, H₃N₂, H₇N₇ e H₃N₈ foram 213,96, 179,42, 231,76 e 231,35UIH/25 µL, respectivamente. As diferenças entre as médias não foram estatisticamente significativas (p>0,05). Conclui-se que os cães apresentaram positividade para ambos vírus influenza humano e equino. O presente estudo sugere, pela primeira vez, evidências de que há circulação do vírus influenza em cães, no Brasil.

ACKNOWLEDGEMENTS

The authors thank Conselho Nacional de Desenvolvimento Científico e Tecnológico for their financial support(CNPq Project 471876/2009/2010).

REFERENCES

1. Alexander DJ, Brown IH. Recent zoonoses caused by influenza A viruses. *Rev Sci Tech*. 2000;19:197-225.
2. Davis LM, Spackman E. Do crocodylians get the flu? Looking for influenza A in captive crocodylians. *J Exp Zool A Ecol Genet Physiol*. 2008;309:571-80.
3. Crawford PC, Dubovi EJ, Castleman WL, Stephenson I, Gibbs EP, Chen I, *et al*. Transmission of equine influenza virus to dogs. *Science*. 2005;310:482-5.
4. Deshpande M, Abdelmagid O, Tubbs A, Jayappa H, Wasmoen T. Experimental reproduction of canine influenza virus H3N8 infection in young puppies. *Vet Ther*. 2009;10:29-39.
5. Harder TC, Wahlenkamp TW. Influenza virus infections in dogs and cats. *Vet Immunol Immunopathol*. 2010;134:54-60.
6. Heinemann MB, Cortez A, Lara MCCSH, Cunha EMS, Nassar AFC, Villalobos EMC, *et al*. Soroprevalência do vírus da influenza equina no Município de Uruará, PA, Brasil, Amazônia Oriental. *Arq Inst Biol*. 2009;76:697-9.
7. Hilling K, Hanel R. Canine influenza. *Compend Contin Educ Vet*. 2010;32(6):E1-9.
8. Mancini DAP, Mendonça RMZ, Cianciarullo AM, Kobashi LS, Trindade HG, Fernandes W, *et al*. Influenza em animais heterotérmicos. *Rev Soc Bras Med Trop*. 2004;37:204-9.
9. Mancini DAP, Mendonça RMZ, Kawamoto AHN, Giorgetti M, Trindade HG, Fernandes W, *et al*. Identification and characterization of influenza virus isolated from Brazilian snakes. In: Méndes-Villas A, editor. *Communicating current research and educational topics and trends in applied microbiology*. Badajoz: Formatex Microbiology Series No. 1; 2007. p. 868-74.
10. Neumann G, Kawaoka Y. Host range restriction and pathogenicity in the context of influenza pandemic. *Emerg Infect Dis*. 2006;12:881-6.
11. Oien N, Mattern S, Brozowski J, Teel J, Salmon S. Cross-reactivity to field isolates of canine influenza virus by a killed canine influenza virus (H3N8, Iowa05) vaccine. *Int J Appl Res Vet Med*. 2012;10:14-8.
12. Paniker CKJ, Nair CMG. Experimental infection of animals with influenza virus types A and B. *Bull World Health Org*. 1972;47:461-3.
13. Said AW, Usui T, Shinya K, Ono E, Ito T, Hikasa Y, *et al*. A sero-survey of subtype H3 influenza A virus infection in dogs and cats in Japan. *J Vet Med Sci*. 2011;73:541-4.
14. Serra VF, Stanzani G, Smith G, Otto CM. Point seroprevalence of canine influenza virus H3N8 in dogs participating in a flyball tournament in Pennsylvania. *J Am Vet Med Assoc*. 2011;238:726-30.
15. Song D, Lee C, Kang B, Jung K, Oh T, Kim H, *et al*. Experimental infection of dogs with avian-origin canine influenza A virus (H3N2). *Emerg Infect Dis*. 2009;15:56-8.
16. Song D, Moon H, Jung K, Yeom M, Kim H, Hans S, *et al*. Association between nasal shedding and fever that influenza A (H3N2) induces in dogs. *Virol J*. 2011;8:1.
17. Taubenberger JK, Morens DM. Influenza: the once and future pandemic. *Public Health Rep*. 2010;125(Suppl 3):16-26.
18. Van Riel D, Munster VJ, de Wit E, Rimmelzwaan GF, Fouchir RA, Osterhaus AD, *et al*. Human and avian influenza viruses target different cells in the lower respiratory tract of humans and other mammals. *Am J Pathol*. 2007;171:1215-23.
19. Yamanaka T, Nemoto M, Tsujimura K, Kondo T, Matsumura T. Interspecies transmission of equine influenza virus (H3N8) to dogs by close contact with experimentally infected horses. *Vet Microbiol*. 2009;139(3-4):351-5.
20. Webster RG, Shortridge KF, Kawaoka Y. Influenza: interspecies transmission and emergence of new pandemics. *FEMS Immunol Med Microbiol*. 1997;18:275-9.

Received: 12 December 2011

Accepted: 15 June 2012