



Universidade de São Paulo

Biblioteca Digital da Produção Intelectual - BDPI

Departamento de Gastroenterologia - FM/MGT

Artigos e Materiais de Revistas Científicas - FM/MGT

2012

Phylogenetic analysis of complete genome sequences of hepatitis B virus from an Afro-Colombian community: presence of HBV F3/A1 recombinant strain

VIROLOGY JOURNAL, LONDON, v. 9, OCT 24, 2012
<http://www.producao.usp.br/handle/BDPI/42594>

Downloaded from: Biblioteca Digital da Produção Intelectual - BDPI, Universidade de São Paulo

RESEARCH

Open Access

Phylogenetic analysis of complete genome sequences of hepatitis B virus from an Afro-Colombian community: presence of HBV F3/A1 recombinant strain

Mónica V Alvarado-Mora^{1*}, Camila M Romano², Michele S Gomes-Gouvêa¹, Maria F Gutierrez³, Flair J Carrilho¹ and João R R Pinho¹

Abstract

Background: Hepatitis B virus (HBV) infection is one of the most prevalent viral infections in humans and represents a serious public health problem. In Colombia, our group reported recently the presence of subgenotypes F3, A2 and genotype G in Bogotá. The aim of this study was to characterize the HBV genotypes circulating in Quibdó, the largest Afro-descendant community in Colombia. Sixty HBsAg-positive samples were studied. A fragment of 1306 bp (S/POL) was amplified by nested PCR. Positive samples to S/POL fragment were submitted to PCR amplification of the HBV complete genome.

Findings: The distribution of HBV genotypes was: A1 (52.17%), E (39.13%), D3 (4.3%) and F3/A1 (4.3%). An HBV recombinant strain subgenotype F3/A1 was found for the first time.

Conclusions: This study is the first analysis of complete HBV genome sequences from Afro-Colombian population. It was found an important presence of HBV/A1 and HBV/E genotypes. A new recombinant strain of HBV genotype F3/A1 was reported in this population. This fact may be correlated with the introduction of these genotypes in the times of slavery.

Keywords: Hepatitis B virus, Subgenotype A1, Genotype E, Subgenotype D3, Bayesian analysis, Recombinant subgenotype F3/A1, Colombia

Introduction

Hepatitis B virus (HBV) infection is a relevant global health problem with 2 billion people that have been infected worldwide, including 350 million of them suffering from chronic HBV infection [1]. In Latin America, the estimated HBsAg seroprevalence ranges from 0.5% to 3.0%, with the total number of HBsAg carriers approaching 11 millions [2]. The highest prevalence surpassing 8.0 % is found among the native populations of the western Amazon basin, which includes Brazil [3], Colombia [4], Peru [5] and Venezuela [6].

HBV genome is a partially double-stranded circular DNA molecule of approximately 3,200 bp that encodes four overlapping open reading frames (ORFs) [7]. A genetic classification based on the comparison of complete HBV genomes has identified nine genotypes, A through I [8], that differ by at least 8% at nucleotide level from each other. Genotype A was initially identified in southern Africa [9]. Phylogenetic analysis of the complete genomes of subgenotype A1 isolates classified it in two clusters (African and Asian) [10]. The introduction of subgenotype A1 into Asia could be the result of movements along the East coast of Africa, from Somalia in the horn of Africa to the Arabic Peninsula in Asia [10]. Subgenotype A2, also denoted Ae, from "European" subgenotype, was isolated from South African carriers and is also found in Northern Europe and Greenland [11].

* Correspondence: monica.viviana@usp.br

¹Laboratory of Gastroenterology and Hepatology, São Paulo Institute of Tropical Medicine and Department of Gastroenterology, School of Medicine, University of São Paulo, São Paulo, Brazil

Full list of author information is available at the end of the article

Subsequently, subgenotype A3 was characterized in Cameroon, Mali and Gambia [12]. Complete genomes of two HBV isolates from Mali (subgenotype A4) have been found to diverge by 6% from A3. Finally, four Nigerian genotype A isolates formed another group, tentatively designated subgenotype A5 [13]. More recently, subgenotypes A6 was detected in African-Belgian patients [14] and subgenotype A7 was reported in Cameroon [15].

Genotypes B and C are predominant in East and Southeast Asia. Subgenotype D1 occurs mostly in the Mediterranean basin and Middle East. D2 has been reported in India, Japan, Europe and the United States. D3 was found in South Africa, Brazil, Rwanda, India, Costa Rica, Iran, Serbia and the United States. Finally, D4 was reported in Australia, South Africa, Somalia, Rwanda and Oceania [11].

Genotype E was first described in West Africa in high prevalence but in a surprisingly low diversity, as the mean diversity over the whole genome is 1.75%. HBV/E is the most prevalent genotype in western and central Africa [10]. In Quibdó, Colombia we previously reported for the first time the presence of this genotype in nine cases [16]. Genotypes F and H are found in populations from Alaska to Central and South America [17,18].

The aim of the present study was to characterize the HBV genotypes circulating in Quibdó city, Colombia. We report for the first time the HBV complete genome sequences from Afro-Colombian infected people and inferred their origin using phylogenetic analyses approaches.

Materials and methods

Study population

To evaluate HBV genotypes distribution in Quibdó, the largest Afro-descendant community in Colombia, 60 positive samples for the Hepatitis B virus surface antigen (HBsAg) were obtained from sera stored at -20°C in a public health laboratory in Quibdó, Colombia in 2007. This protocol was approved by the Ethical Committees from Pontificia Universidad Javeriana, Bogotá Colombia and School of Medicine, University of São Paulo, São Paulo, Brazil. Before

HBV DNA extraction

HBV DNA extraction was carried out from 100 μL serum using the acid guanidinium thiocyanate/phenol/chloroform method [19]. Briefly, 300 μL GT solution was added to each sample. Ice-cold chloroform (50 μL) was added, followed by homogenization and centrifugation. The supernatant was transferred to a conical tube and precipitated with 300 μL cold ethanol. After discarding the ethanol, samples were dried at 94°C for 1 min, resuspended in 50 μL ultrapure MilliQ water and stored at -20°C .

HBV PCR amplification

To characterize HBV genotypes, a fragment of 1306 bp partially comprising HBsAg and DNA polymerase coding regions (S/POL) was amplified by nested PCR using the primers PS3132F/2920R and PS3201F/P1285R [18]. After purification of the PCR product through ChargeSwitch PCR Clean-Up Kit, Sanger sequencing was performed using dideoxynucleotide triphosphates (ddNTPs) in Big Dye Terminator v3.1 Cycle Sequencing Ready Reaction kit – Applied Biosystems, Foster City, CA, USA). The electrophoresis was done in an ABI Prism 377 Automatic Sequencer (Applied Biosystems, Foster City, CA, USA). The quality of each electropherogram was evaluated using the Phred-Phrap software and consensus sequences were obtained by alignment of both sequenced strands (sense and antisense) using CAP3 software available at the web page Electropherogram quality analysis Phred (<http://asparagin.cenargen.embrapa.br/phph>).

Amplification of the whole HBV genome was performed with P1 and P2 primers described previously with slight modifications [20]. The quality of each electropherogram was evaluated as cited above.

HBV genotyping analysis

Sequences were genotyped by phylogenetic reconstructions using reference sequences from all HBV genotypes obtained from GenBank ($n=412$), comprising 1306 bp of partial HBsAg and DNA polymerase coding regions (S/POL). Complete genomes were also obtained from Genbank ($n=192$) and phylogenetic analyses were performed. All sequences were aligned using Muscle software [21] and edited with the SE-AL software (available at <http://tree.bio.ed.ac.uk/software/seal/>). The Bayesian Markov chain Monte Carlo (MCMC) simulation implemented in BEAST v.1.5.4 [22] was done to obtain the best possible estimates under both relaxed uncorrelated log_{normal} and exponential molecular clock and using the model of nucleotide substitution (GTR+G+I). The molecular clock that best fitted the data was chosen by Bayes factor (BF) comparison. After 10 million generations, the maximum credibility tree (MCC) was obtained by summarizing the 10,000 substitution trees after removing a burn-in of 10% using Tree Annotator v.1.5.3 [22]. Phylogenetic trees were visualized and midpoint rooted in FigTree v1.2.2 (<http://tree.bio.ed.ac.uk/software/figtree/>).

Results

Of the 60 HbsAg-positive samples, 29 (48.3%) were positive by nested PCR for S/POL region and among them, 23 were obtained with good quality sequenced for phylogenetic analysis (Figure 1). The distribution of HBV genotypes in these 23 samples was: A1 (52.17%), E (39.13%),



Figure 1 (See legend on next page.)

(See figure on previous page.)

Figure 1 The Maximum Clade Credibility (MCC) tree was estimated by a Bayesian analysis from a larger dataset comprising 413 sequences with 1306 nucleotides of S and Polymerase HBV region of the different subgenotypes. The posterior probabilities of the key nodes are shown above the respective clusters. The cluster containing the strains of other HBV subgenotypes were collapsed.

D3 (4.3%) and F3/A1 (4.3%). Nine genotype E sequences generated in this study from the same community were previously published [16] since they represented an exclusively African HBV genotype circulating in South America. Due to a small sample size, these results were considered inconclusive about the prevalence of these genotypes in the population.

This work shows for the first time globally a recombinant strain of HBV genotype F3/A1 with 1306bp. The break point was at position 941 of HBV genome (POL region). The analysis using the Simplot program and bootscanning analysis confirmed this complex recombination. The sequence was compared with a consensus sequence of each HBV genotype (A–H) in order to identify the breakpoints. The analysis was carried out using a window size of 200 bp, a step size of 20 bp, 100 bootstrap replicates, gapstrip on and neighbour-joining analysis. The break point at position 941 of HBV genome (POL region) was in the codon ATT (C) (HBV/F3) → ACT (C) (HBV/A1) (Figure 2). Furthermore, we obtained for the first time seven HBV Colombian complete genome sequences from A1, E and D3 genotypes (Figure 3). Complete genome HBV sequences were deposited at the GenBank under accession numbers: JQ023660–JQ023666.

Discussion

These results also showed that the distribution pattern of HBV genotypes and/or genotypes variation could be different in Colombia based on the region studied since unlike in Quibdó, a previous study in Bogota found that

the most frequent subgenotype was the subgenotype F3 followed by A2 and G [18]. The presence of genotype E in more than two thirds of cases studied herein highlights the importance to carry out larger studies in Quibdó population to ascertain if this genotype is widespread in this region.

Such results could provide additional knowledge on the history of HBV/E around the world and further clarify molecular chronometer of such viral infections among humans. Based on our recent publication on genotype E in this afro-Colombian community [16], the TMRCA data suggested a recent infection; however such high frequency of infection in a population does not seem to match a recent infection. Further investigations are required to correlate these findings.

In Africa, viruses belonging to five genotypes, A (HBV/A) to E (HBV/E), have been found. Subgenotype A1 was identified in HBV isolates from South Africa using phylogenetic analysis of preS2/S sequences and confirmed by analysis of complete genomes from South Africa and Malawi [9]. This subgenotype has also been found in Somalia, Zimbabwe, Kenya, Rwanda, Philippines, India and Nepal and Yemen [10]. In South America, previous studies have shown that subgenotype A1 was the most frequent in Brazilian population [23]. While in Argentina there is a low prevalence of this subgenotype [24], in Haiti more than 90% of the population descended from African Slaves has the A1 as the most common subgenotype [25]. While this suggests that subgenotype A1 in South America is probably of similar origin, the subgenotype A1 in Colombia reported in this




Figure 2 Nucleotide similarity comparisons with consensus sequences represented each HBV genotype (A-I) and the Colombian recombinant strain HBV F3/A1. The break point of the Colombian strain is at position 941 of HBV genome (POL region).




Figure 3 (See legend on next page.)

(See figure on previous page.)

Figure 3 The Maximum Clade Credibility (MCC) tree was estimated by a Bayesian analysis of 192 complete genome sequences of HBV strains. The posterior probabilities of the key nodes are shown above the respective nodes. The HBV Colombian complete genome sequences (n = 7) were analyzed together with other strains from around the world. The cluster containing the strains of other HBV subgenotypes were collapsed.

study though grouped in the same cluster, seems so suggest otherwise since the posterior probability was low (0.21), thus indicating a distant genetic relatedness.

An important result of this work was the presence of recombinant strain between subgenotype F3 and A1 found in a sixteen years old girl. Since this girl was pregnant when this sample was collected, is probably that sexual contact was the way of HBV transmission. Several HBV recombinants inter genotypes have been reported around the world. In Bolivia, South America, several recombinants A/D, C/B, D/C and F/C have been detected [26]. Furthermore, recombination within and between genotypes created complex patterns and altered the cladistic structure of HBV genotypes [27]. For example, in previous study the B2 subgenotype proved to be a hybrid of genotypes B and C [28]. Unfortunately, we had no success on full-length genome amplification of our recombinant strain. Thus, although we know that the break point was in the polymerase region, we cannot exclude the possibility with of existence of other recombination points in this virus. In sum, this is the first HBV recombinant strain reported in Colombia.

Here, we present a study of the first seven complete HBV genomes of Colombian population. The results obtained from E and A1 genotypes support the theory that HBV may have been introduced into this Afro-descendent community in Colombia in the times of slavery.

Competing interests

The authors declare that they have no competing interests.

Author's contributions

MVAM participated in the design of the study and drafted the manuscript, conducted the sequencing process and phylogenetic analysis. CMR conducted the recombination and evolutionary analysis. MSGG participated in the PCR amplification. MFG and FJC participated in the design of the study. JRRP participated in the elaboration of the manuscript. All authors read and approved the final manuscript.

Acknowledgements

This study was previously presented as a poster in 2011 in the 62nd annual meeting of the American association for the study of liver diseases in San Francisco, United States of America. This work has been supported by CNPq and by Fundação de Amparo à Pesquisa do Estado de São Paulo – FAPESP 2007/53457-7/2011/50562-0 and 2008/50461-6, HCFMUSP and FFM.

Author details

¹Laboratory of Gastroenterology and Hepatology, São Paulo Institute of Tropical Medicine and Department of Gastroenterology, School of Medicine, University of São Paulo, São Paulo, Brazil. ²São Paulo Institute of Tropical Medicine, Department of Infectious and Parasitic Diseases (LIMHC), School of Medicine, University of São Paulo, São Paulo, Brazil. ³Laboratory of Virology, Department of Microbiology, Pontificia Javeriana University, Bogotá, Colombia.

Received: 13 February 2012 Accepted: 22 October 2012

Published: 24 October 2012

References

1. Lee WM: Hepatitis B virus infection. *N Engl J Med* 1997, **337**(24):1733–1745.
2. Tanaka J: Hepatitis B epidemiology in Latin America. *Vaccine* 2000, **18**(Suppl 1):S17–S19.
3. Arboleda M, Castilho MC, Fonseca JC, Albuquerque BC, Saboia RC, Yoshida CF: Epidemiological aspects of hepatitis B and D virus infection in the northern region of Amazonas, Brazil. *Trans R Soc Trop Med Hyg* 1995, **89**:481–483.
4. Alvarado-Mora MV, Fernandez MF, Gomes-Gouveia MS, de Azevedo Neto RS, Carrilho FJ, Pinho JR: Hepatitis B (HBV), hepatitis C (HCV) and hepatitis delta (HDV) viruses in the Colombian population—how is the epidemiological situation? *PLoS One* 2011, **6**(4):e18888.
5. Fay OH: Hepatitis B in Latin America: epidemiological patterns and eradication strategy. The Latin American Regional Study Group. *Vaccine* 1990, **8**:S100–S106.
6. Duarte MC, Cardona N, Poblete F, Gonzalez K, Garcia M, Pacheco M, Botto C, Pujol FH, Williams JR: A comparative epidemiological study of hepatitis B and hepatitis D virus infections in Yanomami and Piaroa Amerindians of Amazonas State, Venezuela. *Trop Med Int Health* 2010, **15**(8):924–933.
7. Stuyver L, De Gendt S, Van Geyt C, Zoulim F, Fried M, Schinazi RF, Rossau R: A new genotype of hepatitis B virus: complete genome and phylogenetic relatedness. *J Gen Virol* 2000, **81**(Pt 1):67–74.
8. Yu H, Yuan Q, Ge SX, Wang HY, Zhang YL, Chen QR, Zhang J, Chen PJ, Xia NS: Molecular and phylogenetic analyses suggest an additional hepatitis B virus genotype "I". *PLoS One* 2010, **5**(2):e9297.
9. Kramvis A, Weitzmann L, Owiredu WK, Kew MC: Analysis of the complete genome of subgroup A' hepatitis B virus isolates from South Africa. *J Gen Virol* 2002, **83**(Pt 4):835–839.
10. Kramvis A, Kew MC: Molecular characterization of subgenotype A1 (subgroup Aa) of hepatitis B virus. *Hepatol Res* 2007, **37**(s1):S27–S32.
11. Norder H, Courouce AM, Coursaget P, Echevarria JM, Lee SD, Mushahwar IK, Robertson BH, Locarnini S, Magnius LO: Genetic diversity of hepatitis B virus strains derived worldwide: genotypes, subgenotypes, and HBsAg subtypes. *Intervirology* 2004, **47**(6):289–309.
12. Hannoun C, Soderstrom A, Norkrans G, Lindh M: Phylogeny of African complete genomes reveals a West African genotype A subtype of hepatitis B virus and relatedness between Somali and Asian A1 sequences. *J Gen Virol* 2005, **86**(Pt 8):2163–2167.
13. Olinger CM, Venard V, Njayou M, Oyefolu AO, Maiga I, Kemp AJ, Omilabu SA, le Faou A, Muller CP: Phylogenetic analysis of the precore/core gene of hepatitis B virus genotypes E and A in West Africa: new subtypes, mixed infections and recombinations. *J Gen Virol* 2006, **87**(Pt 5):1163–1173.
14. Pourkarim MR, Lemey P, Amini-Bavil-Olyae S, Maes P, Van Ranst M: Novel hepatitis B virus subgenotype A6 in African-Belgian patients. *J Clin Virol* 2010, **47**(1):93–96.
15. Hubschen JM, Mbah PO, Forbi JC, Otegbayo JA, Olinger CM, Charpentier E, Muller CP: Detection of a new subgenotype of hepatitis B virus genotype A in Cameroon but not in neighbouring Nigeria. *Clin Microbiol Infect* 2011, **17**(1):88–94.
16. Alvarado Mora MV, Romano CM, Gomes-Gouveia MS, Gutierrez MF, Carrilho FJ, Pinho JR: Molecular epidemiology and genetic diversity of hepatitis B virus genotype E in an isolated Afro-Colombian community. *J Gen Virol* 2010, **91**(Pt 2):501–508.
17. Devesa M, Loureiro CL, Rivas Y, Monsalve F, Cardona N, Duarte MC, Poblete F, Gutierrez MF, Botto C, Pujol FH: Subgenotype diversity of hepatitis B virus American genotype F in Amerindians from Venezuela and the general population of Colombia. *J Med Virol* 2008, **80**(1):20–26.

18. Alvarado Mora MV, Romano CM, Gomes-Gouvea MS, Gutierrez MF, Botelho L, Carrilho FJ, Pinho JR: **Molecular characterization of the Hepatitis B virus genotypes in Colombia: a Bayesian inference on the genotype F.** *Infect Genet Evol* 2011, **11**(1):103–108.
19. Chomczynski P, Sacchi N: **The single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction: twenty- something years on.** *Nat Protoc* 2006, **1**(2):581–585.
20. Gomes-Gouvea MS: *Characterization of complete genomes of the hepatitis B virus of different genotypes isolated in Brazil.* São Paulo: Secretaria da Saúde/ Coordenadoria de Controle de Doenças. Programa de Pós-Graduação em Ciências para obtenção do grau de Mestre; 2005:130.
21. Edgar RC: **MUSCLE: a multiple sequence alignment method with reduced time and space complexity.** *BMC Bioinformatics* 2004, **5**:113.
22. Drummond AJ, Rambaut A: **BEAST: Bayesian evolutionary analysis by sampling trees.** *BMC Evol Biol* 2007, **7**:214.
23. Motta-Castro AR, Martins RM, Yoshida CF, Teles SA, Paniago AM, Lima KM, Gomes SA: **Hepatitis B virus infection in isolated Afro-Brazilian communities.** *J Med Virol* 2005, **77**(2):188–193.
24. Mbayed VA, Pineiro y Leone FG, Pezzano SC, Campos RH: **Molecular characterization of hepatitis B virus genotype A from Argentina and Brazil.** *Arch Virol* 2009, **154**(3):525–529.
25. Andernach IE, Nolte C, Pape JW, Muller CP: **Slave trade and hepatitis B virus genotypes and subgenotypes in Haiti and Africa.** *Emerg Infect Dis* 2009, **15**(8):1222–1228.
26. Simmonds P, Midgley S: **Recombination in the genesis and evolution of hepatitis B virus genotypes.** *J Virol* 2005, **79**(24):15467–15476.
27. Purdy MA, Gonzales AC, Dimitrova Z, Khudyakov Y: **Supragenotypic groups of the hepatitis B virus genome.** *J Gen Virol* 2008, **89**(Pt 5):1179–1183.
28. Sugauchi F, Orito E, Ichida T, Kato H, Sakugawa H, Kakumu S, Ishida, Chutaputti A, Lai CL, Ueda R, *et al*: **Hepatitis B virus of genotype B with or without recombination with genotype C over the precore region plus the core gene.** *J Virol* 2002, **76**(12):5985–5992.

doi:10.1186/1743-422X-9-244

Cite this article as: Alvarado-Mora *et al.*: Phylogenetic analysis of complete genome sequences of hepatitis B virus from an Afro-Colombian community: presence of HBV F3/A1 recombinant strain. *Virology Journal* 2012 **9**:244.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

