



Universidade de São Paulo

Biblioteca Digital da Produção Intelectual - BDPI

Departamento de Epidemiologia - FSP/HEP

Artigos e Materiais de Revistas Científicas - FSP/HEP

2013

Phylogeography of the neotropical *Anopheles triannulatus* complex (Diptera: Culicidae) supports deep structure and complex patterns

Parasites & Vectors. 2013 Feb 22;6(1):47
<http://www.producao.usp.br/handle/BDPI/34972>

Downloaded from: Biblioteca Digital da Produção Intelectual - BDPI, Universidade de São Paulo

RESEARCH

Open Access

Phylogeography of the neotropical *Anopheles triannulatus* complex (Diptera: Culicidae) supports deep structure and complex patterns

Marta Moreno^{1,13*}, Sara Bickersmith¹, Wesley Harlow¹, Jessica Hildebrandt¹, Sascha N McKeon^{1,2}, Teresa Fernandes Silva-do-Nascimento³, Jose R Loaiza⁴, Freddy Ruiz⁵, Ricardo Lourenço-de-Oliveira³, Maria AM Sallum⁶, Eduardo S Bergo⁷, Gary N Fritz⁸, Richard C Wilkerson⁵, Yvonne M Linton⁵, Maria J Dantur Juri⁹, Yadira Rangel¹⁰, Marinete M Póvoa¹¹, Lina A Gutiérrez-Builes¹², Margarita M Correa¹² and Jan E Conn^{1,2}

Abstract

Background: The molecular phylogenetic relationships and population structure of the species of the *Anopheles triannulatus* complex: *Anopheles triannulatus* s.s., *Anopheles halophylus* and the putative species *Anopheles triannulatus* C were investigated.

Methods: The mitochondrial *COI* gene, the nuclear *white* gene and rDNA ITS2 of samples that include the known geographic distribution of these taxa were analyzed. Phylogenetic analyses were performed using Bayesian inference, Maximum parsimony and Maximum likelihood approaches.

Results: Each data set analyzed separately yielded a different topology but none provided evidence for the separation of *An. halophylus* and *An. triannulatus* C, consistent with the hypothesis that the two are undergoing incipient speciation. The phylogenetic analyses of the *white* gene found three main clades, whereas the statistical parsimony network detected only a single metapopulation of *Anopheles triannulatus* s.l. Seven *COI* lineages were detected by phylogenetic and network analysis. In contrast, the network, but not the phylogenetic analyses, strongly supported three ITS2 groups. Combined data analyses provided the best resolution of the trees, with two major clades, Amazonian (clade I) and trans-Andean + Amazon Delta (clade II). Clade I consists of multiple subclades: *An. halophylus* + *An. triannulatus* C; trans-Andean Venezuela; central Amazonia + central Bolivia; Atlantic coastal lowland; and Amazon delta. Clade II includes three subclades: Panama; cis-Andean Colombia; and cis-Venezuela. The Amazon delta specimens are in both clades, likely indicating local sympatry. Spatial and molecular variance analyses detected nine groups, corroborating some of subclades obtained in the combined data analysis.

Conclusion: Combination of the three molecular markers provided the best resolution for differentiation within *An. triannulatus* s.s. and *An. halophylus* and C. The latest two species seem to be very closely related and the analyses performed were not conclusive regarding species differentiation. Further studies including new molecular markers would be desirable to solve this species status question. Besides, results of the study indicate a trans-Andean origin for *An. triannulatus* s.l. The potential implications for malaria epidemiology remain to be investigated.

Keywords: *Anopheles triannulatus* s.s., *Anopheles halophylus*, *Anopheles triannulatus* C, phylogeography, *COI* gene, *White* gene, ITS2

* Correspondence: mmorenoleirana@ucsd.edu

¹New York State Department of Health, Wadsworth Center, Griffin Laboratory, Albany, NY, USA

¹³Present address: Division Infectious Diseases University of California San Diego, George Palade Labs, School of Medicine, 9500 Gilman Drive, MC 0741, La Jolla, CA 92093, USA

Full list of author information is available at the end of the article

Background

Neotropical anophelinae have shown an extraordinary diversity and complexity due, in part, to the capability of dipterans to adapt to and utilize a broad variety of ecological niches [1]. The subgenus *Nyssorhynchus* provides an example of extensive morphological and genetic variation within taxa, and several studies have focused on the processes driving this differentiation. Some hypotheses are related to climatic changes that occurred in different epochs, provoking drastic modifications to the habitats of different organisms [2,3]. However, this remains controversial because of the paucity of available evidence-based data [4-6]. Furthermore, factors affecting speciation and population differentiation, such as ecology, behaviour and genetics, evolve at different rates and are not necessarily congruent [7].

Some of these taxa, estimated to be 10% of all anophelinae, are directly responsible for malaria parasite transmission and their accurate identification is necessary for the implementation of effective control strategies. Understanding the current distribution of species, investigating past or recent demographic events (population growth or contraction), gene flow, as well as human interventions (exploitation of new ecological niches, introduction of non endemic species, among others), can provide powerful tools and valuable predictors for the management of pathogens transmitted by anophelinae.

Anopheles triannulatus s.l., subgenus *Nyssorhynchus*, [8] was first described from adult females in central Brazil and subsequently reported in Central America (Costa Rica, Nicaragua and Panama), in the majority of South American countries [9,10], and recently in some Caribbean islands [11]. This species has been previously described under different names (*syn. bachmanni* Petrocchi, *syn. chagasi* Galvao, *syn. cuyabensis* Neiva and Pinto, *syn. davisii* Paterson and Shannon, *syn. perezii* Shannon and Del Ponte). Morphological variation was later considered intraspecific and attributed to adaptation to different habitats [9,12-18]. However, recent investigations based on morphological characters of the male genitalia and immature stages led to the designation of a new species, *Anopheles halophylus*, and elevated the status from polymorphic species to complex, (i.e. the *Anopheles triannulatus* complex) [18-20].

Genetic distance analysis of allozymes and RAPD detected a third species, *An. triannulatus* C, and showed that *An. halophylus* and *An. triannulatus* species C formed a reciprocally monophyletic group [20]. Apart from this preliminary finding, very little is known about the phylogenetic relationships of members of the *An. triannulatus* complex, except for the results of analysis of sequences of *cpr* and *timeless* genes [21], which confirmed previous findings and suggested that *An. halophylus* and *An. triannulatus* C are in the process of incipient speciation.

Seasonal population density and behavioral differences have also been reported within the complex, for example, potential species-specific preferences for different larval habitats of species that occur in sympatry, e.g., *An. halophylus* and *An. triannulatus* species C [19,22].

An. triannulatus s.l. has been incriminated in human malaria transmission in different regions of Brazil [23-26]), and probably Peru and Venezuela [27,28], although the role of each species within the complex remains unknown. However, zoophilic and exophilic behaviour has been much commonly reported in Brazil [29-32].

The study of sibling species is aggravated by the difficulty of identification based exclusively on morphological characters when key traits, especially in adult females, may exhibit great phenotypic plasticity [33]. Nowadays, DNA sequences are an essential tool for delineating and identifying species, as well as for gathering information about the genetic variation within species complexes. The mitochondrial gene cytochrome oxidase subunit 1 (*COI*) is usually informative in phylogenetic reconstruction and geographic variability [34-36]. To represent the nuclear genome the single-copy *white* gene was chosen [37,38]. The ribosomal DNA (r-DNA) internal transcribed spacer 2 (ITS2) was used because it plays an important role in distinguishing cryptic anopheline species [39,40].

The major objective of this study was to reconstruct and clarify the evolutionary relationships based on mtDNA and nuclear sequences, to illustrate the demographic history at the population level, and to provide information on the distribution of *An. triannulatus* s.l. in several countries across its range. The information obtained would be useful to distinguish the species and help to focus scarce vector control resources on species involved in malaria transmission.

Methods

Sampling

Mosquitoes were collected from 31 localities in eight countries, including Argentina, Bolivia, Brazil, Colombia, Ecuador, Panama, Trinidad-Tobago and Venezuela (Table 1) covering a wide range of the reported species distribution (Figure 1). Adult mosquitoes were collected outdoors by human landing catches (protocols approved by the Institutional Review Board of the New York State Department of Health, University of Antioquia, University of Florida and University of Vermont), except specimens from Argentina and Brazil that were collected by CDC light traps or horse-baited Shannon traps, respectively. Some specimens included in the analysis from Brazil (donated by MAMS) were collected as either larvae or pupae and kept in the laboratory to obtain adults linked with larval and/or pupal exuviae. Two topotype specimens

Table 1 Summary data for collection localities and sample information of the *Anopheles triannulatus* complex

Site No.	Country	Locality	N			Coordinates	Collector
			COI (n = 326)	white (n = 85)	ITS2 (n = 34)		
1	Argentina	Yuto Farm (FY)	23	3	1	23°63'S/64°46'W	MJD
2		Lake Yuto (LY)	3	0	0	23°38'S/64°27'W	MJD
3	Bolivia	Guayaramerin (GU)	4	0	0	10°49'S/65°21'W	JEC
4		Puerto Villaroel (PV)	10	2	2	16°52'S/64°46'W	JEC
5	Brazil	Ceara (CE)	2	2	0	5°05'S/40°23'W	TFS
6		Espirito Santo (ES)	1	0	0	1°15'N/50°54'W	TFS
7		Fazenda S. Joao (MT)	2	1	1	16°57'S/56°36'W	TFS
8		Monte Negro (RO)	5	2	0	10°15'S/63°19'W	MAMS/ESB
9		Aquidauana (Pantanal do Rio Negro) (AQU)	6	4	0	19°29'S/55°36'W	MAMS/ESB
10		Inubia Paulista (SP)	6	2	0	21°39'S/50°56'W	MAMS/ESB
11		Camacan/Santa Luzia (Bahia) (BA)	7	3	0	15°25'S/39°32'W	MAMS/ESB
12		Coronel Pacheco (Minas Gerais) (MG)	6	3	0	21°38'S/43°19'W	MAMS/ESB
13		Lagoa da Confusao (Tocantins) (TO)	6	2	0	10°35'S/49°41'W	MAMS/ESB
14		Itaituba (ITB)	5	1	1	4°15'S/55°59'W	MAMP/JEC
15		Oswaldo Cruz (OC)	19	7	3	8°00'S/35°00'W	MAMP/JEC
16		Salobra (MS)	20	9	4	20°12'S/56°29'W	TFS
17		Silva Jardim (RJ)	2	2	1	22°39'S/42°23'W	TFS
18		Tartarugalzinho (TAR)	23	4	1	1°30'N/50°54'W	MAMP/JEC
19	Colombia	Monitos (MO)	22	6	4	9°13'N/76°08'W	NN/PA
20		Santa Rosa de Lima (SO)	27	5	4	10°26'N/75°21'W	NN/JP
21		Leticia, km12 (COSW)	10	4	3	04°6'S/69°57'W	JFR
22		Tibu (COL)	3	0	0	08°38'N/72°44'W	JFR
23	Ecuador	Juan Montalvo (JM)	6	1	1	0°50'N/78°51'W	GNF
24	Panama	Bayano (BAY)	7	1	1	9°07'N/79°01'W	JRL
25		Gamboia (GAM)	29	4	2	9°07'N/79°42'W	JRL
26		Meteti (MET)	17	2	1	8°30'N/79°58'W	JRL
27	Venezuela	Boconoito (BOC)	6	2	0	8°50'N/69°58'W	JEC/YR
28		Cano Amarillo (CAM)	15	6	2	8°43'N/71°34'W	JEC/YR
29		Casigua Zulia (CAS)	25	4	2	8°44'N/72°30'W	JEC/YR
30		La Veguita (VG)	6	1	0	8°52'N/70°0'W	JEC/EB
31	Trinidad-Tobago	St.Andrew/St.David, Valencia (TRI)	2	0	0	10°39'N/61°09'W	RCW

Salobra (16) and Fazenda S. Joao (7), in Brazil, are the type localities for *An. halophylus* and *An. triannulatus* C, respectively. N, sample size of each of the genes sequenced; MJD, MJ Dantur; JEC, JE Conn; TFS, T Fernandes-da-Silva; MAMS, MA Mureb-Sallum; ESB, ES Bergo; MAMP: MAM Povoia; NN, N Naranjo; PA, P Aviles; JP, J Pinto; JFR, JF Ruiz; GNF, GN Fritz; JRL, JR Loaiza; YR: Y Rangel; EB: E Brown; RCW, RC Wilkerson.

of *Anopheles halophylus* and *Anopheles triannulatus* C (Salobra, Brazil) and *An. triannulatus* s.s. were included in the analysis. Specimens were morphologically identified using the available identification keys [9,17]. Identification of *An. halophylus*, *An. triannulatus* C and some representatives of *An. triannulatus* s.s. was based on morphological characters and the *Mpi* diagnostic allozyme [20]. Mosquitoes were then stored in either 95% ethanol or on silica gel at room temperature until DNA extraction.

DNA extraction and gene amplification

Total genomic DNA was extracted from each specimen using the DNeasy tissue kit (Qiagen, CA, USA). Polymerase chain reaction was used to amplify a 1200 base pair (bp) fragment of the mtDNA *COI* gene using the primers UEA3 and UEA10 described in [41]. The *white* gene was also amplified via PCR using the primers WF and W2R [42,43]. The amplification of the ITS2 was performed following the protocol described in [44] with




Figure 1 *Anopheles triannulatus* s.l. sampled localities. The locality numbers correspond to those in Table 1. Black dots represent *Anopheles triannulatus* s.l., black and white triangles represent *An. halophylus* and *An. triannulatus* C type localities, respectively. The map depicts the $k = 9$ groups yielded by SAMOVA software and supported by AMOVA analysis.

the primers 5.8S and 28S, yielding a 500 bp fragment. Each PCR reaction was carried out using a Ready-To-Go-PCR bead (Amersham Pharmacia/Biotech NJ, USA) and performed in a PTC-200 thermal cycler (BioRad Inc.). PCR products were purified on CentriSpin 40 columns (Princeton Separations, NJ, USA). Following standard PCR reactions, both strands were sequenced at the Applied Genomic Technologies Core (Wadsworth Center) on an ABI PRISM 3700 automated DNA sequencer.

Sequence alignment and phylogenetic analysis

Sequencer 4.1 (Gene Codes Corps.) was used for automatic sequence alignment into contigs and proofreading sequences files; subsequently, all sequences were aligned with Clustal W and then compiled in Bioedit 4.0 [45].

The data sets defined for the subsequent analyses were: 1) the *COI* fragment, 2) the *COI* gene with nt1 + nt2 positions (to avoid redundancy in the nt3 position), 3) the *white* gene (with and without intron), 4) the ITS2 and 5) a combined concatenated data set of the three fragments. For each data set, an appropriate model of nucleotide substitutions was obtained with the software jModeltest 3.5 [46]. The models obtained were GTR + I + G for the *COI* gene (all positions), K81uf + I + G for *COI* gene (nt1 + nt2 positions), GTR + G for the *white* gene and HKY for the ITS2, selected by using the Akaike Information Criterion

(AIC). These model parameters were used as prior information in the subsequent maximum likelihood (ML) and Bayesian analyses.

To reconstruct phylogenetic relationships, four different approaches with each of the data sets were conducted. The ML analyses were performed using the heuristic search algorithm; both analyses were performed on the haplotypes with branch support evaluated by 1000 bootstrap replicates.

Bayesian inference (BI) analysis was performed using MrBayes 3.0.4 [47] allowing, in the combined data set, the specification of a distinct model and parameters for the subset [48]. Different partitioning strategies were defined: P1) is a partition for each gene; P2) a partition for each gene without the 3rd codon position for the mitochondrial marker and P3) the combined sequence without partitions. A Bayes factor ($2\ln B_{F}$) was used to choose the best partitioning strategies. Analyses were initiated with random starting trees and run for 10,000,000 to 20,000,000 generations, sampling every 1000 generations with a 'burn-in' of 25%. Posterior probabilities were used to assess nodal support. FigTree v1.2.1 was employed for visualization of the consensus trees.

Population structure and genetic diversity analysis

Nucleotide and haplotype diversities were calculated within the main mitochondrial lineages, defined by the

previous BI and the minimum spanning network, using DNASP v 5 [49].

A spatial analysis of molecular variance (SAMOVA) was performed to define groups of populations as geographically homogeneous and with a maximum of differentiation from each other, combining genetic and geographic sample information [50]. The random partition of the data was run by defining $K = 2-20$, with K number of groups to maximize the F_{CT} index, i.e., the proportion of total genetic variance due to differences between groups of populations [51].

In addition, the characterization of patterns of genetic diversity variation in the *Anopheles triannulatus* Complex was performed with the analysis of molecular variance (AMOVA), with the option of establishing different hierarchical levels [51]. Groups of variation were defined based on SAMOVA, BI and networks (Figure 2): 1) 9 groups obtained from SAMOVA; 2) SAMOVA groups

but with Ecuador (23; Figure 1) and Amazonian Colombia (21; Figure 1) together; 3) SAMOVA groups but combining Venezuela (29) with those from western Colombia (19, 20); and 4) SAMOVA groups but combining locality 29 with the Venezuelan and Trinidad-Tobago populations.

To describe patterns of population genetic variation that originated from spatially limited gene flow, isolation by distance (IBD) was tested with a nonparametric Mantel test through the web-based computer program IBDWS v3.16 [52]. This analysis was applied only to populations in northwestern Latin America for which we have adequate regional (geographical) sampling. In this case, the distance was calculated as the straight line geographical distance between pairs of sampled localities using the Geographic Distance Matrix Generator tool [53] and the validation of the analysis was provided by 10,000 randomizations. Partial Mantel tests were performed to test for the effect of the Andes on genetic divergence [54].




Figure 2 Tree topology MP, ML and BI of the mitochondrial *COI* gene of the *Anopheles triannulatus* complex. Both procedures produced trees of similar topology, with only branch support values above 70% shown. Bayesian posterior probability is above the branch and Maximum Likelihood bootstrapping (percentage) is below the branch. *Anopheles albimanus*, *Anopheles darlingi*, *Anopheles nuneztovari* and *Anopheles marajoara* were used to root the tree.

Phylogeographic analysis

To infer the haplotype relationships within the data sets, the median-joining network algorithm was performed [55], available in NETWORK v 4.5.1.0 (www.fluxus-engineering.com), which combines the topology of a minimum spanning tree with a parsimonious search for the missing haplotypes. To determine the ancestral network node, *COI* sequences of closely related species, *Anopheles nuneztovari* and *Anopheles darlingi* were included. Because this algorithm was designed for non-recombining molecules, a statistical parsimony network was used for the nuclear *white* gene and the ITS2 data [56] with a 95% confidence level, using the TCS v 1.21 software [57].

Inference of population history

To investigate the historical demography of the *Anopheles triannulatus* Complex, neutrality tests were performed and applied to the SAMOVA groups and to the entire data set. The first two estimates (Tajima's D and R_2) are based on the frequency spectrum of mutation whereas Fu's F_s is based on linkage disequilibrium and haplotype distribution. To further confirm demographic expansion events, mismatch distribution analyses were conducted. The time of expansion was estimated using the formula $\tau = 2ut_e$, where τ is estimated in the mismatch distribution analysis, u is the nucleotide mutation rate in the specific DNA region and t_e is time since expansion. The mutation rate in the mitochondrial gene has been estimated from *Drosophila* as 1×10^{-8} /site/year [58]. The time of the beginning of lineage divergence was calculated following $D_A/2k$, where D_A is the average nucleotide divergence between lineages and $2k$ is the divergence rate [59]. We utilized the *COI* gene substitution rate of 2.3% divergence per million years [60] and estimated these parameters and their significance based on the coalescence process in DNASP v 5.10 [49] and Arlequin v3.11 [61].

Results

A total of 326 specimens of the *Anopheles triannulatus* complex were sequenced for the mtDNA *COI* gene (Table 1). All sequences could be unambiguously aligned and no insertions or deletions were found. Translation into amino acid ruled out the presence of nuclear mitochondrial pseudogenes.

The final alignment of the *COI* gene (partial sequence) had a length of 689 bp, of which 202 characters were variable polymorphic sites and 154 were parsimony informative. For the single-copy nuclear *white* gene, 85 individuals randomly selected were included, representing each of the mtDNA-based lineages, and the *An. halophylus* and *An. triannulatus* C samples. A fragment of 646 bp with an intron of 70 bp was detected within the *white* gene sequences in all individuals, independent of the species. This intron

was excluded for some of the analyses using a final fragment of 576 bp.

Finally, ITS2 was amplified from 34 individuals randomly chosen from the *COI* lineages, and the length varied from 542-570 bp (intra and interspecific differences), with a final alignment length of 483 bp (3' and 5' ends to make sequences equal length). All sequences were deposited in GenBank under Accession numbers *COI* (JN085964-JN086138), *white* gene (JF931140-JF931172) and ITS2 (JF972999-JF973009).

Phylogenetic analysis

Mitochondrial DNA data

The Bayesian, ML and MP trees of the overall analysis are shown in Figure 2. Trees were rooted using Neotropical *Nyssorhynchus* species: *Anopheles darlingi*, *Anopheles marajoara*, *Anopheles nuneztovari* and *Anopheles albimanus* as outgroup [62]. The tree topologies obtained by the 50% majority rule consensus using BL, ML and MP in PAUP were basically in agreement, consistently recovering the *Anopheles triannulatus* complex as a monophyletic lineage. Two main clades were identified based on the geographic distribution of the samples (with some exceptions) and their genetic distances: one encompassing samples from Panama, northern Colombia and trans-Andean Venezuela, and the other (Amazonian) including samples from South America and cis-Andean Venezuela, and *An. halophylus* and *An. triannulatus* C. Within the Amazonian clade further subclades can be described according to tree topology and geographic origin of the samples (Figure 2). Both topologies suggest that the clade, which encompasses samples from Colombia is the most basal within the Complex. The main differences in the tree topologies were the recovery of Panama, Colombia and trans-Venezuela (northern) as a unique clade, although with low posterior probability branch support. The second clade (Figure 2) is shallower when estimated by MP, and not all the subclades are supported.

The samples identified as *An. halophylus* and *An. triannulatus* C were in a clade together with other individuals identified as *An. triannulatus* s.s. from southern Brazil, Bolivia and Argentina. In all tree topologies, *An. halophylus* shares the same haplotype as one individual from eastern Brazil originally identified morphologically and by *Mpi* as *An. triannulatus* s.s.

Analysis excluding the nt3 codon position was performed because of saturation issues [63], but this did not improve the resolution of the topologies. No evidence for saturation at any of the 3 positions was found using the test in [64] (Additional file 1).

Nuclear DNA data

A total of 33 haplotypes were identified among 85 individuals sequenced for the *white* gene. Bayesian tree (Figure 3) was similar to the three generated, employing

MP analyses topology resolved a similar phylogeny, with some of the clades associated to geographic distribution. For instance, one major node includes individuals from central and northwestern South America and another includes samples from eastern and central Brazil (localities 5, 10, 13, Figure 1). In contrast, other individuals have low support or no geographic concordance, such as the genotypes from localities 4, 17 and 28. *An. halophylus* and *An. triannulatus* C clustered together with high support with two other *An. triannulatus* s.l. specimens from localities 1 and 7.


All 33 unique *white* gene genotypes could be connected (Figure 4A). Genotype 1, with individuals from Panama, Colombia and Venezuela, was identified as ancestral [65]: the internal position in the network, the number of lineages that arise from it and the frequency (24 samples). The second most common genotype (number 6) is separated by only 3 mutational steps from genotype 1, and contains 9 samples from NE Brazil. Many of the remaining genotypes were unique ($n = 23$) and only eight were shared. The minimum spanning network depicted a complex pattern of relationships among the genotypes, and there was no support for the BI groups (Figure 2) in the mitochondrial or the nuclear gene genealogy.


Bayesian inference, maximum parsimony and maximum likelihood analysis of the ITS2 marker yielded similar topologies with only subtle differences in bootstrap and posterior probability support values (data not shown).

The tree topologies showed some possibly basal individuals from diverse localities (Colombia, Panama, Brazil, and Venezuela) and two fairly well supported subclades: one of them recovers *An. halophylus* and *An. triannulatus* C as a single monophyletic clade. The other clade contains individuals from central Brazil, southern Colombia, Ecuador and Bolivia. In addition, the ITS2 network (Figure 4B) clearly depicts 3 main groups in the *Anopheles triannulatus* Complex; lineage I recovered individuals from Panama, trans-Andean Venezuela, NE Brazil and SE Brazil whereas lineage II comprised Amazon Basin and trans-Andean Venezuela individuals. Lineage III consisted of two genotypes separated by 1 mutation step with the *An. halophylus* and *An. triannulatus* C specimens, plus two individuals from Brazil and Argentina (originally identified as *An. triannulatus* s.l.), respectively. The deep differentiation of populations separated by the Andes, detected by the *COI* data, is also strongly supported with ITS2 lineages I and II.

Combined mitochondrial and nuclear DNA data

Analysis of partitioning strategies showed the P1 model (independent evolutionary models for each gene) as the most accurate [66]. B_F -based statistics also indicated this partition as the better fit, and the tree derived from these analyses is shown in Figure 5. In general, the combined data presented high posterior probability values. The tree topology recovered two main clades; clade






I consists of multiple subclades: (1) *An. halophylus* + *An. triannulatus* C; (2) trans-Andean Venezuela; (3) central Amazonia + central Bolivia; (4) Atlantic coastal forest; and (5) Amazon delta. Clade II includes three subclades: (1) Panama; (2) cis-Andean Colombia; and cis-Venezuela. The Amazon delta specimens are in both clades (OC9, OC7 in Clade I and OC14 in Clade II), perhaps indicating sympatry of discrete taxa. *Anopheles halophylus* and *An. triannulatus* C were recovered as a single well-supported monophyletic subclade (1) together with some *An. triannulatus* s.l. individuals from the same type locality (Salobra) and from nearby Mato Grosso and Argentina. The other subclades (2–5) generally had a geographic component except that samples from Bolivia are found in both subclades 2 (PV9) and 3 (PV5) (Figure 5).

Population structure and genetic diversity analysis

The 326 sequences of the *COI* gene yielded one hundred and seventy-five haplotypes, with some shared between

species, for instance, for *An. triannulatus* C and *An. triannulatus* s.s (see Table 2). The haplotype network reveals an intricate topology (Figure 6): from Panama, haplotypes A4 ($n = 12$) and A9 ($n = 16$), were the most common; C6 (Venezuela and Brazil) and D12 (NE Brazil) each comprised 11 individuals, whereas F11 (NW Colombia) consisted of 15 individuals. *An. halophylus* (E8 and E53) present haplotypes intermingled with *An. triannulatus* s.s. from eastern Brazil and Mato Grosso. Conversely, both *An. triannulatus* C haplotypes (E7 and E9) were unique.

Overall, the *Anopheles triannulatus* complex network consists of 7 lineages, very similar to the BI, and depicts some relationships between haplotypes and geographic distribution, which may indicate variation in the population histories (Figure 6). Star shaped lineage A, suggestive of population expansion, is restricted to Panama. Other geographically nearby lineages, separated only by 2 mutational steps, are formed by trans-Andean Colombian (F); but



there are >11 mutation steps between A and the western Venezuelan samples (C). In contrast, lineage B, consisting of haplotypes from cis-Andean Venezuela, Colombia, and Trinidad and Tobago, is >21 mutational steps distant. Lineage E is of mixed origin: samples from Ecuador, Bolivia, Amazonian Colombia, southern Brazil, Argentina and a few haplotypes from central and eastern Brazil, mostly as singletons. Lineage D is mainly composed of eastern Brazilian populations with a star shape suggesting an expansion. Lineage G is restricted to populations from SE Brazil, east of the Central Mountain Range.

The SAMOVA F_{CT} values increased with the number of groups. However, the graphical representation of F_{CT} showed that at $k = 9$, a plateau was reached ($F_{CT} = 0.53778$) concordant with the most statistically significant AMOVA groups (Table 3).

A Mantel test showed a significant correlation of pairwise genetic differences between the Panamanian, Colombian and Venezuelan samples with their respective geographic distances ($r = 0.6564$, $p < 0.001$), suggesting that distance accounts for approximately 65% of the genetic differentiation for this region. Furthermore, a partial Mantel test was performed to test the Andes as a barrier to gene flow between mosquito populations from Panama (trans) and Colombia and Venezuela (cis). In this case, the distance still added significant effect to the correlation when the Andean range was primarily considered ($r = 0.6463$; $p < 0.01$) (controlling for geography: $r = -0.0115$; $p = 0.5860$).

Inference of *COI* population history

All the neutrality tests in lineage A were significant, strongly indicating a demographic expansion (Table 4). All the other lineages presented significant F_s and R_2 statistics. In addition, the unimodal mismatch distribution and the nonsignificant raggedness statistics are congruent with a model of sudden expansion in all lineages (Figure 7). A range of early to late Pleistocene divergence was identified in all the lineages, with the oldest estimation for widespread lineage E. However, visually, lineage E mismatch distribution and the haplotype network are more consistent with an older lineage close to equilibrium. Interestingly, if we combined Venezuelan lineages (B and C, excluding the single D1 haplotype), they depicted two different peaks in the mismatch distribution, corresponding to cis- and trans-Andean populations (395,821-327,737 years ago, respectively). The IBD test may be showing positive correlation with the distance as an artifact, because of the demographic expansion of the populations included in the analysis.

A McDonald-Kreitman (MK) test was performed for each lineage to detect departures from neutral theory at the molecular level. There was no evidence of selection on either *COI* or *white* genes in several locations tested, using different species of *Anopheles* as outgroups.

Discussion

Taxonomic status of the *Anopheles triannulatus* complex

This study assessed the phylogenetic relationship of the *An. triannulatus* complex based on mitochondrial and


Table 2 Description of shared COI haplotypes for the *An. triannulatus* complex

Site no.	Country	Locality	N	Haplotypes	Lineages
1	Argentina	Yuto Farm (FY)	23	E1 ,E2,E4,E15, E44 (2),E45,E46,E47, E50 (2),E51, E52 , E57, E66(3), E67 (4),E68,E69	E
2		Lake Yuto (LY)	3	E58 , E67 ,E70	E
3	Bolivia	Guayaramerin (GU)	4	E32(4)	E
4		Puerto Villaroel (PV)	10	E3,E5,E6,E12(2),E13,E49,E62,E63(2)	E
5	Brazil	Ceara (CE)	2	D21,D22	D
6		Espirito Santo (ES)	1	E53	E
7		Fazenda S. Joao (MT)	2	E61, E8	E
8		Monte Negro (RO)	5	E22, E23, E59 (2), E65	E
9		Aquidauana (Pantanal do Rio Negro) (AQU)	6	D23,E17,E18,E19,E20,E21	D,E
10		Inubia Paulista (SP)	6	E1 ,E28,E29,E30, E48 , E58	E
11		Camacan/Santa Luzia (Bahia) (BA)	7	G3,G4,G5,G6,G7,G8,G9	G
12		Coronel Pacheco (Minas Gerais) (MG)	6	G10,G11,G12,G13,G14,E31	G,E
13		Lagoa da Confusao (Tocantins) (TO)	6	E24, E25, E26, E27, E56 , E65	E
14		Itaituba (ITB)	5	E16, E50 , E56 ,E64, E65	E
15		Oswaldo Cruz (OC)	19	D1 ,D5,D6,D7,D8, D12 (7), D13(2),D14(2),D15, D16, C6	C,D
16		Salobra (MS)	20	E1 , E7 , E8 (2),E9,E10,E11,E14(2), E44 , E48 , E52 (4), E53 ,E54,E55, E59 ,E60	E
17		Silva Jardim (RJ)	2	G1,G2	G
18		Tartarugalzinho (TAR)	23	D2,D3,D4,D9,D10(4),D11, D12 (4),D17,D18(4), D19(2),D20, E43(2)	D, E
19	Colombia	Monitos (MO)	23	B3 , F1 ,F2, F3, F4 (2), F8 , F9 , F11 (9),F12(2),F13,F14,F16,F17	B,F
20		Santa Rosa de Lima (SO)	27	B18, F1 , F4 (6),F5(2),F6,F7, F8 , F9 (3), F10, F11 (6),F15, F18,F19,F20	B,F
21		Leticia, km12 (COSW)	10	E36,E37(4),E38,E39,E40,E41,E42	E
22		Tibu (COL)	3	B14 ,B19,C11	B,C
23	Ecuador	Juan Montalvo (JM)	6	E33(4), E34, E35	E
24	Panama	Bayano (BAY)	7	A3 , A4 , A9 ,A14, A16 (2), A18	A
25		Gamboia (GAM)	29	A1,A2, A3 , A4 (3),A5,A6,A7,A8, A9 (10),A10 (4) A11,A12,A13, A16 ,A17	A
26		Meteti (MET)	17	A4 (8), A9 (5), A14,A15, A16 (2), A19	A
27	Venezuela	Boconoito (BOC)	6	B1, B2 , B3 , B8(2), B9	B
28		Cano Amarillo (CAM)	15	B2 , B3 , B4, B5, B9 ,B10(2), B11, B12(2), B13, B14 (4)	B
29		Casigua Zulia (CAS)	25	D1 ,C1(4),C2,C3(2),C4,C5, C6 (10),C7,C8,C9(2),C10	D,C
30		La Veguita (VG)	6	B6, B7 ,B15,B16,B17,B18	B
31	Trinidad-Tobago	St.Andrew/St.David, Valencia (TRI)	2	B7 (2)	B

The number in parentheses indicates the frequency of the haplotype at that site; bold numbers are shared haplotypes, plain numbers are unique haplotypes in that population and underlined numbers are *An. halophylus* (E8, E53) and *An. triannulatus* C (E7, E9).

nuclear data. Results of phylogenetic analysis showed that *An. halophylus* and *An. triannulatus* C do not form separate monophyletic clades, indicating that they are probably very closely related or are incipient species as proposed by Silva-do-Nascimento and collaborators [20,21]. The COI uncorrected pairwise genetic distances detected a 0.2% divergence between these putative taxa,

whereas with those identified as *An. triannulatus* s.s. the range varied from 1.7-2.3%, depending on the sample. Therefore, this study supports the conclusion that *An. halophylus* and *An. triannulatus* C are more closely related to each other than either is to *An. triannulatus* s.s. [20,21]. Furthermore, the combined three gene analysis recovered a highly supported single clade consisting of



An. halophylus and *An. triannulatus* C, whereas the mitochondrial *COI* gene provided a tree with lower resolution (short branches, low support) for the three species.

Diverse processes in the mitochondrial marker such as introgression or incomplete lineage sorting may be responsible for the inconclusive phylogenetic analysis within the complex. In addition, representatives from Argentina and Mato Grosso (Brazil) were recovered in every tree topology clustered together with *An. halophylus* and *An. triannulatus* C, and may represent a more extensive distribution of this taxa. Actually, specimens morphologically similar to *An. halophylus* and *An. triannulatus* C were reported in Western Brazilian Amazon, Paraguay and Bolivia [15,19,22,29,67].

Evidence of shared haplotypes, i.e. between *An. halophylus* and *An. triannulatus* s.s. and between *An. triannulatus* s.s. and *An. triannulatus* C were also found with this gene. The presence or absence of the intron in the *white* gene is an evolutionary event that appears more than once in the history of anophelines [68,69]. Within the *An. triannulatus* complex, all three species

retained this intron and there were no significant nucleotide differences among their sequences.

The divergence of ITS2 lineages was very low within (0.2% in lineage I and II and 0 in lineage III) and between lineages (0.7% between I and II; 1% and 1.1% between I and III and II and III, respectively). Therefore, the ITS2 findings alone did not conclusively differentiate *An. halophylus* and *An. triannulatus* C, which suggest that a low level of gene flow may occur or have taken place in the recent past, as it has been proposed when analyzing sympatric specimens from Brazil with sequences of *timeless* gene alone [21]. On the other hand, the analysis of *cpr* gene sequences of the same specimens revealed fixed differences and considerable genetic differentiation between *An. halophylus* and *An. triannulatus* C [21].

The phylogenetic reconstruction supports monophyly of the *An. triannulatus* complex. The three genes detected different phylogenetic relationships as well as phylogeographic patterns in this complex. Nuclear DNA markers are expected to provide older demographic

Table 3 Analysis of Molecular Variance (AMOVA) using COI in the *An. triannulatus* complex

Groups (localities)	Among groups	Among populations within groups	Within populations
1)	53.81	3.24	42.95
I: 24, 25, 26	$\Phi_{CT} = 0.53^{***}$	$\Phi_{ST} = 0.57^{***}$	$\Phi_{SC} = 0.07^{***}$
II: 19,20			
III: 5, 15, 18			
IV: 29			
V: 22, 27, 28, 30, 31			
VI: 23			
VII: 11, 12, 17			
VIII: 21			
IX: 1,2,3,4,6,7,8,9,10,13,14,16			
2)	53.13	3.90	42.96
VI: 21,23	$\Phi_{CT} = 0.531^{***}$	$\Phi_{ST} = 0.57^{***}$	$\Phi_{SC} = 0.08^{***}$
3)	47.11	10.45	42.44
II: 19,20,29	$\Phi_{CT} = 0.47^{***}$	$\Phi_{ST} = 0.57^{***}$	$\Phi_{SC} = 0.19^{***}$
4)	37.49	19.18	43.33
IV: 22, 27, 28, 29, 30, 31	$\Phi_{CT} = 0.37^{***}$	$\Phi_{ST} = 0.56^{***}$	$\Phi_{SC} = 0.3^{***}$

1) Groups (localities number) derived from SAMOVA k = 9; 2), 3) and 4) with modifications in the groups based on SAMOVA results. ***p < 0.001.

information than mtDNA because dissimilar effective population size can affect estimations of coalescent time [70]. Therefore, the genetic relationships that we obtained with the *white* gene support an original metapopulation, across the entire known geographic distribution, with posterior divergence in some populations, reflected in the results of the *COI* gene. The mitochondrial loci in most species have a shorter expected coalescence time compared with nuclear loci (only one-fourth of the effective population size), and thus there is a greater probability that the mitochondrial gene tree will accurately reflect the species tree [71].

On the other hand, the ITS2 marker did not show different lengths between putative species, but see [44]. The characteristic mutation replacement of this locus (elevated

number of indels and low frequency of replacement mutations) may be more useful for species-specific PCR diagnostics in species complex members [72,73] than for phylogenetic inference.

The *COI* fragment was useful to describe the genetic structure of the *An. triannulatus* complex. However, the genetic variation and divergence within and between lineages might reach a different conclusion if this study had been based on the more conservative “Folmer region” (DNA barcode standard) [74].


Divergence and demographic expansion of the *Anopheles triannulatus* complex

Anopheles triannulatus s.l. contains cryptic and geographically distinct mtDNA lineages, indicative of either

Table 4 Molecular diversity and neutrality test in the main mt-DNA lineages within the *Anopheles triannulatus* complex

COI gene lineage	N	h	π	Hd (SD)	k	F _s	F*	D*	D _T	R ²
A	53	18	0.00275	0.841 (0.035)	1.894	-11.706***	-3.223**	-3.205*	-1.802*	0.162***
B	33	19	0.00776	0.960 (0.018)	5.348	-8.354**	-1.492	-1.29	-1.21	0.160***
C	26	11	0.00457	0.806 (0.071)	3.148	-2.558	0.106	-0.0009	0.305	0.126***
D	43	23	0.00628	0.921 (0.030)	4.328	-11.479***	-1.373	-0.981	-1.483	0.109***
E	109	70	0.0194	0.989 (0.003)	13.383	-41.832***	-1.972	-1.999	-1.173	0.090***
F	48	20	0.00760	0.906 (0.029)	5.238	-6.361*	-2.775*	-2.748*	-1.575	0.109***
G	14	14	0.00636	1 (0.027)	4.385	-11.622***	-1.588	-1.429	-1.265	0.151***
All	326	175	0.0242	0.989 (0.0017)	16.676	-158.48***	-1.924	-2.334*	-0.914	0.075***

A: Panama; B: cis-Andean Venezuela + Colombia + Trinidad-Tobago; C: Venezuela; D: NE and C Brazil + Venezuela; E: Ecuador, Colombia, Bolivia, Argentina, Brazil; F: Colombia; G: SE Brazil; N: sample size; h: number of haplotypes; π : nucleotide diversity; Hd: Haplotype diversity (SD: Standard deviation); k: average number of pairwise differences; F_s: Fu's F_s statistic; F*: Fu and Li F test; D*: Fu and Li's D test; D_T: Tajima's D; R²: Ramos-Onsins & Rozas R² test. *p < 0.05; **p < 0.02; ***p < 0.001.



speciation or substantial isolation and divergence among populations. The most common process of mosquito speciation has been considered to be the allopatric mode ([72] and references therein). Although the Amazon delta has been hypothesized as a natural barrier for populations of *An. triannulatus* and *An. darlingi* [5,43], in the current study populations of *An. triannulatus* located on both shores were part of a single lineage with shared haplotypes for both the mitochondrial and the nuclear markers. We performed AMOVA analysis with our data based on the recent findings of Pedro and collaborators [5] to test

Amazonian groups of *An. triannulatus*. The five groups suggested in the cited study only contributed 21.64% to the variance, whereas our 3 groupings (TAR + CE + OC; MG + RJ + BA; and the remaining Amazonian populations yielded a significant variance of 41.27% ($p < 0.01$).

The high differentiation among populations in the Caribbean Andes region (Colombia-Venezuela, cis-trans Andean) suggests that the Sierra Nevada (west of Lake Maracaibo, Venezuela) acts as a partial barrier to gene flow for *COI* and *white* gene, and as a complete barrier for ITS2. The Andean orogeny seems to be responsible

for this large source of variation, promoting isolation and secondary contact among different lineages in several taxa such as butterflies, birds, bees, sandflies [75], including neotropical anophelines such as *An. darlingi* [3], *An. albimanus* [76] and *An. nuneztovari* [43]. Furthermore, the bimodal *COI* mismatch distribution for all Venezuelan populations supports the hypothesis of two independent migration events [77]. The current study identified the Serranía de Perija in the Eastern Andes cordillera (between Venezuela and Colombia) as a porous genetic barrier to *An. triannulatus*, with some mtDNA haplotypes shared as well as some *white* gene exchange between cis-trans Andean populations. The ITS2 marker depicted the eastern Sierra Nevada Venezuelan populations as more closely related to those from Brazil, Ecuador and Bolivia.

Taking into account lineage divergence estimates in this study, climatic changes during the Pleistocene could have influenced this isolation, creating refugia on both sides of the mountains [2], and might explain better the lineage distributions than vicariant events associated with the older uplift of the North Western Andes (from Late Cretaceous to Holocene) [78]. The absence of shared *COI* haplotypes may indicate lack of gene flow between some of the lineages. Although the Mantel test showed a positive significant IBD pattern in this area, the results may be confounded by the recent expansion events [79].

In southern South America, the Coastal Mountain Range in SE Brazil promotes population structure in *An. darlingi* [80], and incipient speciation in *Anopheles cruzii* [81]. In fact, an extra-Andean glaciation event led to a Pleistocene refugium in Mount Itatiaia and the Serra do Mar [82]. Restricted gene flow and genetic structure of *An. triannulatus* *COI* lineage G in this region support a common phylogeographic pattern in mosquitoes and in some other similarly distributed species, such as sandflies [83]. An alternative explanation is that the most extensive *An. triannulatus* lineage E (Figure 6) could be interpreted as evidence of a Pleistocene range expansion combined with genetic exchange.

Although the origin of the *An. triannulatus* complex has been hypothesized to be south of the Amazon River [5], probably in the Belém or Tapajós refugia [84], our data provide mixed signals. On one hand, both *COI* and *white* gene coalescent and phylogenetic analyses found the most likely ancestral sequences to be in northwestern South America (Colombia, Panama, and/or Venezuela). On the other hand, the lowest nucleotide diversity values were found in these populations, and higher values in localities from the Amazon Basin. The low π value in Panamanian locations has been hypothesized in the malaria vector *An. albimanus* to be the result of a past bottleneck event followed by a demographic expansion dated approximately

~22,000 years ago [85], although recent bottlenecks can also mimic the effects of an expansion [86]. Panamanian populations of *An. triannulatus* included in the present study seem to have a similar demographic history and estimation dates for the proposed expansion (~ 23,000 years ago). Besides, the least polymorphic population may reflect a smaller effective size and relative isolation of a population. However, populations of *An. triannulatus* from the Amazon Basin seem to be in equilibrium, whereas in those from the Amazon Delta a demographic expansion around 15,043 years ago was detected (4265–99,387, 95% CI).

Conclusions

Many malaria control programs focus on local vector management and the incrimination of species involved in malaria transmission is crucial. Molecular taxonomy can accurately identify malaria vectors [87]. Phylogeography has been informative in assessing historic migration and colonization routes for *An. albimanus* [88] and can also be used to trace the origin of accidental introductions such as *Aedes albopictus* [89]. Because *An. triannulatus* has been reportedly incriminated in malaria transmission in eastern Amazonian Peru and Amazonian Brazil and Venezuela [24,26,27,90] we hypothesize that at least lineage E is likely a malaria vector.

Additional file

Additional file 1: Numbers of transitions (X and s) and transversions (Δ and v) at each codon position plotted against Tamura-Nei (Tamura and Nei, 1993) genetic distances for the mitochondrial *COI* gene. The steeper slope of the transitions in nt3 suggests no substitution saturation and is evidence for a constant rate of evolution.

Competing interests

The authors declare they have no competing interests.

Authors' contributions

MM, SB, WH, JH and SNM participated in the design of the study, performed molecular and genetic analysis and wrote the manuscript. TFSN, JRL, FR, RLO, MAMS, ESB, GNF, RW, YML, MJDJ, YR, MMP, LAG, MMC provided samples used in this study, advised on the analysis and assisted and helped to draft the manuscript. JEC conceived and supervised the study, and assisted in the writing of the manuscript. All authors read and approved the final manuscript.

Acknowledgments

This project has been financially supported by NIH grant AI ROI 54139–02 to JE Conn, PAPES IV Fiocruz-CNPq to RLO, and from the Fundação de Amparo à Pesquisa do Estado de São Paulo, FAPESP (Grant 2011/20397-7 to MAMS), CNPq (BPP no. 301666/2011-3 to MAMS).

Disclaimer

This research was performed under a Memorandum of Understanding between the Walter Reed Army Institute of Research and the Smithsonian Institution, with institutional support provided by both organizations. The material to be published reflects the views of the authors and should not be construed to represent those of the Department of the Army or the Department of Defense.

Author details

¹New York State Department of Health, Wadsworth Center, Griffin Laboratory, Albany, NY, USA. ²Department of Biomedical Sciences, School of Public Health, State University of New York, Albany, NY, USA. ³Departamento de Entomologia, Instituto Oswaldo Cruz-Fiocruz, Rio de Janeiro, Brazil. ⁴Instituto de Investigaciones Científicas y Servicios de Alta Tecnología, Clayton, Panamá, República de Panamá. ⁵Division of Entomology, Walter Reed Army Institute of Research, Silver Spring, MD, USA. ⁶Departamento de Epidemiologia, Faculdade de Saúde Pública, Universidade de São Paulo, São Paulo, Brazil. ⁷Superintendência de Controle de Endemias, SUCEN, São Paulo, Brazil. ⁸Department of Biological Sciences, Eastern Illinois University, Charleston, IL, USA. ⁹Instituto Superior de Entomología "Dr. Abraham Willink", Facultad de Ciencias, Naturales e Instituto Miguel Lillo, Universidad Nacional de Tucumán, Tucumán, Argentina. ¹⁰Laboratorio de Biología de Vectores, Instituto de Zoología y Ecología Tropical, Universidad Central de Venezuela, Caracas, Venezuela. ¹¹Instituto Evandro Chagas, Seção de Parasitologia, Belém, Brazil. ¹²Grupo de Microbiología Molecular, Escuela de Microbiología, Universidad de Antioquia, Medellín, Colombia. ¹³Present address: Division Infectious Diseases University of California San Diego, George Palade Labs, School of Medicine, 9500 Gilman Drive, MC 0741, La Jolla, CA 92093, USA.

Received: 26 November 2012 Accepted: 13 February 2013

Published: 22 February 2013

References

- Grimaldi D, Engel MS: *Evolution of the insects*. Cambridge: Massachusetts, Cambridge University Press; 2005.
- Haffer J: *Speciation in Amazonian forest birds*. *Science* 1969, **165**:131–137.
- Conn JE, Mirabello L: *The biogeography and population genetics of neotropical vector species*. *Heredity* 2007, **99**:245–256.
- Moritz C, Patton JL, Schneider CJ, Smith TB: *Diversification of rainforest faunas: an integrated molecular approach*. *Annu Rev Ecol Syst* 2000, **31**:533–563.
- Pedro PM, Uezu A, Sallum MA: *Concordant phylogeographies of 2 malaria vectors attest to common spatial and demographic histories*. *J Hered* 2010, **101**:618–627.
- Loaiza JR, Bermingham E, Sanjurjo OI, Scott ME, Bickersmith SA, Conn JE: *Review of genetic diversity in malaria vectors (Culicidae: Anophelinae)*. *Infect Genet Evol* 2012, **12**:1–12.
- de Queiroz K: *Species concepts and species delimitation*. *Syst Biol* 2007, **56**:956–974.
- Neiva A, Pinto C: *Considerações sobre o gênero Cellia theobald, com a descrição de uma nova espécie*. *Brazil Méd* 1922, **36**:355–357.
- Faran ME: *Mosquito studies (Diptera, Culicidae). XXXIV. A revision of the albimanus section of the subgenus Nyssorhynchus of Anopheles*. *Contrib Amer Entomol Inst* 1980, **15**:1–215.
- Faran ME, Linthicum KJ: *A handbook of the Amazonian species of Anopheles (Nyssorhynchus) (Diptera: Culicidae)*. *Mosq Syst* 1981, **13**:1–81.
- Chadee DD, Wilkerson RC: *Anopheles triannulatus (Neiva and Pinto): a new Anopheles record from Trinidad, West Indies*. *J Am Mosq Control Assoc* 2005, **21**:316–317.
- Pinto C: *Disseminação da malária pela aviação. Biologia do Anopheles gambiae e outros anofelinos no Brasil*. *Mem Inst Oswaldo Cruz* 1939, **34**:293–430.
- Galvão ALA: *Contribuição ao conhecimento dos anofelinos do grupo Nyssorhynchus de São Paulo e regiões vizinhas (Diptera: Culicidae)*. *Arq Zool Sao Paulo* 1940, **1**:399–484.
- Galvão ALA, Lane J: *Observações sobre alguns anofelinos de Salobra, Mato Grosso (Diptera, Culicidae)*. *Rev Biol Hyg* 1941, **11**:10–18.
- Deane LM, Causey OR, Deane MP: *Chave ilustrada para a identificação de ovos de trinta espécies de anofelinos brasileiros, com algumas descrições novas*. *Rev Serv Esp Saúde Públ* 1947a, **1**:293–308.
- Deane LM, Causey OR, Deane MP: *Chave ilustrada para a identificação de trinta e cinco espécies de anofelinos das regiões Nordeste e Amazônica do Brasil pelos caracteres da fêmea, com notas sobre os transmissores da malária (Diptera: Culicidae)*. *Rev Serv Esp Saúde Públ* 1947b, **1**:309–336.
- Forattini OP: *Entomologia médica*, Volume 1. São Paulo, Brazil: Univ São Paulo; 1962.
- Rosa-Freitas MG, Lourenço-de-Oliveira R, Pinto JCC, Flores-Mendoza C, Nascimento TFS: *Anopheline species complexes in Brazil. Current knowledge of those related to malaria transmission*. *Mem Inst Oswaldo Cruz* 1998, **93**:651–655.
- Silva-do-Nascimento TF, Lourenço-de-Oliveira R: *Anopheles halophylus, a new species of the subgenus Nyssorhynchus (Diptera: Culicidae) from Brazil*. *Mem Inst Oswaldo Cruz* 2002, **97**:801–811.
- Silva-do-Nascimento TF, Wilkerson RC, Lourenço-de-Oliveira R, Monteiro FA: *Molecular confirmation of the specific status of Anopheles halophylus (Diptera: Culicidae) and evidence of a new cryptic species within An. Triannulatus in central Brazil*. *J Med Entomol* 2006, **43**:455–459.
- Silva-do-Nascimento TF, Pitaluga LD, Peixoto AA, Lourenço-de-Oliveira R: *Molecular divergence in the timeless and cpr genes among three sympatric cryptic species of the Anopheles triannulatus complex*. *Mem Inst Oswaldo Cruz* 2011, **106**:218–222.
- Silva-do-Nascimento TF, Lourenço-de-Oliveira R: *Diverse population dynamics of the three Anopheles species belonging to the triannulatus complex (Diptera: Culicidae)*. *Mem Inst Oswaldo Cruz* 2007, **102**:975–982.
- Dixon K, Roberts DR, Llewellyn CH: *Contribuição ao estudo epidemiológico da malária em trechos da rodovia transamazônica, Brasil*. *Rev Inst Med Trop São Paulo* 1979, **21**:287–292.
- Galardo A, Arruda M, D'almeida A, Wirtz R, Lounibos P, Zimmerman R: *Malaria vector incrimination in three rural riverine village in the Brazilian Amazon*. *AmJTrop Med Hyg* 2007, **76**:461–469.
- Rosa-Freitas M, Tsouris P, Peterson A, Honorio NA, de Barros FS, de Aguiar DB, Gurgel Hda C, de Arruda ME, Vasconcelos SD, Luitgards-Moura JF: *An ecoregional classification for the state of Roraima, Brazil: the importance of landscape in malaria biology*. *Mem Inst Oswaldo Cruz* 2007, **102**:349–357.
- Tadei WP, Dutary-Thatcher B: *Malaria vectors in the Brazilian Amazon: Anopheles of the subgenus Nyssorhynchus*. *Rev Inst Med Trop S Paulo* 2000, **42**:87–94.
- Aramburú GJ, Ramal AC, Witzig R: *Malaria re-emergence in the Peruvian Amazon region*. *Emerg Infect Dis* 1999, **5**:209–215.
- Benarroch EL: *Studies of malaria in Venezuela*. *Am J Hyg* 1931, **14**:690–693.
- Deane LM, Causey OR, Deane MP: *Notas sobre a distribuição e a biologia dos anofelinos das regiões Nordeste e Amazônica do Brasil*. *Rev Serv Esp Saúde Públ* 1948, **1**:827–965.
- Lourenço-de-Oliveira R, Guimarães AE, Arlê M, Silva TF, Castro MG, Motta MA, Deane LM: *Anopheline species, some of their habits and relation to malaria in endemic areas of Rondônia state, Amazon region of Brazil*. *Mem Inst Oswaldo Cruz* 1989, **84**:501–514.
- Lourenço-de-Oliveira R, Luz SLB: *Simian malaria at two sites in the Brazilian Amazon. II. Vertical distribution and frequency of anopheline species inside and outside the forest*. *Mem Inst Oswaldo Cruz* 1996, **91**:687–694.
- Zimmerman RH, Galardo AK, Lounibos LP, Arruda M, Wirtz R: *Bloodmeal hosts of Anopheles species (Diptera: Culicidae) in a malaria-endemic area of the Brazilian Amazon*. *J Med Entomol* 2006, **43**:947–956.
- Matson R, Rios CT, Chavez CB, Gilman RH, Florin D, Sifuentes VL, Greffa RC, Yori PP, Fernandez R, Portocarrero DV, Vinetz JM, Kosek M: *Improved molecular technique for the differentiation of neotropical anopheline species*. *AmJTrop Med Hyg* 2008, **78**:492–498.
- Avice JC: *Phylogeography: The history and formation of species*. Massachusetts: Harvard University Press; 2000.
- Paredes-Esquivel C, Donnelly MJ, Harbach RE, Townson H: *A molecular phylogeny of mosquitoes in the Anopheles barbirostris subgroup reveals cryptic species: implications for identification of disease vectors*. *Mol Phy Evol* 2009, **50**:141–151.
- Gutierrez LA, Gomez GF, Gonzales JJ, Castro MI, Luckhart S, Conn JE, Correa MM: *Microgeographic genetic variation of the malaria vector Anopheles darlingi (Diptera: Culicidae) from Córdoba and Antioquia, Colombia*. *AmJTrop Med Hyg* 2010, **83**:38–47.
- Krzywinski J, Besansky NJ: *Molecular systematics of Anopheles: from subgenera to subpopulations*. *Annu Rev Entomol* 2003, **48**:111–139.
- Bourke BP, Foster PG, Berge ES, Calado DC, Sallum MA: *Phylogenetic relationships among species of Anopheles (Nyssorhynchus) (Diptera, Culicidae) based on nuclear and mitochondrial gene sequences*. *Acta Trop* 2010, **114**:88–96.
- Bower JE, Downton M, Cooper RD, Beebe NW: *Intraspecific concerted evolution of the rDNA ITS1 in Anopheles farauti sensu stricto (Diptera: Culicidae) reveals recent patterns of population structure*. *J Mol Evol* 2008, **67**:397–411.

40. Zapata MA, Cienfuegos AV, Quiros OI, Quinones ML, Luckhart S, Correa MM: **Discrimination of seven *anopheles* species from San Pedro de uraba, Antioquia, Colombia, by polymerase chain reaction-restriction fragment length polymorphism analysis of its sequences.** *AmJTrop Med Hyg* 2007, **77**:67–72.
41. Lunt DH, Zhang DX, Szymura JM, Hewitt GM: **The insect cytochrome oxidase I gene: evolutionary patterns and conserved primers for phylogenetic studies.** *Insect Mol Biol* 1996, **5**:153–165.
42. Mirabello L: *Molecular population genetics of the malaria vector anopheles darlingi throughout central and south America using mitochondrial, nuclear and microsatellites markers.* State University of New York at Albany, Biomedical Sciences Department: Ph.D. thesis; 2007.
43. Mirabello L, Conn JE: **Population analysis using the nuclear *white* gene detects Pliocene/Pleistocene lineage divergence within *anopheles nuneztovari* in south America.** *Med Vet Entomol* 2008, **22**:109–119.
44. Li C, Wilkerson RC: **Identification of *anopheles (nyssorhynchus) albittarsis* complex species (diptera: culicidae) using rDNA internal transcribed spacer 2-based polymerase chain reaction primers.** *Mem Inst Oswaldo Cruz* 2005, **100**:495–500.
45. Hall TA: **BioEdit: a user-friendly biological sequence alignment editor and analysis program for windows 95/98/NT.** *Nucl Acids Symp Ser* 1999, **41**:95–98.
46. Posada D: **JModelTest: phylogenetic model averaging.** *Mol Biol Evol* 2008, **25**:1253–1256.
47. Huelsenbeck JP, Ronquist F: **MRBAYES: Bayesian inference of phylogenetic trees.** *Bioinformatics* 2001, **17**:754–755.
48. Brandley MC, Schmitz A, Reeder TW: **Partitioned Bayesian analyses, partition choice, and the phylogenetic relationships of scincid lizards.** *Syst Biol* 2005, **54**:373–390.
49. Librado P, Rozas J: **DnaSP v. 5: a software for comprehensive analysis of DNA polymorphism data.** *Bioinformatics* 2009, **25**:1451–1452.
50. Dupanloup I, Schneider S, Excoffier L: **A simulated annealing approach to define the genetic structure of populations.** *Mol Ecol* 2002, **11**:2571–2581.
51. Excoffier L, Smouse PE, Quattro JM: **Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data.** *Genetics* 1992, **131**:479–491.
52. Jensen JL, Bohonak AJ, Kelley ST: **Isolation by distance, web service.** *BMC Genet* 2005, **6**:13. v.3.16.
53. Ersts PJ: *Geographic distance matrix generator (version 1.2.3):* American Museum of Natural History, Center for Biodiversity and Conservation; http://biodiversityinformatics.amnh.org/open_source/gdmg.
54. Smouse PE, Long JC, Sokal RR: **Multiple regression and correlation extensions of the mantel test of matrix correspondence.** *Syst Zool* 1986, **35**:627–632.
55. Bandelt HJ, Forster P, Röhl A: **Median-joining networks for inferring intraspecific phylogenies.** *Mol Biol Evol* 1999, **16**:37–48.
56. Templeton AR, Crandall KA, Sing CF: **A cladistic analysis of phenotypic associations with haplotypes inferred from restriction endonuclease mapping and DNA sequence data III. Cladogram estimation.** *Genetics* 1992, **132**:619–633.
57. Clement M, Posada D, Crandall KA: **TCS: a computer program to estimate gene genealogies.** *Mol Ecol* 2000, **9**:1657–1660.
58. Powell JR, Caccone A, Amato GD, Yoon C: **Rates of nucleotide substitution in *Drosophila* mitochondrial DNA and nuclear DNA are similar.** *Proc Natl Acad Sci USA* 1986, **83**:9090–9093.
59. Heckel G, Burri R, Fink S, Desmet JF, Excoffier L: **Genetic structure and colonization processes in European populations of the common vole, *Microtus arvalis*.** *Evolution* 2005, **59**:2231–2242.
60. Brower AV: **Rapid morphological radiation and convergence among races of the butterfly *Heliconius erato* inferred from patterns of mitochondrial DNA evolution.** *Proc Natl Acad Sci USA* 1994, **91**:6491–6495.
61. Excoffier L, Laval G, Schneider S: **Arlequin ver. 3.0: an integrated software package for population genetics data analysis.** *Evol Bioinform* 2005, **1**:47–50.
62. Sallum MAM, Schultz TR, Foster PG, Aronstein K, Wirtz RA, Wilkerson RC: **Phylogeny of Anophelinae (Diptera: Culicidae) based on nuclear ribosomal and mitochondrial DNA sequences.** *Syst Entomol* 2002, **27**:361–382.
63. Gleeson DM, Rowell DM, Tait NN, Briscoe DA, Higgins AV: **Phylogenetic relationships among Onychophora from Australasia inferred from the mitochondrial cytochrome oxidase subunit I gene.** *Mol Phylogenet Evol* 1998, **10**:237–248.
64. Xia X, Xie Z, Salemi M, Chen L, Wang Y: **An index of substitution saturation and its application.** *Mol Phylogenet Evol* 2003, **26**:1–7.
65. Castollee J, Templeton AR: **Root probabilities for intraspecific gene trees under neutral coalescent theory.** *Mol Phylogenet Evol* 1994, **3**:102–113.
66. Nylander JAA, Ronquist F, Huelsenbeck JP, Nieves Aldrey JL: **Bayesian phylogenetic analysis of combined data.** *Syst Biol* 2004, **53**:47–67.
67. Root FM: **Studies on Brazilian mosquitoes. I. Anophelines of the *Nyssorhynchus* group.** *Am J Hyg* 1926, **6**:684–717.
68. Krzywinski J, Besansky NJ: **Frequent intron loss in the *white* gene: a cautionary tale for phylogeneticists.** *Mol Biol Evol* 2002, **19**:362–366.
69. Merritt TJ, Young CR, Vogt RG, Wilkerson RC, Quattro JM: **Intron retention identifies a malaria vector within the *Anopheles* (Nyssorhynchus) albittarsis complex (Diptera: Culicidae).** *Mol Phylogenet Evol* 2005, **35**:719–724.
70. Avise JC: *Molecular markers, natural history, and evolution.* Massachusetts: Second ed Sinauer; 2004.
71. Moore WS: **Inferring phylogenies from mtDNA variation: mitochondrial gene trees versus nuclear gene trees.** *Evolution* 1995, **49**:718–726.
72. Morgan K, O'Loughlin SM, Mun-Yik F, Linton YM, Somborn P, Min S, Htun PT, Nambanya S, Weerasinghe I, Sochanta T, Prakash A, Walton C: **Molecular phylogenetics and biogeography of the neocellia series of *Anopheles* mosquitoes in the oriental region.** *Mol Phylogenet Evol* 2009, **52**:588–601.
73. Dyer NA, Lawton SP, Ravel S, Choi KS, Lehane MJ, Robinson AS, Okedi LM, Hall MJ, Solano P, Donnelly MJ: **Molecular phylogenetics of tsetse flies (Diptera: Glossinidae) based on mitochondrial (COI, 16S, ND2) and nuclear ribosomal DNA sequences, with an emphasis on the palpalis group.** *Mol Phylogenet Evol* 2008, **49**:227–239.
74. Hebert PDN, Cywinska A, Ball SL, deWaard JR: **Biological identifications through DNA barcodes.** *Proc R Soc Lond B Biol Sci* 2003, **270**:313–321.
75. Arrivillaga JC, Norris DE, Feliciangeli MD, Lanzaro GC: **Phylogeography of the neotropical sand fly *Lutzomyia longipalpis* inferred from mitochondrial DNA sequences.** *Infect Genet Evol* 2002, **2**:83–95.
76. Molina-Cruz A, de Mérida AM, Mills K, Rodríguez F, Schoua C, Yurrita MM, Molina E, Palmieri M, Black WC 4th: **Gene flow among *Anopheles albimanus* populations in central America, south America, and the Caribbean assessed by microsatellites and mitochondrial DNA.** *AmJTrop Med Hyg* 2004, **71**:350–359.
77. Hasan AU, Suguri S, Fujimoto C, Itaki RL, Harada M, Kawabata M, Bugoro H, Albino B, Tsukahara T, Hombhanje F, Masta A: **Phylogeography and dispersion pattern of *Anopheles farauti* sensu stricto mosquitoes in Melanesia.** *Mol Phylogenet Evol* 2008, **46**:792–800.
78. Gregory-Wodzicki KM: **Uplift history of the central and northern Andes: a review.** *Geol Soc Am Bull* 2000, **112**:1091–1105.
79. Telles MPC, Diniz-Filho JAF: **Multiple mantel tests and isolation by distance, taking into account long-term historical divergence.** *Gen Mol Res* 2005, **4**:742–748.
80. Pedro P, Sallum MAM: **Spatial expansion and population structure of the neotropical malaria vector, *Anopheles darlingi*.** *Biol J Linn Soc* 2009, **97**:854–866.
81. Rona LD, Carvalho-Pinto CJ, Peixoto AA: **Molecular evidence for the occurrence of a new sibling species within the *Anopheles (kerteszia) cruzii* complex in south-east Brazil.** *Malaria J* 2010, **26**:9–33.
82. Müller P: *Die herpetofauna der Insel von São Sebastião (Brasilien).* Saarbrücken: Verlag Saarbrücker Zeitung; 1968:1–82.
83. Queiroz Balbino V, Coutinho-Abreu IV, Sonoda IV, Melo MA, de Andrade PP, de Castro JA, Rebêlo JM, Carvalho SM, Ramalho-Ortigão M: **Genetic structure of natural populations of the sand fly *Lutzomyia longipalpis* (Diptera: Psychodidae) from the Brazilian northeastern region.** *Acta Trop* 2006, **98**:15–24.
84. Prance GT: **Forest refuges: evidence from woody angiosperms.** In *Biological diversification in the tropics.* Edited by Prance GT. New York, USA: Columbia University Press; 1982:137–157.
85. Loaiza JR, Scott ME, Bermingham E, Rovira J, Conn JE: **Evidence for Pleistocene population divergence and expansion of *Anopheles albimanus* in southern central America.** *AmJTrop Med Hyg* 2010, **82**:156–164.
86. Tajima F: **Statistical-method for testing the neutral mutation hypothesis by DNA polymorphism.** *Genetics* 1989, **123**:585–595.
87. Cienfuegos AV, Rosero DA, Naranjo N, Luckhart S, Conn JE, Correa MM: **Evaluation of a PCR-RFLP-ITS2 assay for discrimination of *Anopheles* species in northern and western Colombia.** *Acta Trop* 2011, **118**:128–133.
88. Loaiza JR, Scott ME, Bermingham E, Sanjurjo OI, Wilkerson R, Rovira J, Gutiérrez LA, Correa MM, Grijalva MJ, Birnberg L, Bickersmith S, Conn JE: **Late Pleistocene environmental changes lead to unstable demography**

and population divergence of *anopheles albimanus* in the northern neotropics. *Mol Phylogenet Evol* 2010, **57**:1341–1346.

89. Urbanelli S, Bellini R, Carrieri M, Sallicandro P, Celli G: Population structure of *aedes albopictus* (skuse): the mosquito which is colonizing Mediterranean countries. *Heredity* 2000, **84**:331–337.
90. Gabaldon A, Cova Garcia P: Zoogeografía de los anofelinos en Venezuela: I. Los dos vectores secundarios y los no vectores. *Tijeretazos Sobre Malar* 1946, **10**:78–127.

doi:10.1186/1756-3305-6-47

Cite this article as: Moreno *et al.*: Phylogeography of the neotropical *Anopheles triannulatus* complex (Diptera: Culicidae) supports deep structure and complex patterns. *Parasites & Vectors* 2013 **6**:47.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

