



**Universidade de São Paulo**

**Biblioteca Digital da Produção Intelectual - BDPI**

---

Departamento de Geofísica - IAG/AGG

Comunicações em Eventos - IAG/AGG

---

2014

# New paleomagnetic and AMS study of mafic sills from the Rio Branco region - MT (SE of Amazonian Craton)

---

South American Symposium on Isotope Geology, 9, 2014, São Paulo

<http://www.producao.usp.br/handle/BDPI/45872>

*Downloaded from: Biblioteca Digital da Produção Intelectual - BDPI, Universidade de São Paulo*

## NEW PALEOMAGNETIC AND AMS STUDY OF MAFIC SILLS FROM THE RIO BRANCO REGION - MT (SE OF AMAZONIAN CRATON)

**Manoel Souza D'Agrella Filho** – Instituto de Astronomia, Geofísica e Ciências Meteorológicas/USP, **Marcelo Vitor Bravo Queiroz** – Instituto de Astronomia, Geofísica e Ciências Meteorológicas/USP, **Ricardo Trindade** – Instituto de Astronomia, Geofísica e Ciências Meteorológicas/USP, **Wilson Teixeira** – Instituto de Geociências/USP, **Amarildo Salina Ruiz** – Universidade Federal de Mato Grosso.

We present a paleomagnetic and AMS study on mafic sills cutting sedimentary rocks from the Rio Branco region, Mato Grosso state (Amazonian Craton). U-Pb (zircons) dating on a gabbro and a granophyre from the mafic-felsic Rio Branco Suite yielded ages of  $1471 \pm 8$  Ma and  $1427 \pm 10$  Ma, respectively. 108 samples from 11 sites were sampled, from which 3 sites are represented by sedimentary rocks collected close to three sills for a baked contact test. AMS measurements yielded planar fabric for analyzed rocks, which is expected for unmetamorphosed and undeformed sills and sedimentary rocks. After AF and thermal demagnetization southwestern, steep positive inclination characteristic remanent magnetization (ChRM) directions were isolated for the sills  $Dm=208.2^\circ$ ,  $Im=68.5^\circ$  ( $n=8$ ,  $A95=6.4^\circ$ ), which yielded the paleomagnetic pole at  $46.4^\circ S$ ;  $277.0^\circ E$  ( $A95=10.2^\circ$ ). These directions are very close to those found previously on mafic sills in the same region. Consequently, a new mean direction was calculated encompassing 18 sites ( $Dm=197.4^\circ$ ,  $Im=62.9^\circ$ ,  $A95=5.7^\circ$ ) which defined the paleomagnetic pole at  $56.0^\circ S$ ;  $278.5^\circ E$  ( $A95=7.9^\circ$ ). Magnetic mineralogy study indicates PSD-magnetite grains as the main magnetic carrier of ChRM directions. Although the baked contact test was not conclusive, paleomagnetic poles from other contemporary basic intrusions from the same region suggest that the Rio Branco sills' ChRM directions represent a thermoremanent magnetization (TRM) acquired during the emplacement, at about 1430 Ma.

Paleomagnetismo  
Sills  
Rochas máficas  
Rio Branco (AC)