RESEARCH ARTICLE





A survey on clinical presentation and nutritional status of infants with suspected cow' milk allergy

Mário C Vieira¹, Mauro B Morais^{*2}, José VN Spolidoro³, Mauro S Toporovski⁴, Ary L Cardoso⁵, Gabriela TB Araujo⁶, Victor Nudelman⁷ and Marcelo CM Fonseca⁶

Abstract

Background: Cow's milk is the most common food allergen in infants and the diagnosis of cow's milk allergy is difficult, even with the use of several diagnostic tests. Therefore, elimination diets and challenge tests are essential for the diagnosis and treatment of this disorder. The aim of this study is to report the clinical presentation and nutritional status of children evaluated by pediatric gastroenterologists for the assessment of symptoms suggestive of cow's milk allergy.

Methods: An observational cross-sectional study was performed among 9,478 patients evaluated by 30 pediatric gastroenterologists for 40 days in 5 different geographical regions in Brazil. Clinical data were collected from patients with symptoms suggestive of cow's milk allergy. The nutritional status of infants (age \leq 24 months) seen for the first time was evaluated according to z-scores for weight-for-age, weight-for-height, and height-for-age. Epi-Info (CDC-NCHS, 2000) software was used to calculate z-scores.

Results: The prevalence of suspected cow's milk allergy in the study population was 5.4% (513/9,478), and the incidence was 2.2% (211/9,478). Among 159 infants seen at first evaluation, 15.1% presented with a low weight-for-age z score (< -2.0 standard deviation - SD), 8.7% with a low weight-for-height z score (< -2.0 SD), and 23.9% with a low height-for-age z score (< -2.0 SD).

Conclusion: The high prevalence of nutritional deficits among infants with symptoms suggestive of cow's milk allergy indicates that effective elimination diets should be prescribed to control allergy symptoms and to prevent or treat malnutrition.

Background

The incidence of food allergies has increased in several parts of the world, particularly in developed countries [1-6]. Cow's milk is the most common food allergen in infants and the diagnosis of cow's milk allergy is difficult, even with the use of several diagnostic tests. Therefore, elimination diets and challenge tests are essential for the diagnosis and treatment of this disorder [7,8]. Cow's milk and cow's milk-based infant formulas are the most frequently used substitutes for infants who are not breastfed. The growth and development of infants that are allergic to cow's milk may be compromised if an adequate substitute formula is not available [9-12]. The nutritional quality of cow's milk substitutes used for this group of

² Department of Paediatrics, Universidade Federal de São Paulo, R. Botucatu, 598, São Paulo-SP 04023-062, Brazil patients has a crucial role in promoting appropriate growth and development [9-12].

Prior to 1950, the incidence of cow's milk allergy in the first year of life was low and affected about 0.1% to 0.3% of all infants. Prospective studies conducted in 1970 and 1988 showed that the incidence of cow's milk allergy reached 1.8% to 7.5%, a wide range explained by the differences in diagnostic criteria adopted in different studies [1,6]. Symptoms suggestive of cow's milk allergy are reported by parents of 5.0% to 15.0% of all infants [1]. These data draw attention to the importance of an accurate diagnosis of food allergy in order to avoid the use of elimination diets for prolonged periods of time in infants without a confirmed diagnosis.

The aim of this study was to describe the clinical presentation and nutritional status of children evaluated by



© 2010 Vieira et al; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/2.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

^{*} Correspondence: mbmorais@osite.com.br

Full list of author information is available at the end of the article

pediatric gastroenterologists for the assessment of symptoms suggestive of cow's milk allergy.

Methods

Data collection

An observational cross-sectional study was performed applying a questionnaire to 30 pediatric gastroenterologists from 20 different cities in 11 states of five Brazilian geographical regions (North, Midwest, Northeast, Southeast, and South). Data were collected during 40 consecutive days in 2004. Information on the number of patients evaluated for food allergies, demographic data, clinical manifestations, time of onset of symptoms, management, and anthropometric data at birth and at the first consultation was obtained and recorded.

A total of 9,478 children were evaluated over the study period. Five hundred thirteen children were identified as having suspected food allergy. Each specialist evaluated a mean of 2.4 patients with cow's milk allergy per week, including 1.0 new case.

The nutritional status of infants (age ≤ 24 months) seen for the first time was evaluated according to z-scores for weight-for-age, weight-for-height, and height-for-age. Epi-Info (CDC-NCHS, 2000) software was used to calculate z-scores and the National Center for Health Statistics (NCHS) growth charts were used as reference values [13]. Considering that the nutritional needs and growth velocity are variable in the first 2 years of life, weight and length were presented according to 3 age groups: ≤ 6.0 months, 6.1 - 12.0 months, 12.1 - 24.0 months.

According to WHO, a nutritional deficit is defined when z-scores are below -2.0 standard deviations [14]. In a normal population distribution, 2.5% of the values are expected to be below this cut-off point; therefore, a rate > 2.5% suggests that the study population has a nutritional deficit in comparison with reference values.

SigmaStat ^{*} for Windows, version 3.1, was used for statistical analysis. Differences were classified as statistically significant when the p value was < 0.05. The study was approved by the Human Research Ethics Committee at the Hospital Pequeno Príncipe - Curitiba, Brazil.

Results

Cow's milk allergy was suspected in 513 of 9,478 consultations in the pediatric age range. In 211 patients, suspicion of the diagnosis was made at the first medical visit, and in 302 patients the diagnosis was suspected at followup visits. Therefore, the prevalence of diagnosed and suspected cow's milk allergy in the study population was 5.4% (513/9,478), and the incidence was 2.2% (211/9,478). At first consultation, pediatric gastroenterologists agreed with the diagnostic hypothesis of cow's milk allergy made by the referring pediatrician in 82.0% of the cases. Among patients who had suspected cow's milk allergy (n = 211) at first consultation, 49.3% were referred to the pediatric gastroenterologist having already been switched to a substitute infant diet. The milk substitute most frequently prescribed by general pediatricians was a soy formula (58%). Other inappropriate treatments, including lactose-free cow's milk infant formula or goat' milk (11%), were also prescribed. Extensively hydrolyzed formulas were used in 11% of patients, as well as amino acid-based formulas (5%). A diet without milk substitutes was prescribed in 5% of patients.

The following results refer to new cases of infants (≤ 24 months of age) with symptoms suggestive of cow's milk allergy. Weight and length were recorded in 159 (90.8%) of the 175 patients < 24 months of age. One hundred thirty patients (81.7%) were evaluated in the 1st year of life. In this group, 79 (49.7%) patients were evaluated in their 1st six months of life and 51 (32.1%) in their 2nd six months. Twenty-nine (18.2%) patients were evaluated in the 2nd year of life.

Demographic data, weight and length at birth, clinical presentation, and duration of symptoms according to age group are presented in Table 1. The comparisons of gender and weight and height at birth in the three age groups did not show any statistically significant differences. The mean duration of symptoms showed a direct correlation with the child's age.

There was a predominance of gastrointestinal manifestations in all age groups (Table 1). Gastrointestinal symptoms were more frequent in the 1st six months of life than in the other age groups (p < 0.05) whereas respiratory manifestations were more frequent in the 2nd six months of life (p < 0.05).

The clinical manifestations of infants were grouped as follows and are presented in Table 2: systemic (weight loss, anorexia, and irritability); gastrointestinal; cutaneous (atopic dermatitis and urticaria); and respiratory. Exclusive gastrointestinal symptoms were observed in 74 patients (46.5%), exclusive respiratory symptoms in 7 patients (4.4%), exclusive cutaneous manifestations in 5 patients (3.1%), and systemic symptoms only in 2 patients (1.3%). Therefore, a total of 88 (55.3%) of the 159 patients had clinical manifestations in only one of the 4 categories. Gastrointestinal symptoms and 1 (n = 59) or 2 (n = 8) other clinical symptoms in another category (systemic, cutaneous, or respiratory) occurred in 67 patients (42.1%). Only 18 patients (11.3%) had no gastrointestinal manifestations.

The distribution of z-scores for weight-for-age, weightfor-height, and height-for-age of the 159 infants is presented in Figure 1. The analysis of z-score for weight-forheight revealed greater value dispersion, but the mean value was similar to the reference value. The weight-forage curve followed an expected dispersion, but showed a deviation to the left. The height-for-age curve also

		Age group (months)		р
_	≤ 6.0 (n = 79)	6.1 - 12.0 (n = 51)	12.1 - 24.0 (n = 29)	
Gender (male/ female) ¹	39/40	30/21	16/13	0.561
Birth weight (grams) ²	3015 (2821; 3348)	3020 (2793; 3364)	3122 (2955; 3370)	0.663
Birth length (cm) ²	48.5 (47.3; 50.0)	49.0 (48.0; 50.0)	49.0 (47.1; 50.0)	0.901
Duration of symptoms (months) ^{2,3}	1.0 (1.0; 2.0)	5.0 (4.0; 7.0)	12.0 (10.0; 18.0)	<0.001
Systemic symptoms ¹	14 (17.7%)	18 (35.3%)	7 (24.1%)	0.075
Digestive symptoms ^{1,4}	76 (96.2%)	42 (82.4%)	23 (79.3%)	0.011
Cutaneous symptoms ¹	13 (16.5%)	10 (19.6%)	4 (13.8%)	0.789
Respiratory symptoms ^{1,4}	9 (11.4%)	16 (31.4%)	6 (20.7%)	0.019

Table 1: Gender, weight and length at birth, duration of symptoms, and clinical presentation according to age group

1. Chi-square test

2. Median and 25th and 75th percentiles; Kruskal-Wallis test for multiple groups.

3. Dunn test for multiple comparisons for the variable "duration of symptoms"

- p < 0.05: ($\leq 6.0 \text{ months}$) × (6.1 - 12.0 months);

- p < 0.05: ($\leq 6.0 \text{ months}$) \times (12.1 - 24.0 months);

- p < 0.05: 6.1 - 12.0 months) × (12.1 - 24.0 months);

4. Chi-square results:

Digestive symptoms: ≤ 6 months versus 6-12 months, p = 0.008; ≤ 6 months versus 12-24 months, p = 0.005; 6-12 months versus 12-24 months, p = 0.737.

 $\label{eq:respiratory symptoms: $$ 6 months versus 6-12 months, p = 0.005; $$ 6 months versus 12-24 months, p = 0.304; 6-12 months versus 12-24 months, p = 0.216.$

showed a deviation to the left, but greater dispersion in comparison with the reference curve.

The distribution of patients according to anthropometric deficits confirmed by a z-score < -2.0 standard deviations is presented in Table 3. No statistically significant differences were found in the three age groups. The total number of infants with deficits involving weight-for-age, weight-for-height, and height-for-age were 15.1%, 11.3%, and 23.9%, respectively.

Discussion

The incidence and prevalence of food allergies are believed to be increasing in several countries [1,2,15]. However, the comparison of data from different studies is difficult because criteria to diagnose food allergies, as well as the definitions of groups, vary substantially between studies. Therefore, rates vary from 35% when parental reports are used as criteria to define food allergies, to 1% when proper double-blind placebo-controlled food challenges (DBPCFC) are used [5,6].

A study was conducted in the Isle of Wight in the United Kingdom in order to establish the rates of objectively-assessed food allergies in the 1st year of life and to compare this with the rate of parental reports. A cohort of 969 infants was recruited between September 2001 and August 2002. Symptoms of food allergies were reported by 132 parents (14.2%) at 3 months, 83 parents (9.1%) at 6 months, 49 parents (5.5%) at 9 months, and 65 parents (7.2%) at 12 months of age [4]. The cumulative incidence of reported parental perceived food hypersensitivities was 25.8% (250/969; 95% CI, 23.1% to 28.7%) by 12 months of age. Of these, only 14% and 6% were diagnosed with food allergies by means of open food challenges and DBPCFC, respectively. Thus, in this cohort, the incidence of food allergies by the age of 12 months was 2.6% (25/969; 95% CI, 1.7% to 3.8%) on the basis of open food challenge and 1.2% (12/969; 95% CI, 0.6% to 2.2%) on the basis of DBPCFC.

However, the actual occurrence of food allergies may be underestimated because challenge tests may be per-

Clinical manifestation	Ν	%	
Gastrointestinal	141	88.7	
Vomiting and regurgitation	85	53.5	
Colic	54	34.0	
Diarrhea without blood	30	18.9	
Constipation	25	15.7	
Blood in stools with normal consistency	23	14.5	
Bloody diarrhea	10	6.3	
Systemic	39	24.5	
Cutaneous	29	18.2	
Respiratory	31	19.5	

Table 2: Clinical presentation of 159 infants (age ≤ 24 months) with symptoms suggestive of cow's milk allergy (each infant may have one or more clinical manifestation)

formed after the development of tolerance. This is a limitation for the method considered the gold standard for epidemiologic studies concerning food allergies. Therefore, the suggestion that the incidence of food allergies in the 1st year of life ranges from 2% to 3% and symptoms compatible with food allergies found in 5% to 15% of infants may be reasonable and close to reality [1]. As distinct from other studies, we evaluated the prevalence and incidence of symptoms suggestive of cow's milk allergy according to data collected in the offices of pediatric gastroenterologists. The prevalence of suspected cow's milk allergy was 5.4% in all consultations, and the incidence of new cases was 2.2%. The pediatric gastroenterologists agreed with the diagnostic hypothesis made by the referring pediatrician in 82.0% of the consultations. Although the diagnosis of cow's milk allergy was not established by a milk challenge, patients were started on a cow's milkfree diet, which is the initial approach for this disorder. Nearly one-half of the patients had already received a prescription to eliminate cow's milk protein from their diets when they were seen for the first time by their pediatricians. However, many infants were being fed inappropriate substitutes, such as soy-based infant formulas, extracts with soybean proteins, goat's milk, or even lactose-free cow's milk formula. Only 10.0% of the infants received extensively hydrolyzed or amino acid-based formulas, considered adequate substitutes recommended for infants with cow's milk allergy according to the European Society for Pediatric Allergology and Clinical Immunology (ESPACI), the European Society for Pediat-Gastroenterology, Hepatology and Nutrition ric (ESPGHAN), and the American Academy of Pediatrics [7,16].

Approximately one-half of the infants in the study were younger than 6 months of age. In this age group, gastrointestinal manifestations occurred at a greater frequency than in the group of infants older than 6 months of age. As expected, digestive symptoms were the most common (88.7%), including regurgitation and vomiting, colic, diarrhea, and blood in stools. A variety of gastrointestinal allergic disorders typically affect in infants and children. Infants with allergic colitis present small amounts of blood mixed with mucus in their stools. Cow's milk-sensitive enteropathy may present with malabsorption leading to diarrhea and failure to thrive. The most serious form of gastrointestinal food allergy in infants is food protein-induced enterocolitis syndrome which has a symptom complex of profuse vomiting and diarrhoea, and potentially a sepsis-like clinical picture [17]

Cutaneous and respiratory symptoms were less frequent, possibly due to the fact that patients were referred to pediatric gastroenterologists. Although the predominant type of clinical manifestation may depend on the type of specialty care where the study patients are enrolled, there is a consensus that gastrointestinal or cutaneous symptoms are the predominant forms of presentation of cow's milk allergy.

A review of the literature did not yield studies with similar designs for comparisons. In a group of 204 infants with cow's milk allergy studied in the 1950s, the most common symptoms were atopic dermatitis in 43% of the cases, vomiting and regurgitation in 38%, colic in 31%, wheezing in 9%, irritability and anorexia in 22%, and constipation in 6% of the cases [18]. The most remarkable difference from our data is that we observed a greater frequency of allergic colitis and less atopic dermatitis. This Vieira et al. BMC Pediatrics 2010, **10**:25 http://www.biomedcentral.com/1471-2431/10/25



	≤ 6.0 m	6.1 - 12.0 m (n = 51)	12.1 - 24.0 m (n = 29)	р
	(n = 79)			
z-scores (mean \pm SD) ¹				
Weight-for-age	-0.70 ± 1.33	-0.83 ± 1.06	-0.9 ± 1.30	0.671
Weight-for-height	-0.21 ± 1.72	-0.08 ± 1.54	-0.27 ± 1.55	0.854
Height-for-age	-0.93 ± 1.82	-0.71 ± 1.78	-0.68 ± 1.76	0.724
Deficit (z-score<-2.0 SD) ²				
Weight-for-age	13 (16.5%)	5 (9.8%)	6 (20.7%)	0.380
Weight-for-height	11 (13.9%)	4 (7.8%)	3 (13.8%)	0.556
Height-for-age	22 (27.8%)	12 (23.5%)	4 (13.8%)	0.315

Table 3: Mean ± standard deviation (SD) of z-scores for weight-for-age, weight -for-height, and height-for-age, and number and percentage of infants with cow's milk allergy and nutritional deficits (z-score < -2.0 SD) according to age group

1. One-way analysis of variance

2. Chi-square test

current trend is supported by a study conducted in London reporting a reduction in cases of cow's milk allergy associated with intestinal malabsorption due to infectious gastroenteritis [19].

The mean z-score deviations, particularly for weightfor-age and height-for-age, suggest that failure to thrive or malnutrition may occur as a consequence of cow's milk allergy. The analysis of weight and height showed greater deficits (< -2.0 standard deviations) than expected (2.5%) according to the CDC-NCHS reference values (2000): specifically, 15.1% of weight-for-age, 11.3% of weight-forheight, and 23.9% of height-for-age z-scores (Table 3). The World Health Organization has recently issued new weight and height reference charts for infants who were exclusively breastfed [20]. These values were not used in our study because our patients were not breastfed, and this may help compare our data with other studies in the literature. A study reporting on a group of 26 Brazilian children demonstrated that 23% had a deficit in weightfor-age, 7.7% had a deficit in weight-for-height, and 11.5% had a deficit in height-for-age when the same diagnostic criteria and reference values were used. These patients were also shown to be receiving a diet with a lower energy intake and calcium content when compared with controls in the same age group and socioeconomic conditions. A nutritionally-inadequate elimination diet may lead to or aggravate anthropometric deficits of infants with symptoms suggestive of cow's milk allergy according to the few studies that investigated this issue [9-12]. Our results have shown that height-for-age deficit was the most predominant indicator of nutritional impairment in contrast with the expected predominance of weight-for-age deficit. Similar findings were reported in children with constipation secondary to cow's milk allergy, who presented a mean height-for-age deficit (-0.90 \pm 1.24) more pronounced than the weight-for-age deficit (-0.67 \pm 1.30) [21]. It may be possible that due to chronic inflammation secondary to milk allergy, linear growth may be impaired as it is observed in chronic liver disorders [22].

In this study, approximately half of the patients referred to the paediatric gastroenterologists were already switched to a substitute infant diet. However, only 16% of patients were receiving extensively hydrolyzed formulas or amino acid-based formulas. The duration of the substitute diet and the efficacy of treatment were variable. Therefore, the nutritional deficit observed in patients at the time of inclusion in the study may be attributable either to the use of inappropriate milk substitutes or to the insufficient duration of treatment for nutritional recovery.

Cow's milk allergy in infants is usually non-IgE mediated, and the diagnostic hypothesis should be raised using clinical symptoms and, if available, functional and morphological markers of gastrointestinal function. Since there are no effective laboratory methods for the diagnosis of this disorder, an elimination diet without allergenic proteins remains the first essential step to make a diagnosis of cow's milk allergy. Clinical follow-up to evaluate the response to treatment (elimination diet) is an essential step in the management of these patients [23]. After an initial phase of clinical and nutritional recovery, food challenges, when recommended, may provide a definitive diagnosis.

Conclusions

Our data revealed the profile of infants with symptoms suggestive of cow's milk allergy and the presence of nutritional deficits in a considerable percentage of patients. These findings highlight the need to prescribe highlyeffective elimination diets in order to control symptoms, to ensure fast nutritional recovery, and to avoid malnutrition. Further studies should be conducted to develop public healthcare strategies to provide adequate substitute diets and treat infants that have a diagnosis or symptoms suggestive of cow's milk allergy, a current concern in many countries [24].

Competing interests

The authors have received fees from Support Advanced Medical Nutrition - Danone for technical assistance in this article.

Authors' contributions

All authors read and approved the manuscript. MCV: study concept and design; analysis and interpretation of data; drafting of the manuscript; study supervision. MBM: study concept and design; analysis and interpretation of data; drafting of the manuscript; statistical analysis; study supervision. JVNS: study concept and design; analysis and interpretation of data; drafting of the manuscript; study supervision. MST: study concept and design; analysis and interpretation of data; drafting of the manuscript; study supervision. ALC: analysis and interpretation of data; drafting of the manuscript; critical revision of the manuscript for important intellectual content. GTBA: study concept and design: acquisition of data; analysis and interpretation of data; critical revision of the manuscript for important intellectual content; statistical analysis; study supervision. VN: analysis and interpretation of data; drafting of the manuscript; critical revision of the manuscript for important intellectual content. MCMF: study concept and design; acquisition of data; analysis and interpretation of data; critical revision of the manuscript for important intellectual content; statistical analysis; study supervision.

Acknowledgements

We should like to thank all the families and paediatricians who participated in the study. The research was funded by Support Advanced Medical Nutrition - Danone. The views expressed in this publication are those of the authors.

Author Details

¹Centre for Paediatric Gastroenterology, Hospital Pequeno Príncipe, R. Desembargador Motta, 1070, Curitiba-PR 80250-060, Brazil, ²Department of Paediatrics, Universidade Federal de São Paulo, R. Botucatu, 598, São Paulo-SP 04023-062, Brazil, ³Department of Paediatrics - School of Medicine, Pontificia Universidade Católica, Av. Ipiranga, 6690, Porto Alegre-RS 90610-000, Brazil, ⁴Department of Paediatrics - School of Medicine, Santa Casa de São Paulo, R. Cesário Motta Jr., 112 São Paulo-SP 01221-020 Brazil, ⁵Department of Paediatrics, Instituto da Criança, Hospital das Clínicas, Faculdade de Medicina da Universidade de São Paulo, Av. Dr. Enéas Carvalho Aguiar, 647, São Paulo - SP 05403-000, Brazil, ⁶Health Economics, Axia.bio Consulting, R. Setembrino Woitechumas, 38/4, São Paulo-SP 04563-090, Brazil and ⁷Department of Paediatrics, Hospital Albert Einstein, Av. Albert Eisntein, 627, São Paulo - SP 05651-901, Brazil

Received: 6 September 2009 Accepted: 23 April 2010 Published: 23 April 2010

References

- 1. Host A: Frequency of cow's milk allergy in childhood. Ann Allergy Asthma Immunol 2002, 89(6 Suppl 1):33-7.
- 2. Cataldo F, Accomando S, Fragapane ML, Montaperto D, SIGENP and GLNBI Working Groups on Food Intolerances: Are food intolerances and

allergies increasing in immigrant children coming from developing countries? *Pediatr Allergy Immunol* 2006, **17**:364-9.

- Isolauri E, Huurre A, Salminen S, Impivaara O: The allergy epidemic extends beyond the past few decades. *Clin Exp Allergy* 2004, 34:1007-10.
- Venter C, Pereira B, Grundy J, Clayton CB, Roberts G, Higgins B, et al.: Incidence of parentally reported and clinically diagnosed food hypersensitivity in the first year of life. J Allergy Clin Immunol 2006, 117(5):1118-24.
- American College of Allergy, Asthma, & Immunology: Food allergy: a practice parameter. Ann Allergy Asthma Immunol 2006, 96(3 Suppl 2):S1-68.
- Keil T: Epidemiology of food allergy: what's new? A critical appraisal of recent population-based studies. Curr Opin Allergy Clin Immunol 2007, 7:259-63.
- Host A, Koletzko B, Dreborg S, Muraro A, Whan U, Agget P, et al.: Dietary products used in infants for treatment and prevention of food allergy. Joint statement of the European Society for Pediatric Allergology and Clinical Immunology. Committee on hypoallergenic formulas and the European Society for Pediatric Gastroenterology. *Hepatology and Nutrition Arch Dis Child* 1999, 81:80-4.
- Vandenplas Y, Brueton M, Dupont C, Hill D, Isolauri E, Koletzko S, et al.: Guidelines for the diagnosis and management of cow's milk protein allergy in infants. Arch Dis Child 2007, 92:902-8.
- 9. Tiainen JM, Nuutinen OM, Kalavainen MP: Diet and nutritional status in children with cow's milk allergy. *Eur J Clin Nutr* 1995, **49**:605-121.
- Arvola T, Holmberg-Marttila D: Benefits and risks of elimination diets. Ann Med 1999, 31:293-8.
- Medeiros LC, Speridião PG, Sdepanian VL, Fagundes-Neto U, Morais MB: Nutrient intake and nutritional status of children following a diet free from cow's milk and cow's milk by-products. J Pediatr (Rio J) 2004, 80:363-70.
- 12. Noimark L, Cox HE: Nutritional problems related to food allergy in childhood. *Pediatr Allergy Immunol* 2008, **19**:188-95.
- Centers for Disease Control and Prevention and National Center for Health Statistics: 2000 CDC growth charts: United States. 2008 [<u>http://</u><u>www.cdc.gov/growthcharts</u>].
- World Health Organization: Physical status: the use and interpretation of anthropometry. In *Technical Report Series*, 854 Geneve: WHO; 1995.
- The International Study of Asthma and Allergies in Childhood (ISAAC) Steering Committee: Worldwide variation in prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and atopic eczema: ISAAC. Lancet 1998, 351(9111):1225-32.
- Committee on Nutrition. American Academy of Pediatrics: Hypoallergenic infant formulas. *Pediatrics* 2000, 106:346-9.
- 17. Sicherer SH: Food allergy. Lancet 2002, 360:701-10.
- Clein NW: Cow's milk allergy in infants. Pediatr Clin North Am 1954, 4:949-62.
- Walker-Smith J: An eye witness perspective of the changing patterns of food allergy. Eur J Gastroenterol Hepatol 2005, 17:1313-6.
- de Onis M, Garza C, Onyango AW, Borghi E: Comparison of the WHO child growth standards and the CDC 2000 growth charts. J Nutr 2007, 137:144-8.
- Tahan S, Motta MEFA, Goshima S, Daher S, Naspitz CK, Solé D, Fagundes-Neto U, Morais MB: Chronic constipation secondary to cow's milk allergy affects nutritional status in children. J Pediatr Gastroenterol Nutr 2004, 39:S235.
- 22. Sokol RJ, Stall C: Anthropometric evaluation of children with chronic liver disease. *Am J Clin Nutr* 1990, **52**:203-8.
- 23. Walker-Smith JA: Diagnostic criteria for gastrointestinal food allergy in childhood. *Clin Exp Allergy* 1995, **25(Suppl 1)**:20-2.
- 24. Guest JF, Valovirta E: Modeling the resource implications and budget impact of new reimbursement guidelines for the management of cow milk allergy in Finland. *Curr Med Res Opin* 2008, **24**:1167-77.

Pre-publication history

The pre-publication history for this paper can be accessed here: http://www.biomedcentral.com/1471-2431/10/25/prepub

doi: 10.1186/1471-2431-10-25

Cite this article as: Vieira *et al.*, A survey on clinical presentation and nutritional status of infants with suspected cow' milk allergy *BMC Pediatrics* 2010, **10**:25