1	BABY ON BOARD: OLFACTORY CUES INDICATE PREGNANCY
2	AND FOETAL SEX IN A NONHUMAN PRIMATE
3	
4	Jeremy Chase Crawford ^{1,2,3,4} and Christine M. Drea ^{4,5,*}
5	¹ National Evolutionary Synthesis Center, Durham, NC USA
6	² Department of Integrative Biology, University of California, Berkeley, CA USA
7	³ Museum of Vertebrate Zoology, University of California, Berkeley, CA USA
8	⁴ Department of Evolutionary Anthropology, Duke University, Durham, NC USA
9	⁵ Department of Biology, Duke University, Durham, NC USA
10	*Author for correspondence (cdrea@duke.edu)
11	
12	Running headline: Olfactory cues of pregnancy in lemurs
13	

14 **ABSTRACT**

Olfactory cues play an integral, albeit underappreciated, role in mediating vertebrate social and 15 reproductive behaviour. These cues fluctuate with the signaler's hormonal condition, coincident 16 17 with and informative about relevant aspects of its reproductive state, such as pubertal onset, 18 change in season and, in females, timing of ovulation. Although pregnancy dramatically alters a female's endocrine profiles, which can be further influenced by foetal sex, the relationship 19 between gestation and olfactory cues is poorly understood. We therefore examined the effects of 20 21 pregnancy and foetal sex on volatile genital secretions in the ring-tailed lemur (Lemur catta), a strepsirrhine primate possessing complex olfactory mechanisms of reproductive signaling. 22 23 Whilst pregnant, dams altered and dampened their expression of volatile chemicals, with 24 compound richness being particularly reduced in dams bearing sons. These changes were 25 comparable in magnitude to other, published chemical differences among lemurs that are salient to conspecifics. Such olfactory 'signatures' of pregnancy may help guide social interactions, 26 potentially promoting mother-infant recognition, reducing intragroup conflict, or counteracting 27 behavioural mechanisms of paternity confusion; cues that also advertise foetal sex may 28 29 additionally facilitate differential sex allocation.

30

Keywords: olfactory communication, reproductive signal, gestation, sex allocation, hormone,
 chemosignal

33 **1. INTRODUCTION**

Female animals routinely broadcast information about their reproductive state. In general, the 34 35 research emphasis in reproductive signaling has been on cues associated with fecundity, ovulation or mate quality; however, there are also potentially potent cues associated with 36 37 gestation. The physiological changes underlying pregnancy are complex and, compared to 38 ovulation, are more enduring. The corresponding changes to endocrine concentrations, that may guide reproductive cues of pregnancy, can be extreme. Notably, in comparison to maximum 39 40 concentrations outside of pregnancy, concentrations of certain sex steroids during pregnancy can increase >1,600% [1] and vary by foetal sex (reviewed in [2]). Moreover, pregnancy cues may 41 42 serve important functions. In primates, for instance, gestational cues may reduce intragroup 43 conflict in species that compete intensely for mates [3] or may prepare cooperatively breeding 44 males for the energetic and behavioural changes necessary for paternal care [4]. Historically, the conveyance of reproductive information, particularly in primates, has been evaluated primarily 45 46 through visual and behavioural channels (e.g. [5,6]); nonetheless, there is growing recognition of reproductive signaling via olfactory channels [7]. As in several non-primate species [8–10], 47 48 olfactory cues of pregnancy occur in humans, possibly facilitating mother-infant recognition 49 [11]; otherwise, the potential for olfactory gestational cues in primates has been ignored. Here, 50 we investigate pregnancy effects on olfactory cues in a promiscuous, nonhuman primate, the 51 ring-tailed lemur (Lemur catta).

Recently, the ring-tailed lemur has emerged as a useful model for examining the form and function of olfactory signals relevant to reproduction. Among females, that use scent to demarcate birthing sites [12] and for competitive overmarking [7], variations in odorant profiles (whether in compound type, number, or relative concentration, see e.g. diversity indexes in [13]) correspond to natural [14,15] and experimentally induced [16] variations in reproductive 57 hormones. Through behavioural bioassays, genetic profiling and chemical analyses, lemur 58 chemosignals have been shown to convey information about sex, breeding condition and 59 individual identity, as well as genetic quality and relatedness (reviewed in [7,16]). By contrast, 60 dominance status within the sexes, which in lemurs can be nonlinear, non-transitive, and highly variable [17], is recognizable via scent in known individuals only [18], but is not chemically 61 encoded [18,19]. Using a within-subjects sampling design, we extend these prior studies by 62 characterizing chemical changes in lemur genital secretions between preconception and 63 pregnancy, whilst also considering the potential effects of other concurrent physiological or 64 65 environmental changes, such as age, season, litter size, and foetal sex.

66

67 2. MATERIAL AND METHODS

The subjects were 12 captive, sexually mature (1.5-21 years), female ring-tailed lemurs, housed in semi-free-ranging, mixed-sex groups at the Duke Lemur Center in Durham, NC, USA. We monitored reproduction over a six-year period (2004-2010), including during 14 pregnancies ([2]; see Electronic Supplementary Material, ESM). To illustrate any potential olfactoryendocrine relationship, we present accompanying sex steroid concentrations in female subjects, recalculated from [2]. Our animal procedures were approved by the Institutional Animal Care and Use Committee of Duke University (see ESM).

We collected samples of labial secretions near conception (mean \pm SEM: 17.3 \pm 3.7 days preconception) and mid-pregnancy (mean \pm SEM: 88.8 \pm 4.6 days of a 135-day gestation period) for analysis by gas chromatography-mass spectrometry (see [20]). We used richness as a proxy of chemical complexity [13], retaining only compounds that eluted at 8-43 minutes and comprised \geq 0.01% of the chromatogram [20]. We assigned 'foetal sex' following [2], grouping dams carrying singleton or twin females as 'females bearing daughters' (FBD; n=8) and dams
carrying at least one male (singleton male, twin males, or mixed-sex twins) as 'females bearing
sons' (FBS; n=6; ESM).

83 We first addressed and dismissed any potential effects on a female's chemical richness of her age, the seasonal timing of her pregnancy, and her litter size (see ESM). We then tested for 84 effects of reproductive condition (preconceptive versus pregnant) and foetal sex (FBD versus 85 FBS) on her chemical richness and endocrine concentrations, using two-factor ANOVAs 86 (univariate repeated-measures model, JMP PRO 11.0, SAS; ESM). To contextualise the main 87 88 effect of reproductive condition on chemical complexity, we also calculated the average 89 differences in chemical richness between several previously studied groups from this population ([13,14,16,19]; see ESM). Having decided a priori to compare FBD and FBS, we resolved the 90 91 significant interaction using F-tests for simple effects. These data can be accessed at Dryad [21].

92

93 **3. RESULTS**

94 The odorant profiles of female ring-tailed lemurs changed with pregnancy, relative to preconception, as revealed by differences in the gas chromatograms of their genital secretions 95 96 (Figures 1a,b): pregnancy was associated with changes in the relative proportions of the volatile compounds expressed and with a significant decrease in the total number of compounds ($F_{1,10}$ = 97 14.41, P < 0.01; Figure 1c). The change in chemical richness (mean \pm SEM: 9.67 \pm 6.62 98 99 compounds) was comparable in magnitude to the behaviourally salient, chemical differences 100 observed between other groups of individuals, including those found between intact and hormonally contracepted females (Table 1). 101

102 Remarkably, a dam's volatile chemical profile during pregnancy also varied with the sex of her foetus. A significant interaction between reproductive condition and foetal sex ($F_{1, 10}$ = 103 5.21, P < 0.05) owed primarily to a greater loss of chemical richness during pregnancy in FBS 104 105 than in FBD (F_{1, 12,34} = 5.51, P < 0.05; Figure 2*a*). Because there was no effect of a female's eventual foetal sex on the chemical richness of her preconceptive odorants ($F_{1, 12, 34} = 0.74$, P = 106 0.40, n.s.), the differences by foetal sex during pregnancy could not be explained by the scent 107 signatures (see [19]) of individual females. Instead, these findings may relate to underlying 108 109 differences, by foetal sex, in the concentrations of maternal sex steroids, which increase during 110 pregnancy (Figure 2b; Table 2).

111

112 **4. DISCUSSION**

113 We show that odorant expression in lemurs is altered during pregnancy and is further 114 differentiated in accordance with the sex of the dam's developing offspring. The chemical effects induced by pregnancy were comparable in magnitude to other differences in scent that, as 115 116 revealed by behavioural bioassays, are salient to ring-tailed lemurs [16,18,22]. The degree to which compounds are lost during pregnancy and influenced by foetal sex appears to be inversely 117 118 related to the female's changing sex steroid concentrations. Given the established link between 119 sex steroids and olfactory profiles [7], these findings suggest possible endocrine involvement (or competing energy allocation) in the production of olfactory gestational cues. 120

Gestational cues could serve various functions across primate species, from promoting social cohesion [3] to engendering parental investment [4] or kin recognition [11]. To the extent that olfactory cues of pregnancy occur in other primates, our findings could be relevant to these existing hypotheses, as well as to theories on the functionality of multiple mating. Notably, a 125 female's efforts to confuse males about their potential paternity ('paternity confusion') is 126 commonly invoked to explain a female primate's engagement in multiple mating during 127 pregnancy, particularly at times when there are no overt behavioural cues of pregnancy or when 128 visual cues of pregnancy are not yet apparent [23,24] – in other words, when males presumably 129 lack knowledge about female reproductive state. Yet, we have no information about potential scent cues of pregnancy in these species, whose males are nevertheless known to engage in 130 olfactory investigation of vaginal secretions prior to mating [7,25,26]. We must therefore 131 consider the possibility that olfactory cues could provide males with a means of detecting a 132 133 proceptive dam's reproductive state. Discerning males could possibly thwart the female's purported tactics, either by not mating with her or by mating with her without becoming 134 confused about paternity. If so, one might only expect multiple mating to be an effective means 135 136 of paternity confusion in species in which either the female does not produce olfactory pregnancy cues or the male is unable to detect them. 137

To the extent that olfactory advertisement of foetal sex might also occur more broadly 138 139 than in *Lemur*, our findings might help elucidate a potential mechanism of differential sex allocation. Under certain conditions, animals increase fitness by preferentially investing in 140 141 offspring of a particular sex [27]. Although theories addressing sex allocation typically implicate 142 mechanisms operating at conception [28] or post-parturition [29], mechanisms operating during gestation would allow animals to respond to important social or environmental changes whilst 143 144 the ongoing costs of investment are still considerable. An expectant dam could potentially use 145 her own olfactory cues as a self-referent to inform her investment decisions; alternately, conspecifics could use these cues as predictors of imminent sex ratios to guide their own sex-146 147 allocation strategies. Although opportunities to study a putative gestational mechanism of sex

allocation will likely remain limited in this endangered, long-lived species, such limitations do
not preclude investigations of potential sex-specific gestational cues or gestational sex allocation
in more tractable study systems that share key life history characteristics (e.g., [30]).

The observation that pregnancy alters scent profiles in lemurs (this study) and in humans [11] suggests that olfactory cues of pregnancy may be widespread or highly conserved among primates. Such findings highlight the importance of considering multiple sensory modalities when examining reproductive signals, even in taxa historically thought to rely relatively little on olfaction [7,31,32].

156

Figure 1. Volatile chemicals varied among the genital secretions of preconceptive and pregnant ring-tailed lemurs: gas chromatograms depict variation in the relative proportions of compounds present in the labial secretions of a representative ring-tailed lemur before conception (*a*) and whilst pregnant with twin daughters (*b*). Numbers denote: 1, internal standard (hexachlorobenzene); 2, squalene; 3, cholesterol. Bar graphs represent differences by reproductive condition in the chemical richness (mean \pm SEM) derived from the chromatograms of all female subjects (*c*).

164

Figure 2. Mean \pm SEM chemical richness (*a*) and serum steroid concentrations (*b*) varied by reproductive condition and foetal sex in female ring-tailed lemurs. T, testosterone; A₄ androst-4ene-3,17,dione; OHP, 17 α -hydroxyprogesterone; E₂, 17 β -oestradiol.

168

Table 1. Differences in the chemical richness of ring-tailed lemur genital secretions between groups or conditions for which existing behavioural data support the salience of the olfactory signal.

172

Table 2. Differences in the endocrine profiles of female ring-tailed lemurs in relation to reproductive condition and foetal sex. Notably, serum concentrations during gestation were significantly different between females bearing daughters (FBD) and females bearing sons (FBS) for all measured steroids.

177

178 ACKNOWLEDGEMENTS

We thank M. Boulet, D. Brewer, G. Dubay, B. Schopler, J. Taylor, C. Williams, and S. Zehr for
facilitating this work. This research was funded by NSF (BCS-0409367, IOS-0719003 to
C.M.D.). Analysis and write-up were supported by NSF (IOS-1021633 to C.M.D.), and by a
Visiting Scholar's Fellowship from the National Evolutionary Synthesis Center (NSF EF0423641), a Chancellor's Fellowship from the University of California, Berkeley, an NSF
Graduate Research Fellowship, and funding from the Research Council of Norway (to J.C.C.).
DLC publication no: X.

186

187 **REFERENCES**

- Abbassi-Ghanavati M, Greer LG & Cunningham FG. 2009 Pregnancy and laboratory studies: a reference table for clinicians. *Obstet. Gynecol.* 114, 1326–1331.
 (doi:10.1097/AOG.0b013e3181c2bde8)
- Drea CM. 2011 Endocrine correlates of pregnancy in the ring-tailed lemur (Lemur catta):
 Implications for the masculinization of daughters. *Horm. Behav.* 59, 417–427.
 (doi:10.1016/j.yhbeh.2010.09.011)
- Gerald MS, Waitt C & Little AC. 2009 Pregnancy coloration in macaques may act as a
 warning signal to reduce antagonism by conspecifics. *Behav. Processes* 80, 7–11.
 (doi:10.1016/j.beproc.2008.08.001)
- Ziegler TE, Prudom SL, Schultz-Darken NJ, Kurian AV & Snowdon CT. 2006 Pregnancy
 weight gain: marmoset and tamarin dads show it too. *Biol. Lett.* 2, 181–183.
 (doi:10.1098/rsbl.2005.0426)
- 5. Altmann S. 1973 The pregnancy sign in savannah baboons. J. Zoo Anim. Med. 4, 8–12.
- Nunn CL. 1999 The evolution of exaggerated sexual swellings in primates and the graded-signal hypothesis. *Anim. Behav.* 58, 229–246. (doi:10.1006/anbe.1999.1159)
- 203 7. Drea CM. 2014 D'Scent of man: A comparative survey of primate chemosignaling in
 204 relation to sex. *Horm. Behav.* (doi:10.1016/j.yhbeh.2014.08.001)
- 8. Hradecký P. 1986 Volatile fatty acids in urine and vaginal secretions of cows during reproductive cycle. *J. Chem. Ecol.* 12, 187–196.
- Jemiolo B, Andreolini F, Wiesler D & Novotny M. 1987 Variations in mouse (Mus musculus) urinary volatiles during different periods of pregnancy and lactation. *J. Chem. Ecol.* 13, 1941–1956.
- 10. Theis KR, Venkataraman A, Dycus JA, Koonter KD, Schmitt-Matzen EN, Wagner AP,
 Holekamp KE & Schmidt TM. 2013 Symbiotic bacteria appear to mediate hyena social
 odors. *Proc. Natl. Acad. Sci.* **110**, 19832–19837. (doi:10.1073/pnas.1306477110)
- 11. Vaglio S, Minicozzi P, Bonometti E, Mello G & Chiarelli B. 2009 Volatile signals during
 pregnancy: a possible chemical basis for mother–infant recognition. *J. Chem. Ecol.* 35, 131–
 139. (doi:10.1007/s10886-008-9573-5)
- Sauther ML. 1991 Reproductive behavior of free-ranging Lemur catta at Beza Mahafaly
 Special Reserve, Madagascar. *Am. J. Phys. Anthropol.* 84, 463–477.
- 13. Charpentier MJE, Boulet M & Drea CM. 2008 Smelling right: the scent of male lemurs
 advertises genetic quality and relatedness. *Mol. Ecol.* 17, 3225–3233. (doi:10.1111/j.1365-220 294X.2008.03831.x)

- 14. Boulet M, Charpentier MJ & Drea CM. 2009 Decoding an olfactory mechanism of kin
 recognition and inbreeding avoidance in a primate. *BMC Evol. Biol.* 9, 281.
 (doi:10.1186/1471-2148-9-281)
- 15. Boulet M, Crawford JC, Charpentier MJE & Drea CM. 2010 Honest olfactory ornamentation
 in a female-dominant primate. *J. Evol. Biol.* 23, 1558–1563. (doi:10.1111/j.14209101.2010.02007.x)
- 16. Crawford JC, Boulet M & Drea CM. 2011 Smelling wrong: hormonal contraception in
 lemurs alters critical female odour cues. *Proc. R. Soc. B* 278, 122–130.
 (doi:10.1098/rspb.2010.1203)
- Pereira ME. 1995 Development and social dominance among group-living primates. *Am. J. Primatol.* 37, 143–175. (doi:10.1002/ajp.1350370207)
- 18. Scordato ES & Drea CM. 2007 Scents and sensibility: information content of olfactory
 signals in the ringtailed lemur, Lemur catta. *Anim. Behav.* 73, 301–314.
 (doi:10.1016/j.anbehav.2006.08.006)
- 19. Scordato ES, Dubay G & Drea CM. 2007 Chemical composition of scent marks in the
 ringtailed lemur (Lemur catta): glandular differences, seasonal variation, and individual
 signatures. *Chem. Senses* 32, 493–504. (doi:10.1093/chemse/bjm018)
- 238 20. Drea CM, Boulet M, Delbarco-Trillo J, Greene LK, Sacha CR, Goodwin TE & Dubay GR.
 239 2013 The 'secret' in secretions: methodological considerations in deciphering primate
 240 olfactory communication. *Am. J. Primatol.* **75**, 621–642. (doi:10.1002/ajp.22143)
- 241 21. Crawford JC & Drea CM. In review. Data from: Baby on board.
- 242 22. Charpentier MJE, Crawford JC, Boulet M & Drea CM. 2010 Message 'scent': lemurs detect
 243 the genetic relatedness and quality of conspecifics via olfactory cues. *Anim. Behav.* 80, 101–
 244 108. (doi:10.1016/j.anbehav.2010.04.005)
- 245 23. van Schaik C, Hodges J & Nunn C. 2000 Paternity confusion and the ovarian cycles of
 246 female primates. In *Infanticide by Males and Its Implications*, pp. 361–387. Cambridge:
 247 Cambridge University Press.
- 248 24. van Schaik CP, Pradhan GR & van Noordwijk MA. 2004 Mating conflict in primates:
 249 infanticide, sexual harassment and female sexuality. In *Sexual selection in primates: new and*250 *comparative perspectives.*, pp. 131–150. Cambridge: Cambridge University Press.
- 25. Matsumoto-Oda A, Kutsukake N, Hosaka K & Matsusaka T. 2006 Sniffing behaviors in
 Mahale chimpanzees. *Primates* 48, 81–85. (doi:10.1007/s10329-006-0006-1)
- 253 26. Nishida T. 1997 Sexual behavior of adult male chimpanzees of the Mahale Mountains
 254 national park, Tanzania. *Primates* 38, 379–398. (doi:10.1007/BF02381879)

- 27. Wild G & West SA. 2007 A Sex Allocation Theory for Vertebrates: Combining Local
 Resource Competition and Condition-Dependent Allocation. *Am. Nat.* 170, E112–E128.
 (doi:10.1086/518205)
- 258 28. Saragusty J, Hermes R, Hofer H, Bouts T, Göritz F & Hildebrandt TB. 2012 Male pygmy
 259 hippopotamus influence offspring sex ratio. *Nat. Commun.* 3, 697.
 260 (doi:10.1038/ncomms1700)
- 261 29. Robert KA & Braun S. 2012 Milk composition during lactation suggests a mechanism for
 262 male biased allocation of maternal resources in the tammar wallaby (Macropus eugenii).
 263 *PLoS ONE* 7, e51099. (doi:10.1371/journal.pone.0051099)
- 264 30. Cameron EZ & Linklater WL. 2007 Extreme sex ratio variation in relation to change in
 265 condition around conception. *Biol. Lett.* 3, 395–397. (doi:10.1098/rsbl.2007.0089)
- 31. Hohenbrink P, Mundy NI, Zimmermann E & Radespiel U. 2012 First evidence for functional
 vomeronasal 2 receptor genes in primates. *Biol. Lett.* 9. (doi:10.1098/rsbl.2012.1006)
- 32. Krause ET, Kruger O, Kohlmeier P & Caspers BA. 2012 Olfactory kin recognition in a songbird. *Biol. Lett.* 8, 327–329. (doi:10.1098/rsbl.2011.1093)

270