

posters

P – 198

Circulating microRNAs in metastatic colorectal cancer (mCRC) patients (pts) treated with regorafenib

M. Schirripa¹, F. Loupakis¹, C. Cremolini¹, L. Polisenò², L. Salvatore¹, A. Tuccoli³, C. Antoniotti⁴, R. D'Aurizio⁵, F. Marmorino¹, B. Borelli⁶, D. Rossini¹, A. Saettini¹, S. Gini⁴, R. Moretto⁷, I. Rizzo³, E. Dell' Aquila⁸, M. Pellegrini⁹, A. Falcone⁸

¹Oncologia Medica 2, Azienda Ospedaliero Universitaria Pisana, Istituto Toscano Tumori, Pisa, Italy

²Oncogenomics Unit, CRL-Istituto Toscano Tumori, IFC-CNR, Pisa, Italy

³Oncogenomics Unit, CRL-Istituto Toscano Tumori, Pisa, Italy

⁴Oncologia Medica 2, Azienda Ospedaliero-Universitaria Pisana, Istituto Toscano Tumori, Pisa, Italy

⁵Laboratory of Integrative Systems Medicine (LISM), Institute of Informatics and Telematics (IT) and Institute of Clinical Physiology (IFC), CNR, Pisa, Italy

⁶Oncologia Medica 2, Azienda Ospedaliero Universitaria Pisana, Pisa, Italy

⁷Oncologia Medica 2, Azienda Ospedaliero-Universitaria Pisana, Pisa, Italy

⁸Department of Translational Research and New Technologies In Medicine, University of Pisa, Pisa, Italy

Introduction: Regorafenib is indicated for the treatment of mCRC patients who have failed all other therapies. Nevertheless a substantial percentage of patients experiences rapid disease progression (PD) and serious adverse events may occur. For these reasons, clinical and/or molecular markers able to improve the cost/benefit ratio are

urgently needed. Circulating microRNAs (c-miRNAs) have been recognized as possible prognostic and diagnostic markers in mCRC. The aim of this study was to describe the early changes in plasma levels of 10 selected c-miRNAs during the treatment with regorafenib and to investigate their correlation with clinical outcome.

Methods: Plasma samples of patients treated with regorafenib at our Institution were collected at baseline (D1) and after 15 days of treatment (D15). Plasma levels of c-miR-17, c-miR-21, c-miR-29, c-miR-34, c-miR-92, c-miR-126, c-miR-141, c-miR-221, c-miR-601, c-miR-760 were analysed by means of real-time PCR. Paired levels at D1 and D15 were compared by means of Wilcoxon test for each c-miRNA. C-miRNAs showing significant changes were further analysed in order to identify possible correlations with outcome.

Results: Thirty-four patients were included in the present study. Main characteristics were the following: M/F = 50%/50%; median age = 65 (range 48-78 years); ECOG-PS 0/1-2 = 71%/29%; time from diagnosis of metastases </ ≥ 18 months 15%/85%. Median PFS and OS were 2.4 and 6.5 months, respectively. One (3%) patient achieved a response and 16 (47%) had disease stabilization (disease control rate: 50%). As compared to D1, the following c-miRNAs increased at D15: c-miR-601 (p = 0.01), c-miR-141 (p = 0.04) and c-miR-21 (p = 0.06). Despite a median increase in the overall population, 12 (35%) out of 34 patients showed reduced level of c-miR-21 at D15. Nine out of 12 (75%) patients with reduced levels of c-miR-21 achieved disease control, as compared to 8 out of 23 (35%) patients with increased levels (Fisher's Exact Test, p = 0.035). Median PFS of patients with increased and decreased level of levels of c-miR-21 were 2.1 and 3.9 months, respectively (HR = 1.89 95%CI 0.92-4.14 p = 0.08). Data on OS are not yet mature. Early modifications of c-miR-21 levels showed a sensitivity of 82% in predicting benefit from regorafenib.

Conclusion: The early modulation of c-miR-21 levels may predict benefit from regorafenib in terms of disease control. These results need validation in independent series.