

Ammonium Thiosulphate Assisted Phytoextraction of Mercury and Arsenic in Multi-Polluted Industrial Soil

Francesca Pedron^{1,*}, Gianniantonio Petruzzelli¹, Irene Rosellini¹, Meri Barbafieri¹,
Elisabetta Franchi², Roberto Bagatin²

¹Institute of Ecosystem Study, National Council of Research, Pisa, Italy

²Eni S.p.A, Renewable Energy & Environmental Laboratories, S.Donato Milanese (MI), Italy

Abstract The possibility of using ammonium thiosulphate in assisted phytoextraction was evaluated on a greenhouse scale (mesocosm) for the simultaneous removal of mercury and arsenic from multi-polluted industrial soil. The addition of thiosulphate to the soil greatly promoted the uptake and translocation of both contaminants in the aerial parts of *Brassica juncea* and *Lupinus albus*. Thiosulphate showed great potential since it is a common fertilizer used to promote plant growth and is able to promote plant uptake of both Hg and As. Hg concentration in the aerial part of the plants reached 867 mg kg⁻¹ in *B. juncea* and 114 mg kg⁻¹ in *L. albus*. In the aerial parts, As concentration was about 9 mg kg⁻¹ in *B. juncea* and 20 mg kg⁻¹ in *L. albus*. This thus increases the applicability of phytoextraction in terms of cost and time especially if the remedial targets are based on bioavailable metal concentrations.

Keywords Thiosulphate, Phytoextraction, Mercury, Arsenic, Greenhouse experiment, Multi-contaminated soil, *Brassica juncea*, *Lupinus albus*, Bioavailability, Repeated growing cycles

1. Introduction

Mercury and arsenic are widely distributed throughout the environment in the air, water and soil. Both contaminants are present in the atmosphere from volcanoes activity and wildfires. As trace elements, they are naturally present in soils due to their geological origin. Soil weathering can release As and Hg compounds which may be dispersed by wind and rainfall water. However, mercury and arsenic constitute one of the largest environmental problems in former industrial sites, due to the very high concentrations in the soil in comparison to background values [1-3]. Depending on their speciation, which influences their mobility and bioavailability in soil, these two contaminants are particularly dangerous for human health through the direct inhalation of contaminated dust and hand-to-mouth ingestion. Arsenic and mercury are also some of most hazardous contaminants in drinkable and groundwater throughout the world since they can be assimilated and bioaccumulated with occasional biomagnification [4].

Exposure to mercury represents a great hazard for human health, since Hg has no biological function. The most toxic form of Hg, methylmercury, is not present in well-aerated

soil as it is mainly produced in aquatic environments [1]. In the soil, Hg can be involved in adsorption and release from solid phases, complexation with organic and inorganic ligands, oxidation-reduction, and methylation [5]. Depending on the redox conditions, Hg can exist in soil as Hg⁰, Hg²⁺, Hg₂²⁺. Mercury can be present in many organic and inorganic complexed forms depending on pH, ionic strength, and dissolved organic matter in soil [6]. Many compounds of Hg with sulphide are highly stable and insoluble, while inorganic ligands such as Cl⁻ ions promote the release of Hg in the soil solution due to the formation of soluble stable complexes [7].

Long-term exposure to arsenic from the food chain and drinking-water can cause many significant health problems with the development of neurotoxicity and cardiovascular diseases [8-9]. Arsenic and arsenic compounds have also been classified as carcinogenic to humans [10]. Arsenic is present in soil mainly in inorganic forms, the two most commonly found species are As(III) and As(V). In oxidizing conditions, arsenate As(V) species is the most stable, forming strong inner-sphere complexes with the oxides of iron, manganese and aluminium and partly with clay minerals [11-12]. Arsenic interactions with soil organic matter are particularly effective at positively charged sites where it can form both "inner-sphere" and "outer-sphere" complexes [13].

In alkaline conditions, As is only slightly involved in adsorption to soil components, while with decreasing pH values the adsorption and precipitation of this element

* Corresponding author:

francesca.pedron@ise.cnr.it (Francesca Pedron)

Published online at <http://journal.sapub.org/re>

Copyright © 2015 Scientific & Academic Publishing. All Rights Reserved

increase thus reducing the amount in soil solution [14].

Soil properties that determine Hg and As retention by mineral and organic surfaces are mainly pH, oxides, hydroxides and dissolved ions. Clayey and organic soils have the greatest ability to adsorb these elements. If soils are neutral to alkaline, Hg shows stronger adsorption to mineral components, while As is mainly present in the soil liquid phase. A decrease in pH reduces the adsorption processes of Hg [7], but increases As adsorption and precipitation [12]. The different chemical behaviour of the two metals makes the remediation of soils contaminated from both the elements particularly difficult. However, the presence of more than one metal or metalloid in soil is widespread in contaminated sites. Often in these cases excavation and landfilling, soil washing and stabilization / inertization are the most used solutions for remediation.

Among the alternative approaches based on sustainability and the conservation of natural resources, phytoextraction, the use of plants to extract inorganic pollutants from soil, may be a viable solution. Phytoremediation has been intensively investigated due to its environmental friendly approach to remediation, which favours greater social acceptance [15]. Phytoremediation was originally based on the use of metal hyperaccumulator plants, which are characterized by the ability to accumulate high amounts of heavy metals or metalloids [16]. However, the use of hyperaccumulators is often hindered by their reduced production of biomass and by the ability to accumulate only one specific element, which makes them impractical in multi-contaminated soil. As an alternative “assisted phytoextraction” is commonly used. In this technology, high biomass plant species are used together with the modification of soil properties by chemical additives, which increase the bioavailability of the contaminants. Many additives, in particular chelating agents such as ethylenediaminetetra-acetic acid (EDTA), have been commonly used [17-18]. However, these agents persist in the soil with residual toxic effects and there is a risk of metal leaching into the groundwater. New mobilizing agents, which are under study, have no adverse effects on the environment while promoting the bioavailability of contaminants. Among these, thiosulphate appears particularly attractive for mercury remediation since it is a common fertilizer used to promote plant growth and form stable soluble complexes with mercury in soil [19-20-21], thus promoting the uptake of the metal by growing plants. While thiosulphate has been successfully used in many contaminated soils to increase mercury phytoextraction, recent research has shown that thiosulphate competes with arsenate ions for sorption on the same surface of oxides. This competition may, in turn, increase the release of arsenic in the liquid phase of soil and its bioavailability to plant uptake [22]. These results suggested that thiosulphate could be used also to phytoextract arsenic. This solution would be of considerable interest in soils contaminated with both elements.

The aim of this paper was to evaluate the applicability of

phytoextraction technology to reduce the bioavailable fraction of Hg and As from a soil contaminated by the two elements using only one additive: ammonium thiosulphate. Greenhouse experiments were carried out to compare also the efficiency of thiosulphate with phosphate, which is the typical mobilizing agent for arsenic in assisted phytoextraction.

2. Materials and Methods

2.1. Soil Characterization

The soil used in this study was sampled from a former industrial site, where various industrial activities had been carried out. As and Hg contamination was discovered with mean values of 37.6 and 65.8 mg kg⁻¹, respectively. Soil samples were air dried and sieved with a 2 mm sieve for soil analysis. The following parameters were determined: pH, cation exchange capacity (CEC), texture (sand silt and clay) and organic matter [23]. Readily bioavailable mercury and arsenic in the soil solution were determined by H₂O extraction by shaking the soil and extractant (ratio 1:5) for 3h, which is often used as the first step in heavy metal sequential extractions [24]. Hg and As were analyzed in the extracts after centrifugation and filtration. Long-term potential bioavailability was determined using 0.27 M ammonium thiosulphate for Hg and 0.05 M potassium biacid phosphate for As, according to the procedure of Enhanced Bioavailable Contaminant Stripping (EBCS) [20-25]. The extractability of As was also tested with 0.27 M ammonium thiosulphate.

2.2. Mesocosm Experiment

To obtain a representative sample of the field situation, the soil used in these experiments was prepared by eliminating (sieving at 2 cm) the coarser materials, however without sieving to 2 mm. Experiments were carried out in a greenhouse. Mesocosms were polypropylene containers filled with 4 kg of soil. Each pot had a hole in the middle of the base where a plastic tube was inserted to collect the leachate in a plastic bottle [20].

The mesocosm experiments lasted 60 days. The plant species selected for the tests, *Brassica juncea* and *Lupinus albus*, were considered viable candidates for phytoremediation due to their ability to grow in a Mediterranean climate and their relative tolerance to heavy metal soil contamination [18-20]. A total of 1.0 g per pot of *B. juncea* seeds or nine seeds per pot of *L. albus* were used in four replicates for each species. Thirty days after sowing, a solution of ammonium thiosulphate 0.27 M (NH₄)₂S₂O₃ (T treatment) or a solution of potassium biacid phosphate 0.05 M KH₂PO₄ (P treatment) as mobilizing agents for Hg [19-21] and for As [26], respectively. After the first harvest a second growing cycle was performed. On some mesocosms the mobilizing agents were added with the same doses as the first cycle (T/T, P/P), in others no further treatment (nt) was carried out (T/nt, P/nt).

Both additives were added to the soil by splitting the total

dose into five-day applications to minimize the possible toxic effects on the plant species. Control mesocosms (CT) for each species were run simultaneously in untreated soil. At the end of the growing period, plants were harvested and aerial parts and roots were separated. Vegetal samples were washed with deionized water; the roots were further washed in an ultrasound bath (Branson Sonifier 250 ultrasonic processor; Branson, Danbury, Conn.) for 10 min to eliminate any soil particles from the root surfaces. The dry biomass of shoots and roots was gravimetrically determined after maintaining samples at 40°C until a constant weight was achieved. The dry plant samples were ground and digested with an acid mixture (HNO₃+H₂O₂) for As and Hg analysis.

2.3. Determination of As and Hg

As concentration in soil, plant samples, and soil extracts was determined by ICP-OES (Varian AX Liberty) with a method for the generation of hydrides [23]. The mercury concentration in the same samples was determined using atomic absorption spectroscopy with an Automatic Mercury Analyzer (AMA 254, FKV, Bergamo, Italy), according to the SW-846 method 7473 (27 U.S. EPA, 1998).

2.4. Quality Assurance and Quality Control

Quality assurance and quality control were performed by testing standard solution every 10 samples. Certified reference material (BCR n°141) was used to control the quality of analytical system. The detection limits were 2 µg L⁻¹ for Hg and 0.05 mg L⁻¹ for As, respectively.

The recovery of spiked samples ranged from 93 to 101% with a RSD of 1.89 of the mean.

2.5. Statistical Analysis

Statistical analyse was performed using Statistica version 6.0 (Statsoft, Inc.). Effects of mobilizing agents were analyzed using one-way analysis of variance (ANOVA). Differences among means were compared and a post-hoc analysis of variance was performed using the Tukey Honestly and significant differences considered at $p < 0.05$.

3. Results and Discussion

3.1. Soil Characteristics

Soil was characterized by a pH value of 8.06, 1.5% of organic matter and a cation exchange capacity of 15.6 cmol(+) kg⁻¹. The soil can be defined as sandy according to the texture values: 78.9% sand, 13.1% silt and 8.0% clay.

To evaluate the potential efficiency of thiosulphate as a solubilizing agent, Hg and As extractability was determined by 0.27 M (NH₄)₂S₂O₃ extraction. As extractability was also determined by 0.05 M KH₂PO₄ the typical mobilizing agent for arsenic. Results showed that (NH₄)₂S₂O₃ was able to extract the same amounts of As as those extracted by phosphate (Table 1). These results highlighted the effectiveness of thiosulphate as single mobilizing agent for

assisted phytoextraction in this contaminated soil.

Table 1. Extractability of As and Hg. Data are the means of three replicates and are expressed as mg kg⁻¹ dry soil

Extractant	As	Hg
0.05 M KH ₂ PO ₄	7.6±0.86	nd
0.27 M (NH ₄) ₂ S ₂ O ₃	7.4±0.65	12.9±0.88

3.2. Biomass Production and Metal Uptake

The biomass data, expressed as g dry weight, are reported in Table 2, in which for the first growing cycle the controls are indicated by CT, while P and T are the treated mesocosms with phosphate and thiosulphate, respectively. For the second growing cycle the letters after the slash indicates the presence (P/P or T/T) or not (P/nt or T/nt) of a second treatment. Control pots were obviously not treated.

In the first growing cycle, a biomass reduction of the two plant species was observed following thiosulphate treatment (T). In the second growing cycle similar results were obtained following the repeated addition of thiosulphate (T/T treatment) in the case of the aerial part of *L. albus*, while for *B. juncea* data of the biomass production were not statistically different among the treatments. The effect of thiosulphate was more significant in the reduction of biomass of the roots for which, on the contrary, the addition of phosphate produced a slight biomass increase in the case of *B. juncea*.

Table 2. Biomass production (g) in the two growing cycles

	Treatment	Aerial part		Roots	
		<i>B. juncea</i>	<i>L. albus</i>	<i>B. juncea</i>	<i>L. albus</i>
Growing cycle 1	CT	3.22 ^b	4.18 ^b	0.43 ^b	0.72 ^b
	P	3.36 ^b	4.60 ^b	0.46 ^b	0.66 ^b
	T	1.82 ^a	2.85 ^a	0.13 ^a	0.15 ^a
Growing cycle 2	CT/CT	4.01 ^a	6.33 ^b	0.46 ^b	1.04 ^c
	P/nt	4.10 ^a	6.26 ^b	0.73 ^c	0.88 ^c
	P/P	3.20 ^a	6.68 ^b	0.62 ^c	0.97 ^c
	T/nt	3.90 ^a	5.02 ^b	0.24 ^a	0.54 ^b
	T/T	3.88 ^a	2.53 ^a	0.16 ^a	0.16 ^a

Note: Separately for each growing cycles, values in each column with the same letter were not significantly different from each other, according to the Tukey test ($p < 0.05$).

The efficiency of phytoextraction depends on both the contaminant concentration in plants and their biomass. The reduction in plant biomass after thiosulphate treatment may be due to the greater absorption by plants of the mobilized Hg. In fact, mercury can induce toxicity, thus hindering the growth of plants [28]. These effects of yield reduction are not ascribable to As. This is also confirmed by the fact that the phosphate treatment did not influence the yield even increasing the As uptake by plants. In the second growing cycle, when the amount of Hg taken up by the plants is significantly lower, the yields of *B. juncea* from (T/nt) and (T/T) mesocosms and that of *L. albus* from (T/nt) mesocosms were similar to the control plants. The roots yield

showed a different trend to the one of the aerial part of the plants with a reduction following thiosulphate treatments (T/nt and T/T).

Concentrations of contaminants in plant tissues are described separately for Hg and As. The Hg concentration values in plants are reported in Table 3.

Table 3. Hg concentration (mg kg^{-1}) in the selected species in the two growing cycles

	Treatment	<i>B. juncea</i>		<i>L. albus</i>	
		Aerial part	Roots	Aerial part	Roots
Growing cycle 1	CT	1.56±0.87	38.7±10.0	0.96±0.07	10.8±2.4
	P	0.13±0.02	39.9±15.4	0.15±0.07	7.2±1.8
	T	867±99	356±60.1	114±10.2	1469±200
Growing cycle 2	CT/CT	n.d.	32.2±10.1	n.d.	25.1±2.2
	P/nt	n.d.	53.2±12.4	n.d.	28.2±1.9
	P/P	n.d.	55.6±15.8	n.d.	22.4±4.8
	T/nt	43.4±11.0	162±28.3	36.9±6.3	142±10.3
	T/T	181±15.3	722±41.4	57.4±9.8	1015±101

Note: n.d. = not detectable

As expected, in both growing cycles the phosphate treatment did not influence the Hg absorption by plants, due to the formation of stable precipitates among Hg and phosphate ions. In the first growing cycle, the thiosulphate treatment was particularly effective. The concentration of Hg in plants, which was about 1.5 mg kg^{-1} in the controls, increased to around 900 mg kg^{-1} in the aerial part of the *B. juncea*. A similar trend also occurred for *L. albus*, the concentration values after the T treatment increased to 100 mg kg^{-1} , while in untreated plants the Hg concentration was 0.96 mg kg^{-1} . Also in the case of the root portion, the effect of the T treatment produced a very effective increase compared to the controls where the concentrations of Hg were much lower. Regarding *L. albus*, the absorption of mercury after the T treatment resulted in values of concentrations close to 1500 mg kg^{-1} , while the mean Hg concentration in the roots of *B. juncea* was 356 mg kg^{-1} . These values are much higher than in the plants grown in the control mesocosms, where the mean Hg concentrations were 38.7 and 10.8 mg kg^{-1} for *B. juncea* and *L. albus*, respectively. A comparison between the two plant species after T treatment, showed significant differences in the Hg uptake process, in fact *B. juncea* absorbed higher amounts of metal in the aerial part while *L. albus* absorbed more in the root system.

Also in the second growing cycle, plants were able to uptake a further quantity of Hg following treatment with thiosulphate. In the mesocosms treated only once (T/nt), the mean mercury concentration in the aerial part of *B. juncea* was 43.4 mg kg^{-1} , while following a repeated T treatment (T/T) it increased to a mean value of 181 mg kg^{-1} . The metal concentration in the aerial part of the untreated plants (CT/CT) was essentially negligible. A similar trend was also found in the root portion where the mean Hg concentration was 162 mg kg^{-1} in the T/nt treatment and 722 mg kg^{-1} after T/T treatment.

In *L. albus*, the mean Hg concentration in the aerial part increased from a negligible value in the control (CT/CT) to 36.9 mg kg^{-1} (T/nt) and to 57.4 mg kg^{-1} (T/T). In the root portion, the concentrations increased from 25.1 mg kg^{-1} in the control (CT/CT) to 142 mg kg^{-1} (T/nt) and up to 1015 mg kg^{-1} after the repeated treatment (T/T).

These results also confirm the ability of the thiosulphate to increase the bioavailability of Hg in the soil. In the control soil, the high concentration of Hg did not produce any visual symptoms of toxicity, however these symptoms appeared in the treated soil after the third day of adding thiosulphate. As known, the addition of thiosulphate to soil promotes the release of Hg from soil surfaces and the formation of the soluble complex $[\text{Hg}(\text{S}_2\text{O}_3)_2]^{2-}$ which is available for plant uptake [19-20]. Often thiosulphate promotes the translocation of Hg in the aerial parts of the plants [20]. However, in this case study the addition of thiosulphate did not increase the translocation of mercury from the plant roots to the shoots. The Hg concentrations were higher in the roots than in the shoots with the exception of *B. juncea* after thiosulphate treatment in the first growing cycle. This may depend both on the varieties of plants used and on the specific soil characteristics.

The concentration values of As in plants are reported in Table 4.

Table 4. As concentration (mg kg^{-1}) in the selected species in the two growing cycles

	Treatment	<i>B. juncea</i>		<i>L. albus</i>	
		Aerial part	Roots	Aerial part	Roots
Growing cycle 1	CT	3.1±0.87	3.9±0.78	12.0±1.1	6.6±1.4
	P	11.6±2.5	18.6±2.6	18.9±3.1	11.5±1.0
	T	8.7±1.4	14.8±1.9	20.0±2.6	13.5±1.2
Growing cycle 2	CT/CT	n.d.	n.d.	n.d.	n.d.
	P/nt	4.2±0.98	16.2±1.4	n.d.	8.0±1.4
	P/P	10.9±1.1	20.9±1.7	6.3±0.82	8.7±1.1
	T/nt	4.2±0.70	6.0±0.80	1.5±0.50	4.7±0.80
	T/T	9.7±0.83	8.6±1.1	6.3±0.60	8.6±0.99

Note: n.d. = not detectable

In both growing cycles, As concentrations in plants generally increased in the treated mesocosms compared to the control pots. In the first growing cycle, mean As concentrations in the aerial parts of *B. juncea* were 11.6 and 8.7 mg kg^{-1} after the addition of P and T, respectively, with no statistically significant differences between the two treatments. In the roots, these values were higher, 18.6 and 14.8 mg kg^{-1} . Similar trends were found in *L. albus* plants, with an increase in As concentration both in aerial parts and in roots, following the P and T treatments. Also in these cases, no statistically significant differences were observed between the concentrations in plants after P or T addition. Irrespectively of whether being treated with T or P, the mean As concentrations in the aerial parts were about 19 mg kg^{-1} and about 12 mg kg^{-1} in the roots. All these concentration values were higher than those of the plants grown in the

controls.

In the second growing cycle, the As concentration in the control plants was negligible. When the addition of phosphate and thiosulphate was not repeated (P/nt) and (T/nt), the amount of As absorbed by the plants was greater than the controls but generally less than that absorbed during the first cycle. Only with a second addition of phosphate (P/P) or thiosulphate (T/T), and only in *B. juncea* plants the As concentration values in the aerial parts were similar to those obtained in the first cycle. Following the (P/P) treatment, the concentration values in *B. juncea* were similar to those in the first growth cycle both in the aerial parts and in the roots, but only in the aerial part after the (T/T) treatment. In the case of *L. albus*, all the concentration values in the second cycle were lower than those found in the first cycle for both the roots and the shoots.




Figure 1. Hg total uptake (μg) of the two plant species in mesocosm experiment

3.3. Total Accumulation

The product of As and Hg concentration values in shoots and the corresponding biomass, expressed as "total accumulation" [29], provides an estimate of the amount of contaminants removed from the polluted soil and therefore of the efficiency of phytoextraction. The data obtained using the mean values in the calculation are reported in Figure 1 for mercury and Figure 2 for arsenic.

In the case of Hg, the addition of thiosulphate, as expected, increased the amount of bioavailable metal and consequently the accumulation by the plants was promoted. As previously described, this can be ascribed to the formation of soluble complexes in the soil between Hg and thiosulphate ion.




Figure 2. As total uptake (μg) of the two plant species in mesocosm experiment

These complexes are taken up by roots and transported to the above ground parts of the plants [19-21]. In the case of *B. juncea*, the amount of Hg accumulated after T treatment was about 1600 μg in the first growing cycle and around 700 μg in the second after a further treatment with thiosulphate. In the controls and in the pots treated with phosphate, these values were negligible. With regard to *L. albus*, the amount of Hg accumulated in the plants was lower than that found for *B. juncea*, but the trend was similar. In the first growing cycle, after the T treatment the value obtained was 325 μg , while after the second cycle, this decreased to about 160 μg (T/nt and T/T). In the controls and after the P treatment, the accumulation was much lower or even below the detection limit.


Regarding the As accumulated in the aerial parts of the plants, both treatments promoted the absorption of the contaminant compared to the controls, thus confirming the mobilizing action on As not only of phosphate, but also of thiosulphate [22]. In the case of *B. juncea*, in the first growing cycle, plants grown in the control pots showed a mean total accumulation of 10 μg . After treatments with thiosulphate and phosphate this value increased to 15.8 and 39.0 μg , respectively. In the second growing cycle, the highest total accumulation (36 μg) was determined in the plants after a repeated treatment (T/T and P/P), while in the P/nt and T/nt pots the value was about 17 μg .

In the case of *L. albus* in the first growing cycle, the total accumulation was higher than that obtained for *B. juncea* and ranged from a minimum of 50.2 μg in the controls to a


maximum of 86.9 μg after the P treatment. In the second growing cycle, the maximum accumulation was determined after the repeated treatment with phosphate P/P (42.3 μg). A comparison between the two tested plant species should take into account the simultaneous presence of the two contaminants, Hg and As. The greatest amount of Hg was taken up by *B. juncea* following the T treatment, which also promoted a notable increase in As uptake compared to the controls. *L. albus* accumulated higher amounts of As than *B. juncea* but much lower amounts of Hg. Thus, the use of *B. juncea* in combination with T treatment appears to be the best available choice in this contaminated soil.

The data obtained with the use of thiosulphate should be considered very positively since they show that only one mobilizing agent was able to simultaneously increase the absorption of Hg and As by plants.

The addition of thiosulphate has often led to very positive results for Hg phytoextraction in different contaminated soils [19-20-21]. In contrast, the effect of thiosulphate on the bioavailability of As has not been widely investigated and is of great interest. A recent study on the competitive adsorption between arsenate and thiosulphate ions, highlighted the great influence of thiosulphate on As adsorption by hematite [22]. The competition can be ascribed to an inner-sphere complexation, with the possible formation of monodentate non protonated surface complexes similar to those formed by phosphate in the adsorption on hematite [30]. Thiosulphate ion can be considered as a sulphate ion and a sulphide ion linked by a double bond [31]. In soil, thiosulphate decomposes into sulphur and sulphate. Sulphur can give rise to precipitates while sulphate remains in solution according to the following reaction: thiosulphate, tetrathionate, sulphite, sulphate.



The reaction is either abiotic or biotic with kinetics of oxidation depending on the chemical and biological characteristics of the soil. In the presence of plants, which increase the microbiological activity, the transformation of thiosulphate may follow a different pattern [32] without producing tetrathionate as an intermediate:



Thus, thiosulphate may substitute arsenate on iron oxides by the sulphate part of the molecule, with oxygen atoms bonding to the surface Fe ions. This competition is essential in determining the As available for plant uptake, since arsenate in soil is mainly retained by Fe oxides. Thiosulphate ions competing for sorption on the same surface may significantly reduce the arsenate adsorption and enhance the mobility of arsenic in soil, similarly to the influence of phosphate on arsenate. In alkaline soil, the competition between the two anions is very high [33] thus, we can assume that the excess of sulphate ions, deriving from the addition of thiosulphate, released arsenate ions in the soil solution, which were thus bioavailable for plant uptake.

However, it is also important to consider the interactions

between sulphur and arsenic. Sulphur can promote arsenic absorption and transport to the aerial parts of the plants, since sulphur plays an anti-stress role in reducing the toxicity of arsenic [34]. Thiosulphate can thus act as a detoxifying agent by stimulating the defence system of plants while increasing the efficiency of phytoremediation due to competition between arsenate and sulphate ions for the same sites on the soil surfaces.

One of the main drawbacks of assisted phytoextraction is the potential leaching of the contaminants solubilized by the additives used, however in our experimental conditions, no arsenic and mercury were found in the small amounts of leachate produced following the irrigation necessary for the plant growth. However, as a precautionary principle, we determined the Hg and As water extractable in soil after the first and the second growing cycles. We can suppose that the addition of thiosulphate mobilizes contaminants and the plants are able to uptake only a part of this amount.

Following the T treatment, the soluble amount of Hg was around 2.5 mg kg^{-1} after the first growing cycle. Without further T addition, after the second growing cycle, the values decreased to about 0.7 mg kg^{-1} and 1.3 mg kg^{-1} for pots planted with *B. juncea* and *L. albus*, respectively. In the mesocosms treated twice with thiosulphate (T/T) the water soluble Hg amount in soil was about 1.5 mg kg^{-1} for all the pots. The results showed that a certain amount of mercury remained in the soil, bioavailable for further growing cycles essentially in the rhizosphere. In fact, water extractable Hg decreased after the second growth. These results are in accordance with previous findings [21] that reported an increase in soluble mercury in the rhizosphere following thiosulphate addition did not correspond to an increase in the bulk soil. The results were explained as the consequence of mercury complex decomposition with the production of sulphate ions and subsequently a decrease in mobile mercury species [21].

Similar results were obtained for As. Following the T treatment, the water extractable As after the first growing cycle accounted for 1.4 and 0.91 mg kg^{-1} for *B. juncea* and *L. albus* mesocosms, respectively. After the second growing cycle without any further T addition (T/nt), these extractable amounts decreased slightly to 0.80 and 0.65 mg kg^{-1} in the pots planted with *B. juncea* and *L. albus*, respectively. The amounts increased slightly to 1.30 and 0.80 mg kg^{-1} when the T treatment was repeated (T/T) in the pots planted with *B. juncea* and *L. albus*, respectively.

This procedure, which involves H_2O extraction at the end of the growing cycle, may provide a linkage between plant uptake and the residual bioavailable fraction of the two metals. This is useful when planning other growing cycles aimed at reducing and possibly eliminating the amounts of bioavailable Hg and As which are the most dangerous for humans and the environment [25]. Unlike the use of chelating agents such as EDTA, it is not necessary to apply large amounts of thiosulphate to counteract the co-solubilization of Ca due to the low solubility of CaSO_4 . The risk of additive leaching is negligible, in contrast to the

case of poorly biodegradable chelating agents, such as EDTA. Adverse effects on soil quality can be ruled out since the decomposition of thiosulphate produced sulphur and sulphate, which are essential for fertility.

The efficiency of phytoextraction in cleaning up metal-contaminated soils has been questioned due to the long time required to reach the remediation target based on total metal concentration. However, not all the total metals in soils may be involved in environmental processes. There is growing interest in using phytoextraction to reduce only the bioavailable fractions, which are the most dangerous to the environment and human health. The clean-up time can thus be substantially shortened and the imbalance between the potential of phytoextraction and full-scale applicability can be overcome [20].

The potential use of plants for phytoextraction can be derived from the ratio between the concentration of Hg and As in the shoots and the total concentration of the two elements in soil. This relationship is often called the phytoextraction coefficient (PEC) or translocation factor (TF). If the aim of phytoextraction is to remove the bioavailable fraction, this coefficient is calculated as the ratio between the concentration of the metal in the aerial parts of the plants and the bioavailable concentration in the soil. This ratio has been defined bioavailability factor (BF) [35].

Table 5. Effect of P and T treatments and growing cycles on the bioavailability factor (BF) of Hg and As

		Mercury		Arsenic	
		Treatment			
		<i>B. juncea</i>	<i>L. albus</i>	<i>B. juncea</i>	<i>L. albus</i>
First cycle	CT	0.12	0.07	0.39	1.6
	P	0.01	0.01	1.5	2.5
	T	66.8	8.7	1.1	2.7
Second cycle	CT/CT	n.c.	n.c.	n.c.	n.c.
	P/nt	n.c.	n.c.	0.55	n.c.
	P/P	n.c.	n.c.	1.4	0.81
	T/nt	3.3	2.8	0.55	0.15
	T/T	13.8	3.9	1.2	0.85

Note: n.c. = not calculable because Hg and As concentrations in the aerial part of plants were not detectable

Table 5 summarizes the BF data considering the ratio between the concentration in the plant shoots following each treatment and the potential bioavailability of the contaminants determined by the use of 0.27 M thiosulphate. As previously reported, the As extractable by 0.27 M thiosulphate was the same as that extracted by 0.05 M phosphate.

This coefficient can be used to evaluate plants species in a short-growing period in terms of phytoremediation efficiency. When this coefficient is calculated for a remediation target based on the total concentration a value between 0.5 and 60 is considered acceptable [36]. We can

hypothetically assume that the same values might also be indicatively acceptable when the remediation target is based on the bioavailable amounts and the BF coefficient is used.

Our data support the potential for successfully exploiting these plants after the addition of thiosulphate to reduce the bioavailable contaminants concentrations in this soil. The BF coefficient highlights the identical efficiency of the treatment with phosphate or thiosulphate for arsenic and the possibility of removing the bioavailable fraction of As also in presence of Hg in soil.

4. Conclusions

Hg and As are widespread in contaminated sites and are often present simultaneously thus entailing separate remediation strategies, due to their different chemical characteristics and different behaviour in relation to soil properties. Among clean-up procedures, phytoextraction may be a viable clean up procedure based on the sustainability and conservation of natural resources. High biomass plant species could be used together with a modification of soil properties by chemical additives, which increase the bioavailability of the contaminants (assisted phytoextraction). In this study, an innovative approach based on the use of a single mobilizing agent, ammonium thiosulphate, was tested at a greenhouse scale.

Our results show that the addition of thiosulphate to the soil greatly promoted the uptake and translocation of both contaminants in the aerial parts of *B. juncea* and *L. albus*. Thiosulphate, which is specifically used to mobilize Hg in soil, was also particularly efficient in mobilizing As in this soil.

The comparison between phosphate (specific mobilizing agent for As) and thiosulphate in the As extraction efficiency showed that they extracted the same amounts of this contaminant from the soil. Thiosulphate has no adverse effects on the environment since it is a common fertilizer used to promote plant growth and at the same time, it is able, with different mechanisms, to increase the bioavailability of both Hg and As. The use of the same mobilizing agent to clean up soil contaminated simultaneously with both Hg and As could be of great interest in the phytoextraction process by greatly reducing both time and costs.

ACKNOWLEDGEMENTS

This study was supported by Eni S.p.A, Research & Technological Innovation Department, San Donato Milanese (Italy). Contract n°. 3500023724 and fully funded by Syndial S.p.A.

REFERENCES

- [1] AMAP/UNEP, 2013. Technical Background Report for the

- Global Mercury Assessment 2013. Arctic Monitoring and Assessment Programme, Oslo, Norway/UNEP Chemicals Branch, Geneva, Switzerland.
- [2] Garelick, H., Jones, H., Dybowska, A., and Valsami-Jones, E. 2008. Arsenic pollution sources. In Garelick, H. and Jones, H. (Eds.) *Arsenic Pollution and Remediation: an International Perspective*, Springer, New York, pp. 17-60.
- [3] Mukherjee, A.B., Zevenhoven, R., Brodersen, J., Hylander, L.D., Bhattacharya, P., 2004, Mercury in waste in the European Union: sources, disposal methods and risks, *Resour. Conserv. Recycl.*, 42: 155–182.
- [4] Mann, E., Meyer, T., Mitchell, C.P.J. and Wania, F., 2011, Mercury fate in ageing and melting snow: Development and testing of a controlled laboratory system, *J. Environ. Monitor.*, 13: 2695-2702.
- [5] Jing, Y.D., He Z.L., Yang X.E., 2008, Adsorption–Desorption Characteristics of Mercury in Paddy Soils of China, *J. Environ. Qual.*, 37: 680-688.
- [6] Arbestain, M.C., Rodriguez-Lado, L.R., Bao, M., Macias F., 2009, Assessment of Mercury-Polluted Soils Adjacent to an Old Mercury-Fulminate Production Plant, *Applied Environ. Soil Sci.*, 2009: 1-8.
- [7] Schuster, E., 1991, The behaviour of mercury in the soil with special emphasis on complexation and adsorption processes: a review of the literature, *Water Air Soil Pollut.*, 56: 667-680.
- [8] Fernández, M.I., López, J.F., Vivaldi, B. and Coz, F., 2012, Long-term impact of arsenic in drinking water on bladder cancer health care and mortality rates 20 years after end of exposure, *J. Urol.*, 187: 856-61.
- [9] Hughes, M.F., Beck, B.D., Chen, Y., Lewis, A.S., and Thomas, D.J., 2011, Arsenic Exposure and Toxicology: A Historical Perspective, *Toxicol. Sci.*, 123: 305-332.
- [10] Straif, K., Benbrahim-Tallaa, L., Baan, R., Grosse, Y., Secretan, B., El Ghissassi, F., Bouvard, V., Guha, N., Freeman, C., Galichet, L., 2009, A review of human carcinogens—Part C: metals, arsenic, dusts, and fibres, *Lancet Oncol.*, 10: 453–454.
- [11] Catalano, J.G., Park, C., Fenter, P. and Zhang, Z., 2008, Simultaneous inner and outer-sphere arsenate adsorption on corundum and hematite, *Geochim. Cosmochim. Acta*, 72: 1986-2004.
- [12] Dixit, S. and Hering, J., 2003, Comparison of arsenic V and arsenic III sorption onto iron oxide minerals: Implications for arsenic mobility, *Environ. Sci. Technol.*, 37: 4182–4189.
- [13] Bauer, M., and Blodau, C., 2006, Mobilization of arsenic by dissolved organic matter from iron oxides, soils and sediments, *Sci. Tot. Environ.*, 354(2): 179-190.
- [14] Masscheleyn, P.H., Delaune, R.D. and Patrick, W.H., 1991, Effect of Redox Potential and pH on Arsenic Speciation and Solubility in a Contaminated Soil, *Environ. Sci. Technol.*, 25: 1414–1419.
- [15] Barbaferi, M., Japenga, J., Romkens, P., Petruzzelli, G. and Pedron, F. 2013. Protocols for Applying Phytotechnologies. In Gupta, D.K. (Ed.) *Plant-Based Remediation Processes*, Springer-Verlag, Berlin, Heidelberg, pp. 19–37.
- [16] Brooks, R.R., 1998. *Plants that hyperaccumulate heavy metals*. CAB, New York.
- [17] Doumett, S., Lamperi, L., Checchini, L., Azzarello, E., Mugnai, S., Mancuso, S., Petruzzelli, G., Del Bubba, M., 2008, Heavy metal distribution between contaminated soil and *Paulownia tomentosa*, in a pilot-scale assisted phytoremediation study: influence of different complexing agents, *Chemosphere*, 72: 1481–1490.
- [18] Pedron, F., Rosellini, I., Petruzzelli, G., Barbaferi, M., 2014, Chelant Comparison for Assisted Phytoextraction of Lead in Two Contaminated Soils, *Res. Environ.*, 4(5): 209-214.
- [19] Moreno, F.N., Anderson, C.W.N., Stewart, R.B., Robinson, B.H., Ghomshei, M., Meech, J.A., 2005, Induced plant uptake and transport of mercury in the presence of sulphur-containing ligands and humic acid, *New Phytol.*, 166: 445-454.
- [20] Pedron, F., Petruzzelli, G., Barbaferi, M., Tassi, E., 2013, Remediation of a mercury-contaminated industrial soil using bioavailable contaminant stripping, *Pedosphere*, 23 (1): 104-110.
- [21] Wang, J., Feng, X., Anderson, C.W.N., Wang, H., Wang, L., 2014, Thiosulphate-induced mercury accumulation by plants: metal uptake and transformation of mercury fractionation in soil - results from a field study, *Pl. Soil*, 375: 21–33.
- [22] Petruzzelli, G., Pedron, F., Tassi, E., Franchi, E., Bagatin, R., Agazzi, G., Barbaferi, M., Rosellini, I., 2014, Effects of thiosulfate on the adsorption of arsenate on hematite with a view to phytoextraction, *J. Environ. Earth Sci.*, 6 (6): 326-332.
- [23] Sparks, D.L., 1998. *Methods of Soil Analysis. Part 3. Chemical Methods*, Soil Science Society of America Book Series. Soil Science Society of America, Madison, USA.
- [24] Pedron, F., Petruzzelli, G., Barbaferi, M., Tassi, E., 2009, Strategies to use phytoextraction in very acidic soil contaminated by heavy metals, *Chemosphere*, 75: 808-814.
- [25] Petruzzelli, G., Pedron, F., Rosellini, I., Barbaferi M. 2013. Phytoremediation towards the future: focus on bioavailable contaminants. In Gupta, D.K. (Ed.) *Plant-Based Remediation Processes*, Springer-Verlag, Berlin, Heidelberg, pp. 273-289.
- [26] Tassi, E., Pedron, F., Barbaferi, M., Petruzzelli, G., 2004, Phosphate-assisted phytoextraction in As-contaminated soil, *Eng. Life Sci.*, 4: 341-346.
- [27] U.S. EPA 1998 SW-846 Method 7473. Mercury in solids and solutions by thermal decomposition, amalgamation and atomic absorption spectrophotometry. Method 7473, U.S. EPA, Washington, D.C.
- [28] Perez-Sanz, A., Millan, R., Sierra, M.J., Alarcon, R., Garcia, P., Gil-Diaz, M., Vazquez, S., Carmen Lobo, M., 2012, Mercury uptake by *Silene vulgaris* grown on contaminated spiked soils, *J. Environ. Manag.*, 95: 5233-5237.
- [29] Jarrell, W.M. and Beverly, R.B., 1981, Dilution effect in plant nutrition studies, *Advan. Agron.*, 34: 197-224.
- [30] Elzinga, E.J., and Sparks, D.L., 2007, Phosphate adsorption onto hematite: An in situ ATR-FTIR investigation of effects of pH and loading level on the mode of phosphate surface complexation, *J. Colloid Interf. Sci.*, 308: 53-70.
- [31] Benjamin, M.M., and Leckle, M., 1982, Effects of

- complexation by $C1$, SO_4 and S_2O_3 on adsorption behavior of Cd on oxide surfaces, *Environ. Sci. Technol.*, 76: 162-170.
- [32] Barbosa-Jefferson, V.L., Zhao, F.J., McGrath, S.P., Magan, N., 1998, Thiosulphate and tetrathionate oxidation in arable soils, *Soil Biol. Biochem.*, 30: 553-559.
- [33] Myneni, S.C.B., Traina, S.J., Waychunas, G.A. and Logan, T.J., 1998, Vibrational spectroscopy of functional group chemistry and arsenate coordination in ettringite, *Geochim. Cosmochim. Acta*, 62: 3499-3514.
- [34] Duan, G., Liu, W., Xueping, C.X., Ying, H.Y., Zhu, Y., 2013, Association of arsenic with nutrient elements in rice plants, *Metallomics*, 5: 784-792.
- [35] Mandal, A., Purakayastha, T.J., Patra, A.K., Sanyal, S.K., 2012, Phytoremediation of arsenic contaminated soil by *Pteris vittata* influence of phosphatic fertilizers and repeated harvests, *Int. J. Phytorem.*, 14: 978-995.
- [36] Ampiah-Bonnet, R.J., Tyson, J.F. and Lanza, G.R., 2007, Phytoextraction of arsenic from soil by *Leersia oryzoides*, *Int. J. Phytorem.*, 9: 31-40.