

Multidisciplinary Investigations at Panarea (Aeolian Islands) after the Exhalative Crisis of 2002

G. Bortoluzzi¹, S. Aliani², M. Ligi¹, F. D'Oriano¹, V. Ferrante¹, F. Riminucci¹, C. Carmisciano³, L. Cocchi³, F. Muccini³

1, Institute of Marine Sciences, CNR, Bologna, Italy

2, Institute of Marine Sciences, CNR, Pozzuolo di Lerici (SP), Italy

3, National Institute of Geophysics and Volcanology, La Spezia, Italy

g.bortoluzzi@ismar.cnr.it

Abstract

Panarea and surrounding Islets form a volcanic edifice, that is part of the Eastern sector of the Aeolian Arc, Southern Tyrrhenian Sea. It is now considered inactive, since last documented activity is 20 Ka old. However, on 2002-11-03, gas started to flow violently from the seafloor in an area E of the Island, mainly along NE and NW structural lineaments, and lasting up to 2003-2004 with a consistent flux, orders of magnitude larger than 'steady-state' fumarolic activity documented there in historical times. On the same period a strong effusive activity of Stromboli (10 NM to NNE) was present. Since then, several investigations have been conducted at sea and on land, with the aim of focusing on the problem of effusive activity at sea, mainly in the light of volcanic surveillance and risk. Among these investigations, some of which have been repeated over years, we present and discuss some data and results from: (a) visual inspection and sampling by divers and ROV, (b) GPS networks and mapping by multibeam and LIDAR, (c) oceanographical measurements by current meters and CTD, and water flux and dynamics measurements, (d) magnetic and gravimetric surveys, (e) multichannel reflection Seismic with OBS and land station networks. Data were used for compilation of high resolution bathymetric, magnetic and gravimetric maps, including the emerged and submerged portions of the edifice.

1 Introduction and setting

On 2002-11-03 a burst of gas occurred in the marine area E of Panarea, lasting for years with a consistent flux from fractures and sinkholes on the seafloor, mostly near the islet of Bottaro (Figure 1). Investigation started immediately to monitor this event from geological and geochemical point of views, also in the light of volcanological surveillance and risk [1, 2, 3, 4, 5, 6, 7], and on the possible connection to regional tectonics [8, 9, 10, 11].

This paper aims at providing a review of the geophysical investigations carried out in the area of the eruption since 2002. The Aeolian Islands are part of the volcanic arc formed by the convergence of the African and Eurasian plates and by the subduction and southeastward rollback of the Ionian lithosphere [12, 13, 14, 15, 16, 17, 18], and is characterized by compression in the western sector, strike-slip faulting and extension in the central and eastern ones. The archipelago is formed by 7 islands and minor islets, including to-




Figure 1: Panarea volcanic complex. Bathymetries from ISMAR. Also shown position of MCS lines L09, L10, L11.

day's active Stromboli volcano. Panarea is considered inactive, however [19] have shown possible recent volcanic outcrops near Basiluzzo; present deformation patterns are likely connected to NE-SW trending faults [20].

The gas release of 2002-11-03 in the area E of Panarea, known since historical times for fumarolic activities [21], generated 6-7m diameter columns of bubbles from the seafloor to the surface. Several active spots were identified by divers and ROVs' and by repeated multibeam surveys [3, 5]. The

most impressive one was just SW of Bottaro (PEG1, Figure2) with gas reaching the surface from 15m depth, from an elliptic depression produced by the explosive collapse of the seafloor; a plume of suspended sediments was present at the sea surface for days.

During the most active degassing up to mid 2003, the emissions were found to be an emulsion of CO₂-dominated gas phase with suspended sediments, colloidal sulfur; the water was acidified by dissolution of SO₂, HCl and HF [2, 7]. [1] estimated a

RESP	CRUISE	Date	Ship	Mapping	F	C	M	G	D	R
ISMAR	1994	1996-11-01	J.Charcot	Sim.EM1000						X
ISMAR	TIR96	1996-09-01	Gelenzhik	Sim.EM12			X			
ISMAR	TIR96	1999-02-01	Strakhov	Sim.EM12			X			
ISMAR-INGV	P2002-11	2002-11-07	Thetis			X			X	X
ISMAR	P2002-12	2002-12-10	Alea	Res.8125	X		X		X	
ISMAR	P2003-01	2003-01-20	Thetis							
ISMAR	P2003-07	2003-07-27	Alea	Res.8125		X	X			
ISMAR	P2003-09	2003-09-02	L.Sanzo		X	X			X	X
ISMAR	P2003-12	2003-12-10	Alea	Res.8125					X	
ISMAR-NERC	P2004-04	2004-04-20	plane	LIDAR		X				
ISMAR	P2006-01	2006-01-20	Alea	Res.8125					X	
ISMAR-IMM-INGV	P2006-04	2006-04-15	Aretusa	Kon.EM3000			X			
INGV		2006-04-15	land					X		
ISMAR	P2006-05	2006-05-02		SeaInterf.						
ISMAR	PANA07	2007-08-01	Urania	Res.8160						
ISMAR-INGV	CALA08	2008-04-02	Urania	Res.8160				X		
ISMAR-INGV	PANSTR10	2010-02-01	Urania	Kong.EM710			X	X		

Table 1: Data Acquisition Cruises. Measurements: F=Water-gas Fluxes; C=CTD; M=Magnetics, G=Gravity; D=dive; R=ROV.

gas output of $10^9 \text{l}\cdot\text{d}^{-1}$ (November 2002, all emissions) and of 4 to $2 \times 10^7 \text{l}\cdot\text{d}^{-1}$ (May to July 2003, PEG1), orders of magnitude higher than the total gas output of $10^6 \text{l}\cdot\text{d}^{-1}$ measured within the Islets in the 1980's [21]. [22] measured the water fluxes at PEG1, deriving also the gas fluxes. The plumes affected the marine environment with changes in the biota [23, 24] and in the water properties [25]. Figure 2a shows the location of the major emissions (named PEG1,2,3,8).

2 Materials and Methods

Several cruises were performed in the area for obtaining geophysical and oceanographic data and to monitor the geomorphological features of the seabed and the evolution of the gas outflow after the 2002-11 crisis. The investigations were coordinated by the Italian Department of

Civil Protection (DPC) and 'Commissione Grandi Rischi' and were carried out from immediately after the gas burst up to 2008 (Table 1).

Multibeam data from different cruises and instruments were processed with the Kongsberg's Neptune and RESON's PDS2000 software. Gridding was performed in the geographical and UTM33 projections, at spatial resolution ranging from 10-15m for deeper areas, to 0.20-0.25m for local, shallow areas. Some datasets have been processed separately due to large number of points acquired. Furthermore, a particular attention was paid to the analysis of the gas emissions (Figure2) included in the multibeam data [3]. A LIDAR flight was performed by NERC on April 2004 by the *Airborne Remote Sensing Facility* using an Optechh ALTM 3033 laser scanner; the processed data included first and last pulses and ob-

tained data at the resolution of $\sim 0.2\text{m}$ [26]. Magnetic data were acquired during cruises TIR96, TIR99, P2002-12, P2003-07 and PANSTR10, with GEM GSM19D and Marine Magnetics Sea-Spy 'Overhauser' magnetometers; during cruise P2006-04 the Geometrics G-800 'Cesium magnetometer was used [27]. Data underwent filtering, de-spiking, cross-over error reduction, application of IGRF Models with 2005 coefficients for calculating anomalies and reduction to the Pole. Multichannel data (48 active, 12.5 m group interval, 2xGI Harmonic mode), were acquired during cruise PANA_07, complementing MESC2001 ISMAR's cruise (<http://www.ismar.cnr.it/prodotti/reports-campagne>). On the same cruise a seismic network was set on Islets and on the seafloor by deploying seismometers and OBSs from INGV and University of Trieste. Seismic shots sequences were performed along lineaments connecting the instruments. The gravity data on land [27] were sampled using a pair of LaCoste&Romberg microgravimeters (Aliod model) equipped with a digital data acquisition system, GPS tracking and automatic tide corrections, with a nominal resolution of $1\ \mu\text{gal}$. Marine gravity data were acquired with a Lacoste&Romberg 'AirSea' gravimeter, directly interfaced to DGPS, during cruises CALAMARE08 and PANSTR10 (<http://www.ismar.cnr.it/prodotti/reports-campagne>). Data were de-spiked, corrected for drifts and Eötvös effects, and Free air and Bouguer anomalies were calculated.

Flux of water entering the vent at PEG-1 was estimated by geometry of the ascending gas column and by measuring water velocities inside and outside the gas column. Rotor (Aanderaa and Datasonics) and ADCP current meters were positioned 1m above the sea floor in the gas, and few m away, respectively (Figure 4). Estimates of the venting surface was done by divers and high resolution bathymetries, since multi-beam used were able to detect the gas in the water column. Once obtained the water fluxes entering at the base and exiting the vent, the gas fluxes were estimated by applying reduction factors accounting for (a) the bubble sizes and voids, as seen by divers and ROV, and (b) considerations about reduced velocities at the boundary layers of the cell.

R/V Thetis and R/V L.Sanzo deployed ROV systems, on 2002-11, few days after the burst, 2003-01 and 2002-09. During this last deployment the instrument was also moored on bottom in front of the degassing area for hours, aiming at recording flagged poles on seafloor and dye releases able to visualize the water dynamics at the base of the vent. ROV records confirmed the bathymetric and divers investigations about changes in topography and geometry of the gas column.

CTD investigations were performed on 2002-07, 2002-09 and 2004-04, to evaluate possible plumes and modifications of water column properties. Standard sensors mounted on SBE probe were used and pH was found to be particularly effective for tracing the acid fluid releases.




Figure 2: Gas emitting points (a) and 3D rendering of PEG1 multibeam data (b). Photos on top: (c) the divers' positioning the lander with a rotor current meter into the gas column, and (d) the exposed cemented breccia at the vertical borders of the sinkhole. Images from [2,4], modified.




Figure 3: Area E of Panarea (slope shading). Topography by NERC's LIDAR flight.



Figure 4: September 2003 flux-measurement experiment: the ADCP and rotor current meter (within gas, on the frame) are visible.




Figure 5: Bathymetry and diving observations on 2002-12. Also shown the temperature measured at the seafloor close to emitting spots. Sea water temperature was $\sim 17^{\circ}\text{C}$.

3 Results and Discussion

3.1 Bathymetry

ISMAR performed a high resolution multi-beam bathymetric cruise on 1994 [29]. A wide portion of the Aeolian Central sector was imaged down to ~ 800 m depth, except for very shallow waters, including the area of the Islets. The TIR96 and TIR99 cruises [30] mapped the deeper portions. After 2002 crisis several cruises were able to map almost entirely the submerged edifice, collecting also data near the shoreline

of Panarea and Islets.

Panarea and islets (Figure 1, 3, [31]) emerge from a volcanic edifice (diameter of ~ 18 km at the -1000 m isobath), dissected by gullies and channels and largely dismantled by erosion and by neo- and volcano-tectonics. Its flat summit, with edge at about 80-130 m bsl, is almost totally covered by volcanoclastic sedimentation, arranged in its upper part in sequences of terraced, wedge-shaped prograding units [32]. Several, partly buried, primary volcanic features are present and partially outcrop in the southern sector, among them the




Figure 6: Bathymetry 2002-12, 2003-12, one year after. Also shown the difference in depth.

shoal of Secca del Capo [33, 34, 19, 29, 35].

East of the island, a relief of ~ 1.5 km diameter is present at depth < 30 m, partially emerging in the Islets and enclosing depressed areas where intense exhalative activity occurred. The northeastern summit of the edifice presents a NNE-SSW and NE-SW structural lineament (on the ~ 20 m high fault scarps, fresh rocks and mineralization as well as gas venting are present [36, 29, 35]) and further extends from Basiluzzo with NE-SW direction.

The repeated bathymetries over the years on the PEG1 area 5 showed that less than a year after the gas burst, the depression

was already partially infilled by sediments transported from the flanks by the dismantling of the sub-vertical wall to the N, exposing cemented breccias and rocks of holocene age [5] (Figure6). The survey of 2006-01 further confirmed this infilling, producing variation of bathymetries of 2-3m. Pebbles rolling on the seafloor under wave and current dynamics kept on filling the sinkhole and very likely will produce a new non-active meter size depression filled with sandy or gravel materials similar to others discovered in the area by multibeam. On 2006 and 2007 the flux of gas was visibly reduced to a small area, and similarly the transport caused by entrainment at the




Figure 7: Magnetic anomaly data. 2002-12 Alsea cruise.

base of the degassing cell also reduced. The wave and current dynamics will probably be able to continue the filling process on the medium period. On the plateau just N of the depression, a wide area was covered by sediments, very likely ejected during the explosive collapse of 2002-11-03, and the sizes and water depth seem suitable for bedload transport and distribution. ROV on the field few days after revealed sand dunes and ripples crossing at 30-40°.

3.2 Magnetics and Gravity

On 2002-12 ISMAR collected high resolution data within the Islets (Figure7). The same pattern of lines was repeated by Geophysics and Marine Technology Unit of Istituto Nazionale di Geofisica e Vulcanologia (INGV) of Portovenere in Spring 2006, together with gravity data on the islets and on Panarea [27]. Magnetic anomaly pattern of islets area is dominated by the high positive (500-750 nT) north of Panarelli shoal in correspondence of a topographic high. This anomaly seems related with the




Figure 8: 2D Apparent magnetization map. Structural lineaments from [5], exhalative centers from [28].

main tectonic lineaments and it can be interpreted as the signature of an ensemble of magmatic sources. Rock-sampling by [5] showed andesitic lava products correlated to a shallow cryptodome-like structure. A 2D inversion of the data was performed to evaluate the magnetization pattern of the area, using an FFT-algorithm (Parker inverse approach, [37, 38]), applied to a crustal portion of 1Km below sea bottom. Figure 8 suggests a clear separation between the high magnetized region between Dattilo and Basiluzzo and the region among Lisca Bianca, Bottaro and Lisca Nera where the exhalative crisis of 2002 occurred. In this region the magnetiza-

tion pattern decreases with a null-value strongly driven by the hydrothermal alteration which affects the seafloor. During cruise CALAMARE08 marine gravity data around Panarea and Islets were acquired, and were integrated with the above cited survey on land (see Figure 9). The two dataset were collected by different methodologies and instruments, and the merging was achieved without any fictitious grid-knitting process but using the absolute gravity data from a station in Panarea for reducing the offsets. The off-shore gravity mapping shows several gap, making these results preliminary, while awaiting for on-going data processing of




Figure 9: Free Air gravity anomaly map.

cruise PANSTR10.

Multichannel seismic acquired during cruise PANA07 were shot for investigating the regional tectonic and setting, other than for illuminating the OBS and land station seismological network. The data provide important information on plio-quadernary sediment thickness, to be used for proper gravimetric analysis and modelling. Some examples of the lines are on Figures 10, and 11 (positioning data in Figure1).

3.3 Flux estimation and Oceanography

The water and gas fluxes that we estimated for the main emission at PEG1 [22] are of the same order of magnitude of those calculated as in [1]. The experiment of December 2002 encountered problems be-

cause of the gas flowing violently from the depression; divers experimented strong ascensional force and heavy ballasts had been used for securing the instrumentation on bottom. Similarly [1] was able to measure the flux of gas at PEG1 only on 2002-05, due to such difficulties. Figure 12 shows the data obtained during the 2003-09 experiment and the instruments' deployment. The ADCP data have been influenced by tidal components, while rotor current meter data had a rather constant flux, being anchored well into the gas column. The ADCP to the E measured a much higher number of erratic values, and this is probably due to the beams being invested directly by gas bubbles, which are known to strongly reduce the data quality. These errors depend on the entrainment of gases at intermediate depths, in contrast, with data of 2002-12 when the gas ascending




Figure 10: MCS Line L09, neartrace.

speed was higher and diverged at the surface without reaching buoyancy at intermediate depths. The CTD data of 2003-07 were able to depict a plume of acidified water (pH <6.5) centered on the PEG1. The data of 2004 cruise, instead, showed pH anomaly mostly centered on the PEG8 site to the N, which is up to now the lonely active emitting area left.

4 Conclusions

The Panarea area has been largely investigated through a number of geophysical surveys in the area and a comprehensive description of the event was achieved through an intensive and continuous monitoring. A further step in multidisciplinary knowledge of degassing events was put forward and a number of papers has been published. A lot of scientific discussion arose from this large work and many new insights have been highlighted. This brief description summarizes some of the results.

- High resolution bathymetric surveys have imaged important structural lineaments and morphologies, and helped in studying the evolution of gas emissions near the Islets;
- High resolution magnetic and gravity data provided insights into the volcanic complex; a marked decrease of magnetization is present in the exhalative areas, whereas strong values were found just to the N and NW of Dattilo and the Panarelli;
- The importance of measuring the effects of the eruptions on seawater properties and dynamics has been stressed, also as a possible new tool for monitoring the evolution of hydrothermal activity over time. Strong pH anomalies were reported and seawater was largely acidified. The effects of the acidification of seawater is an important issue and shallow vent systems are excellent sites where the effects of ocean acidification can be studied. Diving and morphobathymetric in-




Figure 11: MCS Lines L10 and L11, neartraces.

vestigations reported of the infilling of PEG1 by sediment transport, suggesting a fate similar to other depressions found in the area.

5 Acknowledgments

We thank E. Bonatti, who supported our research and involved ISMAR since the early beginning. The project was funded by INGV/Dipartimento Protezione Civile and CNR. We thank the Coast Guard of Lipari for logistic support, the friends of Coastal Consulting Exploration of Bari, F. De Giosa and S. Lippolis, the Captain and crew of R/V L. Sanzo, Pippo Arena of Are-

nasub. The inhabitants of Panarea, especially Pina, her husband and family made the work a lot easier with their great hospitality and help. The 1994 multibeam survey was funded by Italian Ministry of Industry. We also acknowledge NERC (UK) that kindly provided LIDAR data. Most of the maps and figures were done by the Paul Wessel's GMT package [39]. Dr. Alina Polonia of ISMAR and the Hydrographic Institute of the Italian Navy helped with R/V Urania and Aretusa ship time. Finally, we would like to remember the work that was done in November 2002 and January 2003 at Panarea and Stromboli by R/V Thetis, which sunk dramatically on August 2007.




Figure 12: September 2003 flux-measurement experiment.

References

- [1] S. Caliro, A. Caracausi, G. Chiodini, M. Ditta, F. Italiano, M. Longo, C. Minopoli, P.M. Nuccio, A. Paonita, and A. Rizzo. Evidence of a recent input of magmatic gases into the quiescent volcanic edifice of Panarea, Aeolian Islands, Italy. *Geophys. Res. Lett.*, 31:L07619, 2004. doi:10.1029/2003GL019359.
- [2] B. Capaccioni, F. Tassi, D. Vaselli, D. Tedesco, and P.L. Rossi. The November 2002 degassing event at Panarea Island (Italy): the results of a 5 months geochemical monitoring program. *Annals of Geophysics*, 48(4-5):755–765, 2005.
- [3] M. Anzidei, A. Esposito, G. Bortoluzzi, and F. Degiosa. The high resolution map of the exhalative area of Panarea (Aeolian Is., Italy). *Annals of Geophysics*, 48(6):899–921, 2005.
- [4] A. Caracausi, M. Ditta, F. Italiano, M. Longo, P.M. Nuccio, A. Paonita, and A. Rizzo. Changes in fluid geochemistry and physico-chemical conditions of geothermal systems caused by magmatic input: The recent abrupt outgassing off the island of Panarea (Aeolian Islands, Italy). *Geochim. Cosmochim. Ac.*, 69(12):3045–3059, 2005. doi:10.1016/j.gca.2005.02.011.
- [5] A. Esposito, G. Giordano, and M. Anzidei. The 2002-2003 submarine gas eruption at Panarea volcano Aeolian Islands, Italy: volcanology of the seafloor and implications for the hazard scenario. *Marine Geology*, 227:119–134, 2006.
- [6] G. Chiodini, S. Caliro, G. Caramanna, D. Granieri, C. Minopoli, R. Moretti, L. Perrotta, and G. Ventura. Geochemistry of the submarine gaseous emissions of Panarea (Aeolian Islands, Southern Italy): magmatic vs. hydrothermal origin and implications for volcanic surveillance. *Pure Appl. Geophys.*, 163(4):759–780, 2006. doi:10.1007/s0024-006-0037-y.
- [7] B. Capaccioni, F. Tassi, O. Vaselli, D. Tedesco, and R. Poreda. Submarine gas burst at Panarea Island (Southern Italy) on 3 November 2002: A magmatic versus hydrothermal episode. *J. Geophys. Res.*, 112:B05201, 2007.
- [8] J. Heinicke, F. Italiano, R. Maugeri, B. Merkel, T. Pohl, M. Schipek, and T. Braun. Evidence of tectonic control on active arc volcanism: The Panarea-Stromboli tectonic link inferred by submarine hydrothermal vents monitoring (Aeolian arc, Italy). *Geophys. Res. Lett.*, 36:L04301, 2009. doi:10.1029/2008GL036664.
- [9] V. Acocella, M. Neri, and T.R. Walter. Structural features of Panarea volcano in the frame of the Aeolian Arc (Italy): Implications for the 2002-2003 unrest. *J. Geodyn.*, 47(5):288–292, 2009. doi:10.1016/j.jog.2009.01.004.
- [10] A. Billi and R. Funicello. Concurrent eruptions at Etna, Stromboli, and Vulcano: casualty or causality? *Annals of Geophysics*, 51(4):655–725, 2008.

- [11] T.R. Walter, R. Wang, V. Acocella, M. Neri, H. Grosser, and J. Zschau. Simultaneous magma and gas eruptions at three volcanoes in southern Italy: an earthquake trigger? *Geology*, 37(3):251–254, 2009.
- [12] F. Barberi, P. Gasparini, F. Innocenti, and L. Villari. Volcanism of the Southern Tyrrhenian Sea and its geodynamic implications. *J. Geophys. Res.*, 78(23):5221–5232, 1973.
- [13] F. Barberi, F. Innocenti, G. Ferrara, J. Keller, and L. Villari. Evolution of the Eolian arc volcanism (Southern Tyrrhenian Sea). *Earth Planet. Sc. Lett.*, 21(3):269–276, 1974. doi:10.1016/0012-821X(74)90161-7.
- [14] A. Argnani and C. Savelli. Cenozoic volcanism and tectonics in the Southern Tyrrhenian Sea: space time distribution and geodynamic significance. *J. Geodyn.*, 27(4-5):409–432, 1999.
- [15] A. Argnani. The Southern Tyrrhenian Subduction System: Recent Evolution and Neotectonic Implications. *Annals of Geophysics*, 43(3):585–607, 2000.
- [16] N. Calanchi, A. Peccerillo, C.A. Tranne, F. Lucchini, P.L. Rossi, P. Kempton, W. Barbieri, and T.W. Woo. Petrology and geochemistry of volcanic rocks from the island of Panarea: implications for mantle evolution beneath the Aeolian island arc (Southern Tyrrhenian sea). *J. Volcanol. Geoth. Res.*, 115(3-4):367–395, 2002. doi:10.1016/S0377-0273(01)00333-X.
- [17] G. De Astis, G. Ventura, and G. Vilardo. Geodynamic significance of the Aeolian volcanism (Southern Tyrrhenian Sea, Italy) in light of structural, seismological, and geochemical data. *Tectonics*, 22(4):1040, 2003. doi:10.1029/2003TC001506.
- [18] G. Bortoluzzi, M. Ligi, C. Romagnoli, L. Cocchi, D. Casalbore, T. Sgroi, F. Caratori Contini, M. Cuffaro, F. D’Oriano, V. Ferrante, A. Remia, and F. Riminucci. Interactions between volcanism and tectonics in the Western Aeolian sector, Southern Tyrrhenian Sea. *Geophys. J. Int.*, 183:64–78, 2010. doi: 10.1111/j.1365-246X.2010.04729.x.
- [19] F. Gamberi, M. Marani, and C. Savelli. Tectonic, volcanic and hydrothermal features of a submarine portion of the Aeolian arc (Tyrrhenian Sea). *Marine Geology*, 140:167–181, 1997.
- [20] F. Lucchi, C.A. Tranne, N. Calanchi, and P.L. Rossi. Late Quaternary deformation history of the volcanic edifice of Panarea, Aeolian Arc. *Bull. Volcanol.*, 69:239–257, 2007.
- [21] F. Italiano and P.M. Nuccio. Geochemical investigations of submarine exhalations to the east of Panarea, Aeolian Islands, Italy. *J. Volcanol. Geoth. Res.*, 46:125–141, 1991.

- [22] S. Aliani, G. Bortoluzzi, G. Caramanna, and F. Raffa. Seawater dynamics and environmental settings after november 2002 gas eruption off Bottaro (Panarea, Aeolian Islands, Mediterranean Sea). *Cont. Shelf Res.*, 30(12):1338–1348, 2010.
- [23] C. Gugliandolo, F. Italiano, and T. Maugeri. The submarine hydrothermal system of Panarea (Southern Italy): biogeochemical processes at the thermal fluids - sea bottom interface. *Annals of Geophysics*, 49(2-3), 2006.
- [24] E. Manini, G.M. Luna, C. Corinaldesi, D. Zeppilli, G. Bortoluzzi, G. Caramanna, F. Raffa, and R. Danovaro. Prokaryote diversity and virus abundance in shallow hydrothermal vents of the Mediterranean Sea (Panarea Island) and the Pacific Ocean (North Sulawesi-Indonesia). *Microbial Ecol.*, 55(4):626–639, 2008.
- [25] F. Tassi, B. Capaccioni, G. Caramanna, D. Cinti, G. Montegrossi, L. Pizzino, F. Quattrocchi, and O. Valselli. Low-pH waters discharging from submarine vents at Panarea Island (Aeolian Islands, southern Italy) after the 2002 gas blast: Origin of hydrothermal fluids and implications for volcanic surveillance. *Appl. Geochem.*, 24(2):246–254, 2009.
- [26] M. Fabris, M. Anzidei, P. Baldi, G. Bortoluzzi, A. Pesci, and S. Aliani. The high resolution combined topographic model of Panarea island (Aeolian island, Italy). In *EGU General Assembly*, 2010. Vienna, 02-07 Maggio.
- [27] L. Cocchi, F. Caratori Tontini, C. Carmisciano, P. Stefanelli, M. Anzidei, A. Esposito, C. Del Negro, F. Greco, and R. Napoli. Looking inside the Panarea Island (Aeolian Archipelago, Italy) by gravity and magnetic data. *Annals of Geophysics*, 51(1):25–38, 2008.
- [28] M. Anzidei, P. Baldi, G. Casula, F. Riguzzi, and L. Surace. La rete Tyrgeonet. *Bollettino di Geodesia e Scienze affini*, LIV(2):1–20, 1995.
- [29] F. Gamberi, C. Savelli, M.P. Marani, M. Ligi, G. Bortoluzzi, V. Landuzzi, and M. Costa. Contesto morfo-tettonico e depositi idrotermali di solfuri ed ossidi di ferro in una porzione sommersa dell’arco eoliano (in base ad indagini ad alta definizione). *Boll.Soc.Geol.It.*, 117(1):55–71, 1998.
- [30] M.P. Marani, F. Gamberi, G. Bortoluzzi, G. Carrara, M. Ligi, and D. Penitenti. Tyrrhenian sea bathymetry. In M.P. Marani, F. Gamberi, and E. Bonatti, editors, *From seafloor to deep mantle: architecture of the Tyrrhenian backarc basin*, volume 44 of *Mem. Descr. Carta Geologica d’Italia*. APAT, 2004.
- [31] C. Romagnoli, D. Casalbore, G. Bortoluzzi, A. Bosman, F.L. Chiocci, F. D’Oriano, F. Gamberi, M. Ligi, and M. Marani. Bathy-morphological setting of the Aeolian Islands. In F. Lucchi, A. Peccerillo, J. Keller, C.A. Tranne, and P.L. Rossi, editors, *Geology of the Aeolian Islands*, *Memoirs*. Geological Society, London, 2010. Special Issue, in press.

- [32] F.L. Chiocci and C. Romagnoli. Terrazzi deposizionali sommersi nelle Isole Eolie (Sicilia). *Mem. Descrittive della Carta Geologica d' Italia, APAT*, 58:81–114, 2004.
- [33] G. Gabbianelli, P.Y. Gillot, G. Lanzafame, C. Romagnoli, and P. L. Rossi. Tectonic and volcanic evolution of Panarea (Aeolian Islands, Italy). *Marine Geology*, 92:313–326, 1990.
- [34] G. Gabbianelli, C. Romagnoli, P.L. Rossi, and N. Calanchi. Marine geology of the Panarea-Stromboli area (Aeolian Archipelago, Southeastern Tyrrhenian Sea). *Acta Vulcanol.*, 3:11–20, 1993.
- [35] F. Gamberi, C. Savelli, M.P. Marani, M. Ligi, G. Bortoluzzi, V. Landuzzi, A. Luppi, M. Badalini, and M. Costa. Carta morfo-batimetrica e dei depositi idrotermali in una porzione sommersa dell'arco eoliano. *Boll.Soc.Geol.It.*, 117(1), 1998. Supplement.
- [36] M.P. Marani, F. Gamberi, and C. Savelli. Shallow-water polymetallic sulfide deposits in the Aeolian island arc. *Geology*, 25(9):815–818, 1997.
- [37] L. Cocchi, F. Caratori Tontini, C. Carmisciano, P. Stefanelli, D. Embriaco, and M. Anzidei. Magnetic and Gravimetric model of Panarea (Aeolian islands). In *Convegno Nazionale MGMEESV*, 2006. Catania, 27-29 Settembre 2006.
- [38] F. Caratori Tontini, L. Cocchi, and C. Carmisciano. Potential-field inversion for a layer with uneven thickness: The Tyrrhenian Sea density model. *Phys. Earth Planet. In.*, 166(1-2):105–111, 2008.
- [39] P. Wessel and W.H.F. Smith. New version of the Generic Mapping Tools released. *Trans. Am. Geophys. Union (EOS)*, 76:329, 1995.