



<b>Title</b>	<b>The Involvement of phosphatidylinositol-3 kinase signaling in neurogenesis during <i>Xenopus</i> embryonic development</b>
<b>Author(s)</b>	<b>Peng, Y</b>
<b>Citation</b>	<b>43rd Annual Meeting of the American Society for Cell Biology, San Francisco, California, 13-17 December 2003, v. 14 n. Suppl, p. 515a</b>
<b>Issued Date</b>	<b>2003</b>
<b>URL</b>	<b><a href="http://hdl.handle.net/10722/54211">http://hdl.handle.net/10722/54211</a></b>
<b>Rights</b>	<b>Molecular Biology of the Cell. Copyright © American Society for Cell Biology.</b>

## The Involvement of Phosphatidylinositol-3 Kinase Signaling in Neurogenesis During *Xenopus* Embryonic Development

Y. Peng; Institute of Molecular Biology, The University of Hong Kong, Hong Kong, China

**Presentation Number:** 2888

**Poster Board Number:** B619

**Objective:** Phosphatidylinositol 3-kinase (PI3K) is a signaling molecule with a demonstrated role in numerous cellular activities. In *Xenopus* embryogenesis, PI3K acts in mesoderm induction and in dorsoventral patterning. To investigate the involvement of the PI3K subunits and PI3K signaling pathway in neural development during *Xenopus* embryogenesis, we report the outcome of experiments using loss-or-gain-of-function. **Methods and Results:** The P110 $\alpha$  and  $\Delta$ P85 are two subunits of PI3K, and Akt is downstream of PI3K in the signaling pathway. We show that overexpression of a constitutively active form of the catalytic subunit P110 $\alpha$  of PI3K, or a constitutively active form of Akt, produces double heads and axes and induces neural markers: NACM, Otx2, and Hoxb9 expression; whereas overexpression of  $\Delta$ P85, a truncated form of P85, or DN-Akt or GSK-3 $\beta$ , produces a ventral phenotype and induces ventral markers: Xvent1, Xvent2, BMP-4, and Xmsx1 expression. It has been demonstrated that GSK-3 $\beta$  antagonized Akt signaling. Furthermore, We has shown that  $\Delta$ P85 blacked P110 $\alpha$  activity, whereas DN-Akt and GSK-3 $\beta$  inhibited both P110 $\alpha$  and Akt activities. **Conclusion:** These results demonstrate that PI3K and the PI3k signaling pathway are involved in neural development during *Xenopus* embryogenesis, and that Akt is a downstream target of PI3K in this signaling pathway during neurogenesis.