



Title	Hepatoprotective effects of Coptidis rhizoma aqueous extract on carbon tetrachloride-induced acute liver hepatotoxicity in rats
Author(s)	Ye, X; Feng, Y; Tong, Y; Ng, KM; Tsao, S; Lau, GKK; Sze, C; Zhang, Y; Tang, J; Shen, J; Kobayashi, S
Citation	Journal Of Ethnopharmacology, 2009, v. 124 n. 1, p. 130-136
Issued Date	2009
URL	http://hdl.handle.net/10722/58180
Rights	Creative Commons: Attribution 3.0 Hong Kong License

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

Hepatoprotective effects of Coptidis Rhizoma aqueous extract on carbon tetrachloride-induced acute liver hepatotoxicity in rats

Xinshen Ye^a, Yibin Feng^{a*}, Yao Tong^a, Kwan-Ming Ng^b, SaiWah Tsao^c, George KK Lau^d,
Chowing Sze^a, Yanbo Zhang^a, Jun Tang^a, Jiangang Shen^a, Seiichi Kobayashi^e

*a School of Chinese Medicine, The University of Hong Kong, 10 Sassoon Road, Pokfulam,
Hong Kong.*

*b Department of Chemistry and Open Laboratory of Chemical Biology of the Institute of
Molecular Technology for Drug Discovery and Synthesis, Faculty of Science, The University
of Hong Kong, Pokfulam Road, Hong Kong.*

*c Department of Anatomy, Faculty of Medicine, The University of Hong Kong, 21 Sassoon
Road, Pokfulam, Hong Kong.*

*d Department of Medicine, The University of Hong Kong, 21 Sassoon Road, Pokfulam, Hong
Kong.*

e Faculty of Healthy Science, Hokkaido University, Kita 15, Nishi 7 Kita-ku, Sapporo, Japan.

* Corresponding author

Tel: (852) 2589 0482

Fax: (852) 2872 5476

E-mail address: yfeng@hku.hk

Mail address: *School of Chinese Medicine, The University of Hong Kong, 10 Sassoon Road,
Pokfulam, Hong Kong.*

Abstract

1
2
3
4
5 *Ethnopharmacological relevance:* Coptidis Rhizoma (CR, Chinese name is *Huanglian*) has
6
7 been used in treating infectious and inflammatory diseases for two thousand years in
8
9 Traditional Chinese Medicine (TCM). Its related pharmacological basis for the therapeutics
10
11 has been studied intensively, but CR can be also used for vomiting of “dampness-heat type or
12
13 acid regurgitation” due to “liver-fire attacking stomach” in TCM, which symptoms seem to
14
15 link the hepatic and biliary disorders, yet details in the therapies of liver diseases and
16
17 underlying mechanism(s) remain unclear. *Aim of the Study:* in the present study,
18
19 hepatoprotective effect of Coptidis Rhizoma aqueous extract (CRAE) and its possible
20
21 mechanism were studied in rats intoxicated with carbon tetrachloride (CCl₄). *Materials and*
22
23 *Methods:* Sprague-Dawley (SD) rats aged 7 weeks old were intraperitoneally injected with
24
25 CCl₄ at a dose of 1.0 ml/kg as a 50% olive oil solution. The rats were orally given the CRAE
26
27 at doses of 400, 600, 800 mg/kg and 120 mg/kg berberine body weight (BW) after 6 hours of
28
29 CCl₄ treatment. At 24 hours after CCl₄ injection, samples of blood and liver were collected
30
31 and then biochemical parameters and histological studies were carried out. *Results:* the
32
33 results showed that CRAE and berberine inhibited significantly the activities of alanine
34
35 aminotransferase (ALT) and aspartate aminotransferase (AST) and increased the activity of
36
37 superoxide dismutase (SOD). Observation on the hepatoprotective effect of berberine was
38
39 consistent to that of CRAE. *Conclusion:* the study is the first time to demonstrate that CRAE
40
41 has hepatoprotective effect on acute liver injuries induced by CCl₄, and the results suggest
42
43 that the effect of CRAE against CCl₄-induced liver damage is related to antioxidant property.
44
45
46
47
48
49
50
51
52
53

54
55
56 *Keywords:* Coptidis Rhizoma aqueous extract; Carbon tetrachloride; Aspartate
57
58 aminotransferase; Alanine aminotransferase; Superoxide dismutase; Liver histopathology.
59
60
61
62
63

1
2 *Abbreviations:* CR, Coptidis Rhizoma; CRAE, Coptidis Rhizoma aqueous extract; CCl₄,
3
4 Carbon tetrachloride; AST, Aspartate aminotransferase; ALT, Alanine aminotransferase;
5
6 SOD, Superoxide dismutase; SD rats, Sprague-Dawley rats; H & E staining, Hematoxylin
7
8 and eosin staining.
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Article Outline

1. Introduction

2. Materials and methods

2.1. Plant materials and preparation of extract

2.2. Drugs and chemical test agents

2.3. Test animals

2.4. CCl₄-induced acute liver damage model in rats and CRAE treatment

2.5. Serum ALT, AST analysis

2.6. Measurement of SOD in serum and liver tissues

2.7. Histopathological analyses

2.8. Statistical analysis

3. Results

3.1. Quality control of CRAE

3.2. Liver protective effects of CRAE and berberine on acute liver damage in rats

3.3. Effect of CRAE and berberine on the levels of serum SOD and liver homogenate

SOD activities in acute liver damage of rats

3.4. Effect of CRAE and berberine on histopathological changes of liver in acute liver damage of rats

4. Discussion

Acknowledgements

References

Tables 1-3

Figures 1 - 2

Figure captions

1. Introduction

Coptidis Rhizoma (CR, Chinese name is *Huanglian*) is a Chinese herbal medicine used as a clearing heat and detoxifying agent, and has been used to treat syndromes incurred by *damp-heat, fire or toxicity* in Traditional Chinese Medicine (TCM) for two thousand years, which can be conceived as inflammatory diseases. Extensive studies exhibited that CR has many pharmacological actions with strong clinical implications, including antibacterial, antiviral, antiinflammatory, antineoplastic, antihypertensive, antioxidative, antihyperglycemic and cholesterol-lowering effects (Chang and But, 2004; Choi, et al., 2007; Kim, et al., 2008; Fukuda, et al., 1998; Li, et al., 2000; Sanae, et al., 2001; Yokozawa, et al., 2003, 2004). Traditionally, CR can be used for vomiting with “dampness-heat type” or “acid regurgitation due to liver-fire attacking stomach” in TCM, whose symptoms seem to be linked with the hepatic and biliary disorders, yet its therapeutic potential remains unexplored. In TCM clinical practice, CR is a key component in many TCM formulae. Typically, *Huanglian Jiedu* decoction (or Oren- gedoku-to in Japanese or JT-15, including Coptidis Rhizoma, Radix Scutellariae, Cortex Phellodendri, and Frucuts Gardeniae) has been used for the therapies of hepatitis and liver dysfunction in addition to gastric ulcers, dermatitis, dementia, and cerebrovascular diseases in Japan (Itoh, 2001) and has intensively studied for scientific basis of hepatitis and liver dysfunction in Japan (Ohta, et al., 1997, 1998, 2004), but whether CR has liver protection or not is unknown. Berberine, due to the phytochemical analysis, was the main ingredient in CR (Xu et al., 2004). It was reported that herbs with high content of berberine exhibit preventive effect or curative effect on liver damage (Nadkarni, 1976, Gilani and Janbaz, 1995; Janbaz and Gilani, 2000), indicating that CR is potential for the treatment of liver injury and hepatitis. According to the history of CR use, clinical indications of

1 CR-containing composite formulae, and clinical indications of berberine-containing plant
2 species in other traditional medicine, we have used CR to prescribe for various liver diseases
3 in TCM clinical practice in China (Feng et al., 2008). It is interesting to explore the exact
4 effect of CR on liver damage and underlying mechanisms.
5
6
7
8

9 It was reported that the changes associated with CCl₄-induced liver damage are similar to
10 that of acute viral hepatitis (Rubinstein, 1962), drug/chemicals-induced hepatopathy and
11 oxidative stress (Recknagel et al., 1989; Kadiiska et al., 2000), so CCl₄-induced
12 hepatotoxicity model is frequently used for the investigation of hepatoprotective effects of
13 drugs and plant extracts. CRAE has shown to ameliorate renal oxidative injury in vivo and in
14 vitro (Yokozawa et al 1999, 2004, 2005). Previous study revealed that preventive effect of
15 huanglian-jie-du-tang extract on progression of CCl₄-induced acute liver injury in rats is
16 related to its antioxidant properties (Ohta et al, 1997, 1998, 2001). Whether CRAE
17 contributes to its antioxidant effects on CCl₄-induced acute liver injury in rats is not known
18 yet.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33

34 In the present study, CCl₄ was therefore introduced to induce liver damage on
35 experimental animal model, and the curative effect of CRAE was examined via determining
36 serum Alanine aminotransferase (ALT), aspartate aminotransferase (AST), serum and tissue
37 superoxide dismutase (SOD) level as well as histopathological study. Phytochemistry of
38 CRAE was also analyzed via High performance liquid chromatogram (HPLC) in this study.
39 The experimental results demonstrated the potential effects of CRAE in protecting liver
40 function, reducing oxidative stress, and improving histopathological structures in the rat
41 model of CCl₄-induced liver damage. The study not only provides helpful information for
42 the application of CRAE in liver disease, but also promotes the understanding of the
43 pharmacological mechanisms of CRAE in the acute toxic liver injury.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

2. Materials and methods

2.1. Plant materials and extraction procedures

Crude herb, *Coptidis Rhizoma* (*Coptis chinensis* Franch.) was collected from Sichuan province, China and authenticated under the guidance of The Pharmacopeia of China (2005). Plant materials were dried under shade and cut into small pieces before extraction. For the preparation of CRAE, one gram of crude CR was boiled in 10ml of distilled water at 100°C for 1 hour. Total was 500g of crude CR. The solution was percolated through filter paper (Whatman, pleated filter grade 597 1/2, 4–7 µm) and then sterilized by filtration through a 0.2 µm pore filter (Minisart[®]-plus, Sartorius), while the residue was further extracted under the same condition once. The filtrates collected from the extraction were combined and evaporated to dryness by vacuum at temperature. The dry extract powder obtained (40g) was stored in –20 °C freezer and used in following experiments.

The chemical profile of CRAE was recorded by high performance liquid chromatography (HPLC) with photodiode array (PDA) detection. CRAE powder (26.6 mg) was accurately weighed and dissolved in 9 ml acidified methanol (a mixture of fuming hydrochloric acid and methanol in 1:100 proportion). The solution was heated in a 60 °C water bath for 15 min, followed by ultrasonication for 30 min. The solution was then filtered by using a 0.45 µm Millex Syringe filter unit and subjected to HPLC analysis. The HPLC analytical system is composed of a Waters 600s solvent delivery system coupled with a 717 plus autosampler (with injection volume at 10 µL) and a 996 photodiode array detector. A reverse-phase C₁₈ column (Alltech Alltima HP C18, 250 mm × 4.6 mm, 5 µm) eluting with a mobile phase (acetonitrile:25 mM potassium dihydrogen phosphate in H₂O (25: 75)) in an isocratic manner and at a flow rate of 1 ml/min was employed. The eluate was monitored at the wavelength of

1 345 nm. The column temperature was kept at 24 °C. Berberine is used as the reference
2 standard for identifying and quantifying the major component in CRAE. The whole analysis
3 was duplicated for confirmation.
4
5

6
7 Atomic absorption spectroscopic (AAS) analysis of five toxic heavy metals, including
8 arsenic, cadmium, chromium, lead and mercury, was performed on PerkinElmer Analyst 800
9 atomic absorption spectrometer with autosampler.
10
11
12
13

14 15 16 17 *2.2. Drugs and chemical test agents*

18
19
20
21
22 Alanine aminotransferase (ALT) and aspartate aminotransferase (AST) test kits were
23 purchased from Stanbio, USA. SOD assay kit was from Biovision, USA. Carbon
24 tetrachloride (CCl₄), olive oil and berberine, were purchased from Sigma, USA. Standards of
25 arsenic, cadmium, chromium, lead and mercury were purchased from the Sigma, USA.
26
27
28
29
30
31

32 33 34 *2.3. Test Animals*

35
36 Adult male SD rats 7 weeks weighing 250 ± 20 g were obtained from animal centre of
37 The University of Hong Kong. The animals left for 2 days for acclimatization to animal room
38 conditions were maintained on standard pellet diet and water *ad libitum* at a temperature of
39 20–25 °C under a 12 h light/dark cycle throughout the experiment. The food was
40 withdrawn on the day before the experiment, but free access of water was allowed. The rats
41 were randomly assigned. A minimum of 8 animals were used in each group. All animals
42 received human care and study protocols complied with the guidelines of the animal centre of
43 the University of Hong Kong. Throughout the experiments, animals were processed
44 according to the suggested international ethical guidelines for the care of laboratory animals.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

2.4. CCl₄-induced acute liver damage model in rats and CRAE treatment

The animals were randomly divided into five groups, that is, normal, CCl₄ alone, and 400, 600, 800 mg/kg BW CRAE and 120 mg/kg berberine BW treatment on CCl₄ groups, containing eight rats in each. Rats were intraperitoneally (*i.p.*) injected with CCl₄ at a dose of 1.0 ml/kg as a 50 % olive oil solution according to the previously reported methods with some modification (Moghaddam, et al. 1998; Feng, et al. 2000) and control ones with the same dose of olive oil. Our previous clinical practice showed the therapeutic effect of CRAE on liver injury, infectious hepatitis and even cancer (Feng, et al., 2008), the dosage of raw herb we use in clinic was about 30~50 gram in single use or 9-18 gram in formulation. Excellent effect and no toxicity were observed in our clinical practice. The result and dosage in present study fit our clinical study, that is, if we use conversion table to compute the clinical dosage from the animal dosage, 400~800 mg/kg, it will be 25~50 gram, near our clinical study. On the other hand, lower than 400mg/kg may have no effect in our dose screening in rats. Hence, the CRAE was dissolved in distilled water at a concentration of 400, 600 or 800 mg/kg BW orally administered to rats injected with or without CCl₄ treatment at 6 hours after CCl₄ exposure. The control rats were orally given the same volume of distilled water. These animals were fasted with free access to water throughout the experiment.

Twenty-four hours after the CCl₄ administration, blood samples were withdrawn by cardiac puncture when the animals had been anaesthetized with ketamine /xylazine mixture (ketamine 67mg/kg, xylazine 6mg/kg, *i.p.*). The animals were sacrificed by an overdose of pentobarbitone (Phenobarbital 200mg/kg, *i.p.*) or diethyl ether immediately after blood collection. Blood samples collected in heparinized tubes were centrifuged at 3000 × g for 10 min to obtain serum. Serum samples were used to determine SOD as well as to test ALT and

1 AST activities. On the other hand, the liver of each rat was promptly removed and used to
2 determine the tissue level SOD and for further histopathological study.
3
4
5
6

7 *2.5. Serum ALT and AST analysis*

8
9
10

11 Biocon standard kits and an auto-analyzer (Hitachi 736-60, Tokyo, Japan) (UV-Rate)
12 were used to measure serum ALT and AST activities in serum samples according to the
13 method published before (Wilkinson et al., 1972). Values are derived based on the
14 “absorptivity micromolar extinction coefficient” of NADH at 340 nm. ALT and AST
15 activities expressed in terms of units per liter (U/L) are the amount of enzyme oxidizing one
16 $\mu\text{mol/L}$ of NADH per minute.
17
18
19
20
21
22
23
24
25
26
27

28 *2.6. SOD levels in serum and liver tissues*

29
30
31
32
33

34 The sensitive SOD assay kit (SOD Assay Kit-WST) utilizes mitochondrial activity that
35 produces a water-soluble formazan dye upon reduction with superoxide anion. The rate of the
36 reduction with a superoxide anion is linearly related to the xanthine oxidase (XO) activity,
37 and is inhibited by SOD. Therefore, the inhibition activity of SOD determined by a
38 colorimetric method was used for the determination of the serum and liver SOD level in this
39 study.
40
41
42
43
44
45
46
47

48 Serum and liver homogenate collected from test animals were for measurement of SOD
49 in blood and liver tissues. Procedure was carried out by according to SOD Assay Kit-WST
50 Technical Manual (Dojindo Laboratories, Kumamoto, Japan).
51
52
53
54
55
56
57

58 *2.7. Histopathological studies*

59
60
61
62
63
64
65

1
2 For the histopathological study, the livers of eight animals in each group were
3 immediately removed after autopsy and the tissues were fixed in 10% buffered formaldehyde
4 solution for a period of at least 24 h. The paraffin sections were then prepared (Automatic
5 Tissue Processor, Lipshaw) and cut into 5 µm thick sections by a Leica RM 2016 rotary
6 microtome (Leica Instruments Ltd., Shanghai, China). The sections were stained with
7 hematoxylin and eosin staining (H & E staining) and then mounted with Canada balsam
8 (Sigma, USA). The degree of liver damage was examined under the microscope (Leica
9 Microsystems Digital Imaging, Germany). The images were taken using Leica DFC 280
10 CCD camera at original magnification of 10×10. Through grading the liver sections
11 numerically to assess their histological features, acute liver injury was evaluated by three
12 independent researchers. Vacuolation, nuclei, hepatocyte necrosis, inflammatory cell
13 infiltration and central vein and portal triad were used as criteria, and a combined score of
14 histological features was given for each liver section. The parameters were graded from score
15 0 to 6, with 0 indicating no abnormality, 1–2 indicating mild injury, 3–4 indicating moderate
16 injury and 5–6 with severe liver injury (Wills and Ahsa, 2006; Wang et.al., 2008).
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41 *2.8. Statistical analysis*

42
43
44
45 The data obtained were analyzed by one-way of variance (ANOVA) and
46 Student–Newman–Kelus post hoc tests for the significant interrelation between the groups.
47
48 Data were expressed as mean ± standard error of the mean and were analyzed with SPSS,
49
50 version 11.5 software. Differences between group means were calculated by a one-way
51
52 analysis of variance. Values of $P < 0.05$ were considered to be statistically significant.
53
54
55
56
57

58 **3. Results**

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

3.1. Quality control of CRAE

The crude herb of *huanglian* was identified by morphological characteristics of CR (*Coptis chinensis Franch.*) according to The Pharmacopeia of China (2005) and the voucher specimens (no. CR20060616) was deposited in the Herbarium of School of Chinese Medicine, The University of Hong Kong. The plant and crude herb were shown in Figure 1A and B.

The HPLC chemical profile of CRAE is shown in Fig. 1 C. Nine characteristic chromatographic peaks were recorded at 345 nm. The chemical identity of the peak at 14.98 min was confirmed as berberine by chromatographic peak matching of berberine reference standard at 15.02 min (Fig. 1 D) and the similar UV-Visible spectra with λ_{\max} at 345 nm. A calibration curve of berberine reference standard showing a good linearity over the concentration range from 10 to 160 $\mu\text{g/ml}$ with a regression coefficient at $r^2 = 0.9999$ (Fig. 1 E) was obtained. The content of berberine in CRAE powder was determined to be 20.3 mg in 100 mg of the powder (i.e., 20.3 % weight by weight) with deviation less than 1% in duplicated analysis. Among other eight peaks, five (including the peaks 4, 5, 6, 7 and 8) showed the similar UV spectra of berberine with λ_{\max} at 345 nm.

As shown in Table 1, the contents of five heavy metals (Arsenic, Cadmium, Chromium, Lead, and Mercury) in CRAE determined from the AAS analysis fell in the range less than the maximum limit (20 ppm or $\mu\text{g/g}$) as regulated in China Pharmacopeia (2005) and the World Health Organization (WHO). The amount of the five harmful elements in CRAE is in the safe range for herbal test use.

3.2. Liver protective effects of CRAE and berberine on acute liver damage in rats

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Compared with the normal group, the ALT activities in serum of the control treated with CCl₄ at an *i.p.* dose of 1.0 ml/kg was significantly elevated ($P<0.01$) after 24 h. While treatment with 400 mg/kg BW, CRAE decreased remarkably the levels of serum ALT in rats treated with CCl₄ though higher than the normal rats. The orally administered CRAE at a higher dose of 600 mg/kg BW and 800 mg/kg BW could significantly reduce the serum ALT level and restored them to normal levels when compared with rats treated with CCl₄ control ($P<0.01$). This result showed that the oral treatment with 800, 600 and 400 mg/kg BW of CRAE could inhibit the elevated ALT activities in rats intoxicated with CCl₄ in a dose-dependent manner (Table. 2).

Compared with the normal group, serum AST activities were significantly elevated ($P<0.01$) by 6 h after the CCl₄ treatment. Post-administration of 600 and 800 mg/kg BW CRAE significantly decreased the AST activities in serum in contrast to CCl₄-treated rats ($P<0.05$, $P<0.01$), but 400 mg/kg BW CRAE did not show effect on the AST levels in the CCl₄-treated rats (Table.2).

Significantly reduced serum ALT and AST levels in liver damage rats treated with berberine 120 mg/kg were also observed (Table. 2).

3.3. *Effect of CRAE and berberine on the levels of serum SOD and liver homogenate SOD activities in acute liver damage of rats*

The inhibition rate of serum SOD activities in control rats treated with CCl₄ alone was remarkably decreased after 24 h ($P<0.01$) compared with the normal ones, which showed the injured liver functions by CCl₄. While treated with 800, 600, 400 mg/kg BW CRAE and 120 mg/kg BW berberine, the inhibition rates of SOD were significantly elevated ($P<0.01$)

1 compared with the control rats, especially the high dose at 800 mg/kg BW could restore the
2 value to the normal level (Table 2). Liver homogenate SOD activities were similar to serum
3 expression (Table 2).
4
5
6
7
8
9

10 *3.4. Effect of CRAE and berberine on histopathological changes of liver in acute liver*
11 *damage of rats*
12
13
14
15

16 The histological changes associated with the hepatoprotective activity in three dosages of
17 CRAE and berberine basically supported the measuring of the serum enzyme activities. There
18 was no abnormal appearance or histological changes in the liver of normal control rats, which
19 received olive oil only (Fig. 2 A). CCl₄ administration caused classical damage in the rat liver
20 at 24 h, as demonstrated by severe hepatocyte necrosis, inflammatory cells infiltration, fatty
21 degeneration, hemorrhage, and hydropic degeneration (Fig. 2 B), vacuole generation and
22 microvascular steatosis were frequently observed. The administration of BW at dose of 800,
23 600 and 400 mg/kg could largely rescue the severity of CCl₄-induced liver intoxication, in
24 which the high dose at 800 mg/kg was most effective (Fig. 2 C-E). The histological patterns
25 showed dose dependant improvements for fatty change, necrosis and lymphocyte infiltration
26 in contrast to treatment with CCl₄ showing an obvious formation of necrosis. Improvement
27 results were observed in liver section from animals treated with 120 mg/kg berberine (Fig 2
28 F). The scoring of histological damage was displayed in Table 2.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52

53 **4. Discussion and conclusions**
54
55

56 Three clues led us to study CR against liver diseases: traditional use and our TCM daily
57 practices for liver diseases (Feng et al., 2008), liver protective effects of
58
59
60
61
62
63
64
65

1 Hunaglian-containing composite formula (Huanglian je du decoction) (Ohta et al., 1997,
2 1998, 2004), and berberis aristata (major compound is berberine which is the same as CR) is
3
4 used for liver disease in South Asian areas, only showed preventive, but not curative effects
5
6 on liver damage in rats treated with CCl₄ (Gilani and Janbaz, 1995; Janbaz and Gilani, 2000).
7
8 We assume that CR is a main effective herb in Huanglian je du decoction for liver protection
9
10 and CR should possess curative effects on liver damage. The aim of the present study was to
11
12 investigate the potential hepatoprotective effects of Coptidis Rhizoma aqueous extract
13
14 (CRAE) on the free radical damage of liver caused by CCl₄ in rats.
15
16
17
18
19
20

21 Firstly, quality control was conducted for the raw herb of CR and CRAE to guarantee the
22
23 reliability of our experimental results. The plant materials of Coptidis rhizoma, Chinese
24
25 name is “*Haunglian*”, have three species. Rhizoma of Coptis Chinensis Franch, one of the
26
27 species, was chosen for our study due to its plentiful cultivation in China and most popular in
28
29 TCM practice. Miao et al. (1997) have shown that when *Coptidis rhizoma*, *Scutellariae radix*,
30
31 and *Phellodendri cortex* are extracted with 50% methanol, the *Scutellariae radix* extract has
32
33 much higher O₂⁻-scavenging activity than the *Phellodendri cortex* extract, while the *Coptidis*
34
35 *rhizoma* extract has little O₂⁻-scavenging activity. Yokozawa et al. (1997) have reported that
36
37 although the boiled water extract of *Coptidis rhizoma*, *Gardeniae fructus*, *Scutellariae radix*
38
39 or *Phellodendri cortex* inhibits lipid peroxidation induced by H₂O₂ in rat liver homogenates,
40
41 the *Coptidis rhizoma* extract has the highest inhibitory activity, followed in the order of
42
43 strength by the *Gardeniae fructus* extract > the *Scutellariae radix* extract > the *Phellodendri*
44
45 *cortex* extract. We assume that water extract of CR should possess better bioactivities
46
47 according to the above two reports. In addition, the water extract of CR is clinically applied
48
49 form, and thus being used in our study. The yield amount of CRAE obtained from raw herb
50
51 was similar to previous reports (Li, et al., 2000; Yokozawa, et al., 2004). The consistency of
52
53 chemical composition in the CRAE is important in safeguarding the reliability of the research
54
55
56
57
58
59
60
61
62
63
64
65

1 results. The chemical profile of CRAE was recorded by the RP-HPLC/PDA analysis. The
2 HPLC chemical profile could be delineated by the measurement of relative retention times of
3 major characteristic peaks using berberine as a marker. The resulting chromatogram was used
4 as a standard for assessment of all extracts used in the current study. The HPLC chemical
5 profile of the CRAE was similar to the previous report (Li, et al., 2000). As toxic heavy
6 metals may induce toxicity including liver damage (Duffus, 2002; Wang, et al., 2007) and
7 traditional Chinese medicines have drawn attention by its heavy metals (Cooper K, et al.,
8 2007), five commonly found toxic metals, including arsenic, cadmium, chromium, lead and
9 mercury were analyzed and their contents in CRAE were in safe range set by the
10 Pharmacopeia of China (2005) and WHO.
11
12
13
14
15
16
17
18
19
20
21
22
23
24

25 Many drugs or chemical substances are known to cause hepatic injuries, such as
26 acetaminophen, CCl₄, D-galactosamine (GalN), aflatoxins and dimethylnitrosamine (DMN),
27 among which, liver injury induced by CCl₄ is the best-characterized system of the
28 xenobiotic-induced hepatotoxicity and a commonly used model for screening the drugs with
29 anti-hepatotoxicity and/or hepatoprotective activity (Brattin et al., 1985). CCl₄-induced liver
30 damage, a free radical damage model, results from oxidative stress that could directly injure
31 hepatocellular membrane by lipid peroxidation, followed by a series of cascades of cellular
32 events such as the massive release of inflammatory mediators or cytokines, which eventually
33 lead to liver injuries (Pessayre, 1995; Dizdaroglu, et al., 2002; Higuchi and Gores, 2003).
34 Superoxide dismutase (SOD) is one of the most important antioxidative enzymes, whose
35 activities decrease after CCl₄ injection. Therefore we selected SOD as parameter for the
36 antioxidative effects of CR. The present study showed that CCl₄ administration caused severe
37 acute liver damage in rats, which was demonstrated by significant elevation of serum AST,
38 ALT levels, decreased SOD activities (Table 2), and classic histopathological changes
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 (Fig.2), indicating that CCl₄-induced liver damage in animal model can be used to evaluate
2 the curative effect of CRAE.
3
4

5
6 All data in our study consistently demonstrated that CRAE treatment at a dose of 800,
7
8 600 and 400 mg/kg BW had a potent protective effect against oxidative stress and acute liver
9
10 damage induced by CCl₄ in rats, as revealed by remarkable elevation of SOD activities in the
11
12 liver and serum (Table 2). Additionally, CRAE could ameliorate acute liver damage to a high
13
14 degree, as demonstrated by reduction of serum ALT and AST levels and the improvement of
15
16 the histopathological changes (Table 2, 3 and Fig.2). Apart from mild hydropic degeneration
17
18 of hepatocytes, the liver had a nearly normal appearance in CCl₄-treated rats simultaneously
19
20 treated with CRAE at a dose of 800 mg/kg BW (Fig.2 and Table 3). The results also indicated
21
22 a remarkable elevation of SOD activities in the liver and serum of rats for CRAE treatment at
23
24 24 h after CCl₄ administration. It was previously reported that CRAE effectively scavenged
25
26 the NO radical in-vitro (Yokozawa et al 2000, 2004), so the action mechanisms underlying
27
28 hepatoprotection of CRAE may be related to both its radical scavenging properties and
29
30 indirect effects as a regulator of antioxidative systems in which more details should be
31
32 figured out in the future.
33
34
35
36
37
38
39
40

41
42 Phytochemical analysis indicated that berberine is also a major compound in CRAE (see
43
44 Fig. 1 C and 1 D). Previous studies have demonstrated that berberine showed
45
46 hepatoprotection possibly through inhibitory action on hepatic drug metabolizing enzymes,
47
48 cytochrome P450s (Gilani and Janbaz, 1995; Janbaz and Gilani 2000), but no evident
49
50 curative effect of berberine against CCl₄-induced acute liver damage is investigated before.
51
52 Our present study shows that 120 mg/kg of berberine can significantly decrease the elevation
53
54 of serum AST and SLT level induced by CCl₄ treatment in animal model, and the recovery of
55
56 SOD activity both in serum and tissue indicates berberine as an antioxidant agent in liver
57
58
59
60
61
62
63
64
65

1 protection. This convinces our observation on CRAE's action on liver damage and further the
2 knowledge that berberine may be the major active component in CRAE when it was used for
3 liver diseases. Berberine 120 mg/kg BW which is the equal of CRAE 600 mg/kg BW
4 (calculated by about 20% of berberine in CRAE according to Fig.1) displays good liver
5 protective effect (effect close to CRAE 800 mg/kg BW). Detailed dose-effect relationship
6 and bioavailability of berberine on hepatoprotective effect need to be further studied.
7
8
9
10
11
12
13
14
15

16 In conclusion, the study is the first time to demonstrate that CRAE has an impressive
17 hepatoprotective effect on acute liver injuries induced by CCl₄, which might be considered to
18 be therapeutic effect in clinical situations. Berberine may be the major active component in
19 CRAE for hepatoprotective effect. As a possible mechanism, CRAE could alleviate liver
20 injury through antioxidative effects. On the other hand, the model of CCl₄-induced hepatic
21 injury in the rat is similar to many features of acute hepatitis induced by toxicants and virus,
22 hence our results could partially explicate therapeutic principle for CRAE in TCM clinical
23 application, suggesting that CRAE could be used as a potential new drug for acute liver
24 injury.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

40 **Acknowledgements**

41
42 The study was financially supported by grants from the research council of the
43 University of Hong Kong (Project Codes: 10206540, 10208005), Medical Faculty Research
44 Grant, the University of Hong Kong (Project Code: 21362502), Pong Ding Yueng
45 Endowment Fund for Education & Research (Project Code: 20005274), The University Grant
46 Committee (UGC) of Hong Kong (Project Code: 764708M) and The University Grant
47 Committee of Hong Kong (Area of Excellence Scheme, AoE/P-10/01). The authors are
48 grateful to the support of Professors Yung-Chi Cheng, Chi-Ming Che and Allan SY Lau. The
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 authors would also like to express special thanks to Mr. Keith Wong and Mr. Freddy Tsang
2 for their technical support.
3
4
5

6 7 **References** 8

9 Brattin, W.J., Glende, E.A., Recknagel, R.O., 1985. Pathological mechanisms in carbon
10 tetrachloride hepatotoxicity. *Free Radical Biology and Medicine* 1:27–28.
11

12 Chang, H.M., Paul, P.H But, 2001. *Pharmacology and applications of Chinese Materia*
13 *Medica*. World Scientific Publishing, Hong Kong, 1061-1077.
14

15 Choi, U.K., Kim, M.H., Lee, N.H., 2007. Optimization of antibacterial activity by
16 Gold-Thread (*Coptidis Rhizoma* Franch) against *Streptococcus mutans* using evolutionary
17 operation-factorial design technique. *Journal of microbiology and biotechnology*,
18 17:1880-4.
19

20 Dizdaroglu, M., Jaruga, P., Birincioglu, M., Rodrriguez, H., 2002. Free radical-induced
21 damage to DNA: mechanisms and measurement. *Free Radical Biology and Medicine*
22 32:1102–1115.
23

24 Feng, Y., Nagamatsu, T., Suzuki, Y., Kawada, T., Feng, Y.G., Kobayashi, S., Koike, T.,
25 2000. Pharmacological studies on the diuretic action of Chinese-Japanese formulations: the
26 application and evaluation of its pharmacological screening. *Journal of Traditional*
27 *Medicine* 17, 122-130.
28

29 Feng, Y., Luo, W.Q., Zhu, S.Q., 2008. Explore new clinical application of Huanglian and
30 corresponding compound prescriptions from their traditional use. *China Journal of Chinese*
31 *Materia Medica*, 33, 1221-1225.
32

33 Fukutake, M., Yokota, S., Kawamura, H., Iizuka, A., Amagaya, S., Fukuda, K., Komatsu,
34 Y.,1998. Inhibitory effect of *Coptidis Rhizoma* and *Scutellariae Radix* on azoxymethane-
35
36
37
38
39
40
41
42
43
44
45

1 induced aberrant crypt foci formation in rat colon. *Biological & pharmaceutical bulletin*.
2 21:814-817.
3

4
5 Gilani, A.H., Janbaz, K.H., 1995. Preventive and curative effects of *Berberis aristata* Fruit
6 extract on paracetamol- and CCl₄-induced hepatotoxicity. *Phytotherapy Research* 9:489
7 -494.
8
9

10
11
12 Higuchi, H., Gores, G.J., 2003. Mechanisms of liver injury: an overview. *Current Molecular*
13 *Medicine* 3:483–490.
14
15

16
17 Janbaz, K.H., Gilani, A.H., 2000. Studies on preventive and curative effects of berberine on
18
19 Chemical-induced hepatotoxicity in rodents. *Fitoterapia* 71, 25-33.
20
21

22 Kadiiska, M.B., Gladen, B.C., Baird, D.D., Dikalova, A.E., Sohal, R.S., Hatch, G.E., Jones,
23
24 D.P., Mason, R.P., Barrett, J.C., 2000. Biomarkers of oxidative stress study: are plasma
25
26 antioxidants markers of CCl₄ poisoning? *Free Radical Biology and Medicine* 28:838-845.
27
28

29 Kim, H.Y., Shin, H.S., Park, H., Kim, Y.C., Yun, Y.G., Park, S., Shin, H.J., Kim, K., 2008.
30
31 In vitro inhibition of coronavirus replications by the traditionally used medicinal herbal
32
33 extracts, *Cimicifuga rhizoma*, *Meliae cortex*, *Coptidis rhizoma*, and *Phellodendron cortex*.
34
35
36 *Journal of clinical virology*, 41:122-128.
37
38

39 Li, X.K., Motwani, M., Tong, W., Bornmann, W., Schwartz, G.K., 2000. Huanglian, A
40
41 Chinese herbal extract, inhibits cell growth by suppressing the expression of cyclin B1 and
42
43 inhibiting CDC2 kinase activity in human cancer cells. *Molecular Pharmacology*
44
45 58:1287-1293.
46
47

48 Miao, Z., Kayahara, H. and Tadasa, K., 1997. Superoxide-scavenging and
49
50 tyrosinase-inhibitory activities of the extracts of some Chinese medicines. *Bioscience,*
51
52
53 *Biotechnology, and Biochemistry* 61, 2106–2108.
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Moghaddam, A.P., Eggers J.S., Calabrese E.J., 1998. Evaluation of sex difference in tissue repair following acute carbon tetrachloride toxicity in male and female Sprague-Dawley rats. *Toxicology* 130, 95-105.
- Nadkarni, A.K., 1976. In: *Indian materia medica*, Vol. I Popular Prakashan Private Ltd, Bombay, 187–189.
- Ohta, Y., Kongo-Nishimura, M., Hayashi, T., Kishikawa, T., 2004. Effect of Oren-gedoku-to (*Huanglian-Jie-Du-Tang*) extract on disruption of hepatic antioxidant defense systems in rats treated with D-galactosamine. *Journal of Ethnopharmacology* 94, 323-329.
- Ohta, Y., Sasaki E., Nishida K., Kongo M., Hayashi T., Nagata M., Ishiguro I., 1998. Inhibitory effect of Oren-gedoku-tu (*Huanglian-Jie-Du-Tang*) extract on hepatic triglyceride accumulation with the progression of carbon tetrachloride-induced acute liver injury in rats. *Journal of Ethnopharmacology* 6, 75-80.
- Ohta, Y., Sasaki, E., Nishida, K., Hayashi, T., Nagata, M., Ishiguro, I., 1997. Preventive effect of oren-gedoku-to (*huanglian-jie-du-tang*) extract on progression of carbon tetrachloride- induced acute liver injury in rats. *American Journal of Chinese Medicine*, 25, 57-68.
- Pessayre, D., 1995. Role of reactive metabolites in drug-induced hepatitis, *Journal of Hepatology*, 23:16–24.
- P.J. Wills, V.V. Asha, 2006. Protective effect of *Lygodium flexuosum* (L.) Sw. extract against carbon tetrachloride-induced acute liver injury in rats. *Journal of Ethnopharmacology*, 108: 320–326
- Rubinstein, D., 1962. Epinephrine release and liver glycogen levels after carbon tetrachloride administration. *American Journal of Physiology* 203:1033-1037.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Recknagel, R.O., Glende, E.A.Jr., Dolak, J.A., Waller, R.L., 1989. Mechanisms of carbon tetrachloride toxicity. *Pharmacology & therapeutics* 43:139-154.
- Sanae, F., Komatsu, Y., Chisaki, K., Kido, T., Ishige, A., Hayashi, H. 2001. Effects of San'o-shashin-to and the constituent herbal medicines on theophylline-induced increase in arterial blood pressure of rats. *Biological & pharmaceutical bulletin*. 24:1137-1141.
- Tao Wang, Ning-Ling Sun, Wei-Dong Zhang, Hui-Liang Li, Guo-Cai Lua, Bo-Jun Yuan, Hua Jiang, Jia-Hong She, Chuan Zhang, 2008, Protective effects of dehydrocavidine on carbon tetrachloride-induced acute hepatotoxicity in rats. *Journal of Ethnopharmacology*. 117:300–308;
- Xu, J.T., Wang, L.Q., Xu, B., 2004. Research development of *Coptis chinensis*. *Acta Academia Medicines Sonica* 26:704-707.
- Yasukawa, K., Takido, M., Ikekawa, T., Shimada, F., Takeuchi, M., Nakagawa, S., 1991. Relative inhibitory activity of berberine-type alkaloids against 2-O-tetradecanoylphorbol-13-acetate-induced inflammation in mice. *Chemical & pharmaceutical bulletin (Tokyo)*. 39:1462-1465.
- Yokozawa, T., Dong, E., Liu, Z.W. and Oura, H., 1997. Antiperoxidation activity of traditional Chinese prescriptions and their main crude drugs in vitro. *Natural Medicines* 51, 92–97.
- Yokozawa, T., Satoh, A., Cho, E.J., Kashiwada, Y., Ikeshiro, Y., 2005. Protective role of *Coptidis Rhizoma* alkaloids against peroxynitrite-induced damage to renal tubular epithelial cells. *Journal of Pharmacy and Pharmacology*. 57:367-74.
- Yokozawa, T., Ishida, A., Kashiwada, Y., Cho, E.J., Kim, H.Y., Ikeshiro, Y., 2004. *Coptidis Rhizoma*: protective effects against peroxynitrite-induced oxidative damage and elucidation of its active components. *Journal of Pharmacy and Pharmacology*. 56:547-56.

1 Yokozawa, T., Chen, C.P., Tanaka, T., 2000. Direct scavenging of nitric oxide by traditional
2 crude drugs. *Phytomedicine*. 6: 453-463.
3

4 Yokozawa T, Ishida A, Cho EJ, Nakagawa T. 2003. The effects of *Coptidis Rhizoma* extract
5 on a hypercholesterolemic animal model. *Phytomedicine*. 10:17-22.
6
7
8
9

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Ye, et al. Table 1

Table 1. Content (ppm or $\mu\text{g/g}$) of five selected toxic metals in CRAE

Heavy metals	Mean \pm RSD (%)	Calibration correlation cofactor (%)
Arsenic	-0.26 \pm 32.54%	99.95
Cadimum	0.48 \pm 18.90%	99.91
Chromium	0.01 \pm 8.67%	99.82
Lead	-0.04 \pm 8.84%	99.90
Mercury	0.14 \pm 39.69%	99.40

Table 2 Effect of CRAE and berberine on CCl₄-induced liver damage in rat (Mean±S.D., N=8)

Group	ALT(U, in serum)	AST(U, in serum)	SOD (% in serum)	SOD (% in tissue)
Normal	19.877±7.34	61.759±30.62	77.26±0.11	72.42±5.89
Control	133.27±32.11 ^{##}	342.11±55.27 ^{##}	10.80±0.21 ^{##}	9.32±5.31 ^{##}
CRAE 400 mg/kg	66.26±11.24 ^{**##}	366.22±52.16 ^{##}	26.70±0.23 ^{*#}	21.61±4.97 ^{*#}
CRAE 600 mg/kg	41.17±10.66 ^{**}	155.39±30.24 ^{*#}	40.30±0.15 ^{*#}	46.22±4.46 ^{*#}
CRAE 800 mg/kg	23.29±12.76 ^{**}	20.78±10.77 ^{**}	73.70±0.20 ^{**}	70.63±6.54 ^{**}
Berberine 120 mg/kg	26.06±7.48 ^{**}	23.48±5.91 ^{**}	75.21±1.79 ^{**}	68.42±8.13 ^{**}

#<p<0.05 compared with normal group

##p<0.01 compared with normal group

*p<0.05 compared with CCl₄ control group

**p<0.01 compared with CCl₄ control group

Table 3 Microscopic observation on CRAE and berberine against CCl₄-induced acute liver damage (Mean ± S.D., n=8)

Group	Vacuolation	nuclei	hepatocyte necrosis	inflammatory cell infiltration	central vein and portal triad	combined score
Normal	0.4±0.3	1.0±0.7	0.5±0.2	0.6±0.3	1.1±0.6	0.5±0.3
Control	5.0±0.6##	0.7±0.2	4.2±0.8##	3.9±1.6##	0.5±0.3#	4.9±0.5##
CRAE 400 mg/kg	4.2±1.1*	0.8±0.6	2.2±1.7*	2.8±1.3*	1.3±0.8*	3.6±1.4*
CRAE 600 mg/kg	3.0±1.0**	1.1±0.3	2.1±1.5**	2.1±1.3**	1.1±0.6*	2.7±0.9**
CRAE 800 mg/kg	1.9±0.7**	1.2±0.6	1.5±1.2**	1.3±0.9**	0.9±0.5	1.7±0.6**
Berberine 120 mg/kg	1.7±1.3**	1.8±0.2	1.4±0.4**	1.2±0.5**	1.1±0.4*	1.6±0.8**

#<p<0.05 compared with normal group

##p<0.01 compared with normal group

*p<0.05 compared with CCl₄ control group

**p<0.01 compared with CCl₄ control group

Ye, et al. Fig.1

Figure 1.



A: Plant of *Coptis Chinensis* Franch.



B: Raw herb of *Coptis Chinensis* Franch.






Figure 2.




Figure captions

Figure 1. Huanglian identification and HPLC chemical profile of CRAE. A. The species of CR plant is *Coptis Chinensis* Franch. B. Raw herb of CR. C. The HPLC chromatogram of CRAE D. The HPLC chromatogram of berberine reference standard. E. Standard calibration curve of berberine.

Figure 2. The photomicrography of liver sections from rats treated with CCl₄, the post-doses of CRAE at 400, 600 and 800 mg/kg BW, and olive oil vehicle. A. liver section of normal rat; B. liver section of the control rat treated with CCl₄; C. liver section of the CCl₄-treated rat post-dosed by CRAE at 800 mg/kg BW; D. liver section of the CCl₄-treated rat post-dosed by CRAE at 600 mg/kg BW; E. liver section of the CCl₄-treated rat post-dosed by CRAE at 400 mg/kg BW; F. liver section of the CCl₄-treated rat post-dosed by berberine at 120 mg/kg BW.