



Title	The iron-chelating drug M30 down-regulates carbon tetrachloride (CCl₄)-induced hepatic oxidative stress, inflammation and apoptosis in vitro
Author(s)	Guo, R; Xiao, J; Pan, J; Liang, EC; Fung, ML; So, KF; Youdim, M; Lin, B; Tipoe, GL
Citation	The 23rd Conference of the Asian Pacific Association for the Study of the Liver (APASL 2013), Singapore, 7-10 March, 2013. In <i>Hepatology International</i>, 2013, v. 7 n. 1 suppl., p. S37-S38, abstract no. 2816
Issued Date	2013
URL	http://hdl.handle.net/10722/184896
Rights	Creative Commons: Attribution 3.0 Hong Kong License

The iron-chelating drug M30 down-regulates carbon tetrachloride (CCl₄)-induced hepatic oxidative stress, inflammation and apoptosis in vitro

Rui Guo¹, Jia Xiao^{1,2}, Jing Fei Pan³, Emily C. Liang¹, Man Lung Fung^{3,4}, Kwok Fai So^{1,4}, Moussa Youdim⁵, Bin Lin^{1,4}, George L. Tipoe^{1,4}

¹Anatomy, LKS Faculty of Medicine, The University of Hong Kong, Hong Kong, Hong Kong S.A.R., ²Center for Gene and Cell Engineering, Shenzhen Institute of Advanced Technology, Chinese Academy of Sciences, Shenzhen, China, ³Physiology, ⁴Brain Hormone Healthy Aging Centre, LKS Faculty of Medicine, The University of Hong Kong, Hong Kong, Hong Kong S.A.R., ⁵Faculty of Medicine, Technion-Israel Institute of Technology, Haifa, Israel

BACKGROUND/AIMS: The novel multifunctional brain permeable ironchelator M30 possesses neuroprotective activities against several insults applicable to various neurodegenerative diseases. However, the effect of M30 on CCl₄ induced acute liver damage is still unknown. The aim of this study is to investigate whether the multifunctional drug M30 could ameliorate CCl₄ induced hepatic injury in human HepG2 cell line. **METHODS:** HepG2 cells were grown in DMEM supplemented with 10 %fetal bovine serum and they were divided into control, CCl₄, M30, and CCl₄ + M30 co-treatment groups. M30 was pretreated for 2 h containing a final concentration at 5 μM. Then CCl₄ was added with a final concentration at 2 μl/ml and incubated for 1 h. Finally, the cells were harvested and the cell viability was determined by colorimetric MTT assay based on conversion of MTT to blue formazan crystals by viable cells. In addition, the malondialdehyde levels were determined using a Bioxytech LPO-586™ kit, and real time PCR was also utilized to test the RNA expression levels of antioxidant enzymes, pro-inflammatory mediators and apoptotic markers. **RESULTS:** M30 significantly reduced CCl₄ triggered cell death and MDA levels. Co-treatment of M30 and CCl₄ up-regulated the expression levels of antioxidant enzymes catalase and glutathione peroxidase, which indicated that M30 reduced CCl₄-induced oxidative stress and inhibited lipid peroxidation. In addition, administration of M30 attenuated hepatic inflammation triggered by CCl₄ via inhibiting pro-inflammatory mediators, such as tumor necrosis factor alpha and interleukin-6. M30 also exhibited its anti-apoptotic activity by downregulating proapoptotic protein Bax, up-regulating anti-apoptotic protein Bcl-XL, as well as by recruiting Fas-associated death domain (FADD). **CONCLUSION:** M30 attenuates CCl₄-induced HepG2 cellular damage through its antioxidant, anti-inflammatory and anti-apoptotic properties, which could serve as a potential functional drug to treat acute hepatotoxicity and chronic liver diseases such as non-alcoholic fatty liver disease and steatohepatitis.