



Title	Perceived Neighborhood Environmental Attributes Associated with Walking and Cycling for Transport among Adult Residents of 17 Cities in 12 Countries: The IPEN Study
Author(s)	Kerr, JWC; Emond, J; Badland, H; Reis, R; Sarmiento, O; Carlson, J; Sallis, J; Cerin, E; Cain, K; Conway, T; Schofield, G; Macfarlane, DJ; Christiansen, L; Van Dyck, D; Davey, R; Aguinaga-Ontoso, I; Salvo, D; Sugiyama, T; Owen, N; Mitáš, J; Natarajan, L
Citation	Environmental Health Perspectives, 2015
Issued Date	2015
URL	http://hdl.handle.net/10722/214744
Rights	Creative Commons: Attribution 3.0 Hong Kong License

**Perceived Neighborhood Environmental Attributes
Associated with Walking and Cycling for Transport
among Adult Residents of 17 Cities in 12 Countries:
The IPEN Study**

**Jacqueline Kerr, Jennifer A. Emond, Hannah Badland,
Rodrigo Reis, Olga Sarmiento, Jordan Carlson, James F. Sallis,
Ester Cerin, Kelli Cain, Terry Conway, Grant Schofield,
Duncan J. Macfarlane, Lars B. Christiansen, Delfien Van Dyck,
Rachel Davey, Ines Aguinaga-Ontoso, Deborah Salvo,
Takemi Sugiyama, Neville Owen, Josef Mitáš,
and Loki Natarajan**

<http://dx.doi.org/10.1289/ehp.1409466>

Received: 11 November 2014

Accepted: 15 July 2015

Advance Publication: 17 July 2015

This article will be available in a 508-conformant form upon final publication. If you require a 508-conformant version before then, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Perceived Neighborhood Environmental Attributes Associated with Walking and Cycling for Transport among Adult Residents of 17 Cities in 12 Countries: The IPEN Study

Jacqueline Kerr¹, Jennifer A. Emond², Hannah Badland³, Rodrigo Reis⁴, Olga Sarmiento⁵, Jordan Carlson¹, James F. Sallis¹, Ester Cerin^{6,7}, Kelli Cain¹, Terry Conway¹, Grant Schofield⁸, Duncan J. Macfarlane⁷, Lars B. Christiansen⁹, Delfien Van Dyck¹⁰, Rachel Davey¹¹, Ines Aguinaga-Ontoso¹², Deborah Salvo^{13,14,15}, Takemi Sugiyama¹⁶, Neville Owen¹⁷, Josef Mitáš¹⁸, and Loki Natarajan¹

¹Division of Behavioral Medicine, University of California San Diego, La Jolla, California, USA; ²Community and Family Medicine, Dartmouth College, Hanover, New Hampshire, USA; ³McCaughey VicHealth Centre for Community Wellbeing, The University of Melbourne, Melbourne, Australia; ⁴Pontiff Catholic University of Paraná, Curitiba, Brazil; ⁵Faculty of Medicine, Universidad de los Andes, Bogotá, Colombia; ⁶Centre for Physical Activity and Nutrition Research, Deakin University, Melbourne, Australia; ⁷Institute of Human Performance, The University of Hong Kong, Pokfulam, Hong Kong; ⁸Human Potential Centre, Auckland University of Technology, Auckland, New Zealand; ⁹Department of Sports Science and Clinical Biomechanics, University of Southern Denmark, Odense, Denmark; ¹⁰Department of Movement and Sport Sciences, Ghent University, Ghent, Belgium; ¹¹Centre for Research and Action in Public Health, University of Canberra, Canberra, Australia; ¹²Department of Health Sciences, Public University of Navarra, Pamplona, Spain; ¹³Nutrition and Health Research Center, Instituto Nacional de Salud Publica (National Institute of Public Health), Cuernavaca, Morelos, Mexico;

¹⁴Michael & Susan Dell Center for Healthy Living, The University of Texas Health Science Center at Houston, School of Public Health (Austin Regional Campus). Austin, Texas, USA;

¹⁵Nutrition and Health Sciences, Graduate Division of Biological and Biomedical Sciences, Emory University, Atlanta, Georgia, USA; ¹⁶School of Population Health, University of South Australia, Adelaide, Australia; ¹⁷Behavioural Epidemiology Laboratory, Baker IDI Heart and Diabetes Institute, Melbourne, Australia; ¹⁸Institute of Active Lifestyle, Faculty of Physical Culture, Palacký University, Olomouc, Czech Republic

Address correspondence to Jacqueline Kerr, Department of Family Medicine & Public Health, University of California, San Diego, 9500 Gilman Dr., La Jolla, CA 92093 USA. Telephone: 1 858 534 9305. E-mail: jkerr@ucsd.edu

Running title: International built environments & active travel

Acknowledgments: Australian data collection was supported by National Health and Medical Research Council (NHMRC) of Australia Project Grant #213114. The contributions of Neville Owen were supported by NHMRC Program Grant #569940, NHMRC Senior Principal Research Fellowship #1003960, and by the Victorian Government's Operational Infrastructure Support Program. Data collection in Hong Kong was supported by the HK Research Grants Council GRF grants (#HKU740907H and #747807H) and HKU URC Strategic Research Theme (Public Health). The contributions of Ester Cerin are supported by the ARC Future Fellowship #140100085. US data collection and Coordinating Center processing was supported by the NIH grants R01 HL67350 (NHLBI) and R01 CA127296 (NCI). The Danish study was partly funded by the Municipality of Aarhus. Data collection in the Czech Republic was supported by the grant MEYS (# MSM 6198959221). The study conducted in Colombia was funded by Colciencias

grant 519_2010, Fogarty and CeIBA. Data collection in New Zealand was supported by the Health Research Council of New Zealand grant # 07/356. Data collection in Mexico was supported by the CDC Foundation, which received a training grant from The Coca-Cola Company. The UK study was funded by the Medical Research Council under the National Preventive Research Initiative.

Competing financial interests: All financial support has been declared in the acknowledgements section. There were no competing financial interests.

Abstract

Introduction: Prevalence of walking and cycling for transport is low, varying greatly across countries. Few studies have examined neighborhood perceptions related to walking and cycling for transport in different countries. Therefore it is challenging to prioritize appropriate built environment interventions. The aim of this study was to examine the strength and shape of the relationship between adults' neighborhood perceptions and walking and cycling for transport across diverse environments.

Methods: As part of the International Physical activity and Environment Network (IPEN) adult project, self-report data were taken from 13,745 adults (18 – 65 years) living in physically and socially diverse neighborhoods in 17 cities across 12 countries. Neighborhood perceptions were measured using the Neighborhood Environment Walkability Scale, and walking and cycling for transport were measured using the International Physical Activity Questionnaire – Long Form. Generalized additive mixed models were used to model walking or cycling for transport during the last seven days with neighborhood perceptions. Interactions by city were explored.

Results: Walking for transport outcomes were significantly associated with perceived residential density, land use mix access, street connectivity, aesthetics, and safety. Any cycling for transport was significantly related to perceived land use mix access, street connectivity, infrastructure, aesthetics, safety, and perceived distance to destinations. Between-city differences existed for some attributes in relation to walking or cycling for transport.

Conclusions: Many perceived environmental attributes supported both cycling and walking; however highly walkable environments may not support cycling for transport. People appear to walk for transport despite safety concerns. These findings can guide the implementation of global health strategies.

Introduction

The 2011 *United Nations High-level Meeting on Non-Communicable Diseases* identified increasing physical activity as one of five priority intervention areas to reduce the impact of non-communicable diseases, noting modification of the built environment to support habitual physical activity as a key focus area (Beaglehole et al. 2011). Engaging in active transport (AT) (i.e., walking and cycling for travel purposes) provides opportunities to habitually accumulate physical activity (Badland 2005), and those who engage in AT tend to be more active in duration and frequency than those who do not (Berrigan et al. 2006).

Compared with those who travel by private motor vehicle, those who walk and cycle for transport have been reported to be less likely to be overweight or obese (Badland and Schofield 2008; Bassett et al. 2008). Co-benefits of AT reported by previous studies include greater social inclusion (Currie et al. 2007), improved air quality and reduced traffic congestion, vehicle miles travelled, and road infrastructure expenditure (Haines et al. 2009). The prevalence of walking and bicycling for transport varies worldwide, with estimated bicycling rates ranging from 1-2% in North America and Australasia to 25% in The Netherlands (Bassett et al. 2008;González et al. 2014; Merom et al. 2010; Reis et al. 2013). Because private motor vehicle journeys often cover distances (< 5 km) that are feasible for AT modes, there is great potential to replace automobile trips with AT that will have health benefits (Sugiyama et al. 2012a; European Commission 1999) (World Health Organization 2004b).

The role of environmental and policy strategies to increase AT has recently received attention, with calls for further evidence on the most-relevant and potentially-modifiable environmental attributes (Fraser and Lock 2011). Several studies have explored associations between built

environment attributes and walking or cycling for transport (Badland et al. 2008; Pucher and Buehler 2008; Saelens and Handy 2008). Few studies, however, had sufficient power and variability to assess walking and cycling separately.

We hypothesises that identifying environmental attributes that benefit both modes will be important to maximize health, social, and environmental gains in a fiscally constrained global environment. AT studies thus far have been primarily limited to Australasia, Europe, and North America (Bassett et al. 2008), and associations have been weak or inconsistent, possibly due to limited variability in the samples. Although objective measures of built environment are important, perceptions of environments also are related to behavior and may provide complementary information. Some attributes, such as aesthetics, cannot be measured objectively, and other attributes, such as sidewalks, are simply unavailable as objective data in most cities. International studies using comparable methods can identify the relevant differences and similarities between countries and inform evidence-based international and country-specific interventions to increase AT.

The purpose of the present study was to examine the strength and shape of the relationship of adults' perceptions of several built-environment attributes, selected for a priori theoretical and empirical reasons, with walking and cycling for transport across diverse cities and countries. Analyses controlled for multiple potential confounding variables and interactions by city were explored to assess international generalizability of findings. Understanding these relationships is critical for guiding policy and practice to support walking and cycling for transport.

Methods

Study Design and Locations

The IPEN (International Physical activity and Environment Network) adult study is an observational epidemiologic multi-country cross-sectional study using a common design and comparable methods (Kerr et al. 2013). Seventeen cities from 12 countries participated: Australia (AUS): Adelaide; Belgium (BEL): Ghent; Brazil (BR): Curitiba; China: Hong Kong (CN); Colombia (COL): Bogota; Czech Republic (CZ): Olomouc and Hradec Kralove; Denmark (DEN): Aarhus; Mexico (MEX): Cuernavaca; New Zealand (NZ): North Shore, Waitakere, Wellington, and Christchurch; Spain (SP): Pamplona; the United Kingdom (UK): Stoke-on-Trent; and the United States of America (US): Seattle and Baltimore. In each of the 17 cities, study neighborhoods were chosen first, and then participants were recruited from these neighborhoods (Frank et al. 2010; Kerr et al. 2013).

Neighborhood selection

Neighborhoods were chosen in each city to maximize variability in environmental attributes and socioeconomic status (SES). Neighborhood walkability index (Frank et al. 2010) determined objectively with geographical information systems (GIS) was used for this purpose, except in Spain where neighborhoods were selected based on their construction date (a proxy measure of walkability (Berrigan and Troiano 2002)). For each country, we used the smallest administrative area unit that represented a neighborhood-level geographic sector for the development of the walkability measures (Adams et al., 2014).

Administrative units were ranked into deciles based on the normalized walkability index and on neighborhood-level (socio-economic status) SES data drawn from the census (e.g., household

income, education attainment, or ethnicity) in each city. The walkability index and census-based SES scores were crossed to produce four neighborhood quadrants: higher walkable / higher SES; higher walkable / lower SES; lower walkable / higher SES; and lower walkable / lower SES (Kerr et al. 2013). Equal numbers of neighborhoods were selected from each of the quadrants. The neighborhood selections methods for each country can be found elsewhere (Adams et al. 2014; Kerr et al. 2013).

Participant Recruitment

Participants were selected systematically using addresses from the identified neighborhoods. Four countries recruited and conducted data collection by phone and mail, with the remaining eight countries contacting households in person. Adults living in the selected neighborhoods were contacted and invited to complete surveys on their physical activity behaviors and perceptions of the neighborhood environment. Study dates ranged from 2002 to 2011, with participants' ages ranging between 16-94 years. Analyses were performed on those aged 18-66 years, as only three countries had a wider age range. Six countries used monetary incentives, three countries provided non-monetary incentives (e.g., physical activity feedback), and three countries provided no incentives for recruitment. Participants were recruited across the seasons to control for variations in weather that may affect physical activity. Further details on the participant recruitment response rates across countries can be found elsewhere (Kerr et al. 2013).

Quality control and comparability

All investigators completed the San Diego State University Institutional Review Board training (as the grant was housed there during the data collection phase), the NIH Fogarty International Center ethical requirements, and their own research institution's ethics requirements. All

participants provided signed informed consent for participation in their home country. Participant confidentiality for pooled data was maintained by de-identification using numeric identification codes rather than names.

All survey data were assessed for completeness by sites and double-checked by the single coordinating center at the University of California, San Diego. Study investigators in each country provided back translations of surveys and the comparability of item wording, response options and number of items was assessed by two independent raters, who were experts in the area (B Saelens UW, B Ainsworth ASU). Only comparable items were included in the scales created and employed in the current analyses.

Measures

Physical activity

The self-administered International Physical Activity Questionnaire long form (IPAQ-LF) was used to measure participants' physical activity for recreation and transport purposes. The IPAQ-LF assesses frequency and duration of activities separately across multiple domains (i.e., recreation, transport, occupation, household) (Craig et al. 2003). The IPAQ-LF has been evaluated in 14 studies across 12 countries on five continents and found to have acceptable test-retest reliability (0.8). Validity was tested by correlations with accelerometers (0.3), with the results being comparable to other self-report surveys (Craig et al. 2003). Seven countries collected IPAQ-LF data using interview techniques, while three countries provided an online version in addition to, or instead of, mailing out paper copies.

The IPAQ-LF items used in the present analyses assessed walking and cycling for transport. The items queried the number of days during the last week that were spent walking or cycling for at least 10 minutes to get from place to place and the usual minutes spent doing so per day. Total minutes per week spent walking or cycling for transport (days*minutes per day) were calculated and treated as continuous variables. In addition, dichotomous outcome measures were derived to represent any walking or cycling for transport during the last week (no, yes) that lasted for at least 10 minutes, and whether ≥ 150 minutes of walking or cycling for transport during the last week was accumulated (no, yes). This reflects the current international adult physical activity guidelines (World Health Organization 2011).

Perceived Environment

Many studies have established the independent predictive value of resident perceptions of the neighborhood environment, in addition to the objective assessment of neighborhood attributes (e.g., those based on audits or GIS), as they relate to physical activity (Adams et al. 2009; Bauman et al. 2012; Mason et al. 2013; Saelens and Handy 2008). Perceptions of neighborhood attributes were assessed among US participants using the 67-item Neighborhood Environment Walkability Scale (NEWS), and perceptions among participants in the 10 remaining countries were assessed using original or slightly modified items from the NEWS scale in combination with items from the NEWS-A scale, an empirically derived abbreviated (54-item) version of NEWS (Cerin et al. 2006). See Appendix for a list of common items employed in each city to assess each of the following environmental predictor subscales: neighborhood residential density, land use mix-access, street connectivity, pedestrian infrastructure, aesthetics, traffic safety, and crime safety. Subscale scores ranged from 0-1044 for residential density, and from 1

(strongly disagree) to 4 (strongly agree) for all other items, with higher scores indicating with more favorable environments (Cerin et al. 2013). Scales were scored so that higher numbers corresponded with more favorable environments. In addition, we assessed perceived distances to walk to 13 common neighborhood destinations (also known as mixed-use diversity). Response options for each destination type were 1 = \geq 31 minute walk or don't know; 2 = 21-30 minute walk; 3 = 11-20 minute walk; 4 = 6-10 minute walk; 5 = 1-5 minute walk. The responses were averaged across the 13 destinations to create a score that ranged from 1 to 5, where higher values represented more destinations within a close walking distance. Reliability and validity of NEWS and NEWS-A have been documented in several countries (Cerin et al. 2007; De Bourdeaudhuij et al. 2003; Leslie et al. 2005; Malavasi et al. 2007), with all included scales having test-retest reliability ICCs \geq 0.75.

Demographic variables

Demographic items collected by all countries included age, sex, education, and marital status. While types of education available varied by country, all country data could be categorized into college graduate or not. Marital status was recoded to indicate ‘married or living with a partner’ versus not.

Data analytic plan

Descriptive statistics (means, medians, standard deviations, percentages, and percentages of missing values) were computed, as appropriate, by study city for all relevant variables. Over 8% of participants had missing data on at least one of the examined variables. To avoid potential biases associated with a complete-case-analysis (Rubin 1987) and to improve efficiency, we used multiple imputation methods to impute missing values. Consequently, 10 imputed datasets were

created for the main regression analyses (see below) as recommended (Rubin 1987; van Buuren 2012). Multiple imputations were performed using Markov Chain Monte Carlo methods (Schafer 1997) accounting for within-site administrative-unit-level cluster effects arising from the two-stage stratified sampling strategy employed in each study site. The ten imputed datasets were created in R (R core team 2013) using the package ‘mix’ for multiple imputation of mixed categorical and continuous variables, and following the model-building and diagnostic procedures outlined by van Buuren (2012). Non imputed results yielded the same conclusions (data not shown). The main aim of this paper was to estimate the strength and shape of associations of multiple perceived environmental attributes with walking and cycling for transport for the whole sample, and to examine if these associations varied by city. The built environment variables we focused on have theoretical and empirical support for their inclusion (Saelens and Handy 2008).

Six physical activity outcomes were explored: any bouts (> 10 min) of walking or cycling (dichotomous), walking or cycling for ≥ 150 minutes per week (dichotomous), and total minutes of walking or cycling in those who walked / cycled for transport (continuous). The three different types of outcomes (any, ≥ 150 minutes, and total minutes) were explored because they have different implications for public health. For example, ≥ 150 minutes represents international physical activity guidelines (World Health Organization 2011); yet even small amounts of activity may still be beneficial for health (Blair et al. 2004). Further, examining total minutes spent walking or cycling allows one to investigate what built environmental correlates are related to more walking / cycling among those who do any amount of these activities.

For these analyses, generalized additive mixed models (GAMMs) were used (Wood 2006). GAMMs can model data following various distributional assumptions (e.g., positively-skewed physical activity data), account for dependency in error terms due to clustering (observations sampled from selected administrative units), and estimate complex, dose-response relationships of unknown form (Wood 2006). In our analysis, the shape of dose-response relationships was estimated using thin plate splines (Wood 2006). Random intercepts were specified to account for within-administrative unit correlations. The appropriateness of the GAMMs and their link functions was assessed via residual plots; quasi-Akaike Information Criteria (qAIC) values were used for model selection (e.g., linear vs nonlinear), whereby a lower qAIC is indicative of a better-fitting model (Wood 2006). For the current analysis, absolute differences in qAIC values of 10 or higher were used as the criteria for model selection (Burnham and Anderson 1998). For the dichotomous outcome variables GAMMs used binomial variance and logit link functions. The reported antilogarithms of the regression coefficient estimates of these models represent odds ratios of walking versus not walking, cycling versus not cycling, and meeting the ≥ 150 minute per week activity recommendations or not. For the ‘total minutes of walking’ (or cycling) outcomes, we used a negative binomial regression model for over dispersed count data. The anti-logarithm of the coefficients from the negative binomial models can be used to estimate proportional increases (or decreases) in minutes of walking (or cycling) associated with changes in environmental attributes.

A first set of models estimated the dose-response relationships of the perceived environmental attribute theoretically relevant to walking and cycling for transport with the outcomes, adjusting for study city, socio-demographic covariates, and design variables including neighborhood-level

and socio-economic status. Separate models were run to estimate main associations of each environmental attribute. Quasi-AIC criteria were used to choose: 1) between curvilinear (thin plate splines) versus linear relationships of environmental attributes with outcomes; and 2) whether to include two-way city by environmental attributes interaction effect estimates. Interactions were tested for each model to see if there were significant ($AIC < 10$) differences in the relationships across the 17 cities. For variables where a significant interaction was found, the main associations for each city were presented as forest plots. For significant nonlinear associations the shape of the curve was plotted. All analyses were conducted in R (R core team 2013) using the packages ‘mix’ (Schafer 2013), and ‘mgcv’ (Wood 2006).

Results

Descriptive Results

Table 1 describes the sample in each city. The total sample size was 13,745 adults. The study aimed to balance samples by walkability, SES, and gender, and the percentages demonstrate these goals were achieved. Percentages of the sample with a partner ranged from 44.8-74.2% and those with a university degree ranged from 14.1-67.6%. The average age ranged from 34.0-46.6 years.

Table 2 demonstrates the varying range in the six AT outcomes across the 17 cities. The highest percentages reporting any walking for transport in the previous seven days were found in Pamplona (SP) (92.3%), followed by Cuernavaca (MEX), and Bogota (COL) (90.3%); Aarhus (DEN), the cities from the Czech Republic, and Wellington (NZ) were also over 80%. Levels of any cycling for transport in the last seven days were much lower (in contrast to any walking) ranging from 1.2% in Cuernavaca (MEX) to 62.5% in Aarhus (DEN).

The self-reported built environment perceptions for the neighborhood attributes varied greatly across cities. Residential density scores ranged from 18.2 in Waitakere (NZ) to 439.7 in Hong Kong (CN) (Supplemental Material, Table S1). The differences in means of the other environmental variables across the cities were relatively small, about 0.7 in the variables assessed with a 4-point scale. Larger between-city variability was found for crime safety, from 2.1 in Bogota (COL) to 3.5 in Pamplona (SP). Participants in some cities like Curitiba (BRZ) for example, reported high land use access (3.7) but low traffic safety (2.4).

Results of Regression Analyses

Estimated associations of perceived environment subscales with four of the six outcome variables (≥ 150 min. walking, total minutes walking, any cycling, and total minutes cycling) are shown in Table 3. Associations with any walking for transport (data not shown) were very similar to associations with walking ≥ 150 minutes, which may be more relevant to health outcomes. The low prevalence of participants meeting ≥ 150 minutes cycling for transport outcome led us to present the environmental correlates of any cycling only.

Walking

There was a significant nonlinear association between perceived residential density and ≥ 150 minutes walking for transport that was positive up to a perceived density score of approximately 500, and flat or negative for higher scores (Figure 1A). Perceived land use mix-access, street connectivity, pedestrian infrastructure, aesthetics, and perceived distance to destinations all had significant positive linear associations with ≥ 150 minutes walking for transport during the previous week, while traffic safety had a significant negative association with this outcome (Table 3). In addition, there were significant differences among cities (interactions) for

associations between this outcome and street connectivity and perceived distance to destinations (Figure 2A and B).” Total minutes of walking for transport during the previous week was positively associated with perceived residential density, land use mix access, street connectivity, and aesthetics, and negatively associated with traffic and crime safety (Table 3).

Cycling

There was a significant nonlinear association between perceived residential density and any cycling for transport that was consistently negative in slope (Figure 1B). No other relationship was curvilinear. Any cycling for transport was positively associated with perceived land use mix access, street connectivity, infrastructure, aesthetics, traffic safety, crime safety, and perceived distance to destinations when estimated across all populations (Table 3). However, there were significant interactions of perceived land use mix access and infrastructure with study city, indicating significant variation among the different sites (Figure 2C and D).

Discussion

These analyses explored the strength and shape of the relationship between perceived environment and walking / cycling for transportation and city differences in these associations in a 17-city study. The variation in prevalence across cities for any walking for transport (52% to 92%) and cycling for transport (1% to 63%) demonstrates the value of studying such behaviors globally.

Perceived residential density

Perceived residential density had a significant nonlinear association with walking for transport \geq 150 minutes during the previous week with a positive slope that subsequently plateaued,

suggesting that walking for transport did not continue to increase with perceived density in extremely dense neighborhoods; e.g. in neighborhoods with a perceived residential density score above approximately 500 i.e. above the mean score for Hong Kong (439.7). Perceived residential density has consistently been positively related with walking for transport in many other studies, but densities were not as varied and high as in the current study (Bauman et al. 2012; Eriksson et al. 2012; Saelens and Handy 2008; Sugiyama et al. 2012b; Van Holle et al. 2012; Witten et al. 2012).

In contrast, although the association between perceived density and any cycling for transport was also nonlinear, it was consistently negative across the entire range of perceived density scores, suggesting that increases in perceived density did not support cycling for transport. . Previous studies, mostly in children, have found population density positively related to cycling (Fraser and Lock 2011). Cycling also increased in adult participants who relocated to higher residential density communities (Beenackers et al. 2012). Previous studies in Brazil, Sweden and the US, however, found that neighborhood walkability was not significantly related to cycling (Eriksson et al. 2012; Reis et al. 2013; Sallis et al. 2013). Since cycling trips in adults are normally greater than the 1km neighborhood buffer used in such studies, it is not surprising that neighborhood features alone do not explain these behaviors. Residential density was negatively related to cycling in our study, possibly because highly connected pedestrian streets with cross walks in densely populated areas do not support preferred cycling speeds. Bicycling rates in moderately-dense cities, like Aarhus, Denmark, are partly due to due to specialized infrastructure that supports uninterrupted cycling (Pucher and Buehler 2008). Examples include traffic lights timed to enhance cycling speeds or bikeways that are separate from any pedestrian or road traffic.

Cities like Hong Kong, that are dense, may not support cycling because distances are very short and can be covered by walking without the inconvenience of storing and transporting a bicycle in high rise buildings. Present results may contradict previous studies conducted only in cities with limited variations in density. Present findings also demonstrate the importance of analyzing walking and cycling separately so that different environmental predictors can be detected.

Perceived land use mix access

Perceived land use mix access (having easy access to shops, recreation uses, and transit stops) was significantly and positively related to walking ≥ 150 minutes a week and total minutes of walking. It was also associated with any cycling for transport, but there was a significant city interaction. In two cities (Olomouc and Hong Kong), higher land use mix access was significantly associated with lower odds of doing any cycling, while a positive association was found in seven cities (Van Holle et al. 2012). Stronegger and colleagues (2010) found better access to diverse land uses was related to walking for transport but not to cycling. Many previous studies of cycling found no associations for neighborhoods with multiple land uses and destinations but may have lacked statistical power (Van Holle et al. 2012). It is possible that the positive associations between land use access and both walking and cycling in our multi-city study were evident in part because of the greater variability in exposures and outcomes compared with previous study populations.

Perceived street connectivity

Studies that have reported on intersection density (an objective measure of perceived street connectivity), including in the US, demonstrated intersection density was related to walking for transport (Saelens et al. 2012) but not with cycling (Sallis et al. 2013), potentially due to lack of

statistical power with low cycling rates in these studies. A European review (Van Holle et al. 2012) identified only one study (Van Dyck et al. 2011) showing perceived street connectivity to be positively related to cycling for transport, but not with walking. In the present 17-city study, perceived street connectivity was related to both total minutes walked for transport and any cycling for transport. The significant city interaction for obtaining ≥ 150 minutes of walking for transport revealed that street connectivity was positively related to walking in all but 2 cities, but only significant in 3 cities. This suggests that street connectivity alone may not be sufficient to support AT in most cities.

Perceived pedestrian infrastructure

We found that perceived infrastructure focused on facilities for pedestrians (i.e., street lights, crossings and sidewalks) was positively related to ≥ 150 minutes of walking for transport. Sidewalks have been associated with walking for transport (not recreation) in some previous studies (McCormack et al. 2012; Sugiyama et al. 2012b). In three cities, pedestrian infrastructure was negatively related to cycling for transport. The negative relationships between perceived infrastructure and any cycling in Pamplona, Hong Kong, and Cuernavaca suggest these environments are designed predominantly for pedestrians and are less supportive of cycling (Mosquera et al. 2012).

Perceived aesthetics

Mixed results for aesthetics with AT behaviors were shown in a recent European review (Van Holle et al. 2012), and previous studies have found aesthetics are related to recreational activity, but not AT engagement (Bauman et al. 2012; Saelens and Handy 2008; Sugiyama et al. 2012b; Witten et al. 2012). Aesthetics, however, was found to be significant for both walking and

cycling for transport in this 17-city study. Aesthetics ratings, like safety ratings, were low across all cities, and this may be an area for improvement with less cost implications than other structural changes (Becerra et al. 2013; Beenackers et al. 2012).

Perceived traffic and crime safety

We found that perceived safety from crime and traffic was positively related to any cycling for transport, but negatively related to the amount of walking/cycling among those who walked/bicycled. Safety from traffic was negatively associated with ≥ 150 minutes walked for transport per week. In contrast, previous studies did not find significant associations between perceived safety and walking (Van Holle et al. 2012). It may be that individuals walk regardless of safety issues because they have no other choice (e.g., they do not own a car, travel in areas not serviced by public transport). Further, walkers may be more aware of the threats to safety, compared to those who do not walk (Adams et al. 2009).

Perceived safety is challenging to assess within an international context because participants rate their neighborhoods relative to their own experience, which can differ between countries. For example, a study in New York City, US, found that lower homicide rates were related to higher population-level AT behaviors (Lovasi et al. 2013). Incorporating objective measures of safety (Foster and Giles-Corti 2008) and more nuanced measures of bicycling risk maybe required to improve understanding of likely-complex relation of perceived safety and physical activity.

Perceived distance to local destinations

We found that proximity of perceived destinations was marginally related to total walking for transport, but not associated with minutes spent cycling for transport. Perceiving local

destinations nearby was significantly related to more adults doing any cycling for transport, but there was a significant city interaction for performing ≥ 150 minutes of walking for transport. The relationship between proximity of local destinations and walking was significant for six cities. Many other studies have shown associations between local destinations and walking (Bauman et al. 2012; Saelens and Handy 2008). The negative relationship in Bogota (with the 2nd highest number of destinations) could be explained by trips made by walking being less than 10 minutes per day. The IPAQ-LF includes only trips of at least 10 minutes, therefore may not capture shorter trips which may be common in Bogota.

Policy implications

The negative relationship between perceived residential density and cycling for transport, alongside the two city interactions showing that walkable cities may not support cycling have important policy implications. In highly pedestrianized areas it may be necessary to co-locate cycling facilities. In some cities this could mean locating cyclists on a dedicated path on the road, or in pedestrian centers, providing sidewalks with separation for cyclists from pedestrians. In countries such as the US, where walkability is limited but distances are cycleable, promotion of cycle use may increase cycling rates. Pucher et al. (2008) concluded that substantial increases in cycling for transport requires an integrated package of multiple complementary interventions, including infrastructure provision and pro-bicycle programs, supportive land use planning, and restrictions on car use. The combination of economic, convenience and health arguments in favor of cycling may be important to increase support for cycling investments among policy makers (Jones and Ogilvie 2012; Kahlmeier et al. 2010; Wooller et al. 2012). In some countries, such as Brazil, Mexico and Colombia, cycling is considered a ‘poor man’s’ transport mode, but

environmental supports for cycling and changing social norms could alter such negative perceptions (Mosquera et al. 2012).

Limitations

The cross sectional design of the current study limits evidence of causality that might support policy change. The variability and strength of associations achieved in this study, however, improve on previous cross sectional studies and set the stage for a meaningful prospective study.

The present study focused only on self-reported perceptions of the built environment assessed with scales that had limited variability, even within this international context. Self-report measures are limited when making international comparisons because people tend to make assessments relative to their own experience. Further, thresholds from self-reported scales may not be helpful to policy makers. There were some methodological differences across countries during the neighborhood selection and recruitment phases that reflected local conditions and capacity and may have affected study findings and contributed to variations in associations across countries. The independent variables and outcome measures, however, were collected consistently and checked for comparability. The present analyses did not include assessment of cycling infrastructure, because the NEWS was developed in the US where such infrastructure is mainly absent. A subset of countries deployed additional cycling infrastructure scales, and these will be explored in future analyses. Unfortunately, most cities do not have good GIS data for bicycling infrastructure so future studies will need to employ street audits to assess quality and quantity of these facilities.

The present study relied on self-reported measures of walking and cycling for transport. Total physical activity estimates are often over reported by the IPAQ-LF, but AT is usually more accurately reported than recreation physical activity (Johnson-Kozlow et al. 2006). Future studies should employ GPS devices and/or travel diaries to identify trips in different modes (Carlson et al. 2015; Duncan et al. 2009). Finally, the IPAQ-LF elicits information about trips \geq 10 minutes, potentially underestimating the relationship between the built environment and AT in highly walkable environments.

Conclusions

This 17-city study of perceived environmental correlates of walking and cycling for transport demonstrated the importance of designing a study to capture environmental and behavioral variability. Many environmental attributes supporting both cycling and walking were found. People may walk for transport despite safety concerns. Highly walkable environments may not support cycling for transport. Our study highlights the importance of examining walking and cycling separately and testing neighborhood attributes discretely.

References

- Adams, M.A., Frank, L.D., Schipperijn, J., Smith, G., Chapman, J., Christiansen, L.B., Coffee, N., Salvo, D., du Toit, L., Jan Dygrýn, J., Hino, A.A.F., Lai, P-C., Mavoa, S., Pinzón, J.D., Van de Weghe, N., Cerin, E., Davey, R., Macfarlane, D., Owen, N., and Sallis, J.F. (2014). International variation in neighborhood walkability, transit, and recreation environments using Geographic Information Systems: The IPEN Adult Study. *International Journal of Health Geographics*, 13:43. doi: 10.1186/1476-072X-13-43.
- Adams M, Ryan S, Kerr J, Sallis J, Patrick K, Frank L, et al. 2009. Validation of the neighborhood environment walkability scale (NEWS) items using geographic information systems. *Journal of Physical Activity & Health* 6:S113-S123.
- Badland H, Schofield G. 2008. Health associations with transport-related physical activity and motorized travel to destinations. *International Journal of Sustainable Transport* 2:77-90.
- Badland H, Schofield G, Garrett N. 2008. Travel behavior and objectively measured urban design variables: Associations for adults traveling to work. *Health & Place* 14:85-95.
- Bassett D, Pucher J, Buehler R, Thompson D, Crouter S. 2008. Walking, cycling, and obesity rates in Europe, North America, and Australia. *Journal of Physical Activity & Health* 5:795-814.
- Beaglehole R, Bonita R, Horton R, Adams C, Alleyne G, Asaria P, et al. 2011. Priority actions for the non-communicable disease crisis. *The Lancet* 377:1438-1447.
- Becerra J, Reis R, Frank L, Ramirez-Marrero F, Welle B, Arriaga Cordero E, et al. 2013. Transport and health: A look at three latin american cities. *Cad Saude Publica* 29:654-666.
- Beenackers M, Foster S, Kamphuis C, Titze S, Divitini M, Knuijman M, et al. 2012. Taking up cycling after residential relocation: Built environment factors. *American Journal of Preventive Medicine* 42:610-615.
- Berrigan D, Troiano R. 2002. The association between urban form and physical activity in U.S. Adults. *American Journal of Preventive Medicine* 23:74-79.
- Berrigan D, Troiano R, McNeel T, DiSogra C, Ballard-Barbash R. 2006. Active transportation increases adherence to activity recommendations. *American Journal of Preventive Medicine* 31:210-216.

- Blair S, LaMonte M, Nichaman M. 2004. The evolution of physical activity recommendations: How much is enough? *American Journal of Clinical Nutrition* 79:913S-920S.
- Burnham K, Anderson D. 1998. *Model selection and inference: A practical information-theoretic approach*. Berlin:Springer Verlag.
- Carlson, J.A., Saelens, B.E., Kerr, J., Schipperijn, J., Conway, T.L., Frank, L.D., Chapman, J.E., Glanz, K., Cain, K., and Sallis, J.F. (2015). Association between neighborhood walkability and GPS-measured walking, bicycling and vehicle time in adolescents. *Health and Place*, 32, . Published online January 9, 2015.
- Cerin E, Saelens B, Sallis J, Frank L. 2006. Neighborhood environment walkability scale: Validity and development of a short form. *Medicine & Science in Sports & Exercise* 38:1682-1691.
- Cerin E, Macfarlane D, Ko H, Chan K. 2007. Measuring perceived neighbourhood walkability in Hong Kong. *Cities* 24:209-217.
- Cerin E, Conway T, Cain K, Kerr J, De Bourdeaudhuij I, Owen N, et al. 2013. Sharing good news across the world: Developing comparable scores across 12 countries for the neighborhood environment walkability scale (NEWS). *BMC Public Health* 8.
- Craig C, Marshall A, Sjostrom M, Bauman A, Booth M, Ainsworth B, et al. 2003. International physical activity questionnaire: 12-country reliability and validity. *Medicine & Science in Sports & Exercise* 35:1381-1395.
- Currie G, Stanley J, Stanley J. 2007. *No way to go: Transport and social disadvantage in australian communities*. Melbourne:Monash University ePress.
- De Bourdeaudhuij I, Sallis J, Saelens B. 2003. Environmental correlates of physical activity in a sample of belgian adults. *American Journal of Health Promotion* 18:83-92.
- Duncan M, Badland H, Mummery W. 2009. Applying gps to enhance understanding of transport-related physical activity. *Journal of Science and Medicine in Sport* 12:549-556.
- Erikkson U, Arvidsson D, Gebel K, Ohlsson H, Sundquist K. 2012. Walkability parameters, active transportation and active physical activity: Moderating and mediating effects of motor vehicle ownership in a cross-sectional study. *International Journal of Behavioral Nutrition and Physical Activity* 9.

- European Commission. 1999. *Cycling: The way ahead for towns and cities*. Belgium:European Commission.
- Foster S, Giles-Corti B. 2008. The built environment, neighborhood crime and constrained physical activity: An exploration of inconsistent findings. *Preventive Medicine* 47:241-251.
- Frank L, Sallis J, Saelens B, Leary L, Cain K, Conway T, et al. 2010. The development of a walkability index: Application to the neighborhood quality of life study. *British Journal of Sports Medicine* 44:924-933.
- González S, Sarmiento O, Lozano O, Ramírez A, Grijalba C. 2014. Niveles de actividad física de la población colombiana: Desigualdades por sexo y condición socioeconómica. *Biomédica* doi: 10.1186/1471-2458-12-474.
- Haines A, McMichael A, Smith K, Roberts I, Woodcock J, Markandya A, et al. 2009. Public health benefits and strategies to reduce greenhouse-gas emissions: Overview and implications for policy makers. *The Lancet* 374:2104-2114.
- Johnson-Kozlow M, Sallis J, Gilpin E, Rock C, Pierce J. 2006. Comparative validation of the ipaq and the 7-day par among women diagnosed with breast cancer. *International Journal of Behavioral Nutrition and Physical Activity* 3.
- Jones C, Ogilvie D. 2012. Motivations for active commuting: A qualitative investigation of the period of home or work relocation. *International Journal of Behavioral Nutrition and Physical Activity* 9.
- Kahlmeier S, Racioppi F, Cavill N, Rutter H, Oja P. 2010. 'Health in all policies' in practice: Guidance and tools to quantifying the health effects of cycling and walking. *Journal of Physical Activity & Health* 7:S120-S125.
- Kerr J, Sallis J, Owen N, De Bourdeaudhuij I, Cerin E, Reis R, et al. 2013. Advancing science and policy through a coordinated international study of physical environments: IPEN methods. *Journal of Physical Activity & Health* 10:581-601.
- Leslie E, Saelens B, Frank L, Owen N, Bauman A, Coffee N, et al. 2005. Residents' perceptions of walkability attributes in objectively different neighbourhoods: A pilot study. *Health & Place* 11:227-236.

- Malavasi L, Duarte M, Both J, Reis R. 2007. Escala de mobilidade ativa no ambiente comunitario—news brasil: Retraducao e reprodutibilidade. (neighborhood walkability scale—NEWS Brazil: Back translation and reliability) *Revista Brasileira de Cineantropometria & Desempenho Humano* 9:339-350.
- Mason P, Kearns A, Livingston M. 2013. "Safe going": The influence of crime rates and perceived crime and safety on walking in deprived neighbourhoods. *Social Science & Medicine* 91:15-24.
- McCormack G, Friedenreich C, Sandalack B, Giles-Corti B, Doyle-Baker P, Shiell A. 2012. The relationship between cluster-analysis derived walkability and local recreational and transportation walking among canadian adults. *Health & Place* 18:1079-1087.
- Merom D, van der Ploeg H, Corpuz G, Bauman A. 2010. Public health perspectives on household travel surveys: Active travel between 1997 and 2007. *American Journal of Preventive Medicine* 39:113-121.
- Mosquera J, Parra D, Gomez L, Sarmiento O, Schmid T, Jacoby E. 2012. An inside look at active transportation in Bogotá: A qualitative study. *Journal of Physical Activity & Health* 9:776-785.
- Owen N, Humpel N, Leslie E, Bauman A, Sallis J. 2004. Understanding environmental influences on walking: Review and research agenda. *American Journal of Preventive Medicine* 27:67-76.
- Pucher J, Buehler R. 2008. Making cycling irresistible: Lessons from the netherlands, denmark, and germany. *Transport Reviews* 28.
- R core team. 2013. *A language and environmetn for statistical computing*. Vienna:R Foundation for Statistical Computing.
- Reis R, Hino A, Parra D, Hallal P, Brownson R. 2013. Bicycling and walking for transportation in three brazilian cities. *American Journal of Preventive Medicine* 44:e9-e17.
- Rubin D. 1987. *Multiple imputation for non-response in surveys*. New York:Wiley J & Sons.
- Saelens BE, Handy SL. Built environment correlates of walking: a review. *Med Sci Sports Exerc*. 2008 Jul;40(7 Suppl):S550-66
- Schafer J. 1997. *Analysis of incomplete multivariate data*. London:Chapman & Hall.

- Schafer J. 2013. Estimation / multiple imputation for mixed categorical and continuous data. Schafer, JL.
- Stronegger W, Titze S, Oja P. 2010. Perceived characteristics of the neighborhood and its association with physical activity behavior and self-rated health. *Health & Place* 16:736-743.
- Sugiyama T, Merom D, van der Ploeg H, Corpuz G, Bauman A, Owen N. 2012a. Prolonged sitting in cars: Prevalence, socio-demographic variations, and trends. *Preventive Medicine* 55:315-318.
- van Buuren S. 2012. Flexible imputation of missing data. Boca Raton, Florida: Chapman & Hall.
- Van Dyck D, Cardon G, Deforche B, Giles-Corti B, Sallis J, Owen N, et al. 2011. Environmental and psychosocial correlates of accelerometer-assessed and self-reported physical activity in Belgian adults. *International Journal of Behavioral Medicine* 18:235-245.
- Van Dyck D, Cerin E, Conway T, De Bourdeaudhuij I, Owen N, Kerr J, et al. 2012. Perceived neighborhood environmental attributes associated with adults' transport-related walking and cycling: Findings from the USA, Australia and Belgium. *International Journal of Behavioral Nutrition and Physical Activity* 9.
- Van Holle V, Deforche B, Van Cauwenberg J, Goubert L, Maes L, Van de Weghe N, et al. 2012. Relationship between the physical environment and different domains of physical activity in European adults: A systematic review. *BMC Public Health* 2012, 12:807
- Wood S. 2006. Generalized additive models: An introduction with r. Boca Raton, Florida: Chapman & Hall.
- Wooller L, Badland H, Schofield G. 2012. Pedestrianisation: Are we reading from the same page? Perspectives from key stakeholders in Takapuna, Auckland. *Graduate Journal of Sport, Exercise, and Physical Education Research* 1:16-30.
- World Health Organization. 2004a. Transport, environment and health. 89. Austria: Regional Office for Europe of the World Health Organization.
- World Health Organization. 2004b. Global strategy on diet, physical activity and health. Geneva: World Health Organization.
- World Health Organization Centre for Health Development. 2011. Healthy urban planning: Report of a consultation meeting. Kobe: World Health Organization Centre for Health Development.

Table 1: Demographic characteristics of sample by city

City	N	% low walk NH	% low SES NH	% female	% with partner	% college grad	Mean age \pm SD
AUS: Adelaide	2650	51.4	47.9	64.0	56.5	46.3	44.5 \pm 12.3
BEL: Ghent	1166	50.0	49.7	52.1	73.4	60.9	42.7 \pm 12.6
BRZ: Curitiba	697	49.8	50.2	52.9	58.1	38.7	41.1 \pm 13.2
CN: Hong Kong	493	47.1	48.7	58.9	59.0	40.0	42.8 \pm 11.7
COL: Bogota	963	44.8	59.5	63.7	53.4	22.2	40.0 \pm 13.7
CZ: Hradec Kralove	167	53.3	31.7	60.5	47.4	26.1	34.0 \pm 13.1
CZ: Olomouc	330	32.1	40.6	62.7	58.4	32.2	37.9 \pm 14.7
DEN: Aarhus	642	46.6	43.9	56.7	65.4	48.0	39.0 \pm 13.9
MEX: Cuernavaca	677	50.5	49.8	55.4	64.7	27.6	42.1 \pm 12.6
NZ: Christchurch	495	50.3	50.3	55.8	55.4	32.0	41.7 \pm 12.6
NZ: North Shore	511	50.3	33.3	63.9	70.4	38.3	41.1 \pm 11.8
NZ: Waitakere	512	48.6	59.0	60.7	74.2	30.7	40.8 \pm 11.8
NZ: Wellington	496	49.4	50.0	51.2	56.7	52.2	39.2 \pm 12.7
SP: Pamplona	904	32.0	56.9	55.2	53.0	57.8	38.7 \pm 14.2
UK: Stoke-on-Trent	843	77.5	47.1	56.1	44.8	14.1	43.0 \pm 13.3
US: Baltimore	912	50.8	47.5	52.3	60.5	67.6	46.6 \pm 10.7
US: Seattle	1287	49.4	48.7	45.2	63.2	63.2	44.0 \pm 11.0

Key: grad = graduate; NH = neighborhood; SD = standard deviation; SES = socioeconomic status; walk = walkability

Australia (AUS), Belgium (BEL), Brazil (BR), Colombia (COL), Czech Republic (CZ), Denmark (DEN), China (CN), Mexico (MEX), New Zealand (NZ), Spain (SP), the United Kingdom (UK) and the United States (US).

Table 2: Prevalence of walking and cycling for transport outcomes by city assessed in previous week by IPAQ-LF

City	Any walking for transport N (%)	Any cycling for transport N (%)	≥ 150 minutes walking for transport N (%)	≥ 150 minutes cycling for transport N (%)	Mean ± SD total minutes walking for transport ^a	Mean ±SD total minutes cycling for transport ^b
AUS: Adelaide	1998 (77.1)	304 (11.8)	973 (36.7)	130 (4.9)	200.2 ±412.7	27.7 ±174.0
BEL: Ghent	608 (52.1)	504 (43.2)	194 (16.6)	194 (16.6)	79.3 ±155.9	63.9 ±119.7
BRZ: Curitiba	538 (77.3)	52 (7.5)	195 (27.9)	19 (2.7)	153.3 ±300.0	14.7 ±78.2
CN: Hong Kong	377 (78.9)	44 (9.7)	249 (50.5)	22 (4.4)	288.9 ±881.2	18.8 ±83.1
COL: Bogota	870 (90.3)	89 (9.2)	490 (50.9)	39 (4.1)	303.7 ±490.9	21.7 ±125.0
CZ: Hradec Kralove	139 (83.2)	59 (35.3)	87 (52.1)	26 (15.6)	298.1 ±411.1	85.3 ±233.7
CZ: Olomouc	272 (83.2)	60 (18.2)	203 (61.5)	24 (7.3)	401.5 ±591.9	35.7 ±141.0
DEN: Aarhus	514 (86.1)	401 (62.5)	229 (35.7)	188 (29.3)	190.8 ±331.4	136.1 ±222.5
MEX: Cuernavaca	611 (90.3)	8 (1.2)	303 (44.7)	1 (0.2)	325.7 ±584.9	0.8 ±8.5
NZ: Christchurch	278 (56.2)	66 (13.3)	84 (17.0)	29 (5.9)	79.7 ±177.4	26.4 ±110.5
NZ: North Shore	334 (65.5)	31 (6.1)	98 (19.1)	11 (2.2)	86.1 ±154.8	14.1 ±127.3
NZ: Waitakere	319 (62.3)	37 (7.2)	70 (13.7)	13 (2.5)	88.7 ±279.8	12.6 ±81.0
NZ: Wellington	422 (85.1)	40 (8.1)	210 (42.3)	20 (4.0)	180.0 ±220.5	18.6 ±128.0
SP: Pamplona	810 (92.3)	111 (12.4)	560 (61.9)	42 (4.6)	322.0 ±353.1	23.1 ±108.4
UK: Stoke-on-Trent	553 (65.8)	35 (4.2)	287 (34.0)	25 (3.0)	218.4 ±426.3	13.9 ±109.1
US: Baltimore	620 (68.1)	60 (6.6)	305 (33.4)	19 (2.1)	171.4 ±302.8	8.5 ±43.1
US: Seattle	877 (68.3)	116 (9.0)	405 (31.5)	49 (3.8)	173.9 ±359.4	17.0 ±99.2

Australia (AUS), Belgium (BEL), Brazil (BR), Colombia (COL), Czech Republic (CZ), Denmark (DEN), China (CN), Mexico (MEX), New Zealand (NZ), Spain (SP), the United Kingdom (UK) and the United States (US).

^a total minutes in those who reported any walking ^b total minutes in those who reported any cycling

Table 3: Estimated associations between perceived environment attributes and walking and cycling for transport assessed in the previous week by IPAQ-LF ^a

Environmental Attributes	≥ 150 minutes walking for transport OR (95% CI); p value N=13745	Total minutes walking for transport in those who report any walking exp(β) (95% CI); p value N=4939	Any cycling for transport OR (95% CI); p value N=13745	Total minutes cycling for transport in those who report any cycling exp(β) (95% CI); p value N=851
Residential density	NA ^b	1.001 (1.000, 1.001); p<0.001	NA ^b	1.00 (0.999, 1.001); p=0.805
Land use mix-access	1.33 (1.24, 1.42); p<0.001	1.08 (1.03, 1.14); p=0.001	1.24 (1.13, 1.36); p<0.001 ^c	1.05 (0.95, 1.15); p=0.359
Street connectivity	1.15 (1.09, 1.21); p<0.001 ^c	1.06 (1.02, 1.11); p=0.003	1.14 (1.06, 1.22); p=0.001	1.00 (0.92, 1.08); p=0.945
Pedestrian infrastructure	1.12 (1.04, 1.21); p=0.002	1.04 (0.98, 1.10); p=0.193	1.22 (1.10, 1.36); p<0.001 ^c	0.94 (0.84, 1.05); p=0.267
Aesthetics	1.19 (1.11, 1.27); p<0.001	1.05 (1.00, 1.11); p=0.032	1.15 (1.05, 1.26); p=0.003	1.01 (0.92, 1.11); p=0.814
Traffic safety	0.92 (0.86, 0.97); p=0.005	0.93 (0.89, 0.98); p=0.002	1.14 (1.05, 1.24); p=0.001	0.91 (0.83, 0.99); p=0.033
Crime safety	0.99 (0.93, 1.05); p=0.667	0.94 (0.90, 0.99); p=0.010	1.17 (1.07, 1.28); p=0.001	0.87 (0.80, 0.95); p=0.002
Distance to local destinations	1.19 (1.12, 1.27); p<0.001 ^c	1.05 (1.00, 1.10); p=0.052	1.16 (1.06, 1.27); p=0.001	1.07 (0.99, 1.17); p=0.108

Key: CI = Confidence Interval; OR = odds ratio;

^a All models adjusted for participant sociodemographics, site, and study design variables (neighborhood-area unit and socio-economic status)

^b Association significant but not linear. Shape of relationship presented in Figure 1.

^c Significant interaction by city, see Figure 2.

Appendix. Neighborhood Environment Walkability Scale-Abbreviated (NEWS-A): IPEN Subscales and Items

Residential density (weighted rating of housing types in neighborhood) *How common are...*

1. Detached single-family residences
2. Townhouses or rows of 1-3 stories houses
3. Apartments or condos with 1-3 stories
4. Apartments or condos with 4-6 stories
5. Apartments or condos with 7-12 stories
6. Apartments or condos with >12 stories
7. Apartments or condos with >20 stories

Land use mix – access

1. Stores are within easy walking distance of my home.
2. There are many places to go within easy walking distance of my home.
3. It is easy to walk to a transit stop (bus, train) from my home.

Street connectivity

1. The distance between intersections in my neighborhood is usually short (100 yards or less; the length of a football field or less).
2. There are many alternative routes for getting from place to place in my neighborhood (I don't have to go the same way every time).

Pedestrian infrastructure

1. There are sidewalks on most of the streets in my neighborhood.
2. My neighborhood streets are well lit at night.
3. Walkers and bikers on the streets in my neighborhood can be easily seen by people in their homes.
4. There are crosswalks and pedestrian signals to help walkers cross busy streets in my neighborhood.

Aesthetics

1. There are trees along the streets in my neighborhood.
2. There are many interesting things to look at while walking in my neighborhood.
3. There are many attractive natural sights in my neighborhood (such as landscaping, views).
4. There are attractive buildings/homes in my neighborhood.

Traffic safety

1. There is so much traffic along nearby streets that it makes it difficult or unpleasant to walk in my neighborhood.
2. The speed of traffic on the street I live on is usually slow (30 mph or less).
3. Most drivers exceed the posted speed limits while driving in my neighborhood.

Crime safety

1. There is a high crime rate in my neighborhood.
2. The crime rate in my neighborhood makes it unsafe to go on walks during the day.
3. The crime rate in my neighborhood makes it unsafe to go on walks at night.

Perceived distance to local destinations *About how long would it take to walk from your home to the nearest...*

1. Supermarket
2. Other food/grocery, small grocery/convenience, fruit/veg market, bakery, butchery
3. Post office
4. Any school, elementary, other, nursery
5. Transit stop
6. Any restaurant, fast food, non-fast food, café/coffee place
7. Park/other public open space
8. Gym/fitness facility, recreation center, swimming pool
9. Library

10. Video store

11. Drug store/pharmacy

12. Bookstore

13. Other shops and services

Figure Legends

Figure 1a: Shape of significant non-linear relationships between perceived residential density and 150 minutes walking for transport during the last week.

Figure 1b: Shape of significant non-linear relationships between perceived residential density and any cycling for transport during the last week. The solid lines represent point estimates (and dashed lines their 95% confidence intervals). The grey lines are the median (and CIs) of the imputed point estimates. The tick lines above the x-axis represent the number of participants reporting this level of residential density. Residential density was the only variable with a significant non linear association.

Figure 2: Forest plots of city-specific effects for any cycling for transport and 150 minutes walked for transport during the last week. The variables plotted demonstrated a significant interaction in the main analyses, highlighted by ^c in table 3. Interactions that were not significant are not plotted. Analyses adjusted for participant sociodemographics and study design variables (neighborhood-area unit and socio-economic status)




Figure 1 (A, B)




Figure 2 (A-D)