

Discussion

Discussion on the article "Remarks on the Permian-Triassic transition in Central and Eastern Lombardy (Southern Alps, Italy)" by G. Cassinis, M. Durand and A. Ronchi

Discusión sobre el artículo "Apuntes sobre el tránsito Pérmico-Triásico en Lombardía central y oriental (Alpes Meridionales, Italia)" de G. Cassinis, M. Durand y A. Ronchi

R. Posenato¹, D. Sciunnach², E. Garzanti³

¹ Dipartimento di Scienze della Terra, Università di Ferrara, Via Saragat 1, 44100 Ferrara (Italy)
psr@unife.it

² Regione Lombardia, Infrastruttura per l'Informazione Territoriale, Via Sasseti 32/2, 20124 Milano (Italy)

³ Dipartimento di Scienze Geologiche e Geotecnologie, Università di Milano-Bicocca, Piazza della Scienza 4, 20126 Milano (Italy)

Received: 17/1/08 / Accepted: 23/01/08

In their paper "Remarks on the Permian-Triassic transition in Central and Eastern Lombardy (Southern Alps, Italy)", Cassinis *et al.* (2007) propose regional correlations in the Southalpine area (Northern Italy) around the P/Tr boundary. Their reconstructions are based on the integration of paleontological, stratigraphic, and sedimentological data from published works and original analyses on the studied succession, spanning from the Early Permian (Cisuralian) to the Olenekian. The authors attempt to integrate their observations in a coherent scenario, but their conclusions are largely biased by the incorrect preconception that climatically-controlled sedimentary features should be necessarily synchronous, even if in contrast with biostratigraphic evidence.

Specifically, the occurrence of ventifacts (wind-worn pebbles, interpreted as markers of arid climate)

is not only assigned an undemonstrated chronostratigraphic significance, but even claimed to merit predominance on the occurrence of the cosmopolitan marker *Claraia* (Valsassina, Orobic Alps, Posenato *et al.*, 1996). The *Claraia* horizon of the Servino Formation, representing the Lower Triassic in the western Southern Alps (Sciunnach, 2005), is thus arbitrarily shifted upwards in the stratigraphic chart, to comply with preconceived ideas based on equivocal sedimentological evidence.

The Valsassina *Claraia* horizon is located in the middle Servino Formation about 7 m above the basal Prato Solaro Member (Posenato *et al.*, 1996; Sciunnach *et al.*, 1996, 1999), which contains the ventifacts of Cassinis *et al.* (2007). The fossiliferous horizon only yields the bivalve *Claraia*, an opportunistic crisis progenitor taxon (*sensu* Kauffman and Harries, 1996), which is characterised here, as everywhere, by a wide range of morphologi-

cal variability. The following morpho-species have been recognized: *Claraia intermedia* (Bittner), *C. radialis* (Leonardi), *C. tesidea* (Leonardi), *C. cf. bittneri* (Ichikawa), *C. cf. clarai* (Emmrich) and *C. cf. aurita* (Hauer). But for the latter, all these taxa have been classified, applying a species-population concept, into *C. intermedia* (Posenato et al., 1996), a species belonging to the *Claraia stachei* group (Ichikawa, 1958), which is characterised by well developed radial ornamentation. Specimens of this group have been reported in the *Claraia clarai* subzone of Dolomites (Broglia Loriga et al., 1983, p. 559), while *Claraia intermedia* and *C. aurita* dominate the *Claraia* beds located between the *C. clarai* and *C. aurita* subzones of the Dolomites (Leonardi, 1960). Therefore the *Claraia intermedia* horizon of Valsassina was correlated to the transitional beds between the *C. clarai* and *C. aurita* subzone (*sensu* Broglia Loriga et al., 1983; 1990) of the Dolomites (Posenato et al., 1996), which range in age from late Griesbachian to Dienerian (Broglia Loriga et al., 1983, 1990).

Cassinis et al. (2007), adducing the high morphological variability of *Claraia*, disregard its chronostratigraphical value and cite some old papers where *C. claraia* and *C. aurita* were also quoted in Smithian units of the Southern Alps (e.g. Yin, 1985). Such younger position, however, has never been demonstrated, and all recent papers report a *Claraia* range in the western Tethys (Dolomites and Hungary) restricted to the Induan Mazzin and middle Siusi members of the Werfen Formation (Broglia Loriga et al., 1983, 1986, 1990; Neri, 2007) and lower-middle Aracs Marl and Alcsutdoboz formations of the Transdanubia (Hungary) (Kozur, 1987; Broglia Loriga et al., 1990). In the Dolomites, this dating is constrained by conodont biostratigraphy (e.g. Perri, 1991), magnetostratigraphy (Scholger et al., 2000), and chemostratigraphy (Horacek et al., 2007a). The Induan age of the *Claraia* beds has been recently also re-proposed in Iran (Horacek et al., 2007b) and South China (Tong and Yin, 2002; Zhao et al., 2007), where the genus appears in the late Changhsingian (He et al., 2007). Smithian occurrences of *Claraia* in the Tethys concern the *Claraia decidens* (Bittner) group, for which the new genus *Crittendenia* has been proposed (Newell and Boyd, 1995), or species with only dense, concentric lines (e.g. *Claraia hubeiensis* Chen: Tong et al., 2006, pl. 2, figs. 11-14).

Ventifact-bearing conglomerates of the French Buntsandstein have “pratically no element for direct dating” and the “correlation between NE France and northern Germany...are well still debated”, although the Volpriehausen Formation of the Germanic basin is “the best candidate” (Durand, 2006, p. 290) to record the arid

phase which “may correlate to the late Dienerian to early Smithian interval” (Durand, 2006, p. 291). However, a younger dating for the Volpriehausen Formation is known in literature, e.g. a middle-late Smithian age has been proposed by Kozur and Bachmann (2005). In the Germanic basin, another, older arid episode is recorded in the uppermost Nordhausen Fm (now Calvörde Fm) which was correlated to the Andraz Horizon (Werfen Formation) of the Dolomites (Kozur, 1989), a prevailing evaporitic unit located between the *Claraia wangi-griesbachi* and *C. clarai* subzones (Broglia Loriga et al., 1990). The Prato Solaro Member may thus be correlated with the Andraz Horizon, consistently with the correlation between the *Claraia* beds of the Valsassina and the transitional *Claraia claraia* - *C. aurita* beds (Siusi Member) of the Dolomites proposed by us (Posenato et al., 1996). This part of the Siusi Member records the MFS of the second Scythian sequence (Sc 2) of the Dolomites (Gianolla et al., 1998; Neri, 2007).

In summary, we consider the hypothesis that ventifacts should necessarily occur in a single interval during the Early Triassic, as proposed by Cassinis et al. (2007), as unjustifiable and biased by an unscientific approach. Incidentally, we note that because of incomplete fossil record of central Europe continental successions, and consequently uncertain chronostratigraphic framework, biostratigraphy, magnetostratigraphy, and sequence stratigraphy of continental Buntsandstein (e.g. Kozur, 1993; Kozur and Bachmann, 2005; Szurlies, 2007) were calibrated by marine Lower Triassic successions (e.g. *Claraia* beds of Western Tethys).

References

- Broglia Loriga, C., Masetti, D., Neri, C. (1983): La Formazione di Werfen (Scitico) delle Dolomiti occidentali. *Rivista Italiana di Paleontologia e Stratigrafia*, 88(4): 501-598.
- Broglia Loriga, C., Neri, C., Posenato, R. (1986): The Lower Triassic in the Dolomites and Cadore. In: Italian IGCP Proj. no. 203 (ed.), *Permian and Permian-Triassic boundary in the South-Alpine segment of the Western Tethys*, Field guidebook, 29-34.
- Broglia Loriga, C., Goczan, F., Haas, J., Lenner, K., Neri, C., Oravec Sheffer, A., Posenato, R., Szabo, I., Toth Makk, A. (1990): The Lower Triassic sequences of the Dolomites (Italy) and Transdanubian Mid-Mountains (Hungary) and their correlation. *Memorie di Scienze Geologiche, Università di Padova*, 42: 41-103.
- Cassinis, G., Durand, M., Ronchi, A. (2007): Remarks on the Permian-Triassic transition in Central and Eastern Lombardy (Southern Alps, Italy). *Journal of Iberian Geology*, 33(2): 143-162.

- Durand, M. (2006): The problem of the transition from the Permian to the Triassic Series in southeastern France: comparison with other Peritethyan regions. In: Lucas, S.G., Cassinis, G., Schneider, J.W. (eds.), *Non-marine Permian Biostratigraphy and Biochronology*. Geological Society, London, *Special Publications*, 265: 281-296.
- Gianolla, P., De Zanche, V., Mietto, P. (1998): Triassic sequence stratigraphy in the southern Alps (Northern Italy): definition of the sequences and basin evolution. In: de Graciansky, P.C., Hardenbol, J., Jacquin, T., Vail, P. (eds.), *Mesozoic and Cenozoic sequence stratigraphy of European basins*, *SEPM Special Publications*, 60: 719-747.
- He, W., Feng, Q., Weldon, E., Gu, S., Meng, Y., Zhang, F., Wu, S. (2007): A Late Permian to Early Triassic bivalve fauna from the Dongpan section, southern Guangxi, South China. *Journal of Paleontology*, 81(5): 1009-1019.
- Horacek, M., Brandner, R., Abart, R. (2007a): Carbon isotope record of the P/T boundary and the Lower Triassic in the Southern Alps: Evidences for rapid changes in storage of organic carbon. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 252: 347-354.
- Horacek, M., Richo, S., Brandner, R., Krystyn, L., Spötl, C. (2007b): Evidences for recurrent changes in Lower Triassic oceanic circulation of the Tethys: The $\delta^{13}\text{C}$ record from marine sections in Iran. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 252: 355-369.
- Ichikawa, K. (1958): Zur Taxonomie und Phylogenie der Triadischen "Pteriidae" (Lamellibranch.). *Palaeontographica, Abt. A*, 111(5/6): 131-212.
- Kauffman, E.G., Harries, P.J. (1996): The importance of crisis progenitors in recovery from mass extinction. In: Hart M.B. (ed.), *Biotic Recovery from Mass Extinction Events*, Geological Society, London, *Special Publications*, 102: 15-39.
- Kozur, H.W. (1987): Stratigraphisch wichtige Fossilfunde in der Werfen-Gruppe des Balaton-Hochlandes (Vorläufige Mitteilung). *Geologisch-Paläontologische Mitteilungen Innsbruck*, 14(7): 157-167.
- Kozur, H.W. (1989): The Permian-Triassic boundary in marine and continental sediments. *Zentralblatt für Geologie und Paläontologie*, Teil 1 (1988), 11/12: 1245-1277.
- Kozur, H.W. (1993): Range charts of conchostracans in the Germanic Buntsandstein. In: Lucas, S.G., Morales, M. (eds.), *The Nonmarine Triassic*, *New Mexico Museum of Natural History & Science Bulletin*, 3: 249-253.
- Kozur, H.W., Bachmann, G. H. (2005): Correlation of the Germanic Triassic with the international scale. *Albertiana*, 32: 21-35.
- Leonardi, P. (1960): Studio statistico-sedimentologico di alcune faune werfeniane della Valle di Fiemme nel Trentino. *Studi Trentini di Scienze Naturali*, 37(1): 17-29.
- Neri, C. (2007): Formazione di Werfen. In: Cita M.B. et al. (eds.), *Carta Geologica d'Italia 1:50000. Catalogo delle formazioni-unità tradizionali (2)*. *Quaderni del Servizio Geologico d'Italia*, ser. 3, 7/7, 83-96.
- Newell, N.D., Boyd, D.W. (1995): Pectinoid bivalves of the Permian-Triassic crisis. *Bulletin of the American Museum of Natural History*, 227: 95 pp.
- Perri, M.C. (1991): Conodont biostratigraphy of the Werfen Formation (Lower Triassic), Southern Alps, Italy. *Bollettino della Società Paleontologica Italiana*, 30(1): 23-46.
- Posenato, R., Sciunnach, D., Garzanti, E. (1996): First report of *Claraia* (Bivalvia) in the Servino Formation (Lower Triassic) of the Western Orobic Alps, Italy. *Rivista Italiana di Paleontologia e Stratigrafia*, 102(2): 201-210.
- Scholger, R., Mauritsch, H.J., Brandner, R. (2000): Permian-Triassic boundary magnetostratigraphy from the Southern Alps (Italy). *Earth and Planetary Science Letters*, 176: 495-508.
- Sciunnach, D. (2005): Servino. In: Cita M.B. et al. (eds.), *Carta Geologica d'Italia 1:50000. Catalogo delle formazioni - unità tradizionali*. *Quaderni del Servizio Geologico d'Italia*, ser. 3, 7/6, 33-41.
- Sciunnach, D., Garzanti, E., Confalonieri, M. (1996): Stratigraphy and petrography of Upper Permian to Anisian terrigenous wedges (Verrucano Lombardo, Servino and Bellano Formations; western Southern Alps). *Rivista Italiana di Paleontologia e Stratigrafia*, 102(1): 27-48.
- Sciunnach, D., Garzanti, E., Posenato, R., Rodeghiero, F. (1999): Stratigraphy of the Servino Formation (Lombardy Southern Alps): towards a refined correlation with the Werfen Formation of the Dolomites. *Memorie di Scienze Geologiche, Università di Padova*, 51(1): 103-118.
- Szurliès, M. (2007): Latest Permian to Middle Triassic cyclo-magnetostratigraphy from the Central European Basin, Germany: Implications for the geomagnetic polarity timescales. *Earth and Planetary Science Letters*, 261: 602-619.
- Tong, J., Yin, H.F. (2002): The Lower Triassic in South China. *Journal of Asian Earth Sciences*, 20: 803-815.
- Tong, J., Wu, S., Li, Z., Guo, G., Zhang, J. (2006): Lower Triassic bivalves from Chaohu, Anhui Province, China. *Albertiana* 34: 42-51.
- Yin, H.F. (1985): Bivalves near the Permian-Triassic boundary in South-China. *Journal of Paleontology*, 59(3): 572-600.
- Zhao, L., Orchard, M.J., Tong, J., Sun, Z., Zuo, J., Zhang, S., Yun, A. (2007): Lower Triassic conodont sequence in Chaohu, Anhui Province, China and its global correlation. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 252: 24-38.