

REVISTA MATEMÁTICA COMPLUTENSE

Volumen 11, número 1: 1998

[http://dx.doi.org/10.5209/rev\\_REMA.1998.v11.n1.17291](http://dx.doi.org/10.5209/rev_REMA.1998.v11.n1.17291)

## **P-convexity of Musielak-Orlicz function spaces of Bochner type.**

Paweł KOLWICZ\* and Ryszard PŁUCIENNIK\*\*

### **Abstract**

It is proved that the Musielak-Orlicz function space  $L_{\Phi}(\mu, X)$  of Bochner type is  $P$ -convex if and only if both spaces  $L_{\Phi}(\mu, R)$  and  $X$  are  $P$ -convex. In particular, the Lebesgue-Bochner space  $L^p(\mu, X)$  is  $P$ -convex iff  $X$  is  $P$ -convex.

## **1 Introduction**

Relationships between various kinds of convexities of Banach spaces and the reflexivity of them were developed by many authors. The earliest result concerning that problem was obtained by D. Milman in 1938 (see [17]). Milman proved that every uniformly convex Banach space is reflexive. D. Giesy [3] and R.C. James [9] raised the question whether Banach spaces which are uniformly non- $l_n^1$  with some positive integer  $n \geq 2$  (i.e.  $B$ -convex spaces) are reflexive. Although there were some affirmative results in particular cases, the answer in general case was negative [10]. In 1970 C.A. Kottman [14] introduced a slightly stronger than  $B$ -convexity geometric property implying reflexivity and called it  $P$ -convexity. Ye Yining, He Miaohong and R. Płuciennik [22] proved that for Orlicz spaces reflexivity is equivalent to  $P$ -convexity. The same result for Musielak-Orlicz sequence and function spaces were obtained by Ye Yining and Huang Yafeng [23] and by P. Kolwicz and R. Płuciennik [11] respectively.

---

1991 Mathematics Subject Classification: 46E30, 46E40, 46B20.

\* Supported by Foundation for Polish Science, scholarship'97.

\*\* Supported by KBN grant 2 PO3A 031 10.

Servicio Publicaciones Univ. Complutense. Madrid, 1998.

In this paper we consider Musielak-Orlicz function spaces of Bochner type  $L_\Phi(\mu, X)$ . The question of whether or not a geometrical property is inherited from  $X$  into  $L_\Phi(\mu, X)$  is one of the fundamental problems here. Considerations of that type for various kinds convexities for  $L^p(\mu, X)$  were done by many authors ( see for instance [4], [5], [15], [16], [19], [20], [21]). In [12] it is proved that the Orlicz-Bochner function space  $L_\Phi(\mu, X)$  is  $P$ -convex iff both  $L_\Phi(\mu)$  and  $X$  are  $P$ -convex. We showed that the same fact is true in the case of Musielak-Orlicz sequence spaces of Bochner type (see [13]). Although such result was expected, the proof turned out to be nontrivial and essentially different from the previous one in the case of Orlicz-Bochner spaces. Moreover, our result presented below and the main result from [13] have some interesting consequences. For example, using this result in the case  $X = \mathcal{R}$ , we get immediately the characterization of  $P$ -convexity of Musielak-Orlicz spaces of real valued functions and sequences. Such characterization was proved in a very long and complicated way in the paper [11] and [23]. It is worth to mention that some similar criteria for  $B$ -convexity of Musielak-Orlicz spaces of Bochner type were obtained in [1].

Denote by  $\mathcal{N}$ ,  $\mathcal{R}$  and  $\mathcal{R}_+$  the sets of natural, real and positive real numbers, respectively. Let  $(T, \Sigma, \mu)$  be a measure space with a  $\sigma$ -finite, complete and non-atomic measure  $\mu$ . Denote by  $L^0 = L^0(T)$  the set of all  $\mu$ -equivalence classes of real valued measurable functions defined on  $T$ .

A function  $\Phi : T \times \mathcal{R} \rightarrow [0, \infty)$  is said to be a *Musielak-Orlicz function* if  $\Phi(\cdot, u)$  is measurable for each  $u \in \mathcal{R}$ ,  $\Phi(t, u) = 0$  iff  $u = 0$  and  $\Phi(t, \cdot)$  is convex, even, not identically equal zero and  $\frac{\Phi(t, u)}{u} \rightarrow 0$  as  $u \rightarrow 0$  for  $\mu$ -a.e.  $t \in T$ . Define on  $L^0$  a convex modular  $I_\Phi$  by

$$I_\Phi(x) = \int_T \Phi(t, x(t)) d\mu$$

for every  $x \in L^0$ . By the *Musielak-Orlicz space*  $L_\Phi$  we mean

$$L_\Phi = \{x \in L^0 : I_\Phi(cx) < \infty \text{ for some } c > 0\},$$

equipped with so called *Luxemburg norm* defined as follows

$$\|x\|_\Phi = \inf \left\{ \epsilon > 0 : I_\Phi \left( \frac{x}{\epsilon} \right) \leq 1 \right\}.$$

For every Musielak-Orlicz-function  $\Phi$  we define *complementary function in the sense of Young*  $\Phi^* : T \times \mathcal{R} \rightarrow [0, \infty)$  by the formula

$$\Phi^*(t, v) = \sup_{u>0} \{u|v| - \Phi(t, u)\}$$

for every  $v \in \mathcal{R}$  and  $t \in T$ .

We say that *Musielak-Orlicz-function*  $\Phi$  satisfies the  $\Delta_2$ -condition (write  $\Phi \in \Delta_2$ ) if there exist a constant  $k > 2$  and a measurable non-negative function  $f$  such that  $I_\Phi(f) < \infty$  and

$$\Phi(t, 2u) \leq k\Phi(t, u) \tag{1}$$

for  $\mu$ -a.e.  $t \in T$  and every  $u \geq f(t)$ .

For more details we refer to [18].

Now let us define the type of spaces to be considered in this paper. For a real Banach space  $\langle X, \|\cdot\|_X \rangle$ , denote by  $L^0(T, X)$ , or just  $L^0(X)$ , the family of strongly measurable functions  $f : T \rightarrow X$  identifying functions which are equal  $\mu$ -almost everywhere in  $T$ . Define a new modular  $\tilde{I}_\Phi : L^0(X) \rightarrow [0, \infty)$  by the formula

$$\tilde{I}_\Phi(f) = \int_T \Phi(t, \|f(t)\|_X) d\mu$$

for every  $f \in L^0(X)$ . Let

$$L_\Phi(\mu, X) = \left\{ f \in L^0(X) : \|f(\cdot)\|_X \in L_\Phi \right\}.$$

Then  $L_\Phi(\mu, X)$  becomes a Banach space with the norm

$$\|f\| = \|\|f(\cdot)\|_X\|_\Phi$$

and it is called a *Musielak-Orlicz space of Bochner type*.

Linear normed space  $X$  is called *P-convex* if there exist  $\epsilon > 0$  and  $n \in \mathbb{N}$  such that for all  $x_1, x_2, \dots, x_n \in S(X)$

$$\min_{i \neq j; i, j \leq n} \|x_i - x_j\|_X \leq 2(1 - \epsilon),$$

where  $S(X)$  denotes the unit sphere of  $X$ .

The notion of  $P$ -convexity can be characterized as follows.

**Lemma 1.** *A Banach space  $X$  is  $P$ -convex iff there exist  $n_0 \in \mathbb{N}$  and  $\delta_0 > 0$  such that for any elements  $x_1, x_2, \dots, x_{n_0} \in X \setminus \{0\}$  integers  $i_0, j_0$  can be found such that*

$$\left\| \frac{x_{i_0} - x_{j_0}}{2} \right\|_X \leq \frac{\|x_{i_0}\|_X + \|x_{j_0}\|_X}{2} \left( 1 - \frac{2\delta_0 \min\{\|x_{i_0}\|_X, \|x_{j_0}\|_X\}}{\|x_{i_0}\|_X + \|x_{j_0}\|_X} \right).$$

For the proof see [12] or [13].

## 2 Auxiliary lemmas

To prove the main result, we need the following.

**Lemma 2.** *Assume that  $\Phi$  and  $\Phi^*$  satisfy the  $\Delta_2$ -condition. Then for every  $\epsilon \in (0, 1)$  there are a measurable function  $h_\epsilon : T \rightarrow \mathcal{R}_+$  with  $I_\Phi(h_\epsilon) < \epsilon$ , numbers  $a(\epsilon) \in (0, 1)$  and  $\gamma = \gamma(a(\epsilon)) \in (0, 1)$  such that for  $\mu$ -a.e.  $t \in T$  the inequality*

$$\Phi\left(t, \frac{u+v}{2}\right) \leq \frac{1-\gamma}{2} [\Phi(t, u) + \Phi(t, v)] \quad (2)$$

holds true for every  $u \geq h_\epsilon(t)$  and  $|\frac{v}{u}| < a$ .

**Proof.** Repeating the same argumentation as in the proof of Theorem 1.3 from [7] and Lemma 1.1 from [2] it can be proved that the conditions  $\Phi \in \Delta_2$  and  $\Phi^* \in \Delta_2$  imply an existence of a number  $\xi_1 > 1$  and a measurable function  $z : T \rightarrow \mathcal{R}_+$  such that  $I_\Phi(z) < \infty$  and

$$\Phi\left(t, \frac{\xi_1}{2}u\right) \leq \frac{1}{2\xi_1}\Phi(t, u)$$

for  $\mu$ -a.e.  $t \in T$ , for every  $u \geq z(t)$ .

Take an arbitrary number  $\epsilon > 0$ . We can find a number  $\lambda = \lambda_\epsilon > 0$  such that  $I_\Phi(\lambda z) < \frac{\epsilon}{2}$ . Define

$$A_k = \left\{ t \in T : \sup_{\lambda z(t) \leq u \leq z(t)} \frac{2(1 + \frac{1}{k})\Phi\left(t, \frac{1}{2}(1 + \frac{1}{k})u\right)}{\Phi(t, u)} \leq 1 \right\}.$$

It is easy to verify that  $A_i \subset A_{i+1}$  for every  $i \in \mathcal{N}$ . Since  $\frac{\Phi(t,u)}{u} \rightarrow 0$  as  $u \rightarrow 0$  for  $\mu$ -a.e.  $t \in T$ ,  $\Phi(t, \cdot)$  is not linear in a certain neighborhood of 0 for  $\mu$ -a.e.  $t \in T$ . Hence  $\mu(T \setminus \bigcup_{i=1}^{\infty} A_i) = 0$ . Then the Beppo-Levi theorem implies that there exists a number  $l = l_\epsilon \in \mathcal{N}$  such that

$$\int_{T \setminus A_l} \Phi(t, z(t)) d\mu < \frac{\epsilon}{2}.$$

Define

$$h_\epsilon(t) = \lambda z(t) \chi_{A_l}(t) + z(t) \chi_{T \setminus A_l}(t).$$

Obviously  $I_\Phi(h_\epsilon) < \epsilon$ . Moreover, denoting  $\xi = \xi(\epsilon) = \min\{\xi_1, 1 + \frac{1}{l}\}$ , we have

$$\Phi\left(t, \frac{\xi}{2}u\right) \leq \frac{1}{2\xi} \Phi(t, u)$$

for  $\mu$ -a.e.  $t \in T$ , and for all  $u \geq h_\epsilon(t)$ . Taking a number  $a = a(\epsilon) \in (0, 1)$  such that  $1 + a \leq \xi$  and putting  $\gamma = \gamma(a(\epsilon)) = \frac{a}{a+1} \in (0, 1)$ , we get

$$\begin{aligned} \Phi\left(t, \frac{1+a}{2}u\right) &\leq \frac{1}{2(1+a)} \Phi(t, u) \leq \frac{1}{2(1+a)} (\Phi(t, u) + \Phi(t, au)) = \\ &= \frac{1}{2}(1-\gamma) (\Phi(t, u) + \Phi(t, au)) \end{aligned} \tag{3}$$

for  $\mu$ -a.e.  $t \in T$  and for all  $u \geq h_\epsilon(t)$ . Furthermore, the convexity of  $\Phi(t, \cdot)$  implies that

$$\frac{2\Phi\left(t, \frac{1+a}{2}u\right)}{\Phi(t, u) + \Phi(t, au)}$$

is a non-decreasing function of  $a$  for  $\mu$ -a.e.  $t \in T$  and for all  $u \geq h_\epsilon(t)$ . Consequently, for every  $a_0 < a$  inequality (3) holds with the same  $\gamma$ . Hence we obtain the thesis.

Moreover, the following two lemmas will be useful.

**Lemma 3** (Lemma 3 in [11]). *If  $\Phi$  satisfies the  $\Delta_2$ -condition, then for every  $\alpha \in (0, 1)$  there exists a non-decreasing sequence  $(B_m^\alpha)$  of measurable sets of finite measure such that*

$$\mu\left(T \setminus \bigcup_{m=1}^{\infty} B_m^\alpha\right) = 0$$

and for every  $m \in \mathcal{N}$  a number  $k_m^\alpha > 2$  can be found such that

$$\Phi(t, 2u) \leq k_m^\alpha \Phi(t, u) \quad (4)$$

for  $\mu$ -a.e.  $t \in B_m^\alpha$  and for every  $u \geq \alpha f(t)$ , where  $f$  is from the  $\Delta_2$ -condition.

**Lemma 4** (Lemma 4 in [11]). *If  $\Phi$  satisfies the  $\Delta_2$ -condition, then for every  $\epsilon \in (0, 1)$  there exist a measurable function  $g_\epsilon : T \rightarrow \mathcal{R}_+$  and  $k_\epsilon > 2$  such that*

$$I_\Phi(g_\epsilon) < \epsilon \quad \text{and} \quad \Phi(t, 2u) \leq k_\epsilon \Phi(t, u) \quad (5)$$

for  $\mu$ -a.e.  $t \in T$ , whenever  $u \geq g_\epsilon(t)$ .

### 3 Main result

**Theorem 1.** *Let  $\Phi$  be a Musielak-Orlicz function and let  $X$  be a Banach space. Then the following statements are equivalent:*

- (a)  $L_\Phi(\mu, X)$  is  $P$ -convex.
- (b) Both  $L_\Phi$  and  $X$  are  $P$ -convex.
- (c)  $L_\Phi$  is reflexive and  $X$  is  $P$ -convex.
- (d)  $X$  is  $P$ -convex,  $\Phi \in \Delta_2$  and  $\Phi^* \in \Delta_2$ .

**Proof.** (a)  $\Rightarrow$  (b). Since the spaces  $L_\Phi$  and  $X$  are embedded isometrically into  $L_\Phi(\mu, X)$  and  $P$ -convexity is inherited by subspaces,  $L_\Phi(\mu)$  and  $X$  are  $P$ -convex.

(b)  $\Rightarrow$  (c). Every  $P$ -convex Banach space is reflexive (see Theorem 3.2 in [14]). Hence  $L_\Phi$  is reflexive.

(c)  $\Rightarrow$  (d). The reflexivity of Musielak-Orlicz function space  $L_\Phi$  is equivalent to the fact that  $\Phi \in \Delta_2$  and  $\Phi^* \in \Delta_2$  (see [8]).

(d)  $\Rightarrow$  (a). Suppose that  $\Phi \in \Delta_2$  and  $\Phi^* \in \Delta_2$  and  $X$  is  $P$ -convex. Let  $n_0$  be a natural number from Lemma 1. For every  $t \in T$  define

$$f(t) = \max \left\{ h_{\frac{1}{4n_0}}(t), g_{\frac{1}{4n_0}}(t) \right\},$$

where functions  $h_{\frac{1}{4n_0}}$  and  $g_{\frac{1}{4n_0}}$  are respectively from Lemma 2 and Lemma 4 with  $\epsilon = \frac{1}{4n_0}$ . Hence  $I_\Phi(f) < \frac{1}{2n_0}$ . Put in Lemma 3  $\alpha = a$ ,

where  $a$  is from Lemma 2. We have  $I_{\Phi} \left( \frac{1}{a} f \right) < \infty$ , because  $\Phi \in \Delta_2$ . Take a set  $B_{m_0}^a$  from Lemma 3 with  $m_0$  large enough to satisfy

$$\int_{T \setminus B_{m_0}^a} \Phi \left( t, \frac{f(t)}{a} \right) d\mu < \frac{1}{2n_0}. \quad (6)$$

Let  $l \in \mathcal{N}$  be such that  $\frac{1}{a} \leq 2^l$ . Then, by Lemma 3, there exists a number  $k_{m_0}^a > 2$  such that

$$\Phi \left( t, \frac{1}{a} v \right) \leq (k_{m_0}^a)^l \Phi(t, v)$$

for  $\mu$ -a.e.  $t \in B_{m_0}^a$ , whenever  $v \geq af(t)$ . Putting

$$\frac{v}{a} = u \text{ and } \frac{1}{(k_{m_0}^a)^l} = \beta(a, m_0) = \beta_{m_0},$$

we get

$$\Phi(t, au) \geq \beta_{m_0} \Phi(t, u) \quad (7)$$

for  $\mu$ -a.e.  $t \in B_{m_0}^a$  and for every  $u \geq f(t)$ . Moreover, by Lemma 4, we have

$$\Phi \left( t, \frac{1}{a} v \right) \leq (k_{\epsilon})^l \Phi(t, v)$$

for  $\mu$ -a.e.  $t \in T$ , and  $v \geq f(t)$ , where  $k_{\epsilon} = k_{\frac{1}{4n_0}}$ . Analogously, we can obtain

$$\Phi(t, au) \geq \beta \Phi(t, u) \quad (8)$$

for  $\mu$ -a.e.  $t \in T$ , and for every  $u \geq \frac{f(t)}{a}$ .

Now, we will show that there exists a number  $r_1 \in (0, 1)$  such that for any elements  $x_1, x_2, \dots, x_{n_0}$  of Banach space  $X$  and for  $\mu$ -a.e.  $t \in T_M$  we have

$$\sum_{i=1}^{n_0} \sum_{j=i}^{n_0} \Phi \left( t, \left\| \frac{x_i - x_j}{2} \right\|_X \right) \leq \frac{n_0 - 1}{2} r_1 \sum_{i=1}^{n_0} \Phi(t, \|x_i\|_X), \quad (9)$$

where  $T_M = \left\{ t \in T : \max_{1 \leq i \leq n_0} \{ \|x_i\|_X \} \geq \frac{f(t)}{a} \right\}$ .

Take  $x_1, x_2, \dots, x_{n_0} \in X$ . Let  $k$  be an index such that  $\|x_k\|_X = \max_{1 \leq i \leq n_0} \{\|x_i\|_X\}$ . For the clarity of the proof, we will divide it into two parts.

I. Suppose that there is  $i_1 \in \{1, 2, \dots, n_0\} \setminus \{k\}$  such that

$$\frac{\|x_{i_1}\|_X}{\|x_k\|_X} < a.$$

Since  $\|x_k\| \geq \frac{f(t)}{a} > f(t)$  for  $\mu$ -a.e.  $t \in T_M$ , by inequality (2), we obtain

$$\begin{aligned} \Phi\left(t, \left\| \frac{x_{i_1} - x_k}{2} \right\|_X\right) &\leq \Phi\left(t, \frac{\|x_{i_1}\|_X + \|x_k\|_X}{2}\right) \\ &\leq \frac{1}{2}(1 - \gamma) (\Phi(t, \|x_{i_1}\|_X) + \Phi(t, \|x_k\|_X)). \end{aligned}$$

Hence, by the convexity of  $\Phi(t, \cdot)$  for  $\mu$ -a.e.  $t \in T$ , we get

$$\begin{aligned} &\sum_{i=1}^{n_0} \sum_{j=i}^{n_0} \Phi\left(t, \left\| \frac{x_i - x_j}{2} \right\|_X\right) \leq \\ &\leq \frac{n_0 - 1}{2} \sum_{i=1}^{n_0} \Phi(t, \|x_i\|_X) - \frac{\gamma}{2} (\Phi(t, \|x_{i_1}\|_X) + \Phi(t, \|x_k\|_X)) \\ &\leq \frac{n_0 - 1}{2} \sum_{i=1}^{n_0} \Phi(t, \|x_i\|_X) - \frac{\gamma}{2n_0} (n_0 \Phi(t, \|x_k\|_X)) \\ &\leq \frac{n_0 - 1}{2} \sum_{i=1}^{n_0} \Phi(t, \|x_i\|_X) - \frac{\gamma}{2n_0} \sum_{i=1}^{n_0} \Phi(t, \|x_i\|_X) \\ &= \frac{n_0 - 1}{2} \left(1 - \frac{\gamma}{n_0(n_0 - 1)}\right) \sum_{i=1}^{n_0} \Phi(t, \|x_i\|_X) \end{aligned} \quad (10)$$

for  $\mu$ -a.e.  $t \in T_M$ .

II. Assume that for all  $i \neq k$  we have

$$\frac{\|x_i\|_X}{\|x_k\|_X} \geq a. \quad (11)$$



Then  $\|x_i\| > 0$  for every  $i \neq k$ . Let  $i_0, j_0$  be from Lemma 1. We may assume that

$$a < \frac{\|x_{i_0}\|_X}{\|x_{j_0}\|_X} \leq \frac{1}{a}. \tag{12}$$

Really, otherwise we have

$$a > \frac{\|x_{i_0}\|_X}{\|x_{j_0}\|_X} \geq \frac{\min\{\|x_{i_0}\|_X, \|x_{j_0}\|_X\}}{\max\{\|x_{i_0}\|_X, \|x_{j_0}\|_X\}} \geq \frac{\min\{\|x_{i_0}\|_X, \|x_{j_0}\|_X\}}{\|x_k\|_X},$$

which contradicts to inequality (11). Hence, applying Lemma 1 and inequality (12), we get

$$\left\| \frac{x_{i_0} - x_{j_0}}{2} \right\|_X \leq \left( 1 - \frac{2\delta a}{1+a} \right) \frac{\|x_{i_0}\|_X + \|x_{j_0}\|_X}{2}.$$

Therefore, by the convexity of  $\Phi(t, \cdot)$  for  $\mu$ -a.e.  $t \in T$ , we obtain

$$\Phi \left( t, \left\| \frac{x_{i_0} - x_{j_0}}{2} \right\|_X \right) \leq \frac{1}{2}(1 - \alpha) (\Phi(t, \|x_{i_0}\|_X) + \Phi(t, \|x_{j_0}\|_X)), \tag{13}$$

where  $\alpha = \frac{2\delta a}{1+a} \in (0, 1)$ . Consequently, by inequalities (13) and (8), we have

$$\begin{aligned} & \sum_{i=1}^{n_0} \sum_{j=i}^{n_0} \Phi \left( t, \left\| \frac{x_i - x_j}{2} \right\|_X \right) \\ & \leq \frac{n_0 - 1}{2} \sum_{i=1}^{n_0} \Phi(t, \|x_i\|_X) - \frac{\alpha}{2} (\Phi(t, \|x_{i_0}\|_X) + \Phi(t, \|x_{j_0}\|_X)) \\ & \leq \frac{n_0 - 1}{2} \sum_{i=1}^{n_0} \Phi(t, \|x_i\|_X) - \alpha \Phi(t, a \|x_k\|_X) \\ & \leq \frac{n_0 - 1}{2} \sum_{i=1}^{n_0} \Phi(\|x_i\|_X) - \frac{\alpha\beta}{n_0} (n_0 \Phi(\|x_k\|_X)) \\ & \leq \frac{n_0 - 1}{2} \sum_{i=1}^{n_0} \Phi(t, \|x_i\|_X) - \frac{\alpha\beta}{n_0} \sum_{i=1}^{n_0} \Phi(t, \|x_i\|_X) \\ & = \frac{n_0 - 1}{2} \left( 1 - \frac{2\alpha\beta}{n_0(n_0 - 1)} \right) \sum_{i=1}^{n_0} \Phi(t, \|x_i\|_X) \end{aligned} \tag{14}$$

for  $\mu$ -a.e.  $t \in T_M$ . Define

$$r_1 = \max \left\{ 1 - \frac{\gamma}{n_0(n_0 - 1)}, 1 - \frac{2\alpha\beta}{n_0(n_0 - 1)} \right\}.$$

Combining inequalities (10) and (14), we get inequality (9). Repeating the same argumentation as in the proof of inequality (9), a number  $r_2 \in (0, 1)$  can be found such that the inequality

$$\sum_{i=1}^{n_0} \sum_{j=i}^{n_0} \Phi \left( t, \left\| \frac{x_i - x_j}{2} \right\|_X \right) \leq \frac{n_0 - 1}{2} r_2 \sum_{i=1}^{n_0} \Phi(t, \|x_i\|_X) \quad (15)$$

holds for every  $x_1, x_2, \dots, x_{n_0}$  elements from Banach space  $X$  and for  $\mu$ -a.e.  $t \in B_{m_0}^a$  satisfying  $\max_{1 \leq i \leq n_0} \{\|x_i\|_X\} \geq f(t)$ . Using (7) in place of (8) one can find that (15) is true with

$$r_2 = \max \left\{ 1 - \frac{\gamma}{n_0(n_0 - 1)}, 1 - \frac{2\alpha\beta m_0}{n_0(n_0 - 1)} \right\}.$$

Let  $f_1, \dots, f_{n_0} \in S(L^\Phi(\mu, X))$ . Define

$$E = \left\{ t \in T : \sum_{i=1}^{n_0} \Phi(t, \|f_i(t)\|_X) \geq n_0 \Phi(t, f(t)) \right\}.$$

Obviously  $\max_{1 \leq i \leq n_0} \{\|f_i(t)\|_X\} \geq f(t)$  for every  $t \in E$ . Divide the set  $E$  into two following subsets:

$$E_1 = \left\{ t \in E : \max_{1 \leq i \leq n_0} \{\|f_i(t)\|_X\} \geq \frac{f(t)}{a} \right\},$$

$$E_2 = \left\{ t \in E : f(t) \leq \max_{1 \leq i \leq n_0} \{\|f_i(t)\|_X\} < \frac{f(t)}{a} \right\}.$$

Next decompose the set  $E_2$  into two subsets  $E_{21}$  and  $E_{22}$  defined by

$$E_{21} = E_2 \cap B_{m_0}^a,$$

$$E_{22} = E_2 \setminus B_{m_0}^a.$$

By inequalities (9) and (15), we have

$$\sum_{i=1}^{n_0} \sum_{j=i}^{n_0} \Phi \left( t, \left\| \frac{f_i(t) - f_j(t)}{2} \right\|_X \right) \leq \frac{1}{n_0} \binom{n_0}{2} \sum_{i=1}^{n_0} \Phi(t, \|f_i(t)\|_X) \quad (16)$$

for  $\mu$ -a.e.  $t \in E_1 \cup E_{21}$ , where  $r = \max\{r_1, r_2\}$ . Obviously  $r \in (0, 1)$ . Moreover, by the definitions of the set  $E$  and the function  $f$ , we have

$$\sum_{i=1}^{n_0} \tilde{I}_\Phi (f_i \chi_{T \setminus E}) < \frac{1}{2}. \quad (17)$$

Now, let  $t \in E_{22}$ . Then, by inequality (6), we have

$$\begin{aligned} \sum_{i=1}^{n_0} \tilde{I}_\Phi (f_i \chi_{E_{22}}) &= \sum_{i=1}^{n_0} \int_{E_2 \setminus B_{m_0}^a} \Phi(t, \|f_i(t)\|_X) d\mu \\ &\leq \int_{E_2 \setminus B_{m_0}^a} n_0 \Phi \left( t, \max_{1 \leq i \leq n_0} \{ \|f_i(t)\|_X \} \right) d\mu \\ &< \int_{E_2 \setminus B_{m_0}^a} n_0 \Phi \left( t, \frac{f(t)}{a} \right) d\mu \leq \int_{T \setminus B_{m_0}^a} n_0 \Phi \left( t, \frac{f(t)}{a} \right) d\mu < \frac{1}{2}. \end{aligned} \quad (18)$$

Hence, by inequalities (17) and (18), we get

$$\sum_{i=1}^{n_0} \tilde{I}_\Phi (f_i \chi_{T \setminus (E_1 \cup E_{21})}) = \sum_{i=1}^{n_0} \tilde{I}_\Phi (f_i \chi_{T \setminus E}) + \sum_{i=1}^{n_0} \tilde{I}_\Phi (f_i \chi_{E_{22}}) < 1.$$

Since  $\|f_i\| = 1$  for  $i = 1, 2, \dots, n_0$  and  $\Phi \in \Delta_2$ , we have  $\tilde{I}_\Phi (f_i) = 1$  for  $i = 1, 2, \dots, n_0$ . Consequently,

$$\sum_{i=1}^{n_0} \tilde{I}_\Phi (f_i \chi_{E_1 \cup E_{21}}) \geq n_0 - 1. \quad (19)$$

Therefore, by inequalities (16) and (19), we have

$$\sum_{i=1}^{n_0} \sum_{j=i}^{n_0} \tilde{I}_\Phi \left( \frac{1}{2} (f_i - f_j) \right) =$$

$$\begin{aligned}
&= \sum_{i=1}^{n_0} \sum_{j=i}^{n_0} \tilde{I}_\Phi \left( \frac{1}{2} (f_i - f_j) \chi_{T \setminus (E_1 \cup E_{21})} \right) + \sum_{i=1}^{n_0} \sum_{j=i}^{n_0} \tilde{I}_\Phi \left( \frac{1}{2} (f_i - f_j) \chi_{E_1 \cup E_{21}} \right) \\
&\leq \frac{1}{n_0} \binom{n_0}{2} \sum_{i=1}^{n_0} \tilde{I}_\Phi (f_i \chi_{T \setminus (E_1 \cup E_{21})}) + \frac{r}{n_0} \binom{n_0}{2} \sum_{i=1}^{n_0} \tilde{I}_\Phi (f_i \chi_{E_1 \cup E_{21}}) \\
&= \frac{1}{n_0} \binom{n_0}{2} \left( \sum_{i=1}^{n_0} \tilde{I}_\Phi (f_i) - \sum_{i=1}^{n_0} \tilde{I}_\Phi (f_i \chi_{E_1 \cup E_{21}}) \right) \\
&\quad + \frac{r}{n_0} \binom{n_0}{2} \sum_{i=1}^{n_0} \tilde{I}_\Phi (f_i \chi_{E_1 \cup E_{21}}) \\
&= \binom{n_0}{2} \left( 1 - \frac{1-r}{n_0} \sum_{i=1}^{n_0} \tilde{I}_\Phi (f_i \chi_{E_1 \cup E_{21}}) \right) \\
&\leq \binom{n_0}{2} \left( 1 - \frac{(1-r)(n_0-1)}{n_0} \right) \leq \binom{n_0}{2} (1-p),
\end{aligned}$$

where  $p = \frac{1-r}{2}$ . So, there exist  $i_1, j_1 \in \{1, 2, \dots, n_0\}$  such that

$$\tilde{I}_\Phi \left( \frac{1}{2} (f_{i_1} - f_{j_1}) \right) \leq 1-p.$$

Finally, by the  $\Delta_2$ -condition for  $\Phi$ , we obtain that

$$\left\| \frac{1}{2} (f_{i_1} - f_{j_1}) \right\| \leq 1 - q(p), \quad 0 < q(p) < 1 \text{ (cf. [6])}.$$

Thus, by Lemma 1, the space  $L_\Phi(\mu, X)$  is  $P$ -convex. This completes the proof.

Theorem 1 is a generalization of Theorem 1 from [12]. Moreover, the following characterization of  $P$ -convex Musielak Orlicz spaces of real functions  $L_\Phi$ , proved directly in [11] in a very complicated way, is an immediate consequence of Theorem 1.

**Corollary 1.** *The following statements are equivalent:*

- (a)  $L_\Phi$  is  $P$ -convex.
- (b)  $L_\Phi$  is reflexive.
- (c)  $\Phi \in \delta_2$  and  $\Phi^* \in \Delta_2$ .

**Proof.** It is enough to apply Theorem 1 with  $X = \mathcal{R}$ .

**Corollary 2.** *The Lebesgue-Bochner function space  $L^p(u, X)$  ( $1 < p < \infty$ ) is P-convex iff  $X$  is P-convex.*

**Proof.** The Lebesgue space  $L^p$  is a Musielak-Orlicz space generated by the Orlicz function  $\Phi(t, u) = |u|^p$  for every  $t \in T$  satisfying all the assumptions of Theorem 1.

## References

- [1] H. Algherik and H. Hudzik, *Uniformly non- $l_n^1$  Musielak-Orlicz spaces of Bochner type*, Forum Math. 1 (1989), 403-410.
- [2] S. Chen and H. Hudzik, *On some convexities of Orlicz and Orlicz-Bochner spaces*, Comm. Math. Univ. Carolinae, 29,1 (1988), 13-29.
- [3] D.P. Giesy, *On a convexity condition in normed linear spaces*. Trans. Amer. Math. Soc. 125 (1966), 114-146.
- [4] P. Greim, *A note on strong extreme and strongly exposed points in Bochner  $L^p$  spaces*, Proc. Amer. Math. Soc. 93 (1985), 65-66.
- [5] P. Greim, *Strongly exposed points in Bochner  $L^p$  spaces*, Proc. Amer. Math. Soc. 88 (1983), 81-84.
- [6] H. Hudzik, *Uniform convexity of Musielak-Orlicz spaces with Luxemburg norm*, Commentationes Math., 23 (1983), 21-32.
- [7] H. Hudzik, A. Kaminska, W. Kurc, *Uniformly non- $l_n^{(1)}$  Musielak-Orlicz spaces*, Bull. Acad. Polon. Sci. Math. 35, 7-8 (1987), 441-448.
- [8] H. Hudzik and A. Kamińska, *On uniformly convexifiable and B-convex Musielak-Orlicz spaces*, Commentat. Math. 25 (1985), 59-75.
- [9] R.C. James, *Uniformly non-square Banach spaces*, Ann. of Math. 2 80 (1964), 542-550.
- [10] R.C. James, *A nonreflexive Banach space that is uniformly nonoctahedral*, Israel J. Math. 18 (1974), 145-155.

- [11] P. Kolwicz and R. Pluciennik, *On  $P$ -convex Musielak-Orlicz spaces*, Comment. Math. Univ. Carolinae **36**, 4 (1995), 655-672.
- [12] P. Kolwicz and R. Pluciennik,  *$P$ -convexity of Bochner-Orlicz spaces*, Proc. Amer. Math. Soc. –to appear.
- [13] P. Kolwicz and R. Pluciennik,  *$P$ -convexity of Musielak-Orlicz Sequence Spaces of Bochner Type*, Collect. Math. (1997) –to appear.
- [14] C.A. Kottman, *Packing and reflexivity in Banach spaces*, Trans. Amer. Math. Soc. **150** (1970), 565-576.
- [15] B.L. Lin and P.K. Lin, *Denting points in Bochner  $L^p$  spaces*, Proc. Amer. Math. Soc. **97.4** (1986), 629-633.
- [16] B.L. Lin, P.K. Lin and S.L. Troyanski, *Property (H) in Lebesgue-Bochner function spaces*, Proc. Amer. Math. Soc. **95.4** (1985), 581-584.
- [17] D. Milman, *On some criteria for the regularity of spaces of type (B)*, (Russian) Doklady Akad. Nauk SSSR **20** (1938), 243-246.
- [18] J. Musielak, *Orlicz spaces and modular spaces*. Lecture Notes in Math. **1034** (1983), 1-222.
- [19] M.A. Smith, *Strongly extreme points in  $L^p(\mu, X)$* , Rocky Mnt. J. Math. **16** (1986), 1-5.
- [20] M.A. Smith and B. Turett, *Rotundity in Lebesgue-Bochner function spaces*, Trans. Amer. Math. Soc. **257** (1980), 105-118.
- [21] K. Sundaresan, *Extreme points of the unit cell in Lebesgue-Bochner function spaces*, Colloq. Math. **22** (1970), 111-119.
- [22] Ye Yining, He Miaohong and R. Pluciennik,  *$P$ -convexity and reflexivity of Orlicz spaces*, Comment. Math. **31** (1991), 203-216.
- [23] Ye Yining and Huang Yafeng,  *$P$ -convexity property in Musielak-Orlicz sequence spaces*, Collect. Math. **44** (1993), 307-325.

Paweł KOLWICZ, Ryszard PLUCIENNIK  
Institute of Mathematics,  
Poznań University of Technology  
ul. Piotrowo 3a, 60-965 POZNAŃ,  
Poland

Recibido: 14 de Octubre de 1996