

Confirmed record of *Monodactylus argenteus* (Linnaeus, 1758 (Family Monodactylidae) from Jubail, Saudi Arabia, Arabian Gulf

L. A. Jawad

Jawad, L. A., 2013. Confirmed record of *Monodactylus argenteus* (Linnaeus, 1758) (Family Monodactylidae) from Jubail, Saudi Arabia, Arabian Gulf. *Arxius de Miscel·lània Zoològica*, 11: 158–162.

Abstract

Confirmed record of Monodactylus argenteus (Family Monodactylidae) from Jubail, Saudi Arabia, Arabian Gulf.— The first record of *M. argenteus* from the Arabian Gulf coasts of Saudi Arabia is confirmed, based on a sample measuring 158 mm in SL. Morphometric and meristic data are provided for this specimen.

Key words: Arabian Gulf, Saudi Arabia, New record, *Monodactylus argenteus*, Range distribution.

Resumen

Confirmación del registro de Monodactylus argenteus (familia Monodactylidae) en Jubail, Arabia Saudita, golfo Pérsico.— Se confirma el primer registro de *M. argenteus* en las costas de Arabia Saudita en el golfo Pérsico con un ejemplar de 158 mm SL. Se indican los datos morfométricos y merísticos del espécimen.

Palabras clave: Golfo Pérsico, Arabia Saudita, Nuevo registro, *Monodactylus argenteus*, Área de distribución.

Received: 4/11/13; Conditional acceptance: 20/01/14; Final acceptance: 22/01/14

Laith A. Jawad, Manukau, Auckland, New Zealand.

E-mail: laith_jawad@hotmail.com

Introduction

Publication of fish taxonomy in the Arabian Gulf started when Blegvad published his book (Blegvad & Løppenthin, 1944) on fishes in the Arabian Gulf in 1944. Since then, several works have appeared on the ichthyofauna of the gulf (White & Barwani, 1971; Kuronoma & Abe, 1972, 1986; Relyea, 1981; Fischer & Bianchi, 1984; Hussain et al., 1988; Randall, 1995).

The fish fauna of the Arabian Gulf coast of Saudi Arabia have been poorly studied and complete, extensive works are lacking. However, a few significant publications have been published from parts of the Saudi coasts of the Arabian Gulf (McCain et al., 1984; Coles & Tarr, 1990; Krupp & Anegay, 1993; Krupp & Müller, 1995; Krupp & Al-Marri, 1996).

The silver moony *M. argenteus* is reported from the Red Sea and East Africa (Heemstra, 1984) to Samoa, north to the Yaeyamas, and south to New Caledonia and Australia (Blaber, 1980). In the Arabian Gulf area, the only record of this species is that of Krupp et al. (2000).

The present paper confirms the presence of *Monodactylus argenteus* in the Saudi Arabian coast of the Arabian Gulf waters following the capture of the present specimen, described in a recent taxonomic work.

Studied material

On 28 May 2013, a single specimen of the silver moony *M. argenteus* (fig. 1) was collected at Jubail City on the Arabian Gulf coast of Saudi Arabia. The specimen was caught by fishermen using 30 x 10 m drifting gill nets of 25 mm mesh size. Its total length was measured with dial calipers to the nearest 0.1 mm and it was weighed to the nearest kilo. It was made available by M. Ibrahim of the Fish Welfare Branch, Al-Jubail, Ministry of Agriculture, Saudi Arabia. The fishermen usually make three to five hauls per hour. The morphometric and meristic details of the species were recorded according to Heemstra (1984) (table 1). The specimen was fixed in 10% formalin and later preserved in 70% ethanol for deposit in the fish collection of the Fish Welfare Branch, Ministry of Agriculture, Al-Jubail, Saudi Arabia.



Fig. 1. *Monodactylus argenteus*, 200 mm TL.

Fig. 1. *Monodactylus argenteus*, 200 mm TL.

Table 1. Morphometric and meristic characters of *Monodactylus argenteus* collected from Jubail, Saudi Arabia, Arabian Gulf: TL. Total length; HL. Head length; SL. Standard length.

Tabla 1. Caràcteres morfològics i mèrics de *Monodactylus argenteus* recollido en Jubail (costas de Arabia Saudita, golfo Pèrsico): TL. Longitud total; HL. Longitud de la cabeza; SL. Longitud estàndar.

| Morphometric characters | in mm |
|----------------------------------|-------|
| Total length (TL) | 200 |
| Standard length (% in TL) | 158 |
| Head length (% in SL) | 50 |
| Eye diameter (% in HL) | 18 |
| Preorbital length (% in HL) | 12 |
| Posorbital length (% in HL) | 30 |
| Predorsal fin length (% in SL) | 63 |
| Postdorsal fin length (% in SL) | 136 |
| Prepectoral fin length (% in SL) | 47 |
| Preanus length (% in SL) | 82 |
| Preanal fin length (% in SL) | 90 |
| Postanal fin length (% in SL) | 144 |
| Maximum body depth (% in SL) | 111 |
| Caudal peduncle depth (% in SL) | 16 |
| Pectoral fin length (% in SL) | 30 |
| Meristic characters | |
| Dorsal fin spines | 7 |
| Dorsal fin rays | 28 |
| Anal fin spines | 3 |
| Anal fin rays | 29 |
| Pectoral fin rays | 17 |

Remarks

The total length was 200 mm and it weighed 33 g. *M. argenteus* can attain lengths of 270 mm (Allen et al., 2002) and it is found in the pelagic–neritic zone of a marine, freshwater and brackish environments (Riel & Baensch, 1991). As far as the author is aware, the only specimen of *M. argenteus* given with SL are in Fish Base (Froese & Pauly, 2013) reported from the Japanese Archipelago, and given as 93 mm SL. *M. argenteus* has five synonyms, *Chaetodon argenteus* Linnaeus, 1758, *Psettus argenteus* (Linnaeus, 1758), *M. argenteus*, *Scomber rhombeus* Forsskål, 1775 and *Psettus rhombeus* (Forsskål, 1775). This species has not been recorded in the Arabian Gulf area under any of its synonyms. The importance of the present record of *M. argenteus* comes as a result of ichthyological collections in the area and the rarity of the members of the family Monodactylidae in Saudi waters. The species reported in this study has not been reported previously in the fish fauna literature from the

Arabian Gulf in general or from the Saudi coasts of the Arabian Gulf (Fischer & Bianchi, 1984; McCain et al., 1984; Coles & Tarr, 1990; Krupp & Anegay, 1993; Krupp & Müller, 1995; Krupp & Al-Marri, 1996; Carpenter et al., 1997). Krupp et al. (2000) obtained their specimen in Dubai, at a Deraa fish landing site at the entrance of Khor Dubai. The Dubai landing site is very large, receiving fish stock mainly from the Sea of Oman and southern coasts of Oman. Such practice sheds doubts on the origin of the specimen obtained by Krupp et al. (2000). It is highly likely the specimen originated somewhere in the Omani waters. Therefore, the record of Krupp et al. (2000) can not confirm the presence of this species in the Arabian Gulf waters and the present record is considered the first confirmation for the presence of this species in the gulf area, indicating a significant range extension of its previously known distribution. The reason why *M. argenteus* was not recorded before in the Saudi Arabian coasts of the Arabian Gulf might be because it is rare in the area or due to lack of adequate sampling. The use of diverse fish sampling techniques in the area, including scuba diving and underwater photography, has already ended the reputed rarity of several species (Jawad & Al-Mamry, 2009). It is premature to assess whether the present population is represented by only a few visitors of *M. argenteus*, exploring the new area or whether it is a well-established population, probably due to a lack of ichthyological expeditions and fishery surveys.

Thus, there is a need to investigate further the frequency of occurrence and to study the biological characteristics of this species to determine whether it has established a sustainable population in its new surroundings.

Acknowledgements

My sincere thanks to M. Ibrahim of the Fish Welfare Branch, Al-Jubail, Ministry of Agriculture, Saudi Arabia for presenting the fish specimen and the Fish Welfare Branch at Jubail City, Saudi Arabia for giving me the opportunity to report this species from the Saudi waters

References

- Allen, G. R., Midgley, S. H. & Allen, M., 2002. *Field guide to the freshwater fishes of Australia*. Western Australian Museum, Perth, Western Australia.
- Blaber, S. J. M., 1980. Fish of the Trinity Inlet System of North Queensland with notes on the ecology of fish faunas of tropical Indo-Pacific estuaries. *Australian Journal of Marine and Freshwater Researches*, 31: 137–146.
- Blegvad, H. & Løppenthin, B., 1944. *Fishes of the Iranian Gulf*. Einar Munksgaard, Copenhagen, 247 pp, XII pls (1999, translation into Farsi by E. Etemad and B. Mokayyer with supplement, Tehran University Publications, 1744: 26 + 416 pp, XI pls).
- Carpenter, K. E., Krupp, F., Jones, D. A. & Zajonz, U., 1997. *FAO species identification guide for fishery purposes. The living resources of Kuwait, eastern Saudi Arabia, Bahrain, Qatar, and the United Arab Emirates*. FAO, Rome.
- Coles, S. L. & Tarr, A. B., 1990. Reef fish assemblages in the western Arabian Gulf: A geographically isolated population in an extreme environment. *Bulletin of Marine Science*, 47: 696–720.
- Fischer, W., Bianchi, G. (Eds.), 1984. *FAO species identification sheets for fishery puposes. Western Indian Ocean (Fishing Area 51)*. Prepared and printed with support of the Danish International Development Agency (DANIDA). FAO, Rome, Vol.1–6.
- Froese, R. & Pauly, D. (Eds.), 2013. *FishBase*. World Wide Web electronic publication. www.fishbase.org, version (07/2013).
- Heemstra, P. C., 1984. Monodactylidae. In: *FAO species identification sheets for fishery*

- purposes: *Western Indian Ocean (Fishing Area 51)*, Vol. 3 (W. Fischer & G. Bianchi, Eds.). FAO, Rome.
- Hussain, N. A., Na'ama, A. K. & Al-Hassan, L. A. J., 1988. Annotated Check-list of the fish fauna of Khor al-Zubair, North west of the Arabian Gulf, Iraq. *Acta Ichthyologica et Piscatoria*, 18: 17–24.
- Jawad, L. A. & Al-Mamry, J. M., 2009. First record of *Antenarius coccineus* (Lesson, 1831) from Gulf of Oman and second record of *Antenarius indicus* Schultz, 1964 from the Arabian Sea coasts of Oman. *Marine Biodiversity Record*, 2 (e163): 1–3. (Published online: doi:10.1017/S1755267209990923.)
- Kuronoma, K. & Abe, Y., 1972. *Fishes of Kuwait*. KISR, Kuwait.
- 1986. *Fishes of the Arabian Gulf*. KISR, Kuwait.
- Krupp, F. & Al-Marri, M.A., 1996. Fishes and fish assemblages of the Jubail marine wildlife sanctuary. In: *A marine wildlife sanctuary for the Arabian Gulf: environmental research and conservation following the 1991 Gulf war oil spill*: 339–350 (F. Krupp, A. H. Abuzinada & I. A. Nader, Eds.). NCWCD, Riyadh and Senckenberg, Research Institute, Frankfurt a.M.
- Krupp, F., Al-Marri, M. A., Zajonz, U., Carpenter, K., Almatar, S. & Zetzsche, H., 2000. Twelve new records of fishes from the Gulf. *Fauna of Saudi Arabia*, 18: 323–325.
- Krupp, F. & Aneqay, F., 1993. Habitats and species composition in subtidal areas north of Abu Ali, Saudi Arabia. In: *The R/V MT. Mitchell Cruise in the ROPME Sea Area 2*: 174–184 (F. Frupp, Ed.). ROPME, Kuwait.
- Krupp, F. & Müller, T., 1995. The status of fish populations in the northern Arabian Gulf two years after the 1991 Gulf War oil spills. *Courier Forschungsinstitute Senckenber*, 166: 67–75.
- McCain, J. C., Tarr, A. B., Carpenter, K. E. & Cole, S. L., 1984. Marine ecology of Saudi Arabia. A survey of coral reefs and reef fishes in the northern area, Arabian Gulf, Saudi Arabia. *Fauna of Saudi Arabia*, 6: 102–126.
- Randall, J. E., 1995. *Coastal fishes of Oman*. Crawford house Publishing Pty Ltd., Bathurst, Australia.
- Relyea, K., 1981. *Inshore Fishes of the Arabian Gulf*. George Allen & Unwin, London.
- Riel, R. & Baensch, H. A., 1991. *Aquarien Atlas. Band. 1 Melle: Mergus*. Verlag Für Natur- und Heimtierkunde, Germany.
- White, A. E. & Barwani, M. A., 1971. *Common sea fishes of the Arabian Gulf and Gulf of Oman*. Dubai, Trucial States.