

Title	Reactions of organolithiums with dialkyl oxalates. A flow microreactor approach to synthesis of functionalized α -keto esters.
Author(s)	Nagaki, Aiichiro; Ichinari, Daisuke; Yoshida, Jun-ichi
Citation	Chemical communications (2013), 49(31): 3242-3244
Issue Date	2013-04-21
URL	http://hdl.handle.net/2433/187101
Right	This journal is © The Royal Society of Chemistry 2013
Type	Journal Article
Textversion	author

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

COMMUNICATION

Reactions of Organolithiums with Dialkyl Oxalates. A Flow Microreactor Approach to Synthesis of Functionalized α -Keto Esters†Aiichiro Nagaki,^a Daisuke Ichinari,^a and Jun-ichi Yoshida^{*a}

Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

DOI: 10.1039/b000000x

Reactions of functionalized aryllithiums with dialkyl oxalates were achieved using a flow microreactor to give α -keto esters with highly selectivity by virtue of fast 1:1 micromixing.

α -Keto carboxylic acids and esters play an important role in organic synthesis as useful building blocks for synthesizing a various of complex molecules such as biologically active natural products.^{1,2} Various methods for synthesizing α -keto carboxylic acids and esters have been reported so far,³ and the most commonly used method is the reaction of organometallics with dialkyl oxalates or oxalyl chloride. However, the method often suffers from low yields of the desired products because the initial products react with another molecule of organometallics to produce byproducts such as alcohols and a diketones even if one equiv of the organometallics is used (the problem of competitive consecutive reactions).⁴ Therefore, alkyl oxalyl chloride, alkyl cyanofornate, and alkyl α -oxo-1H-imidazole-1-acetate are often used to suppress such side reactions.⁵ However, these reagents are more expensive and therefore, are not suitable for large-scale synthesis. These methods using organometallics suffer from another problem, *i.e.* functional group compatibility in preparation and reactions of organometallics.⁶ Therefore, it is highly desired to develop a method for synthesizing α -keto esters bearing functional groups on a large scale. However, to the best of our knowledge, no study to solve these problems has yet been reported in the literature. We envisaged that the use of a flow microreactor system^{7,8} would be effective in solving these problems.

We have already reported that the problem of disguised chemical selectivity⁹ in competitive consecutive reactions is solved by extremely fast 1:1 micromixing based on short diffusion paths and that product selectivity close to kinetically based one is obtained.¹⁰ We have also reported that highly reactive and unstable reactive intermediates could be generated and used before they decompose by virtue of short residence times in flow microreactors¹¹ (flash chemistry¹²). For example, aryllithiums bearing electrophilic functional groups such as cyano, nitro, alkoxy carbonyl and ketone carbonyl groups were successfully

generated and used for reactions with electrophiles using flow microreactor systems.¹³

With such information in hand, we set out to investigate reactions of functionalized organometallics with dialkyl oxalates using flow microreactor systems and found that selective formation of α -keto esters was achieved.

Before studying flow microreactor reactions, we examined the reaction in a flask (50 mL round bottom glass flask with a magnetic stirrer). Thus, phenyllithium, which was generated by the halogen-lithium exchange reaction of bromobenzene with *n*-BuLi, was allowed to react with one equiv of diethyl oxalate (Figure 1). The yields of ethyl 2-oxo-2-phenylacetate (**2**, mono-addition product), 1,2-diphenylethane-1,2-dione (**3**, di-addition product), ethyl 2-hydroxy-2,2-diphenylacetate (**4**, di-addition product), and 2-hydroxy-1,2,2-triphenylethanone (**5**, tri-addition product) were determined by GC. The addition of diethyl oxalate to a stirred solution of phenyllithium at -78 °C gave **2** and **3** in low yields (Table 1). In contrast, the simultaneous addition and the reverse addition gave better results, but the yield of **2** and selectivity were not enough. The selectivity and yield decreased with an increase in the temperature, and a significant amount of **3**, and **5** were produced. It is interesting to note that di-addition product **4** was not obtained at all indicating that the ester carbonyl group of **2** is more reactive than the ketone carbonyl group presumably because of the steric effect. In addition, it should be kept in mind that the batch method cannot be applied to functionalized organolithium species because they decompose very quickly.




Fig. 1 Reaction of diethyl oxalate and phenyllithium generated from bromobenzene.

Table 1 Reaction of diethyl oxalate and phenyllithium using a conventional macro batch system

T ² (°C)	method of addition	conversion of 1 (%)	yield (%)			
			2	3	4	5
-78	addition of 1 to PhLi	59	25	21	0	1
-78	simultaneous addition of 1 and PhLi	66	56	17	0	6
-78	addition of PhLi to 1	70	63	9	0	7

^a Department of Synthetic and Biological Chemistry, Graduate School of Engineering, Kyoto University Nishikyo-ku, 615-8510 Kyoto, Japan.

Fax: +81-75-383-2725; Tel: +81-75-383-2726; E-mail: yoshida@sbchem.kyoto-u.ac.jp

† Electronic Supplementary Information (ESI) available: Experimental section. See DOI: 10.1039/b000000x/

-60	addition of PhLi to 1	71	66	13	0	3
-40	addition of PhLi to 1	81	62	15	0	5
-20	addition of PhLi to 1	83	55	18	0	9
0	addition of PhLi to 1	82	53	15	0	7
20	addition of PhLi to 1	72	43	18	0	10

In the next step, we examined the reaction using a flow microreactor system composed of two T-shaped micromixers (**M1** and **M2**) and two microtube reactors (**R1** and **R2**) as shown in Figure 2.




Fig. 2 Flow microreactor system for reaction of diethyl oxalate and phenyllithium generated from bromobenzene.

The results obtained with varying the temperature ($T^1 = T^2 = -78 \sim 20$ °C), the inner diameter of **M2**, and the flow rate. The effect of the temperature is interesting. As shown in Table 2, the yield of **2** and selectivity were much higher than those for flask reactions, and the highest yield and selectivity was obtained at -60 °C. The yield and selectivity gradually decreases with an increase in T. It is noteworthy that satisfactory yield and selectivity were obtained at -20 °C and hereafter we carried out the reactions at this temperature from a practical point of view because it difficult to achieve lower temperatures using brine, which is a common coolant in industry.

Table 2 Reaction of diethyl oxalate with phenyllithium generated from bromobenzene using a flow microreactor system. The effect of the temperature.

T^1, T^2 (°C)	t^{R1} (s)	Conversion of 1 (%)	yield (%)				
			2	3	4	5	
-78	9.42	96	87	3	0	0	
-60	9.42	100	96	1	0	0	
-40	2.35	100	96	2	0	0	
-20	2.35	98	93	5	0	0	
0	2.35	95	88	7	0	0	
20	2.35	93	82	9	0	1	

The reactions were carried out under the following conditions: The flow rates: PhBr solution (8 mL/min), BuLi solution (2 mL/min), diethyl oxalate solution (4 mL/min). The inner diameter of **M2** (250 μ m).

To examine the effect of mixing, we carried out the reactions with varying the total flow rate at **M2** because it is well known that mixing efficiency of micromixers strongly depends on the flow rate.¹⁴ As shown in Table 3, the yield of **2** and the selectivity decreased with a decrease in the flow rate, indicating the importance of mixing efficiency. The fact that the yield of **2** and selectivity decreased with an increase in the diameter of micromixer **M2** also indicates the importance of mixing. Therefore, it is reasonable to conclude that extremely fast 1 : 1 mixing is responsible for selective formation of α -ketoester **2**.

Table 3 Reaction of diethyl oxalate and phenyllithium generated from bromobenzene using a flow microreactor system. The effect of mixing.

inner diameter of M2 (μ m)	total flow rate at M2 (mL/min)	conversion of 1 (%)	yield (%)				
			2	3	4	5	
250	17.5	98	95	5	0	0	
250	14.0	98	93	5	0	0	
250	10.5	96	90	5	0	0	
250	7.0	92	80	8	0	2	
250	3.5	60	29	10	0	5	
500	14.0	88	77	17	0	2	
800	14.0	72	47	22	0	8	

The reactions were carried out at $T = -20$ °C


40

With the optimized conditions for selective synthesis of α -ketoester **2** ($T^1 = T^2 = -20$ °C, inner diameter of **M2** = 250 μ m, total flow rate at **M2** = 14.0 mL/min) in hand, we next examined reactions of various aryllithiums generated from functionalized halobenzenes. In cases of bromobenzenes *n*-BuLi was used as a lithiating agent. In cases of iodobenzenes PhLi was used as a lithiating agent. The halogen-lithium exchange was conducted with the residence times (t^{R1}) optimized for each halobenzenes.¹⁴ As shown in Table 4, dimethyl oxalate and di-*tert*-butyl oxalate in addition to diethyl oxalate could be used and the corresponding α -keto esters were obtained in good yields and selectivity. Notably, various aryllithiums bearing electrophilic functional groups such as cyano, nitro, and alkoxy groups at ortho-, meta- and para-positions reacted with diethyl oxalate to give the corresponding α -ketoesters in high yields and selectivity. In the case of aryllithiums bearing an alkoxy group, the reactions with diethyl oxalate were carried out at $T^1 = T^2 = -60$ °C in order to avoid their decomposition. Such transformations are very difficult or impossible to achieve using conventional batch reactors. Moreover, the mono-selective Br-Li exchange reaction of *p*-dibromobenzene followed by the reaction with diethyl oxalate gave the corresponding bromo-substituted α -keto ester in a good yield. Highly sterically hindered mesityllithium could be also used for the present transformation.

65

Table 4 Selective reaction of dialkyl oxalate with organometallics using flow microreactors

Halobenzene	T^1, T^2 (°C)	t^{R1} (s)	dialkyl oxalate	product	yield (%)
	-20	4.71			93
	-20	4.71			90
	-20	4.71			75
	-20	2.35			91
	-20	2.35			95
	-20	2.35			94
	-20	0.28			75
	-20	0.28			87
	-20	0.28			63
	-20	0.041			89



In conclusion, we have developed an efficient method for introduction of α -ketoester group into aromatic ring based on the generation and reaction of organometallics such as aryllithiums at temperatures that are easily accessible in industry by virtue of fast mixing in flow microreactor systems. The generation of various highly unstable functionalized aryllithiums followed by the selective formation of α -ketoesters can be achieved by using integrated flow microreactor systems. The method adds a new dimension in the selective synthesis of complex molecules having α -ketoester group.

Notes and references

- (a) E. J. Corey, D. H. Hua, B. C. Pan and S. P. Seitz, *J. Am. Chem. Soc.*, 1982, **104**, 6818; (b) T. Takahashi, T. Okano, T. Harada, K. Imamura and H. Yamada, *Synlett*, 1994, 121; (c) S. Alvarez, R. Alvarez, H. Khanwalkar, P. Germain, G. Lemaire, F. R. Barrios, H. Gronemeyer and A. R. Lera, *Bioorg. Med. Chem.*, 2009, **17**, 4345.
- (a) M. R. Angelastro, S. Mehdi, J. P. Burkhardt, N. P. Peet and P. Bey, *J. Med. Chem.*, 1990, **33**, 11; (b) V. G. Oleksandr, V. K. Pavel, V. S. Oleg, K. M. Pavel, A. Z. Olga and A. Andrey, *Synthesis*, 2011, 1633.
- (a) E. J. Corey and B. W. Erickson, *J. Org. Chem.*, 1971, **36**, 3553; (b) K. Yamada, M. Kato, M. Iyoda and Y. Hirata, *J. Chem. Soc., Chem. Commun.*, 1973, 499; (c) J. M. Photis, *Tetrahedron Lett.*, 1980, **21**, 3539; (d) A. J. L. Cooper, J. Z. Ginos and A. Meister, *Chem. Rev.*, 1983, **83**, 321; (e) J. P. Burkhardt, N. P. Peet, and P. Bey, *Tetrahedron Lett.*, 1988, **29**, 3433; (f) H. H. Wasserman and W. B. Ho, *J. Org. Chem.*, 1994, **59**, 4364; (g) L-S. Li and Y-L. Wu, *Tetrahedron Lett.*, 2002, **43**, 2427; (h) M. Ma, C. Li, L. Peng, F. Xie, X. Zhang and J. Wang, *Tetrahedron Lett.*, 2005, **46**, 3927; (i) J. Zhuang, C. Wang, F. Xie and Wanbin Zhang, *Tetrahedron*, 2009, **65**, 9797; (j) A. Raghunadh, S. B. Meruva, N. A. Kumar, G. S. Kumar, L. V. Rao and U. K. S. Kumar, *Synthesis*, 2012, 283.

- F. Hollwedel and G. Koßmehl, *Synthesis*, 1998, 1241.
- (a) J. Nimitz and H. S. Mosher, *J. Org. Chem.*, 1981, **46**, 211; (b) F. Babudri, V. Fiandanese, G. Marchese and A. Punzi, *Tetrahedron*, 1996, **52**, 13513 and references cited therein; (c) G. Guercio, S. Bacchi, A. Perboni, C. Leroi, F. Tinazzi, I. Bientinesi, M. Hourdin, M. Goodyear, S. Curti, S. Provera and Z. Cimarosti, *Org. Process Res. Dev.*, 2009, **13**, 1100.
- P. Knochel, *Handbook of Functionalized Organometallics*, Wiley-VCH, Weinheim, 2005.
- Microreactor: (a) P. Watts and S. J. Haswell, *Chem. Soc. Rev.*, 2005, **34**, 235; (b) K. Geyer, J. D. C. Codee and P. H. Seeberger, *Chem. Eur. J.*, 2006, **12**, 8434; (c) G. Whitesides, *Nature*, 2006, **442**, 368; (d) A. J. deMello, *Nature*, 2006, **442**, 394; (e) J. Kobayashi, Y. Mori and S. Kobayashi, *Chem. Asian J.*, 2006, **1**, 22; (f) M. Brivio, W. Verboom and D. N. Reinhoudt, *Lab Chip*, 2006, **6**, 329; (g) B. P. Mason, K. E. Price, J. L. Steinbacher, A. R. Bogdan and D. T. McQuade, *Chem. Rev.*, 2007, **107**, 2300; (h) B. Ahmed-Omer, J. C. Brandt and T. Wirth, *Org. Biomol. Chem.*, 2007, **5**, 733; (i) P. Watts and C. Wiles, *Chem. Commun.*, 2007, 443; (j) T. Fukuyama, M. T. Rahman, M. Sato and I. Ryu, *Synlett*, 2008, 151; (k) W. Lin, Y. Wang, S. Wang and H. Tseng, *Nano Today*, 2009, **4**, 470; (l) K. Geyer, T. Gustafsson and P. H. Seeberger, *Synlett*, 2009, 2382; (m) S. Marrea and K. F. Jensen, *Chem. Soc. Rev.*, 2010, **39**, 1183; (n) D. Webb and T. F. Jamison, *Chem. Sci.*, 2010, **1**, 675; (o) J. Yoshida, H. Kim and A. Nagaki, *ChemSusChem*, 2011, **4**, 331.
- Recent examples: (a) A. Nagaki, K. Kawamura, S. Suga, T. Ando, M. L. Sawamoto and J. Yoshida, *J. Am. Chem. Soc.*, 2004, **126**, 14702; (b) L. Ducry and D. M. Roberge, *Angew. Chem. Int. Ed.*, 2005, **44**, 7972; (c) P. He, P. Watts, F. Marken and S. J. Haswell, *Angew. Chem. Int. Ed.*, 2006, **45**, 4146; (d) K. Tanaka, S. Motomatsu, K. Koyama, S. Tanaka and K. Fukase, *Org. Lett.*, 2007, **9**, 299; (e) H. R. Sahoo, J. G. Kralj and K. F. Jensen, *Angew. Chem. Int. Ed.*, 2007, **46**, 5704; (f) C. H. Hornung, M. R. Mackley, I. R. Baxendale and S. V. Ley, *Org. Process Res. Dev.*, 2007, **11**, 399; (g) T. Fukuyama, M. Kobayashi, M. T. Rahman, N. Kamata and I. Ryu, *Org. Lett.*, 2008, **10**, 533; (h) D. L. Browne, M. Baumann, B. H. Harji, I. R. Baxendale and S. V. Ley, *Org. Lett.*, 2011, **13**, 3312; (i) C. F. Carter, H. Lange, D. Sakai, I. R. Baxendale and S. V. Ley, *Chem. Eur. J.*, 2011, **17**, 3398; (j) N. Zaborenko, M. W. Bedore, T. F. Jamison and K. F. Jensen, *Org. Process Res. Dev.*, 2011, **15**, 131; (k) Y. Tomida, A. Nagaki and J. Yoshida, *J. Am. Chem. Soc.* 2011, **133**, 3744 (l) A. C. Gutierrez and T. F. Jamison, *Org. Lett.*, 2011, **13**, 6414.
- P. Rys, *Acc. Chem. Res.*, 1976, **9**, 345.
- (a) A. Nagaki, M. Togai, S. Suga, N. Aoki, K. Mae and J. Yoshida, *J. Am. Chem. Soc.*, 2005, **127**, 11666; (b) S. Suga, A. Nagaki and J. Yoshida, *Chem. Commun.* 2003, 354 (c) A. Nagaki, N. Takabayashi, Y. Tomida and J. Yoshida, *Org. Lett.*, 2008, **10**, 3937; (d) A. Nagaki, Y. Tomida and J. Yoshida, *Macromolecules*, 2008, **41**, 6322; (e) A. Nagaki, N. Takabayashi, Y. Tomida and J. Yoshida, *Beilstein J. Org. Chem.* 2009, **5**, No16, 1-11; (f) J. Yoshida, A. Nagaki, T. Iwasaki and S. Suga, *Chem. Eng. Tech.*, 2005, **3**, 259.
- (a) H. Usutani, Y. Tomida, A. Nagaki, H. Okamoto, T. Nokami and J. Yoshida, *J. Am. Chem. Soc.*, 2007, **129**, 3047; (b) A. Nagaki, Y. Tomida, H. Usutani, H. Kim, N. Takabayashi, T. Nokami, H. Okamoto and J. Yoshida, *Chem. Asian J.*, 2007, **2**, 1513; (c) A. Nagaki, E. Takizawa and J. Yoshida, *J. Am. Chem. Soc.*, 2009, **131**, 1654; (d) A. Nagaki, A. Kenmoku, Y. Moriawaki, A. Hayashi and J. Yoshida, *Angew. Chem., Int. Ed.*, 2010, **49**, 7543; (e) Y. Tomida, A. Nagaki and J. Yoshida, *J. Am. Chem. Soc.*, 2011, **133**, 3744; (f) A. Nagaki, S. Tokuoka, S. Yamada, Y. Tomida, K. Oshiro, H. Amii and J. Yoshida, *Org. Biomol. Chem.*, 2011, **9**, 7559; (g) A. Nagaki, C. Matsuo, S. Kim, K. Saito, A. Miyazaki and J. Yoshida, *Angew. Chem. Int. Ed.* 2012, **51**, 3245.
- (a) J. Yoshida, *Flash Chemistry. Fast Organic Synthesis in Microsystems*: Wiley-Blackwell, 2008; (b) J. Yoshida, *Chem. Commun.*, 2005, 4509; (c) J. Yoshida, A. Nagaki and T. Yamada, *Chem. Eur. J.*, 2008, **14**, 7450; (d) J. Yoshida, *Chem. Record*, 2010, **10**, 332.
- (a) A. Nagaki, H. Kim and J. Yoshida, *Angew. Chem., Int. Ed.*, 2008, **47**, 7833; (b) A. Nagaki, H. Kim and J. Yoshida, *Angew. Chem., Int. Ed.*, 2009, **48**, 8063; (c) A. Nagaki, H. Kim, Y. Moriawaki, C. Matsuo and J. Yoshida, *Chem. Eur. J.*, 2010, **16**, 11167; (d) A. Nagaki, H. Kim, C. Matsuo, H. Usutani and J. Yoshida, *Org. Biomol. Chem.* 2010, **8**, 1212; (e) H. Kim, A. Nagaki and J. Yoshida, *Nature Commun.*, 2011, **2**, 264.
- W. Ehrfeld, K. Golbig, V. Hessel, H. Lowe and T. Richter, *Ind. Eng. Chem. Res.*, 1999, **38**, 1075.