

AXIOMATIZATIONS OF SIGNED DISCRETE CHOQUET INTEGRALS

MARTA CARDIN, MIGUEL COUCEIRO, SILVIO GIOVE,
AND JEAN-LUC MARICHAL

ABSTRACT. We study the so-called signed discrete Choquet integral (also called non-monotonic discrete Choquet integral) regarded as the Lovász extension of a pseudo-Boolean function which vanishes at the origin. We present axiomatizations of this generalized Choquet integral, given in terms of certain functional equations, as well as by necessary and sufficient conditions which reveal desirable properties in aggregation theory.

1. INTRODUCTION

This paper deals with the so-called “signed (discrete) Choquet integral” (also called non-monotonic Choquet integral) which naturally generalizes the Choquet integral [1]. Traditionally, the Choquet integral is defined in terms of a capacity (also called fuzzy measure [10, 11]), i.e., a set function $\mu: 2^{[n]} \rightarrow \mathbb{R}$ such that $\mu(\emptyset) = 0$ and $\mu(S) \leq \mu(T)$ whenever $S \subseteq T$. Dropping the monotonicity requirement in the definition of μ , we obtain what is referred to as a signed capacity (also called non-monotonic fuzzy measure). The signed Choquet integral is then defined exactly the same way but replacing the underlying capacity by a signed capacity. This extension has been considered by several authors, e.g., [3, 7, 8].

A convenient way to introduce the signed Choquet integral is via the notion of Lovász extension. Indeed, the signed Choquet integral can be thought of as the Lovász extension of a pseudo-Boolean function $f: \{0, 1\}^n \rightarrow \mathbb{R}$ which vanishes at the origin. Moreover, we retrieve the classical Choquet integral by further assuming that $f: \{0, 1\}^n \rightarrow \mathbb{R}$ is nondecreasing.

In this paper we consider the latter approach to the signed Choquet integral. In Section 2 we recall the basic notions and terminology concerning Choquet integrals and Lovász extensions needed throughout the paper. In Section 3 we present various characterizations of the signed Choquet integral. First, we recall the piecewise linear nature of Lovász extensions which particularizes to the signed Choquet integral (Theorem 3.1). Then we generalize Schmeidler’s axiomatization of the signed discrete Choquet integral

Date: October 21, 2010.

2010 Mathematics Subject Classification. Primary 39B22, 39B72; Secondary 26B35.

Key words and phrases. Signed discrete Choquet integral, signed capacity, Lovász extension, functional equation, comonotonic additivity, homogeneity, axiomatization.

given in terms of continuity and comonotonic additivity, showing that positive homogeneity can be replaced for continuity (Theorem 3.2). The main result of this paper, Theorem 3.3, presents a characterization of families of signed Choquet integrals in terms of necessary and sufficient conditions which:

- (1) reveal the linear nature of these generalized Choquet integrals with respect to the underlying signed capacities,
- (2) express properties of the family members defined on the standard basis of signed capacities, and
- (3) make apparent the meaningfulness with respect to interval scales of signed Choquet integrals.

We also discuss the independence of axioms given in Theorem 3.3.

Throughout this paper, the symbols \wedge and \vee denote the minimum and maximum functions, respectively.

2. CHOQUET INTEGRALS AND LOVÁSZ EXTENSIONS

A *capacity* on $[n]$ is a set function $\mu: 2^{[n]} \rightarrow \mathbb{R}$ such that $\mu(\emptyset) = 0$ and $\mu(S) \leq \mu(T)$ whenever $S \subseteq T$. A capacity μ on $[n]$ is said to be *normalized* if $\mu([n]) = 1$.

Definition 2.1. Let μ be a capacity on $[n]$ and let $\mathbf{x} \in [0, \infty]^n$. The *Choquet integral* of \mathbf{x} with respect to μ is defined by

$$C_\mu(\mathbf{x}) = \sum_{i=1}^n (\mu_i^\pi - \mu_{i+1}^\pi) x_{\pi(i)},$$

where π is a permutation on $[n]$ such that $x_{\pi(1)} \leq \dots \leq x_{\pi(n)}$ and $\mu_i^\pi = \mu(\{\pi(i), \dots, \pi(n)\})$ for $i \in [n+1]$, with the convention that $\mu_{n+1}^\pi = \mu(\emptyset)$.

The concept of Choquet integral can be formally extended to more general set functions and n -tuples of \mathbb{R}^n as follows. A *signed capacity* (or *game*) on $[n]$ is a set function $v: 2^{[n]} \rightarrow \mathbb{R}$ such that $v(\emptyset) = 0$.

Definition 2.2. Let v be a signed capacity on $[n]$ and let $\mathbf{x} \in \mathbb{R}^n$. The *signed Choquet integral* of \mathbf{x} with respect to v is defined by

$$C_v(\mathbf{x}) = \sum_{i=1}^n (v_i^\pi - v_{i+1}^\pi) x_{\pi(i)},$$

where π is a permutation on $[n]$ such that $x_{\pi(1)} \leq \dots \leq x_{\pi(n)}$ and $v_i^\pi = v(\{\pi(i), \dots, \pi(n)\})$ for $i \in [n+1]$, with the convention that $v_{n+1}^\pi = v(\emptyset)$.

The more general concept of a set function $v: 2^{[n]} \rightarrow \mathbb{R}$ (without any constraint) leads to the notion of the Lovász extension of a pseudo-Boolean function, which we now briefly describe. For general background, see [4, 9].

Let S_n denote the symmetric group on $[n]$ and, for each $\pi \in S_n$, define

$$P_\pi = \{\mathbf{x} \in \mathbb{R}^n : x_{\pi(1)} \leq \dots \leq x_{\pi(n)}\}.$$

Let $v: 2^{[n]} \rightarrow \mathbb{R}$ be a set function and let $f: \{0, 1\}^n \rightarrow \mathbb{R}$ be the corresponding pseudo-Boolean function, that is, such that $f(\mathbf{1}_S) = v(S)$. The *Lovász extension* of f is the continuous function $\hat{f}: \mathbb{R}^n \rightarrow \mathbb{R}$ which is defined on each P_π as the unique affine function that coincides with f at the $n + 1$ vertices of the standard simplex $[0, 1]^n \cap P_\pi$ of $[0, 1]^n$. In fact, \hat{f} can be expressed as

$$(1) \quad \hat{f}(\mathbf{x}) = f(\mathbf{0}) + \sum_{i=1}^n (f_i^\pi - f_{i+1}^\pi) x_{\pi(i)} \quad (\mathbf{x} \in P_\pi).$$

where $f_i^\pi = f(\mathbf{1}_{\{\pi(i), \dots, \pi(n)\}}) = v(\{\pi(i), \dots, \pi(n)\})$ for $i \in [n]$ and $f_{n+1}^\pi = f(\mathbf{0})$. Thus \hat{f} is a continuous function whose restriction to each P_π is an affine function.

It follows from (1) that the Lovász extension of a pseudo-Boolean function $f: \{0, 1\}^n \rightarrow \mathbb{R}$ is a signed Choquet integral if and only if $f(\mathbf{0}) = 0$. Its restriction to $[0, \infty]^n$ is a Choquet integral if, in addition, f is nondecreasing.

It was also shown [6] that the Lovász extension \hat{f} can also be written as

$$(2) \quad \hat{f}(\mathbf{x}) = \sum_{S \subseteq [n]} m(S) \bigwedge_{i \in S} x_i \quad (\mathbf{x} \in \mathbb{R}^n),$$

where the set function $m: 2^{[n]} \rightarrow \mathbb{R}$ is the Möbius transform of v , given by $m(S) = \sum_{T \subseteq S} (-1)^{|S|-|T|} v(T)$. Thus, a signed Choquet integral has the form (2) with $m(\emptyset) = 0$.

3. AXIOMATIZATIONS OF LOVÁSZ EXTENSIONS

We have a first characterization that immediately follows from the definition of Lovász extensions.

Theorem 3.1. *A function $g: \mathbb{R}^n \rightarrow \mathbb{R}$ is a Lovász extension if and only if*

$$(3) \quad g(\lambda \mathbf{x} + (1 - \lambda) \mathbf{x}') = \lambda g(\mathbf{x}) + (1 - \lambda) g(\mathbf{x}') \quad (0 \leq \lambda \leq 1)$$

for all comonotonic vectors $\mathbf{x}, \mathbf{x}' \in \mathbb{R}^n$. The function g is a signed Choquet integral if additionally $g(\mathbf{0}) = 0$.

Proof. The condition stated in the theorem means that g is affine (since it is both convex and concave) on each P_π . Hence, it is continuous on \mathbb{R}^n and thus it is a Lovász extension. \square

The following theorem is inspired from a characterization of the Choquet integral by de Campos and Bolaños [2].

Theorem 3.2. *A function $g: \mathbb{R}^n \rightarrow \mathbb{R}$ is a Lovász extension if and only if the function $h: \mathbb{R}^n \rightarrow \mathbb{R}$, defined by $h = g - g(\mathbf{0})$,*

- (i) *is comonotonic additive.*
- (ii) *is continuous or satisfies $h(r\mathbf{x}) = rh(\mathbf{x})$ for all $r > 0$.*

The function g is a signed Choquet integral if additionally $g(\mathbf{0}) = 0$.

Proof. It is not difficult to see that the conditions are necessary. So let us prove the sufficiency. Fix $\pi \in S_n$ and $\mathbf{x} \in P_\pi$. Then we have

$$\mathbf{x} = x_{\pi(1)} \mathbf{1}_{[n]} + \sum_{i=2}^n (x_{\pi(i)} - x_{\pi(i-1)}) \mathbf{1}_{\{\pi(i), \dots, \pi(n)\}}.$$

By comonotonic additivity, we get

$$h(\mathbf{x}) = h(x_{\pi(1)} \mathbf{1}_{[n]}) + \sum_{i=2}^n h((x_{\pi(i)} - x_{\pi(i-1)}) \mathbf{1}_{\{\pi(i), \dots, \pi(n)\}}).$$

Also by comonotonic additivity, we have

$$0 = h(\mathbf{0}) = h(\mathbf{1}_{[n]} - \mathbf{1}_{[n]}) = h(\mathbf{1}_{[n]}) + h(-\mathbf{1}_{[n]})$$

and hence $h(-\mathbf{1}_{[n]}) = -h(\mathbf{1}_{[n]})$. Moreover, if $h(r\mathbf{x}) = rh(\mathbf{x})$ for all $r > 0$ (and even for $r = 0$ since $h(\mathbf{0}) = 0$), then $h(r\mathbf{1}_{[n]}) = rh(\mathbf{1}_{[n]})$ for all $r \in \mathbb{R}$ and hence

$$\begin{aligned} h(\mathbf{x}) &= x_{\pi(1)} h(\mathbf{1}_{[n]}) + \sum_{i=2}^n (x_{\pi(i)} - x_{\pi(i-1)}) h(\mathbf{1}_{\{\pi(i), \dots, \pi(n)\}}) \\ &= \sum_{i=1}^n (h_i^\pi - h_{i+1}^\pi) x_{\pi(i)} \end{aligned}$$

where $h_i^\pi = h(\mathbf{1}_{\{\pi(i), \dots, \pi(n)\}})$ for $i \in [n]$ and $h_{n+1}^\pi = h(\mathbf{1}_\emptyset)$.

Let us now show that h satisfies the positive homogeneity property as soon as it is continuous. Comonotonic additivity implies that $g(n\mathbf{x}) = ng(\mathbf{x})$ for every $\mathbf{x} \in \mathbb{R}^n$ and every positive integer n . For any positive integers n, m , we then have

$$\frac{m}{n} h(\mathbf{x}) = \frac{m}{n} h\left(n \frac{\mathbf{x}}{n}\right) = m h\left(\frac{\mathbf{x}}{n}\right) = h\left(\frac{m}{n} \mathbf{x}\right)$$

which means that $h(r\mathbf{x}) = rh(\mathbf{x})$ for every positive rational r and even for every positive real r by continuity. \square

In the following characterization of the signed Choquet integral, we will assume that the function to axiomatize is constructed from a signed capacity. More precisely, denoting the set of signed capacities on $[n]$ by Σ_n , we now regard our function as a map $f: \mathbb{R}^n \times \Sigma_n \rightarrow \mathbb{R}$, or equivalently, as the class $\{f_v: \mathbb{R}^n \rightarrow \mathbb{R} : v \in \Sigma_n\}$. We will adopt the latter terminology to state our result, which is inspired from a characterization given in [5].

For every $T \subseteq [n]$, let $v_T \in \Sigma_n$ be the *unanimity game* defined by $v_T(S) = 1$, if $S \supseteq T$, and 0, otherwise. Note that the v_T ($T \subseteq [n]$) form a basis (actually, the standard basis) for Σ_n . Indeed, for every $v \in \Sigma_n$, we have

$$v = \sum_{T \subseteq [n]} m_v(T) v_T,$$

where m_v is the Möbius transform of v .

Theorem 3.3. *If the class $\{f_v: \mathbb{R}^n \rightarrow \mathbb{R} : v \in \Sigma_n\}$ satisfies the following properties*

(i) *There exist 2^n functions $g_T: \mathbb{R}^n \rightarrow \mathbb{R}$ ($T \subseteq [n]$) such that*

$$f_v = \sum_{T \subseteq [n]} v(T) g_T;$$

(ii) *For every $S \subseteq [n]$, we have $f_{v_S}(\mathbf{x}) = 0$ whenever $x_i = 0$ for some $i \in S$;*

(iii) *For every $S \subseteq [n]$, $r > 0$, $s \in \mathbb{R}$, and $\mathbf{x} \in \mathbb{R}^n$, we have*

$$f_{v_S}(r\mathbf{x} + s\mathbf{1}_{[n]}) = rf_{v_S}(\mathbf{x}) + s;$$

then and only then $f_v = C_v$ for all $v \in \Sigma_n$.

Proof. The sufficiency is straightforward, so let us prove the necessity. Given the relation between v and m_v , condition (i) is equivalent to assuming the existence of 2^n functions $h_T: \mathbb{R}^n \rightarrow \mathbb{R}$ ($T \subseteq [n]$) such that

$$f_v = \sum_{T \subseteq [n]} m_v(T) h_T.$$

Thus $f_{v_T} = h_T$. Therefore, it suffices to prove the following claim.

Claim. For any fixed $T \subseteq [n]$, if the function $f_{v_T}: \mathbb{R}^n \rightarrow \mathbb{R}$ satisfies conditions (ii) and (iii), then $f_{v_T}(\mathbf{x}) = \bigwedge_{i \in T} x_i$ for all $\mathbf{x} \in \mathbb{R}^n$.

Let $\mathbf{x} \in \mathbb{R}^n$. If $x_1 = \dots = x_n$, then

$$f_{v_T}(\mathbf{x}) = f_{v_T}\left(\left(\bigwedge_{i \in [n]} x_i\right)\mathbf{1}_{[n]}\right) = \bigwedge_{i \in [n]} x_i,$$

since $f_{v_T}(\mathbf{0}) = 0$ by (iii).

Otherwise, if $\bigvee_{i \in [n]} x_i - \bigwedge_{i \in [n]} x_i \neq 0$, then by (iii) we have

$$(4) \quad f_{v_T}(\mathbf{x}) = \left(\bigvee_{i \in [n]} x_i - \bigwedge_{i \in [n]} x_i\right) f_{v_T}(\mathbf{x}') + \bigwedge_{i \in [n]} x_i,$$

where

$$\mathbf{x}' = \frac{\mathbf{x} - (\bigwedge_{i \in [n]} x_i)\mathbf{1}_{[n]}}{\bigvee_{i \in [n]} x_i - \bigwedge_{i \in [n]} x_i} \in [0, 1]^n.$$

By (iii) and (ii),

$$f_{v_T}(\mathbf{x}') = f_{v_T}\left(\mathbf{x}' - (\bigwedge_{i \in T} x'_i)\mathbf{1}_{[n]}\right) + \bigwedge_{i \in T} x'_i = \bigwedge_{i \in T} x'_i.$$

By (4), $f_{v_T}(\mathbf{x}) = \bigwedge_{i \in T} x_i$. □

Note that the conditions of Theorem 3.3 are independent. Indeed,

(i), (iii) $\not\Rightarrow$ (ii): Consider the class $\{f_v: \mathbb{R}^n \rightarrow \mathbb{R} : v \in \Sigma_n\}$ given by the weighted arithmetic mean functions

$$f_v(\mathbf{x}) = \sum_{T \subseteq [n]} m_v(T) \left(\frac{1}{|T|} \sum_{i \in T} x_i\right),$$

where m_v is the Möbius transform of v .

(i), (ii) \nrightarrow (iii): Consider the class $\{f_v: \mathbb{R}^n \rightarrow \mathbb{R} : v \in \Sigma_n\}$ given by the multilinear polynomial functions

$$f_v(\mathbf{x}) = \sum_{T \subseteq [n]} m_v(T) \prod_{i \in T} x_i,$$

where m_v is the Möbius transform of v .

(ii), (iii) \nrightarrow (i): Define the normalized capacity $v^* \in \Sigma_3$ by $v^*(\{1, 2\}) = v^*(\{3\}) = 0$ and $v^*(\{1, 3\}) = v^*(\{2, 3\}) = 1/2$ and consider the class $\{f_v: \mathbb{R}^3 \rightarrow \mathbb{R} : v \in \Sigma_3\}$ given by $f_v = C_v$ for every $v \in \Sigma_3 \setminus \{v^*\}$, and

$$f_{v^*}(x_1, x_2, x_3) = \left(\frac{x_1 + x_2}{2}\right) \wedge x_3.$$

- Remark 1.* (a) The conditions in Theorem 3.3 can be justified as follows. Condition (i) expresses the fact that the aggregation model is linear with respect to the underlying signed capacities. Condition (ii) expresses minimal requirements on the functions defined on the standard basis $\{v_S : S \subseteq [n]\}$ of Σ_n . Condition (iii) expresses the fact that f_{v_S} is meaningful with respect to interval scales.
- (b) The characterization given in Theorem 3.3 does not use the fact that $v(\emptyset) = 0$. Therefore they can be immediately adapted to Lovász extensions by redefining Σ_n as the set of set functions on $[n]$.

REFERENCES

- [1] G. Choquet. Theory of capacities. *Ann. Inst. Fourier, Grenoble*, 5:131–295 (1955), 1953–1954.
- [2] L. M. de Campos and M. J. Bolaños. Characterization and comparison of Sugeno and Choquet integrals. *Fuzzy Sets and Systems*, 52(1):61–67, 1992.
- [3] A. De Waegenaere and P. Wakker. Nonmonotonic Choquet integrals. *J. Mathematical Economics*, 36:45–60, 2001.
- [4] L. Lovász. Submodular functions and convexity. In *Mathematical programming, 11th int. Symp., Bonn 1982*, 235–257. 1983.
- [5] J.-L. Marichal. An axiomatic approach of the discrete Choquet integral as a tool to aggregate interacting criteria. *IEEE Trans. Fuzzy Syst.*, 8(6):800–807, 2000.
- [6] J.-L. Marichal. Aggregation of interacting criteria by means of the discrete Choquet integral. In *Aggregation operators: new trends and applications*, pages 224–244. Physica, Heidelberg, 2002.
- [7] T. Murofushi, M. Sugeno, and M. Machida. Non-monotonic fuzzy measures and the Choquet integral. *Fuzzy Sets and Systems*, 64:73–86, 1994.
- [8] D. Schmeidler. Integral representation without additivity. *Proc. Amer. Math. Soc.*, 97(2):255–261, 1986.
- [9] I. Singer. Extensions of functions of 0-1 variables and applications to combinatorial optimization. *Numer. Funct. Anal. Optimization*, 7:23–62, 1984.
- [10] M. Sugeno. *Theory of fuzzy integrals and its applications*. PhD thesis, Tokyo Institute of Technology, Tokyo, 1974.
- [11] M. Sugeno. Fuzzy measures and fuzzy integrals—a survey. In M. M. Gupta, G. N. Saridis, and B. R. Gaines, editors, *Fuzzy automata and decision processes*, pages 89–102. North-Holland, New York, 1977.

DEPARTMENT OF APPLIED MATHEMATICS, UNIVERSITY CA' FOSCARI OF VENICE,
DORSODURO 3825/E-30123, VENICE, ITALY
E-mail address: `mcardin[at]unive.it`

MATHEMATICS RESEARCH UNIT, FSTC, UNIVERSITY OF LUXEMBOURG, 6, RUE COUDENHOVE-
KALERGI, L-1359 LUXEMBOURG, LUXEMBOURG
E-mail address: `miguel.couceiro[at]uni.lu`

DEPARTMENT OF APPLIED MATHEMATICS, UNIVERSITY CA' FOSCARI OF VENICE,
DORSODURO 3825/E-30123, VENICE, ITALY
E-mail address: `sgiove[at]unive.it`

MATHEMATICS RESEARCH UNIT, FSTC, UNIVERSITY OF LUXEMBOURG, 6, RUE COUDENHOVE-
KALERGI, L-1359 LUXEMBOURG, LUXEMBOURG
E-mail address: `jean-luc.marichal[at]uni.lu`