

# Power spectrum features of the near-Earth atmospheric electric field in Kamchatka

Yuri M. Mikhailov <sup>(1)</sup>, Galina A. Mikhailova <sup>(1)</sup>, Olga V. Kapustina <sup>(1)</sup>,  
Alexandr V. Buzevich <sup>(2)</sup> and Sergei E. Smirnov <sup>(2)</sup>

<sup>(1)</sup> Institute of Terrestrial Magnetism, Ionosphere and Radiowave Propagation (IZMIRAN),  
Russian Academy of Science, Troitsk (Moscow Region), Russia

<sup>(2)</sup> Institute of Space Physics Researches and Radiowave Propagation, Far-East Branch  
of Russian Academy of Science, Petropavlovsk-Kamchatski (Kamchatka Region), Russia

## Abstract

Power spectrum of the diurnal variations of the *quasi*-electrostatic field  $E_z$  in the near-Earth atmosphere have been presented for the first time. The  $E_z$  power spectrum variations in the period of fine weather have been shown to exhibit two bands of the periods of natural atmospheric oscillations with  $T = 1-5$  and  $6-24$  h. These oscillations are the modes of the internal gravity and tidal waves in the lower atmosphere. On the days under atmospheric precipitation, the spectral power of  $E_z$  increases by an order of magnitude. During the pre-earthquake period, when the diurnal  $E_z$  variation had an anomaly, the intensity of harmonics with  $T = 1.8, 2.2,$  and  $3.8$  h increased by an order of magnitude or more in comparison with the  $E_z$  spectra in fine weather. Two additional spectral bands with  $T = 0.6$  and  $1$  h have appeared simultaneously.

**Key words** atmospheric quasi-static electric field – atmospheric precipitation – atmospheric waves


## 1. Introduction

Anomalous variations in the *quasi*-static electrical field  $E_z$  in the near-Earth atmosphere before the  $M \geq 4$  earthquakes have been recorded in many seismo-active regions of the Earth. They usually occur at an interval of a few hours to a few days before the main shock and prove to be either a bay-like field intensity decrease (even to a sign reversal) or an oscillation train lasting a few hours. These observations were summarized and analysed by Rulenko (2000) and made it possible to establish that both types of  $E_z$  anomalies are caused by the deformation

processes in the subsurface layers of the Earth's crust during the earthquake preparation phase. The former type of anomalies were observed on Kamchatka for a few years and were referred to the radon (Rn) emission from subsurface gases into the atmosphere, resulting in variations of atmospheric conductivity and, therefore in  $E_z$  magnitude (Rulenko *et al.*, 1992, 1996; Buzevich *et al.*, 1998; Smirnov, 2001). These results, as well as the observations in other seismo-active regions, have made it possible to assume that the *quasi*-static electric field in near-Earth atmosphere can be responsible for ionospheric disturbances occurring before seismic shocks (Morgunov, 1988; Pulinets *et al.*, 1998; Mikhailov *et al.*, 1999). Research of electrical field intensity in temporary series has shown that the field value is affected by meteorological characteristics (precipitation, winds) (Smirnov, 2001; Mikhailov *et al.*, 2002). This fact makes it difficult to detect some anomalies in  $E_z$  behaviour before the earthquake.

The object of this paper is the investigation of the power spectrum in the near-Earth atmos-

*Mailing address:* Dr. Yuri M. Mikhailov, Institute of Terrestrial Magnetism, Ionosphere and Radiowave Propagation (IZMIRAN), Russian Academy of Science, 142190 Troitsk (Moscow Region), Russia; e-mail: yumikh@izmiran.rssi.ru



**Fig. 1.** Typical diurnal variations in a *quasi*-static electric field  $E_z$  in the near-Earth atmosphere: 1-3 – days without atmospheric precipitation; 2 – a day before the earthquake; 3 – for the earthquake day; 4 – day with heavy atmospheric precipitation.

pheric *quasi*-static electric field  $E_z$  in a period of the fine weather, under atmospheric precipitation and during the pre-earthquake period of anomalous diurnal variations  $E_z$ .

## 2. Experimental data and processing procedure

The Pole-2M electrostatic flux-meter, whose output data are transmitted into the information processing centre in the analogue form, is used at Paratunka ( $\varphi = 53^\circ\text{N}$ ,  $\lambda = 158.3^\circ\text{E}$ ) to measure the vertical component of the atmospheric electric field ( $E_z$ ). The  $E_z$  values are digitized with sampling rate  $\Delta t = 1$  min. The results of the investigations of the power spectrums of the near-Earth atmospheric electric field in September-October 1999 are presented below. Figure 1 shows the typical diurnal  $E_z$  variations in the fine weather (curve 1), on the day before the earthquake (curve 2, September 17), on the day of the earthquake (curve 3, September 18, 21.28:33.17 UT,  $\varphi = 51.21^\circ\text{N}$ ,  $\lambda = 157.56^\circ\text{E}$ ,  $h = 60$  km,  $M_p = 6.0$ ) and on the day with intense precipitation (curve 4, September 19). The character of the diurnal  $E_z$  variations in the fine weather is described qualitatively by physical processes, occurring in the near-Earth atmosphere. The increase in  $E_z$  magnitude at 19-24 UT (06.00-11.00 LT) is related to the convective processes in the atmosphere due to the air temperature variations in the morning. In the fine weather in the daytime and at night the electrical fields are  $\sim 100$ -120 V/m, varying in the range  $\sim \pm 20$  V/m.

At high and durable precipitation, the electric field value decreased to  $-1$  kV/m. On the rainy day (September 19), which carries to the Earth's surface volume changes, the atmospheric conductivity increases and  $E_z$  decreases, even to a sign reversal. Other examples of the diurnal  $E_z$  variations in fine weather and at high precipitation were shown in the paper (Mikhailov *et al.*, 2002).

On the day before the earthquake (curve 2), the bay of decreased  $E_z$  magnitude was observed at 15.00-24.00 UT against a background of quiet meteorological conditions. Similar effects were observed in the Paratunka during pe-


riod 1997-2000 in 1-36 h before earthquakes (Smirnov, 2001).

By example of 29 September 1999, fig. 2 illustrates the spectral processing of the diurnal  $E_z$  data. The upper curve 1 represents the diurnal variations  $E_z$ . The next curve below 2 is the same data obtained by subtracting the daily average  $E_z$  value from curve 1. Curve 2 shows the well-marked semidiurnal wave, whose daytime amplitude is less than the nighttime one. These values have been added by zeros to have 2048 points, needed for using the fast Fourier transform algorithm (FFT). The power spectrum of the augmented  $E_z$  time series was computed by the method of periodograms with rectangular time «window» at frequencies of  $f_k = k \cdot \Delta f$  with a corresponding frequency step of  $\Delta f = 1/2048 \cdot \Delta t = 8.14 \cdot 10^{-6}$  Hz and with periods of  $T_k = 1/f_k$ , where  $k = 1, 2, \dots, 1024$ . This spectrum is shown in the panel 4 (solid line, left ordinate axis). The period of the predominant harmonic is  $T = 12$  h and less intensive harmonics with periods of 1-5 h are also present. To distinguish them we excluded periods longer than 5 h from the spectrum (the dash-dotted line, right ordinate axis). The time form of these filtered series is shown by curve 3. The spectra labelled 5 are shown in the bottom of fig. 2 as a function of the periods varying to 24 h (on the left) and  $T < 5$  h (on the right).


## 3. Main results

The method described above was used to make the spectral analysis of the diurnal variations in  $E_z$  for September-October 1999. Naturally, the spectrum forms varied from day to day. To obtain statistically reliable spectra and to distinguish stable maximums the data have been averaged over days with specific  $E_z$  behaviour. We used days without atmospheric precipitation and earthquakes, when the spectral variations may be caused by global effects in the Earth's crust in the given region and, consequently, in the near-Earth atmosphere. Figure 3 shows the control atmospheric precipitation data for the observation period.

The resulting averaged spectra for the days without (22 days, curve 1) and with (11 days,




**Fig. 2.** The method for spectral processing of the diurnal  $E_z$  variations.




**Fig. 3.** Variations in the atmospheric precipitation level measured twice a day during September-October 1999.

curve 2) atmospheric precipitation are shown in fig. 4 ( $T < 5$  h) together with the individual spectrum for September 17 (curve 3). The vertical bars in curves 1 and 2 are the mean-square deviations. Curve 1 indicates the spectrum weakening from  $\sim 2.5 \cdot 10^9$   $V^2/m^2$  Hz at  $T \sim 4$  h to  $\sim 2 \cdot 10^8$   $V^2/m^2$  Hz at  $T \sim 1$  h without pronounced peaks at a level of 0.5 relative to the maximum value. The character of the spectrum shown by

curve 2 has changed insignificantly, namely, the energy enhancements at  $T \sim 4$  h and  $\sim 2$  h, but the power level have increased twofold compared to curve 1, with very large mean-square deviations indicating that individual spectra are widely variable. Finally, curve 3 represents the individual  $E_z$  power spectrum for 17 September 1999 without atmospheric precipitation, but with anomalous diurnal variations (see fig. 1).



**Fig. 4.** The averaged  $E_z$  power spectrum in the band of periods  $T < 5$  h on the days without precipitation (curve 1) and with precipitation (curve 2) and 17 September with the anomalous diurnal variations  $E_z$  (curve 3).



**Fig. 5.** The averaged  $E_z$  power spectrum with periods of  $2 \text{ h} < T < 30 \text{ h}$  for the fine weather days from 1 September to 29 October 1999.

The following features are clearly defined in this curve:

- i) There are the spectral bands (at 0.5 level of the maximum) at  $T = 0.6, 1, 1.8, 2.2$  and  $3.8 \text{ h}$ .
- ii) The intensity at maxima of these bands exceeds the corresponding values for the days without atmospheric precipitation by an order of magnitude or more.

The  $T > 5 \text{ h}$  harmonics were distinguished by calculating the spectra of longer initial date series with a time step of  $\Delta t = 1 \text{ h}$  only for days of the fine weather, *i.e.* September 1-4, September 20-24, October 20-24 and October 25-29. Figure 5 represents the resulting spectrum, averaged over these days, which shows the clear maximums with  $T \sim 8, 10, 12$  and  $24 \text{ h}$ .

During September-October 1999 (except September 18), two more Kamchatka earthquakes occurred, whose preparation zone included the  $E_z$  measurement point. The earthquakes occurred on October 5 (05.01:35.94 UT,  $\varphi = 51.21^\circ\text{N}$ ,  $\lambda = 157.61^\circ\text{E}$ ,  $h = 76 \text{ km}$ ,  $M = 5.6$ ) and on October 18 (20.49:47.92 UT,  $\varphi =$

$= 51.30^\circ\text{N}$ ,  $\lambda = 157.12^\circ\text{E}$ ,  $h = 138 \text{ km}$ ,  $M = 4.8$ ). From fig. 3 it is seen, however, that rainfalls were recorded during these days, which probably screened the  $E_z$  effects caused by other sources.

#### 4. Discussion

The continuous measurements of *quasi-static* electric field  $E_z$  with the sampling rate  $\Delta t = 1 \text{ min}$  have made it possible to study the fine structure of the power spectra with periods from 4 min and longer. Two spectral bands with periods  $T = 1-5 \text{ h}$  and  $6-24 \text{ h}$  have been found by analysing the  $E_z$  diurnal variations for September-October 1999. Under fine weather, clearly defined maximums (at a level of 0.5) were absent, but the intensity trended to weaken when the periods decreased from 5 to 1 h. These periods are known to be the periods of internal gravity waves, which are clearly seen in the seismo-gravitational oscillations of the

Earth and related with pressure disturbances in the near-surface atmosphere (Garmash *et al.*, 1989; Petrova, 1999). The coincidence of the  $E_z$  power spectra with those of the Earth's seismo-gravitational oscillations, allows us to state that the  $E_z$  variations are of seismic nature. If we accept the «piston» mechanism of generation of air pressure fluctuations by the Earth's surface oscillations (Garmash *et al.*, 1989), then we may expect similar variations in the radon concentration spectra in subsurface gases.

The  $T = 6-24$  h spectral band shows the clear harmonics with  $T = 8, 12$  and  $24$  h. These are the tidal waves in the lower atmosphere, related to the variations in the temperature of underlying surface. Their intensity is of the same order of magnitude as that of internal gravity waves.

One day before the earthquake on 18 September 1999, the  $E_z$  power spectrum showed three distinct maximums (relative to the half-maximum level) at  $T = 1.8, 2.2,$  and  $3.8$  h. Their intensity increased by an order of magnitude and more, especially at  $T < 3$  h, as compared with the  $E_z$  spectra during the days of the fine weather. Moreover, spectral components at  $T = 0.6$  and  $1$  h appeared too. The similar spectral structures within the  $T = 1-5$  h band were observed by Lin'kov *et al.* (1990) and their enhancement in the variations in the Earth's seismo-gravitational oscillations and in the simultaneous pressure variations in the near-Earth atmosphere before a strong earthquake. The agreement between the  $E_z$  spectra and the results of those papers indicate that not only the quiet diurnal variations in  $E_z$ , but also anomalies before earthquakes are of the same seismo-gravitational nature.

## 5. Conclusions

The fine spectral analysis of diurnal variations in the quasi-static electric field  $E_z$  in the near-Earth atmosphere has been carried out for the first time and has shown the following:

i) In fine weather and on the days with atmospheric precipitation the natural frequencies of atmospheric oscillations have been distinguished in two bands with periods of  $T = 1-5$  h and  $T = 6-24$  h in the  $E_z$  power spectrum. The

former band corresponds to internal gravity wave modes of the lower atmosphere, while the second band to the tidal waves.

ii) During the earthquake preparation period, the intensity of bands with  $T = 1.8, 2.2,$  and  $3.8$  h increased by an order of magnitude and more compared with spectra obtained in fine weather. During this period, two additional spectral bands with  $T = 0.6$  and  $1.0$  h were distinguished.

iii) The nature of the  $E_z$  spectral anomalies from the Earth's seismo-gravitational oscillations has been confirmed experimentally. However, the question has still to be answered as to whether the pre-earthquake bay-like weakening anomalies of  $E_z$  are caused by enhancement of conductivity in the lower atmosphere as a result of the additional ionization by a radioactive gas, or this weakening is a purely seismic electrical signal caused by the rock failure in a hypocenter.

## Acknowledgements

The authors are deeply indebted to L.N. Petrova for useful discussions of our report at the conference «Problems of Geocosmos» held in St. Petersburg and to the referees for their kind help and useful suggestions.

This work was supported by the Russian Foundation for Basic Research, project No. 00-05-65020, No. 00-05-79047, and No. 00-05-65380.

## REFERENCES

- BUZEVICH, A.V., G.I. DRUZHIN, P.P. FIRSTOV, E.F. VERSHININ, S.E. SMIRNOV and V.I. FILIMONOV (1998): The heliogeophysical effects prior to the Kronotsk earthquake of December 5, 1997,  $M = 7.7$ , in *Kronotsk Earthquake of the December 5, 1997, in Kamchatka: Precursors, Specific Features and Aftereffects*, edited by E.V. GORGEEV (Petropavlovsk-Kamchatsky, Ross. Acad. Nauk), 177-188.
- GARMASH, S.V., E.M. LIN'KOV, L.N. PETROVA and G.M. SHVED (1989): Excitation of atmospheric oscillations by seismo-gravitational oscillations of the Earth, *Izv. Akad. Nauk SSSR*, **25** (12), 1290-1299.
- LIN'KOV, E.M., L.N. PETROVA and K.S. OSIPOV (1990): Seismo-gravitational oscillations of the Earth and atmospheric disturbances as possible precursors of strong earthquakes, *Dokl. Akad. Nauk SSSR, Phys. Atmos. Ocean*, **313** (5), 1095-1098.



- MIKHAILOV, YU.M., G.A. MIKHAILOVA and O.V. KAPUSTINA (1999): ULF-ELF electromagnetic emissions over the fault in Kangra Valley of India and their relation with radon emanation (Intercosmos 24 satellite data), *Geomagn. Aeron.*, **39** (4), 62-68.
- MIKHAILOV, YU.M., G.A. MIKHAILOVA, O.V. KAPUSTINA, A.KH. DEPUEVA, A.V. BUZEVICH, G.I. DRUZHIN, S.E. SMIRNOV and P.P. FURSTOV (2002): Variations of different atmospheric and ionospheric parameters in preparing phase of earthquakes at Kamchatka: preliminary results, *Geomagn. Aeron.*, **42** (6), 805-813.
- MORGUNOV, V.A. (1988): To the nature of lithosphere-ionosphere coupling, *Izv. Akad. Nauk SSSR, Phys. Earth*, **5**, 80-87.
- PETROVA, L.N. (1999): Seismic process in the 0.05-0.5 mHz frequency band: regularities and specific features, *Volcanol. Seismol.*, 4-5, 116-124.
- PULINETS, S.A., V.V. KHEGAI, K.A. BOYARCHUK and A.M. LOMONOSOV (1998): Atmospheric electric field as a source of ionospheric variability, *Uspekhi Fiz. Nauk*, **168** (5), 582-589.
- RULENKO, O.P. (2000): Effective earthquake precursors in the electricity of the near-Earth atmosphere, *Volcanol. Seismol.*, **4**, 57-68.
- RULENKO, O.P., A.V. IVANOV and A.V. SHUMEIKO (1992): The short-term atmospheric-electricity precursor of the Kamchatka earthquake of 6 March 1992,  $M = 6.1$ , *Dokl. Akad. Nauk SSSR*, **326** (6), 980-982.
- RULENKO, O.P., G.I. DRUZHIN and E.F. VERSHININ (1996): Measurements of an atmospheric electrical field and natural electromagnetic emission before the Kamchatka earthquake of 11 November 1993,  $M = 7.0$ , *Dokl. Akad. Nauk SSSR*, **348** (6), 814-816.
- SMIRNOV, S.E. (2001): Anomalies behaviour of electric field intensity before the earthquake according to the observations of magnetic observatory «Paratunka» in 1997-2000, in *Solar-Terrestrial Coupling and Electromagnetic Precursors of Earthquakes. Theses of the Repts. at the 2nd International Meeting at Paratunka (Kamchatka Region)*, 14-19 August 2001, edited by YU.S. SHUMILOV and G.I. DRUZHIN (Petropavlovsk-Kamchatsky, IKIR, Far-Eastern Branch of RAS), pp. 48-49.