



# *Wood Anderson Magnitude Scale for Mt. Vesuvius*

- A revised ML scale for VT events at Mt. Vesuvius -

Edoardo Del Pezzo<sup>1,2</sup> and Simona Petrosino<sup>1</sup>

1 Osservatorio Vesuviano

2 Physics Department - University of Salerno- Italy

**A Mathcad-8 Professional Program**

Osservatorio Vesuviano  
Open file report  
1999 n° 3

# Wood Anderson Magnitude Scale for Mt. Vesuvius

## A revised ML scale for VT events at Mt. Vesuvius

Edoardo Del Pezzo<sup>1,2</sup>, Simona Petrosino<sup>1</sup>

1 - Osservatorio Vesuviano

2 - Physics Department, University of Salerno. Italy

**A Mathcad-8 Professional Program**

### Abstract

A Mathcad-8 program to calculate a revised magnitude scale is presented. An application to Mt. Vesuvius is included as a program test. Wood-Anderson seismograms for 131 local earthquakes recorded at station BKE (Osservatorio Vesuviano seismic network) were synthesized to estimate local magnitude from the original definition:

$$Ml = \log A_{max}(\Delta) - \log A_o(\Delta)$$

The distance correction  $\log A_o(\Delta)$  was empirically determined simulating a wave packet which propagates in a structure with assigned Q.

Moment magnitude (calculated both with Kanamori and Thatcher-Hanks formulas) was also determined from the displacement spectra.

Finally a relation between Wood-Anderson magnitude and duration magnitude was derived, allowing the estimate of local magnitude from the duration of the earthquake.

### Theory

#### Local Magnitude definition

The definition of local magnitude is:

$$Ml = \log A_{max}(\Delta) - \log A_o(\Delta) \quad (1)$$

where  $A_{max}$  is the Wood Anderson maximum amplitude, and  $A_o$  is the Wood Anderson maximum amplitude for the reference earthquake. This scale uses as reference the earthquake of Magnitude 3 which in California, where the scale was set up, takes the max amplitude of 1 mm at a distance  $\Delta$  of 100 km. For California the formula giving the local Magnitude as a function of distance is:

$$Ml_{cal} = \log A_{max}(\Delta) + 2.76 \log(\Delta) - 2.48 + C \quad (2)$$

where  $C$  is a correction term taking into account the deviation of the scale at the station of the network.

We normalize the scale for Mt Vesuvius in such a way that an earthquake at  $\Delta=10$  km has the same local Magnitude as in California. This means that at 10 km from the source an earthquake of a given Magnitude in California, would have the same maximum amplitude as at Mt. Vesuvius. This allows a comparison of the Magnitude values at Mt. Vesuvius with those for California. A similar normalization for a distance close to the source was proposed by Hutton and Boore (1987) for local earthquakes. In this way the above authors eliminated the strong regional attenuation anomalies for S wave propagation.

The empirical formula for the attenuation of the maximum amplitude with distance at Mt. Vesuvius was calculated using a numerical simulation. First we generate a synthetic S-wave packet, with a flat spectrum at a distance close to the source (0.1 km). A sequence of 125 random numbers between 1 and -1 with a uniform distribution simulate the wave packet. Then we multiply the sequence by a Hanning window. The signal represents the S-wave packet sampled at 1/125 sps.

### Simulation of the synthetic wave packet

k := 0.. 100

$R_k := \frac{k + 1}{10}$  This is the distance range in km


$rr_k := \log(R_k)$

$v(k) := \text{runif}(125, -1, 1)$  This is the vector of 125 samples, uniformly distributed

$hn := \text{hanning}(125)$  This is the hanning window

$i := 1..124$

$vv_{k,i} := hn_i \cdot v(k)_i \cdot 1000$  This is the wave packet. The amplitude is arbitrary



$\beta := 2$  This is the S-wave wave velocity

$VV(k) := \text{CFFT}\left[\left(vv^T\right)^{\langle k \rangle}\right]$  This is the Fourier transform of the synthetic signal

Now we apply the attenuation operator with  $Q=60$  as measured at Mt Vesuvius


$f_i := i$

$fc := 10$  This is the corner frequency of the source spectrum

$$\text{ampl}_{i,k} := \frac{|VV(k)_i| \cdot 0.1}{R_k \cdot \left[1 + \left(\frac{f_i}{fc}\right)^2\right]} \cdot \left(\exp\left(\frac{-\pi \cdot f_i \cdot R_k}{\beta \cdot 60}\right)\right)$$

$\text{fase}_i := \arg(VV(k)_i)$

$VVatt_{i,k} := \text{ampl}_{i,k} \cdot \exp(i \cdot \text{fase}_i)$




$\text{Attensismo}^{\langle k \rangle} := \text{ICFFT}(VVatt^{\langle k \rangle})$  This is the synthetic seismogram at different distances from the

source. The next plot shows the seismogram recorded at 0.1 and 0.3 km distance from the source.


$$t_i := \frac{i}{125}$$

$$R_1 = 0.2$$

$$R_{100} = 10.1$$



$\text{massimo}_k := \max(\text{Re}(\text{Attensismo}^{<k>}))$  This is the vector of the maximum amplitudes at different distances.



## Best fit with a relation of the form of (2)

$$y_k := -\log(\text{massimo}_k)$$

$$\text{ter}_k := 1$$

$$G^{<0>} := \text{rr}$$

G is the matrix of coefficients

$$G^{<1>} := \text{ter}$$

$$\text{par} := (G^T \cdot G)^{-1} \cdot G^T \cdot y \quad \text{This is the least square fit for the coefficients of relation (2)}$$

$$\text{par} = \begin{bmatrix} 1.3422631 \\ -1.0849864 \end{bmatrix} \quad \text{This is the solution}$$

The relation for Mt Vesuvius is  $1.28 \log(\Delta) + b$ .  $b$  has to be determined by the normalization at 10 km distance. The normalization is given by the amplitude of a  $M=3$  earthquake at 10 km. For this earthquake:

$$Ml = \log A_{max} + 2.76 \log 10 - 2.48 = 3. \rightarrow \log A_{max} = 2.72$$

For Mt. Vesuvius,  $Ml = 2.72 + 1.28 \log 10 + b = 3$ , which gives  $b = -1.1$ , then the formula is:

$$Ml = \log A_{max} + 1.28 \log (\Delta) - 1.1$$

## Application to an example

These are the input traces:

:=   
A:\mag\03010947.e

:=   
A:\mag\03010947.n

$$\text{trace} := \text{trac} \cdot \text{volt} \cdot 10^{-3}$$

The original signal is in mV. This is the correction to Volts

$$\text{trace2} := \text{trac2} \cdot \text{volt} \cdot 10^{-3}$$

$$\text{last}(\text{trace}) = 2499$$

$$\text{max}(\text{trace}) = 0.007872 \text{ V}$$

Traces in Volts. Note that Mathcad automatically checks the units.

$$\text{last}(\text{trace2}) = 2499$$

$$\text{max}(\text{trace2}) = 0.00776 \text{ V}$$



$$j := 1.. \text{last}(\text{trace})$$

$$t_j := j \cdot \frac{1}{125} \cdot \text{s}$$

$$T := \text{last}(\text{trace}) \cdot \frac{1}{125} \cdot \text{sec}$$

$$T = 19.992 \text{ s}$$

T is the seismogram window duration



$$f_0 := 1 \cdot \text{Hz}$$

$\gamma$  is the damping coefficient of the Lennartz portable station which recorded the event shown above

$$\gamma := 0.68$$

$$G_a := \frac{4}{2.4 \cdot 0.01} \cdot \frac{\text{V}}{\frac{\text{m}}{\text{sec}}}$$

$$G_a = 166.6666667 \text{ s} \cdot \text{m}^{-1} \cdot \text{V} \quad \text{This is the internal damping main coil motor constant}$$

$$G := 125 \text{ V} \cdot \text{s} \cdot \text{m}^{-1} \quad G \text{ is the motor constant at } \gamma$$

traspecvel := cfft (trace - mean(trace))      These are the Fourier Transforms of the detrended traces

traspecvel2 := cfft (trace2 - mean(trace2))

last(traspecvel) = 2499      The number of points in the Fourier Transform

last(traspecvel2) = 2499

$$f_j := \left( \frac{1}{125} \cdot \text{last}(\text{trace}) \right)^{-1} \cdot j \cdot \text{Hz} \quad \text{The frequency in Hz}$$

Anti-aliasing Filter Lennartz:

$$a_0 := 1 \quad b_1 := 0.3887$$

$$a_1 := 1.2217 \quad b_2 := 0.3505$$

$$a_2 := 0.9686 \quad b_3 := 0.2756$$

$$a_3 := 0.5131 \quad f_c := 25 \cdot \text{Hz}$$

$$\text{filtro}(f) := \frac{a_0}{\prod_{i=1}^3 \left[ 1 + a_i \cdot \frac{f}{f_c} \cdot i + b_i \cdot \left( \frac{f}{f_c} \cdot i \right)^2 \right]} \quad \text{The transfer function of the anti-alias filter}$$


$$H(f) := \frac{(G \cdot (2 \cdot \pi \cdot f)^2) \cdot \text{filtro}(f)}{(2 \cdot \pi \cdot f_0)^2 - (2 \cdot \pi \cdot f)^2 + 2 \cdot i \cdot (2 \cdot \pi \cdot f) \cdot \gamma \cdot (2 \cdot \pi \cdot f_0)} \quad \text{This is the velocity response curve in V/m/s}$$

Hspo(f) := H(f) · i · 2 · π · f      This is the displacement response curve, obtained multiplying for frequency

The the Wood Anderson magnification curve is:

$$\text{HWA}(f) := \frac{2800 \cdot (2 \cdot \pi \cdot f)^2}{(-2 \cdot \pi \cdot f)^2 + \left( 2 \cdot \pi \cdot \frac{1}{0.8 \cdot \text{sec}} \right)^2 + 2 \cdot i \cdot (2 \cdot \pi \cdot f) \cdot 0.8 \cdot \left( 2 \cdot \pi \cdot \frac{1}{0.8 \cdot \text{sec}} \right)}$$

$$\left| \text{Hspo}(f_6) \right| = 21.2881184 \text{ m}^{-1} \cdot \text{V} \quad \text{Note the physical dimensions in V/m}$$



In the plot the amplitude response for Velocity, Displacement and Wood Anderson is shown.

## Transformation of the Signal in Wood Anderson Equivalent


$j := 1.. \text{last}(\text{trace})$

$\text{spostamento}_j := \frac{\text{traspecvel}_j}{H_{\text{spo}}(f_j)}$  The real ground displacement spectrum for E-W component

$\text{velocità}_j := \frac{\text{traspecvel}_j}{H(f_j)}$  The real ground velocity spectrum

$\text{spostamento2}_j := \frac{\text{traspecvel2}_j}{H_{\text{spo}}(f_j)}$  The same for N-S component

$\text{velocità2}_j := \frac{\text{traspecvel2}_j}{H(f_j)}$




$\text{woodandersonspec}_j := HWA(f_j) \cdot \text{spostamento}_j$  Wood Anderson

$$\text{woodandersonspec2}_j := \text{HWA}(f_j) \cdot \text{spostamento2}_j$$

woodandersontrace := icfft(woodandersonspec)      Wood Anderson converted trace for the two  
 woodandersontrace2 := icfft(woodandersonspec2)      ground motion components

$$\text{spostamento\_trace} := \text{icfft}(\text{spostamento})$$


$$\text{velocità\_trace} := \text{icfft}(\text{velocità})$$



$$\text{integral}_0 := 0 \cdot V \cdot \text{sec}$$

$\text{integral}_j := \text{integral}_{j-1} + \text{trace}_j \cdot (t_j - t_{j-1})$       The trace is integrated in time domain too, to check the results

$$\text{integral}_2 = -0.0000014s \cdot V \quad \text{chek of the units}$$



$$\max(\text{Re}(\text{spostamento})) = 0.0000266\text{m}$$

$$\max(\text{Re}(\text{woodandersontrace})) = 0.0328461\text{m}$$

$$\max(\text{Re}(\text{woodandersontrace2})) = 0.0409515\text{m}$$

$$\text{AWAspe} := \frac{\sqrt{\max(\text{Re}(\text{woodandersontrace}))^2 + \max(\text{Re}(\text{woodandersontrace2}))^2}}{2}$$

$$\sqrt{\max(\text{Re}(\text{woodandersontrace}))^2 + \max(\text{Re}(\text{woodandersontrace2}))^2} = 0.0524966\text{m}$$



# Local Magnitude Evaluation

Q := 60      Quality factor measured at Mt. Vesuvius (Bianco et al.,1999). It is independent of frequency  
 v := 2      Seismic wave velocity.

California(r, AWAspe) := log(2·AWAspe) - 2.48 + 2.76·log(r)      Local Magnitude for California

Vesuvio(r, AWAspe) := log(2·AWAspe) + 1.28·log(r) - 1.1      Local Magnitude for Mt. Vesuvius

Moment Magnitude (Kanamori):

fo := 1·Hz

$\rho := 2.700 \frac{\text{gm}}{\text{cm}^3}$

$v_s := 2 \cdot 10^5 \frac{\text{cm}}{\text{sec}}$


$\text{norm} := \frac{2 \cdot T}{\sqrt{\text{last}(\text{trace})}}$

norm = 0.79984s

$\text{Mo}(R, \omega) := \frac{\left( \omega \cdot \exp\left(\frac{\pi \cdot R \cdot f_0}{v_s \cdot Q}\right) \cdot 4 \cdot \pi \cdot \rho \cdot v_s^3 \cdot R \right)}{0.85}$

$\text{mum}_j := \frac{\left| \text{spostamento}_j \cdot \frac{1}{\text{m}} + \text{spostamento2}_j \cdot \frac{1}{\text{m}} \right|}{2}$

musm := medsmooth(mum, 11)



$$\omega := \frac{10^2 \cdot \text{norm} \cdot \text{cm}}{500} \cdot \sum_{k=100}^{600} \text{musm}_k$$

Automatical evaluation of  $\omega$  in cm\*s:

$$\omega = 0.0000104 \text{ s} \cdot \text{cm}$$

$$\text{Mo}(364000 \text{ cm}, \omega) = 1.331880610^{18} \text{ dyne} \cdot \text{cm}$$

Moment Magnitude (Kanamori) and (Tatcher) Formulas:

$$M_w := \frac{\log\left(\text{Mo}(364000 \text{ cm}, \omega) \cdot \frac{1}{\text{dyne} \cdot \text{cm}}\right)}{1.5} - 10.73$$

$$\text{Tatcher} := \frac{\log\left(\text{Mo}(364000 \text{ cm}, \omega) \cdot \frac{1}{\text{dyne} \cdot \text{cm}}\right)}{1.5} - \frac{16}{1.5}$$

$$M_w = 1.3529769$$

**Moment Magnitude (Kanamori)**

$$\text{Tatcher} = 1.4163102$$

**Formula**


**Moment Magnitude with Tatcher and Hanks**

$$\text{California}\left(3.64, 1000 \cdot \text{AWAspe} \frac{1}{\text{m}}\right) = 0.7887708 \quad \text{California W A Magnitude}$$

$$\text{Vesuvio}\left(3.64, \text{AWAspe} 10^3 \cdot \frac{1}{\text{m}}\right) = 1.3383408 \quad \text{Vesuvius W A Magnitude}$$

We calculated the Vesuvius Magnitude, the Kanamori Magnitude and the Tatcher and Hanks Magnitude for 181 earthquakes recorded in 1996 by a Lennartz seismic station (BKE) of the Osservatorio Vesuviano Seismic Network. For 131 events it was possible to compare the Vesuvius Wood-Anderson Magnitude with the Duration magnitude estimated for the seismic station OVO using the empirical formula  $M_D = 2.75 \log \tau - 2.35$ . The Wood Anderson Magnitude can be related to the Duration Magnitude performing a linear fit (see the next plot), which provides the following relation:

$$M_{WA} = 0.682 + 0.655 M_D$$



As we know the relation between the Duration Magnitude and  $\log \tau$ , we can combine the two formulas to obtain a final relation between the Wood Anderson Magnitude and the duration of the earthquake:

$$M_{WA} = 0.682 + 0.655 * (2.75 \log \tau - 2.35) = 1.8 \log \tau - 0.9$$