



**Comparative proteome analysis of metabolic proteins from seeds of durum wheat (cv. Svevo) subjected to heat stress**

Journal:	<i>PROTEOMICS</i>
Manuscript ID:	Draft
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	
Complete List of Authors:	Laino, Paolo; University of Tuscia, Agrobiology and Agrochemistry Shelton, Dale; DTU, Systems Biology Finnie, Christine; DTU, Systems Biology De Leonardis, Anna Maria; CRA, Cereal Research Centre Mastrangelo, Anna Maria; CRA, Cereal Research Centre Svensson, Birte; DTU, Systems Biology Lafiandra, Domenico; University of Tuscia, Agrobiology and Agrochemistry Masci, Stefania; University of Tuscia, Agrobiology and Agrochemistry
Key Words:	Durum wheat, Heat stress, Wheat kernel proteome, Wheat kernel metabolic proteins



1  
2  
3 **Comparative proteome analysis of metabolic proteins from seeds of durum**  
4  
5  
6 **wheat (cv. Svevo) subjected to heat stress**  
7  
8  
9

10  
11  
12 *Laino Paolo*<sup>1,2</sup>, *Shelton Dale*<sup>2φ</sup>, *Finnie Christine*<sup>2</sup>, *De Leonardis Anna Maria*<sup>3</sup>, *Mastrangelo Anna*  
13  
14 *Maria*<sup>3</sup>, *Svensson Birte*<sup>2</sup>, *Lafiandra Domenico*<sup>1</sup>, *Masci Stefania*<sup>1\*</sup>  
15  
16  
17

18  
19  
20  
21  
22 <sup>1</sup> Department of Agrobiological and Agrochemistry, University of Tuscia, Via S. Camillo De Lellis  
23  
24 snc, 01100 Viterbo, Italy  
25  
26

27 <sup>2</sup>Enzyme and Protein Chemistry, Department of Systems Biology, Technical University of  
28  
29 Denmark, Søtofts Plads Building 224, DK-2800 Kgs. Lyngby, Denmark  
30  
31

32  
33 <sup>3</sup> C.R.A. - Cereal Research Centre, S.S. 16 km 675, 71122 Foggia, Italy  
34  
35

36 <sup>φ</sup>Present address: Institute for Plant Biology, Faculty of Life Science, University of Copenhagen,  
37  
38 Thorvaldsensvej 40, 1841 Frederiksberg C, Denmark  
39  
40  
41

42  
43  
44  
45 **\*Corresponding author:** Prof. Stefania Masci, DABAC, University of Tuscia, Via S. C. de Lellis  
46  
47 snc, 01100 Viterbo (Italy), e-mail: [masci@unitus.it](mailto:masci@unitus.it), fax: 0039 0761 357238, tel. 0039 0761 357255  
48  
49  
50

51  
52  
53  
54 **Abbreviations:** FDR, false discovery rate; GAPDH, glyceraldehyde-3-phosphate dehydrogenase;  
55  
56 LEA, Late Embryogenesis Abundant; NDPK, Nucleoside diphosphate kinase  
57  
58  
59  
60

**Keywords:** Durum Wheat; Heat Stress; Wheat Kernel Proteome; Wheat Kernel Metabolic Proteins

**ABSTRACT**

In Central and Southern Italy, where durum wheat represents one of the most widely cultivated crops, grain filling occurs during Spring, a period characterised by sudden increases in temperature.

Wheat grain proteins are classified into albumins, globulins, and prolamins. The non-prolamin fractions include proteins with metabolic activity or structural function.

In order to investigate the consequences of heat stress on the accumulation of non-prolamin proteins in mature durum wheat kernels, the Italian cultivar Svevo was subjected to two thermal regimes (heat stress *vs.* control) during grain filling. The 2D patterns of non-prolamin proteins were monitored to identify polypeptides affected by heat stress. This study shows that heat stress alters significantly the durum wheat seed proteome, although the fold changes range only between 1.2 and 2.2. This analysis revealed 132 differentially expressed polypeptides, 47 of which were identified by MALDI TOF and MALDI-TOF-TOF MS and included heat shock proteins, proteins involved in the glycolysis and carbohydrate metabolism, as well as stress related proteins. Many of the heat induced polypeptides are considered to be allergenic for sensitive individuals.

The differences observed with previously reported data regarding bread wheat may be explained by the absence of the D genome in durum wheat.

## INTRODUCTION

1  
2  
3  
4  
5  
6  
7  
8  
9  
10 Wheat is a widely used cereal for human consumption. The two most important cultivated wheat  
11 species are *Triticum aestivum* L. (bread wheat) and *T. durum* Desf. (durum wheat). *T. aestivum* is a  
12 hexaploid species possessing the genomes AABBDD, whereas *T. durum* is tetraploid with genome  
13 composition AABB. This last species is extremely important, in the Mediterranean areas, where  
14 represents the most widespread crop, finding large utilization in the production of a wide range of  
15 end products, such as pasta, leavened and unleavened breads, cous cous etc. Genes present on D  
16 chromosomes, such as those determining flour texture [1] or those encoding specific high molecular  
17 weight glutenin subunits [2], missing in durum wheat, confer the typical bread-making properties to  
18 bread wheat flour. In addition, the absence of the D genome in durum wheat may affect also the  
19 response to biotic and abiotic stresses, because it has been shown that several stress-related genes  
20 are located on the D genome [3,4].

21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36 Abiotic stresses, such as drought, salinity, extreme temperatures, chemical toxicity, and oxidative  
37 stress are serious threats to agriculture because they affect quality properties, including yield, and  
38 the characteristics of the final product. These effects stem from altered synthesis and functionality  
39 of specific protein [5]. Proteome analysis is an effective tool for investigation of changes in protein  
40 accumulation in wheat kernel in response to heat stress [6].

41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
The seed protein content is the most important factor determining wheat quality. Wheat kernel  
proteins are divided according to their solubility properties into prolamins (gliadins and glutenins,  
collectively known as gluten proteins), soluble in diluted acid or alkali or alcohol-water mixtures,  
and albumins and globulins, which are water and salt soluble, respectively [7]. Gluten proteins  
represent about 80% of wheat seed proteins, and are the most important determinant of the dough  
properties.

1  
2  
3 High temperature during the grain filling has been reported to alter the yield and the dough quality  
4 of bread wheat [8,9]. Heat stress, in fact, modifies the ratio between the different gluten proteins in  
5 the seed by affecting mainly the composition of the polymeric fraction (soluble/insoluble  
6 polypeptides) [10]. High temperatures thus favor synthesis of gliadins, whereas glutenin synthesis  
7 decreases. An explanation of the gliadin increase after heat stress might be related to the presence of  
8 heat stress elements (HSE) in the upstream regions of some gliadin genes [10]. Because glutenins,  
9 however, are the most important polypeptides in determining quality properties, their decrease,  
10 coupled to the relative increase of gliadins, might explain the negative performance of bread wheat  
11 exposed to high temperatures.  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

24 Water-soluble albumins and salt-soluble globulins constitute about 20% of total flour protein  
25 [11,12]. The non-prolamin proteins have mainly metabolic activity or structural functions [13].  
26  
27  
28

29 Certain wheat globulins, however, are reported to be storage proteins [14,15]. In contrast to gliadins  
30 and glutenins, albumins and globulins have been less thoroughly characterized, most probably due  
31 to their minor role in wheat quality compared to gluten proteins, although the ratio of albumin to  
32 globulin has been reported to correlate with bread-making quality [11]. A relationship between the  
33 water-soluble  $\alpha$ -amylase/trypsin inhibitors and pasta quality has also been suggested [16,17]. The  
34 soluble proteins from wheat seeds are now receiving increasing attention because several  
35 polypeptides from this fraction have been identified as human allergens [18].  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45

46 Effects of heat stress on metabolic proteins in wheat kernel were previously studied in bread wheat  
47 kernels. One of the consequences was a reduction in starch accumulation and activity of soluble  
48 starch synthase [19-21]. More recently, proteome analysis of bread wheat showed the involvement  
49 of enzymes in starch biosynthesis, e.g. glucose-1-phosphate adenylyltransferase and the granule  
50 bound starch synthase in heat stress response. In addition, up-regulation of a large number of  
51 cytoplasmic heat shock proteins (HSP) was confirmed [6].  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Very recent investigations have shown that environmental changes, such as heat, cold, drought, and  
4  
5 waterlogging mainly affect storage proteins of the 7S globulin (vicilin-like) and R-globulin  
6  
7 families, along with defense proteins such as serpins and chitinases [22].  
8  
9

10 Durum wheat is mostly grown in the Mediterranean region where heat stress during grain filling is a  
11  
12 likely event. In order to understand the response of the metabolic protein fraction to heat stress and  
13  
14 make a comparison with bread wheat, we have used proteome analysis to identify the durum wheat  
15  
16 soluble seed proteins that are influenced by heat stress and which may affect the nutritional and  
17  
18 technological quality of the derived products.  
19  
20  
21  
22  
23  
24

## 25 MATERIALS AND METHODS

### 26 27 28 29 **Plant material and heat treatment**

30  
31  
32  
33 Durum wheat (*cv* Svevo) was grown in a climate chamber in a medium composed of soil, sand, peat  
34  
35 (6:3:1) at 10°C (9 h day)/ 7°C (15 h night) with 60% relative humidity and photon flux of 500  $\mu\text{mol}$   
36  
37  $\text{m}^{-2} \text{s}^{-1}$  until appearance of the third leaf. At this point, conditions were gradually (according to the  
38  
39 development stage) switched to 20°C (13 h day)/ 17°C (11 h night) with 55% relative humidity and  
40  
41 photon flux of 500  $\mu\text{mol} \text{m}^{-2} \text{s}^{-1}$ . These conditions were maintained until five days after anthesis.  
42  
43  
44

45 Then, while control plants were maintained in the same conditions, stressed plants were subject to  
46  
47 heat-shock treatment, carried out at 37/17 °C (13 h day/11 h night) with 55% relative humidity for 5  
48  
49 days. Following the heat shock, the temperature was decreased to 28°C for 4 h, and then the  
50  
51 growing cycle was set up at 20°C (13 h day)/ 17°C (11 h night) with 55% relative humidity and 500  
52  
53  $\mu\text{mol} \text{m}^{-2} \text{s}^{-1}$  photon flux. Starting from the milk maturity stage, both control and stressed plants  
54  
55 were brought to complete maturity at 25°C (16 h day)/ 20°C (8 h night) with 45% relative humidity.  
56  
57  
58  
59  
60 In both control and stress treatments, seeds were collected from four biological replicas.

## Protein content analysis and selective extraction

The seed protein content was determined by Kjeldahl nitrogen analysis (Nx5.7). Metabolic seed proteins were extracted according to [23], but amounts were scaled up in order to perform larger 2D gels. Briefly flour (150 mg) was suspended in 600  $\mu$ l of cold KCl buffer (50 mM Tris-HCl, 100 mM KCl, 5 mM EDTA, pH 7.8). The suspension was incubated on ice for 5 min with intermittent mixing and centrifuged (14,500g, 15 min, 4 °C). The KCl-soluble fraction was collected and 5 volumes (v/v) of cold 0.1 M ammonium acetate in methanol were added at room temperature. Following incubation overnight at -20 °C, the methanol-insoluble fraction was pelleted by centrifugation as above. The pellet (containing metabolic proteins) was rinsed with cold acetone, dried down and stored at -20°C until further use.

## 2-DE

Metabolic proteins were dissolved in 800  $\mu$ l of strip rehydration buffer containing 7 M Urea, 2 M thiourea, 2% (w/v) CHAPS, 2% (v/v) Triton X100, 1.2% (v/v) Destreak reagent (GE Healthcare), and 0.5% (v/v) IPG buffer pH 3–10. IEF linear IPG strips (18 cm, GE Healthcare) pH 3–10 were used as first dimension. Strips were rehydrated overnight at 20°C with 200  $\mu$ L of dissolved proteins (about 300  $\mu$ g protein) and 150  $\mu$ l of rehydration buffer was added. The same extract volume (200  $\mu$ L) was loaded. Focusing was performed at 20°C for 80 kVh (200 V 4 h, 500 V 2.5 h, 1000 V 3 h, 5000 V 2 h, 8000 V 9 h). The gel strips were subsequently equilibrated for 25 min in 0.1 M Tris-HCl pH 8.8 containing 6 M Urea, 30% (w/v) glycerol, 2% (w/v) SDS, 1% (w/v) DTT and bromophenol blue as tracking dye. For the second dimension, the strips were transferred onto 18  $\times$  20 cm polyacrylamide gels (15%T, 1.28%C) (Protean II X-Cell, BioRad) and run at 40 mA *per gel* for 3–4 h at 11°C until the dye front left the gel.

1  
2  
3 After electrophoresis, proteins were visualized by CBB staining according to [24]. All gels were  
4 stained for 16 h and destained for 1 h with distilled water before image acquisition. Three technical  
5 replicas were performed for each of the four biological replicas from each treatment, giving a total  
6 of 24 gels. Destained gels were scanned with Image Master LabScan (GE Healthcare) and analyzed  
7 using the software SameSpots Progenesis (vers. 1.0.2602.33289, Nonlinear Dynamics, UK). This  
8 software includes statistical analyses such as ANOVA ( $p \leq 0.05$ ), determination of False Discovery  
9 Rate (FDR,  $q \leq 0.05$ ), and Principle Component Analysis (PCA) calculated according to [25].  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19

### 20 21 22 **Protein identification by MALDI TOF-TOF**

23  
24  
25  
26  
27 Protein identification was done by picking gel spots from two independent gels to minimize  
28 technical variability. CBB-stained spots were excised from the gel, cut to pieces and washed twice  
29 with 40% ethanol until colorless. The destained gel pieces were dehydrated with ACN, treated with  
30 10 mM DTT in 50 mM  $\text{NH}_4\text{HCO}_3$  for 1 h at 56°C and finally alkylated with 55 mM iodoacetamide  
31 in 50 mM  $\text{NH}_4\text{HCO}_3$  for 30 min in the dark. Gel pieces were dehydrated with ACN, rehydrated  
32 with 1 pmol trypsin (Sigma) solution in 50 mM  $\text{NH}_4\text{HCO}_3$  and incubated at 37°C overnight for  
33 digestion. The peptides were extracted twice from gel slices with 5% formic acid in 50% ACN. The  
34 peptide solution was then desalted, concentrated and applied to an Anchorchip target™ (Bruker  
35 Daltonics) using  $\alpha$ -cyano-4-hydroxycinnamic acid as matrix, according to the manufacturer's  
36 instructions and [26]. Mass spectrometric analysis was performed on a MALDI TOF-TOF Ultraflex  
37 II in positive ion reflector mode and spectra were processed and analysed using the software  
38 FlexAnalysis and BioTools (Bruker Daltonics). Database searching was carried out using an in-  
39 house MASCOT server (Matrix Science, London, UK) to search NCBIInr  
40 (<ftp://ftp.ncbi.nih.gov/blast/db/>) and the Wheat Gene Index  
41 (<http://compbio.dfci.harvard.edu/tgi/tgipage.html>) release 10.  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 Proteins were identified by PMF. Matrix contaminants and predominant keratin peaks were  
4 removed from peak lists using PeakErazor  
5  
6  
7  
8 (<http://www.protein.sdu.dk/gpmaw/Help/PeakErazor/peakerazor.html>). In order to identify a protein  
9  
10 unambiguously the following criteria were used: MASCOT score with  $p \leq 0.05$ ; protein sequence  
11 coverage  $> 15\%$ ; at least five independent peptides matching with a mass tolerance of 50 ppm and  
12  
13 maximum one trypsin miscleavage site. The Oxidation (M) and the Gln->pyro-Glu were selected as  
14  
15 variable fragment modifications. Where the PMF-based identification was uncertain, fragment ion  
16  
17 spectra were obtained for at least three peaks with a signal:noise ratio  $> 5$  and  $m/z > 1400$ . Each  
18  
19 fragment ion spectrum was checked against the same database as used for PMF and the  
20  
21 identification was confirmed if correspondence was found.  
22  
23  
24  
25  
26  
27  
28  
29  
30

## 31 RESULTS AND DISCUSSION

32  
33  
34  
35  
36  
37 Heat stress can be an important factor affecting yield and quality of durum wheat Although quality  
38 is mostly determined by gluten proteins, that are the major protein components of wheat seeds, also  
39 the soluble metabolic protein fraction (albumins and globulins) plays a role, especially in terms of  
40  
41 nutritional/antinutritional properties. Most of these polypeptides, in fact, show allergenic properties  
42  
43 in sensitive individuals [18]. Albumins and globulins are distributed mainly in the outer layers of  
44  
45 wheat kernels, and thus are important components in whole-wheat flours and semolina, towards  
46  
47 which there is an increasing consumer interest, because they have a higher amount of fibers,  
48  
49 proteins and functional components [27].  
50  
51  
52  
53  
54  
55

56 In the present paper the durum wheat cultivar Svevo, widely grown in Italy and moderately resistant  
57  
58 to cold stress, was subjected to two thermal regimes, *i.e.* heat stress and control conditions during  
59  
60

1  
2  
3 grain filling. The effect on the metabolic protein fraction was analysed by comparison of the  
4  
5 corresponding proteomes.  
6

7  
8 Firstly, protein contents were compared between stressed and control samples, which revealed that  
9  
10 heat stressed samples accumulated significantly higher amounts of protein. The protein content  
11  
12 being 19.30% ( $\pm 1.43$ ) in stressed and 13.85% ( $\pm 1.87$ ) in control samples ( $p=0.004$ ). A correlation  
13  
14 between heat stress and increase in protein content was previously reported in bread wheat [28]. In  
15  
16 our case, the increase in protein content was due mostly to gluten proteins, that were about 50%  
17  
18 more abundant in heat stressed samples with respect to control samples, whereas the metabolic  
19  
20 fraction increased by about 20% (data not shown).  
21  
22

23  
24 The highly reproducible 2-D gels showed about 1000 spots in the pI range 3–10 and comparison  
25  
26 revealed differential expression of proteins ( $1.2 < \text{fold change} < 2.2$ ) in 132 spots (Fig. 1), 65% were  
27  
28 up-regulated in heat-treated samples (Fig. 2). Because only those spots satisfying ANOVA ( $p < 0.05$ )  
29  
30 and FDR ( $q < 0.05$ ) values were chosen, and the Principal Component Analysis (PCA) performed on  
31  
32 the differentially expressed spots [25] indicated two separate groups corresponding to the two  
33  
34 thermal regimes (Fig. 3), we are confident that the observed differences are significant.  
35  
36  
37  
38  
39  
40

### 41 **Protein identification**

42  
43  
44 MALDI TOF analysis identified 47 proteins from the picked varying spots (Table 1; Fig. 4).

45  
46 Among the identified proteins, 85% were up-regulated and 15% down-regulated. Proteins identified  
47  
48 by mass spectrometry were classified based on to their main activity although most are involved in  
49  
50 different pathways or signaling.  
51  
52

53  
54  
55 Noteworthy is the observation that the fold changes observed, are in a narrow range ( $1.2 > \text{fold}$   
56  
57  $\text{change} > 2.2$ ). This is in agreement with previous findings relative to the bread wheat cultivar Butte  
58  
59 86 [29].  
60

1  
2  
3 As expected, heat stress increased expression of many proteins related to desiccation and oxidation  
4 stress, e.g. Late Embryogenesis Abundant (LEA)-proteins, the heat shock proteins HSP70 and  
5 HSP26 [5, 30-32]. These proteins, through binding or interaction with other proteins, prevent  
6 damage of proteins and cell membranes (reviewed in [33]). Also glyoxalase I, one of the enzymes  
7 of the glyoxalase pathway related to the detoxification pathway of methylglyoxal in plants [34] was  
8 up-regulated.  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21

### 22 *Carbohydrate metabolism or energy related proteins*

23  
24 Nucleoside diphosphate kinase (NDPK), which is required for the synthesis of nucleotide  
25 triphosphate precursors of DNA and RNA, was up-regulated along with some housekeeping  
26 enzymes involved in glycolysis and the pentose phosphate pathway (glyceraldehyde 3-phosphate  
27 dehydrogenase, phosphoglycerate kinase, and glucose and ribitol dehydrogenase), in agreement also  
28 with that reported in [29], in which the effect of high temperature on soluble proteins of developing  
29 bread wheat grains has been studied. Altered expression pattern of NDPK was previously found in  
30 response to abiotic (including heat stress) and biotic stresses in rice and other plant species [35-36],  
31 suggesting that NDPK plays a regulatory role in addition to its primary metabolic functions.  
32  
33 Five spots (427, 444, 460, 503, 514) which were up-regulated by heat stress contained  
34 glyceraldehyde 3-phosphate dehydrogenase (GADPH), which is involved in glycolysis and has been  
35 identified as one of the allergens implicated in bakers' asthma [18]. Bustos and Iglesias [37]  
36 reported that wheat endosperm and shoot GAPDH undergoes posttranslational phosphorylation  
37 enabling interaction with 14-3-3 family proteins, thus exerting a regulation aimed at maintaining the  
38 levels of energy and reductants in the cytoplasm. Recently, it was established using a proteomics  
39 approach that GAPDH activity in *Arabidopsis* was inhibited by H<sub>2</sub>O<sub>2</sub>, suggesting that GAPDH is a  
40 direct target of H<sub>2</sub>O<sub>2</sub> and might have a role in mediating ROS signaling in plants [38].  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Three up-regulated spots (436, 470, 502) belonged to the glucose and ribitol dehydrogenase protein  
4 family.  
5

6  
7  
8 Among the polypeptides down-regulated after heat shock, ATP synthase  $\beta$  subunit (spot 110) was  
9 identified, in agreement with observations in bread wheat [6]. The decrease in amount of ATP  
10 synthase may affect energy-dependent processes involved in heat stress resistance. It has been  
11 reported that energy consuming processes are the primary cellular targets for a decreasing of ATP  
12 demand in response to stresses, such as anoxia [39], and it is known that heat shock and anoxia are  
13 abiotic stresses eliciting similar cellular responses [40].  
14

15 Also phosphoglycerate kinase (spot 109), glycoside hydrolase family 85 (spot 173), and fructose 6  
16 phosphate-phosphotransferase (spot 299), all involved in glycolysis, were down-regulated. The  
17 glycoside hydrolase family is present in essentially all living organisms and has been implicated in  
18 a diversity of roles, such as biomass conversion in microorganisms and activation of defense  
19 compounds, phytohormones, lignin precursors, aromatic volatiles, and metabolic intermediates [41].  
20 Phosphoglycerate kinase was previously found to be over-expressed in the nuclear proteome of  
21 Arabidopsis in the cold stress response [42].  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41

### 42 *Stress related*

43  
44  
45 Two members of the heat shock protein family, HSP70 (spots 70, 426) and HSP26 (spot 497) were  
46 up-regulated after the heat stress. HSP70s have been linked to the development of acquired  
47 thermotolerance in heat stress, although they seem also to correlate with tolerance to low  
48 temperature stress [43]. HSP26 belongs to the family of small heat shock proteins (sHSP) exhibiting  
49 chaperone activity and thought to protect proteins from irreversible aggregation [44].  
50  
51

52  
53  
54 A polypeptide of the 14-3-3 protein family was slightly down-regulated by heat stress (spot 341).  
55 Also Hurkman *et al* [29] observed down-regulation of members of this protein family after heat  
56 stress in bread wheat. Transcripts encoding proteins belonging to the 14-3-3 family accumulate in  
57  
58  
59  
60

1  
2  
3 barley after biotic stresses [45,46] and in wheat a region of chromosome 4AL containing genes  
4  
5 coding for 14-3-3 is associated with a resistance QTL against specific fungal diseases [47]. The up-  
6  
7 regulated polypeptides after heat treatment moreover included flavonoid *O*-methyltransferase (spot  
8  
9 445), a 14-3-3-binding protein involved in production of antimicrobial secondary metabolites ,thus  
10  
11 showing a potential role in response to a pathogen attack in barley [48]. Furthermore these  
12  
13 compounds have antioxidants properties and could thus be involved in oxidative stress.  
14  
15

16  
17  
18 Serpins (*serine protease inhibitors*) were identified in multiple spots (89, 94, 395, 456) as up-  
19  
20 regulated in response to high temperature. The same observation was made by Hurkman *et al* [29].  
21

22  
23 Serpins, found in the soluble fraction of wheat seeds, undergo differential regulation in response to  
24  
25 environmental stress [49]. The serpins are widespread in the plant kingdom and represent up to 4%  
26  
27 of the total protein in the mature endosperm of cereal grains [50,51]. While the precise  
28  
29 physiological role of serpins remains unclear, their activity suggests that they are involved in  
30  
31 inhibition of endogenous proteases, or proteases from grain pests. Because of their high  
32  
33 concentration in the endosperm, serpins have potential to influence grain quality traits [52-54].  
34  
35

36  
37 Other proteins with a defense role have a modified expression profile in response to heat shock.  
38  
39 Such proteins are typically identified in multiple forms on 2-D gels [55] and include: tritins (spots  
40  
41 140, 222),  $\alpha$ -amylase inhibitors (spot 362), and some 14-3-3 related or binding proteins. Most of  
42  
43 these proteins besides having a metabolic role, are also considered as storage proteins. Furthermore,  
44  
45 some are also considered wheat grain allergens along with the serpins [56].  
46  
47

48  
49 Another enzyme found in response to the heat treatment is glyoxalase I (spot 508). The glyoxalase  
50  
51 system is a set of enzymes that carry out detoxification of methylglyoxal and other reactive  
52  
53 aldehydes that are produced during normal metabolism. In plants it has been demonstrated that  
54  
55 different kinds of stress, such as salt and metal stress, elicits enhance the expression of glyoxalase I  
56  
57 [57-59].  
58  
59  
60

1  
2  
3 LEA proteins (spots 483 and 484) were up-regulated and are involved in stress responses. They are  
4  
5 typically correlated with cellular dehydration in response to cold stress. Other LEA functions  
6  
7 include roles as antioxidants and membrane and protein stabilisers during water stress [60].  
8  
9 Finally, 1-Cys peroxiredoxin (spot 14), also observed to be up-regulated, has antioxidant and  
10  
11 chaperone activity [61].  
12  
13  
14  
15  
16  
17  
18

### 19 *Storage proteins*

20  
21 Up-regulated proteins with a storage role here identified in the heat response mainly belong to the  
22  
23 globulin-like protein family (spots 110, 232 350, 509, 511) or protein homologous to embryo  
24  
25 globulin (spots 58, 239, 468, 505). Spot 239 (Homologue to Embryo Globulin) was found to be  
26  
27 down-regulated by heat stress. A role for these proteins in thermotolerance is not known. They may  
28  
29 be directly related to heat stress response by an unknown mechanism or, indirectly, be a target for  
30  
31 other proteins involved in the heat shock response. This is in agreement with [29].  
32  
33  
34  
35  
36  
37

### 38 **CONCLUDING REMARKS**

39  
40  
41 The results here reported offer a picture of the consequences of heat stress occurring during grain  
42  
43 filling on the accumulation of the soluble proteins in the mature seeds of the widely grown Italian  
44  
45 durum wheat *cv* Svevo. The results provide a basis for understanding how this environmental  
46  
47 change influences on protein synthesis and consequently the metabolic and quality traits of the  
48  
49 durum wheat kernel. Moreover, since the durum wheat kernel is of primary interest because the  
50  
51 semolina obtained by crushing mature seeds is the basis of many products of common use, among  
52  
53 which pasta is the most important, it is critical to understand if the different types of stresses that  
54  
55 wheat plants can potentially undergo, may alter protein composition, and, consequently, qualitative  
56  
57 and nutritional properties of the derived products.  
58  
59  
60

1  
2  
3 In the present study 47 metabolic proteins were identified from the soluble seed fraction of durum  
4 wheat which were induced or repressed by heat shock. The fold changes observed were between 1.2  
5 and 2.2, and this was also in agreement with [29].  
6  
7

8  
9  
10 In agreement with previous studies performed in bread wheat cultivars [6,29,62] both HSP70 and  
11 LEA proteins were up-regulated in response to heat stress. Other proteins, such as ATP synthase  $\beta$   
12 subunit and nucleoside diphosphate kinase (NDPK) related to the energy metabolism also  
13 responded to heat treatment. This agrees also with the findings of Hurkman *et al* [29]. Interestingly,  
14 we did not find any difference in the level of expression of starch related enzymes, although this  
15 was a finding of Majoul *et al* [6] in bread wheat. This might be a result either of the different  
16 protein extraction procedures, or of a different protein turnover in the plant material, or the enzyme  
17 forms that are regulated in bread wheat may be coded by genes present on the D genome, that is  
18 absent in durum wheat.  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

31 GAPDH, a housekeeping enzymes involved in glycolysis, was found to be up-regulated. Others  
32 recently suggested a relationship between this enzyme and certain abiotic stresses [63]. Proteins  
33 reported to be influenced by abiotic stresses, e.g. oxidative and drought stresses, and also found as  
34 differentially regulated in the present work include 14-3-3 proteins, serpins, LEA proteins, 1-Cys  
35 peroxiredoxin, glyoxalase I, and proteins with a storage function (e.g. globulin-like proteins).  
36  
37  
38  
39  
40  
41  
42

43 It is noteworthy that some of the differentially regulated proteins are considered as allergens ( $\alpha$ -  
44 amylase inhibitors, serpins, tritins, GAPDH) and were found to be up-regulated after heat stress,  
45 which obviously represents a disadvantage for sensitive individuals.  
46  
47  
48  
49

50 These results illustrate that there is a common network “response” to different types of abiotic  
51 stress, such as drought, oxidative, cold and heat stress.  
52  
53  
54  
55  
56  
57  
58  
59  
60

## ACKNOWLEDGMENTS

1  
2  
3  
4  
5  
6  
7  
8 This work was supported by the National project “Proteine e geni per la protezione delle piante  
9 dagli stress biotici e abiotici (PROTEO-STRESS), MIPAF. Paolo Laino was the recipient of a PhD  
10 fellowship in Plant Biotechnology funded by MIUR, "Fondo per il sostegno dei giovani e per  
11 favorire la mobilità degli studenti" in the framework “Valorizzazione dei prodotti tipici  
12 dell’agroalimentare e sicurezza alimentare attraverso nuovi sistemi di caratterizzazione e garanzia  
13 di qualità”, and the recipient of a STMS award from the COST Action FA0603 Plant Proteomics.  
14  
15 Birgit Andersen (Technical University of Denmark) is thanked for assistance with mass  
16 spectrometry. Dale Shelton was funded by a Large Multi-Disciplinary Research Grant from the  
17 Danish Research Council. The authors wish to acknowledge Dr. Federico Scossa (ENEA-Casaccia,  
18 Rome, Italy) for having performed protein content measurements.  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



**Conflict of interest statement**

The authors declare that there is no financial/commercial conflict of interest

For Peer Review

## REFERENCES

- 1  
2  
3  
4  
5  
6  
7 [1] Morris C.F., Puroindolines: the molecular genetic basis of wheat grain hardness. *Plant Molec.*  
8  
9 *Biol.* 2002, 48, 633–647.
- 10  
11 [2] Shewry, P.R., Halford, N.G., Tatham, A.S., Popineau, Y. *et al.*, The high molecular weight  
12  
13 subunits of wheat glutenin and their role in determining wheat processing properties. *Adv. Food*  
14  
15 *Nutr. Res.* 2003, 45, 219-302.
- 16  
17  
18 [3] Boyko E.V., Gill K. S., Mickelson-Young L., Nasuda S. *et al.*, A high-density genetic linkage  
19  
20 map of *Aegilops tauschii*, the D-genome progenitor of bread wheat. *Theor. Appl. Genet.* 1999,  
21  
22 99, 16-26.
- 23  
24  
25 [4] Aprile, A., Mastrangelo, A.M., De Leonardis, A.M., Galiba, G. *et al.*, Transcriptional profiling  
26  
27 in response to terminal drought stress reveals differential responses along the wheat genome.  
28  
29 *BMC Genomics* 2009, 10, art. no. 279.
- 30  
31  
32 [5] Timperio, A.M., Egidi, M.G., Zolla, L., Proteomics applied on plant abiotic stresses: Role of  
33  
34 heat shock proteins (HSP). *J. Proteomics* 2008, 71, 4, 391-411.
- 35  
36  
37 [6] Majoul, T., Bancel, E., Triboï, E., Hamida, J.B., Branlard, G., Proteomic analysis of the effect of  
38  
39 heat stress on hexaploid wheat grain: Characterization of heat-responsive proteins from non-  
40  
41 prolamins fraction. *Proteomics* 2004, 4, 505-513.
- 42  
43  
44 [7] Osborne, T.B., The vegetable proteins. Longmans, Green & Co., London, 1924.
- 45  
46  
47 [8] Stone, P.J., Nicolas, M.E., Wheat cultivars vary widely in their responses of grain yield and  
48  
49 quality to short periods of post- anthesis heat stress. *Aust. J. Plant Physiol.* 1994, 21, 887-900.
- 50  
51  
52 [9] Wardlaw, I.F., Blumenthal, C., Larroque, O., Wrigley, C.W., Contrasting effects of chronic heat  
53  
54 stress and heat shock on kernel weight and flour quality in wheat. *Funct. Plant Biol.* 2002, 29,  
55  
56 25–34.
- 57  
58  
59 [10] Blumenthal, C.S, Batey, I.L., Wrigley, C.W., Barlow, E.W.R., Involvement of a novel peptide  
60  
in the heat shock response of Australian wheats. *Aust. J. Plant Physiol.* 1990, 17, 441-449.

- 1  
2  
3 [11] Pence, J.W., Weinstein, N.E., Mecham, D., The albumin and globulin contents of wheat flour  
4 and their relationship to protein quality. *Cereal Chem.* 1954, *31*, 303–311.  
5  
6  
7  
8 [12] Singh, J., Skerritt, J. H., Chromosomal control of albumins and globulins in wheat grain  
9 assessed using different fractionation procedures. *J. Cereal. Sci.* 2001, *33*, 163-181.  
10  
11  
12 [13] Goesaert, H., Gebruers, K., Courtin, C.M., Brijs, K., Delcour, J.A., Enzymes in breadmaking.  
13 In “Bakery products: science and technology”, Hui Y.H. Ed. 2006, pp. 337-364.  
14  
15  
16 [14] Singh, N.K., Shepherd, K.W., The structure and genetic control of a new class of disulphide-  
17 linked proteins in wheat endosperm. *Theor. Appl. Genet.* 1985, *71*, 79–92.  
18  
19  
20 [15] Singh, N.K., Shepherd, K.W., Solubility behavior synthesis, degradation and subcellular  
21 location of a new class of disulfide linked proteins in wheat endosperm. *Aust. J. Plant Physiol.*  
22 1987, *14*, 245–252.  
23  
24  
25 [16] Kobrehel, K., Alary, R., The role of a low molecular weight glutenin fraction in the cooking  
26 quality of durum wheat pasta. *J. Sci. Food Agric.* 1989, *47*, 487–500.  
27  
28  
29 [17] Kobrehel, K., Alary, R., Isolation and partial characterization of two low molecular weight  
30 durum wheat (*Triticum durum*) glutenins. *J. Sci. Food Agric.* 1989, *48*, 441–452.  
31  
32  
33 [18] Tatham, A.S., Shewry, P.R., Allergens to wheat and related cereals. *Clin. Exp. Allergy*, 2008,  
34 38, 1712-1726.  
35  
36  
37 [19] Shy, Y.C., Seib, P.A., Bernardin, J.E., Effects of temperature during grain-filling on starches  
38 from six wheat cultivar. *Cereal Chem.* 1994, *71*, 369-383.  
39  
40  
41 [20] Hawker, J.S., Jenner, C.F., High temperature affects the activity of enzymes in the committed  
42 pathway of starch synthesis in developing wheat endosperm. *Austr. J. Plant Physiol.* 1993, *20*,  
43 197-209.  
44  
45  
46 [21] Keeling, P.L., Bacon, P.J., Holt, D.C., Elevated temperature reduces starch deposition in wheat  
47 endosperm by reducing the activity of soluble starch syntase. *Planta* 1993, *191*, 342-348.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- [22] Sancho, A.I., Gillibert, M., Tapp, H., Shewry, P.R., *et al.*, Effect of environmental stress during grain filling on the soluble proteome of wheat (*Triticum aestivum*) dough liquor. *Agr. Food Chem.* 2008, 56, 5386-5393.
- [23] Hurkman, W.J., Tanaka, C.K., Improved methods for separation of wheat endosperm proteins and analysis by two-dimensional gel electrophoresis. *J. Cereal Sci.* 2004, 40, 295–299.
- [24] Neuhoff, V., Arold, N., Taube, D., Ehrhardt, W., Improved staining of proteins in polyacrylamide gels including isoelectric focusing gels with a clear background at nanogram sensitivity using Coomassie Brilliant Blue G-250 and R-250. *Electrophoresis* 1988, 9, 255–262.
- [25] O’Gorman, M., Beauvallet, C., Lepercq, P. David, O., *et al.*, An investigation into Crohn’s disease using the Progenesis SameSpots analysis platform. *24<sup>th</sup> Journée Françaises de Spectrométrie de Masse, PAU 2007.*
- [26] Zhang, X., Shi, L., Shu, S., Whang, Y., *et al.*, An improved method of sample preparation on AnchorChip™ targets for MALDI-MS and MS/MS and its application in the liver proteome projects. *Proteomics* 2007, 7, 2340-2349.
- [27] Dewettinck, K., Van Bockstaele, F., Kühne, B., Van de Walle, D., *et al.*, Nutritional value of bread: influence of processing, food interaction and consumer perception. *J. Cereal Sci.* 2008, 48, 243-257.
- [28] Shewry, P.R., Napier, J.A., Tatham, A.S., Seed storage proteins: structures and biosynthesis. *Plant Cell* 1995, 7, 945-956.
- [29] Hurkman, W.J., Vensel, W.H., Tanaka, C.K., Whitehand, L., Altenbach, S.B., Effect of high temperature on albumin and globulin accumulation in the endosperm proteome of the developing wheat grain. *J. Cereal Sci.* 2009, 49, 12-23.
- [30] Treglia, A., Spano, G., Rampino, P., Giangrande, E., *et al.*, Identification by in vitro translation and Northern blot analysis of heat shock mRNAs isolated from wheat seeds exposed to different temperatures during ripening. *J. Cereal Sci.* 1999, 30, 33-38.

- 1  
2  
3 [31] Renaut, J., Hausman, J.-F., Wisniewski, M.E., Proteomics and low-temperature studies:  
4  
5 bridging the gap between gene expression and metabolism. *Physiol. Plant.* 2006, *126*, 97-109.  
6  
7  
8 [32] Qureshi, M.I., Qadir, S., Zolla, L., Proteomics-based dissection of stress-responsive pathways  
9  
10 in plants. *J. Plant Physiol.* 2007, *164*, 1239-1260.  
11  
12  
13 [33] Buitink, J., Hoekstra, F.A., Leprince, O., Biochemistry and biophysics of tolerance systems.  
14  
15 Black, M and Pritchard, H.W. Eds., CAB International, Wallingford, UK., 2002.  
16  
17 [34] Hossain, M.A., Hossain, M.Z., Fujita, M., Stress-induced changes of methylglyoxal level and  
18  
19 glyoxalase I activity in pumpkin seedlings and cDNA cloning of glyoxalase I gene. *Aust. J.*  
20  
21 *Crop Sci.* 2009, *3*, 53-64.  
22  
23  
24 [35] Salekdeh, Gh.H., Siopongco, J., Wade, L.J., Ghareyazie, B., Bennett, J. Proteomic analysis of  
25  
26 rice leaves during drought stress and recovery. *Proteomics* 2002, *9*, 1131-1145.  
27  
28  
29 [36] Hajheidari, M., Abdollahian-Noghabi, M., Askari, H., Heidari, M., *et al.*, Proteome analysis of  
30  
31 sugar beet leaves under drought stress. *Proteomics* 2005, *4*, 950-960.  
32  
33  
34 [37] Bustos, D.M., Iglesias, A.A., Phosphorylated Non-Phosphorylating Glyceraldehyde-3-  
35  
36 Phosphate Dehydrogenase from heterotrophic cells of wheat interacts with 14-3-3 proteins.  
37  
38 *Plant Physiol.* 2003, *133*, 2081-2088.  
39  
40  
41 [38] Hancock, J.T., Henson, D., Nvirenda M., Desikan R., *et al.*, Proteomic identification of  
42  
43 glyceraldehyde 3-phosphate dehydrogenase as an inhibitory target of hydrogen peroxide in  
44  
45 *Arabidopsis*. *Plant Physiol. Biochem.* 2005, *43*, 828–835.  
46  
47  
48 [39] Boutilier, R.G., St-Pierre, J., Surviving hypoxia without really dying. *Comp. Biochem. Physiol.*  
49  
50 2000, *126A*, 481-490.  
51  
52  
53 [40] Wu, B.S., Lee, J.K., Thompson, K.M., Walker, V.K., *et al.*, Anoxia induces thermotolerance in  
54  
55 the locust flight system. *J. Exp. Biol.* 2002, *205*, 815–827.  
56  
57  
58 [41] Opassiri, R., Pomthong, B., Onkoksoong, T., Akiyama, T., *et al.*, Analysis of rice glycosyl  
59  
60 hydrolase family 1 and expression of Os4bglu12  $\beta$ -glucosidase. *BMC Plant Biol.* 2006, *6*, art.  
no. 33.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- [42] Bae, M.S., Cho, E.J., Choi, E.Y., Park, O.K., Analysis of the *Arabidopsis* nuclear proteome and its response to cold stress. *Plant J.*, 2003, *36*, 652–663.
- [43] Guy, C.L. Freezing tolerance of plants: Current understanding and selected emerging concepts. *Can. J. Bot.* 2003, *81*, 1216-1223.
- [44] Nguyen, H.T., Weng, J., Joshi, C.P., A wheat (*Triticum aestivum*) cDNA clone encoding a plastid-localized heat-shock protein. *Plant Physiol.* 1993, *103*, 675-676.
- [45] Brandt, J., Thordal-Christensen, H., Vad, K., Gregersen, P.L. and Collinge, D.B., A pathogen-induced gene of barley encodes a protein showing high similarity to a protein kinase regulator. *Plant J.* 1992, *2*, 815–820.
- [46] Gregersen, P.L., Thordal-Christensen, H., Forster, H., Collinge, D.B., Differential gene transcript accumulation in barley leaf epidermis and mesophyll in response to attack by *Blumeria graminis* f.sp. *hordei* (syn. *Erysiphe graminis* f.sp. *hordei*). *Physiol. Mol. Plant Pathol.* 1997, *51*, 85–97.
- [47] Faris, J.D., Li, W.L., Liu, D.J., Chen, P.D., Gill, B.S., Candidate gene analysis of quantitative disease resistance in wheat. *Theor. Appl. Genet.* 1999, *98*, 219–225.
- [48] Christensen, A.B., Gregersen, P.L., Olsen, C.E., Collinge, D.B., A flavonoid 7-O-methyltransferase is expressed in barely leaves in response to pathogen attack. *Plant Mol. Biol.* 1998, *36*, 219-227.
- [49] Sancho, A.I., Gillabert, M., Tapp, H., Shewry, P.R., *et al.*, Effect of environmental stress during grain filling on the soluble proteome of wheat (*Triticum aestivum*) dough liquor. *J. Agric. Food Chem.* 2008, *56*, 5386–5393.
- [50] Rasmussen, S.K., Klausen, J., Hejgaard, J., Svensson, B., Svendsen, I., Primary structure of the plant serpin BSZ7 having the capacity of chymotrypsin inhibition. *Biochim. Biophys. Acta* 1996, *1297*, 127-130.
- [51] Oøstergaard, H., Rasmussen, S.K., Roberts, T.H., Hejgaard, J., Inhibitory serpins from wheat grain with reactive centers resembling glutamine-rich repeats of prolamin storage proteins.

- 1  
2  
3 Cloning and characterization of five major molecular forms. *J. Biol. Chem.* 2000, 275, 33272-  
4 33279.  
5  
6  
7  
8 [52] Roberts, T.H., Marttila, S., Rasmussen, S.K., Hejgaard, J., Differential gene expression for  
9 suicide-substrate serine proteinase inhibitors (serpins) in vegetative and grain tissues of barley.  
10  
11 *J. Exp. Bot.* 2003, 54, 2251-2263.  
12  
13  
14 [53] Salt, L.J., Robertson, J.A., Jenkins, J.A., Mulholland, F., Mills, E.N.C., The identification of  
15 foam-forming soluble proteins from wheat (*Triticum aestivum*) dough. *Proteomics* 2005, 5,  
16 1612-1623.  
17  
18  
19 [54] Roberts, T.H., Hejgaard, J., Serpins in plants and green algae. *Funct. Integr. Genom.* 2008, 1,  
20 1-27.  
21  
22  
23 [55] Østergaard, O., Finnie, C., Laugesen, S., Roepstorff, P., Svensson, B., Proteome analysis of  
24 barley seeds: Identification of major proteins from two-dimensional gels (pI 4-7). *Proteomics.*  
25 2004, 4, 2437-2447  
26  
27 [56] Šotkovský, P., Hubálek, M., Hernychová, L., Novák, P., *et al.*, Proteomic analysis of wheat  
28 proteins recognized by IgE antibodies of allergic patients. *Proteomics* 2008, 8, 1677-1691.  
29  
30 [57] Veena, Reddy V.S., Sopory S.K., Glyoxalase I from *Brassica juncea*: molecular cloning,  
31 regulation and its over-expression confer tolerance in transgenic tobacco under stress. *Plant J.*  
32 1999, 17, 385-395.  
33  
34 [58] Singla-Pareek, S.L., Reddy, M.K., Sopory, S.K., Genetic engineering of the glyoxalase  
35 pathway in tobacco leads to enhanced salinity tolerance. *PNAS* 2003, 100, 14672-14677.  
36  
37 [59] Yadav, S.K., Singla-Pareek, S.L., Reddy, M.K., Sopory, S.K., Transgenic tobacco plants  
38 overexpressing glyoxalase enzymes resist an increase in methylglyoxal and maintain higher  
39 reduced glutathione levels under salinity stress. *FEBS Letters* 2005, 579, 6265-6271.  
40  
41 [60] Tunnacliffe, A., Wise, M.J. The continuing conundrum of the LEA proteins.  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

[61] Bystrova, M.F., Budanova, E.N., Novoselov, V.I., Fesenko, E.E., Analysis of the oligomeric state of rat 1-Cys peroxiredoxin. *Biophysics* 2007, 52, 277–281.

[62] Majoul, T., Bancel, E., Triboï, E., Ben Hamida, J., Branlard, G., Proteomic analysis of the effect of heat stress on hexaploid wheat grain: Characterization of heat-responsive proteins from total endosperm. *Proteomics* 2003, 3, 175–183.

[63] Bustos, D.M., Bustamante C.A., Iglesias A.A., Involvement of non-phosphorylating glyceraldehyde-3 phosphate dehydrogenase in response to oxidative stress. *J. Plant Physiol.* 2008, 165, 456-461.

For Peer Review



**Tab.1** Proteins responsive to heat shock identified by PMF and MS-MS analysis. The following criteria were used: MASCOT score ( $p < 0.05$ ), minimal coverage 13%, at least 6 independent peptides should match with a mass tolerance of 50 ppm and 1 miss cleavage site. The database search was applied in NCBI nr and TIGR wheat. For all the identifications three peptide fragmentations were done (MS-MS) and used in combination in the database search. The ANOVA value for each spot, the fold change of the normalized volumes between the spots in the control and heat stressed maps, the proteins names, accession numbers, the MASCOT scores, the protein coverage and the theoretical molecular weight of the proteins identified are reported.

Spot	Anova	Fold Change	Protein	Accession	Species	Score/coverage%	Theor Mw (kDa)
<b>Up-regulated proteins</b>							
14	1.58E-05	1.7	1-Cys peroxiredoxin PER 1	gi/1710077	<i>T. durum</i>	87/35	24.1
42	0.004	1.2	Isoflavone reductase homolog	TC235506		148/30	42.8
58	1.21E-08	2.5	Homologue to Embryo globulin	TC234172		135/27	71.3
70	3.58E-05	1.6	HSP70	gi/2827002	<i>T. aestivum</i>	83/21	71.4
89	0.001	2.2	Serpin	gi/5734506	<i>T. aestivum</i>	96/21	43.3
94	4.61E-04	2	Serpin	gi/5734506	<i>T. aestivum</i>	109/33	43.3
114	0.035	1.3	Nucleoside diphosphate Kinase	TC262718		96/26	19
222	2.45E-04	1.4	Tritin	gi/147744620	<i>T. aestivum</i>	111/38	29.5
232	0.002	1.9	Globulin-like protein	TC246874		356/45	71.1


1								
2	336	0.015	1.3	Globulin-like protein 57%	TC246703		133/22	68.2
3								
4	350	0.002	1.4	Globulin-like protein	TC246874		76/18	71.1
5								
6	362	0.008	1.5	Endogenous $\alpha$ amylase inhibitor (WASI)	gi/123975	<i>H. vulgare</i>	232/85	14.7
7								
8								
9	390	0.033	1.2	Globulin-like protein	TC246703		257/27	68.2
10								
11	395	0.033	1.3	Serpin	gi/5734506	<i>T. aestivum</i>	74/29	43.3
12								
13	413	0.008	1.4	Globulin-like protein	TC246703		133/19	68.2
14								
15	426	0.014	1.2	HSP70	gi/476003	<i>H. vulgare</i>	87/30	71.4
16								
17	427	0.002	1.2	GAPDH cytosolic	gi/32478662		103/53	18.2
18								
19								
20	436	0.023	1.5	Glucose and ribitol dehydrogenase	TC233140		113/17	31.6
21								
22								
23	444	0.007	1.3	GAPDH	gi/148508784	<i>T. aestivum</i>	125/42	36.6
24								
25	445	0.027	1.5	Flavonoid 7-O-Methyltransferase-like (52%)	TC252404		141/26	49.7
26								
27								
28	447	0.026	1.3	Rubisco large subunit binding protein	gi/2493650	<i>Secale</i>	141/38	53.7
29								
30								
31	453	0.014	1.2	Globulin-like protein	TC246874	<i>T. aestivum</i>	383/47	71.1
32								
33	456	0.049	1.2	Serpin	TC236181		87/34	34.8
34								
35	460	0.01	1.2	GAPDH cytosolic	gi/32478662	<i>T. aestivum</i>	103/53	18.2
36								
37	468	0.047	1.4	Embryo globulin	TC234134		94/16	76.5
38								
39	470	0.022	1.3	Glucose and ribitol dehydrogenase	gi/7431022	<i>H. vulgare</i>	89/28	31.6
40								
41								
42								
43								
44								
45								
46								
47								

1								
2	483	0.032	1.3	Late embryogenesis abundant (LEA)	TC268629		97/32	30.5
3								
4								
5	484	0.003	1.3	Late embryogenesis abundant (LEA)	TC268629		126/34	30.5
6								
7								
8	497	0.004	1.6	HSP26	gi/147225072	<i>T. aestivum</i>	110/45	26.6
9								
10	502	0.006	1.2	Glucose and ribitol dehydrogenase homologue	gi/7431022	<i>H. vulgare</i>	95/31	31.6
11								
12								
13	503	0.005	1.2	GAPDH	gi/148508784	<i>T. aestivum</i>	123/61	36.6
14								
15	505	0.01	1.4	Embryo-specific protein	TC235043		98/14	37.6
16								
17								
18	508	0.012	1.4	Glyoxalase I	TC264636		91/36	44
19								
20	509	0.006	1.7	Globulin-like protein	TC246874		290/50	71.1
21								
22	511	0.009	1.3	Globulin-like protein	TC246703		112/26	68.2
23								
24	513	0.033	1.3	Hypothetical protein Oryza with Enolase Domain	gi/115451911	<i>O. sativa</i>	69/22	51.1
25								
26								
27	514	0.046	1.2	GAPDH cytosolic	gi/120680 TC264316		119/28	36.6
28								
29								
30								
31	<b>Down regulated proteins</b>							
32	109	0.018	-1.3	Phosphoglycerate Kinase	gi/129916	<i>T. aestivum</i>	176/43	42.1
33								
34	110	0.016	-1.6	ATP Synthase $\beta$ subunit	gi/525291 and TC264886	<i>T. aestivum</i>	123/21	77.5
35								
36								
37	110	0.016	-1.6	Globulin-like protein	TC246874		104/23	71.1
38								
39								
40	140	8.82E-04	-1.6	Tritin	gi/391929	<i>T. aestivum</i>	112/24	29.5
41								
42								
43								
44								
45								
46								
47								


1								
2	158	0.002	-1.4	ATP Syntase $\beta$ subunit	gi/525291	<i>T. aestivum</i>	81/13	59.3
3								
4	173	0.007	-1.3	Glycosyl hydrolase 85	gi/30695320	<i>A. thaliana</i>	85/29	40.7
5								
6	239	0.022	-1.2	Homologue to Embryo	TC234045		163/41	77.3
7				globulin				
8								
9	245	0.008	-1.3	Single stranded nucleic acid	gi/974605,		100/19	19.3
10				binding protein	TC249148			
11								
12	299	0.003	-1.2	Fructose-6-P 1	TC248170		71/13	82.3
13				phosphotransferase				
14								
15	341	0.007	-1.4	14-3-3 homologue	gi/22607 and	<i>H. vulgare</i>	172/47	29.4
16					TC233195			
17								
18								
19								
20								
21								
22								
23								
24								
25								
26								
27								
28								
29								
30								
31								
32								
33								
34								
35								
36								
37								
38								
39								
40								
41								
42								
43								
44								
45								
46								
47								

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60


For Peer Review



**Fig. 1.** 2-D PAGE map of metabolic proteins in the 3–10 pH range: circles indicate the differential spots




**FIG 2** 2D-PAGE of the metabolic proteins that are differentially regulated after the heat treatment: the yellow selection is referred to spots down-regulated, whereas the black one to up-regulated (fold change between 1.2 and 2.2). In the map all other proteins not involved in the heat response have been removed.



**Fig. 3.** PCA representation in which differential spots relative to each gel analysed are reported.

Circles: heat stressed samples; squares: control samples. Numbers represent the differentially expressed spots.





**Fig. 4.** 2-D PAGE map of metabolic proteins in the 3-10 pH range: circles indicate the identified proteins by MALDI-TOF and MALDI-TOF-TOF