

**Lower bounds for the life span of
solutions of semilinear wave equations
with data of non compact support**

Kimitoshi Tsutaya

Series #229. February 1994

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- # 199: A. Arai, N. Tominaga, Quantization of angle-variables, 31 pages. 1993.
- # 200: S. Izumiya, Y. Kurokawa, Holonomic systems of Clairaut type, 17 pages. 1993.
- # 201: K.-S. Saito, Y. Watatani, Subdiagonal algebras for subfactors, 7 pages. 1993.
- # 202: K. Iwata, On Markov properties of Gaussian generalized random fields, 7 pages. 1993.
- # 203: A. Arai, Characterization of anticommutativity of self-adjoint operators in connection with Clifford algebra and applications, 13 pages. 1993.
- # 204: J. Wierzbicki, An estimation of the depth from an intermediate subfactor, 7 pages. 1993.
- # 205: N. Honda, Vanishing theorem for the tempered distributions, 11 pages. 1993.
- # 206: T. Hibi, Betti number sequences of simplicial complexes, Cohen-Macaulay types and Möbius functions of partially ordered sets, and related topics, 25 pages. 1993.
- # 207: A. Inoue, Regularly varying correlations, 23 pages. 1993.
- # 208: S. Izumiya, B. Li, Overdetermined systems of first order partial differential equations with singular solution, 9 pages. 1993.
- # 209: T. Hibi, Hochster's formula on Betti numbers and Buchsbaum complexes, 7 pages. 1993.
- # 210: T. Hibi, Star-shaped complexes and Ehrhart polynomials, 5 pages. 1993.
- # 211: S. Izumiya, G. T. Kossioris, Geometric singularities for solutions of single conservation laws, 28 pages. 1993.
- # 212: A. Arai, On self-adjointness of Dirac operators in Boson-Fermion Fock spaces, 43 pages. 1993.
- # 213: K. Sugano, Note on non-commutative local field, 3 pages. 1993.
- # 214: A. Hoshiga, Blow-up of the radial solutions to the equations of vibrating membrane, 28 pages. 1993.
- # 215: A. Arai, Scaling limit of anticommuting self-adjoint operators and nonrelativistic limit of Dirac operators, 35 pages. 1993.
- # 216: Y. Giga, N. Mizoguchi, Existence of periodic solutions for equations of evolving curves, 45 pages. 1993.
- # 217: T. Suwa, Indices holomorphic vector fields relative to invariant curves, 10 pages. 1993.
- # 218: S. Izumiya, G. T. Kossioris, Realization theorems of geometric singularities for Hamilton-Jacobi equations, 14 pages. 1993.
- # 219: Y. Giga, K. Yama-uchi, On instability of evolving hypersurfaces, 14 pages. 1993.
- # 220: W. Bruns, T. Hibi, Cohen-Macaulay partially ordered sets with pure resolutions, 11 pages. 1993.
- # 221: S. Jimbo, Y. Morita, Ginzburg Landau equation and stable solutions in a rotational domain, 32 pages. 1993.
- # 222: T. Miyake, Y. Maeda, On a property of Fourier coefficients of cusp forms of half-integral weight, 12 pages. 1993.
- # 223: I. Nakai, Notes on versal deformation of first order PDE and web structure, 34 pages. 1993.
- # 224: I. Tsuda, Can stochastic renewal of maps be a model for cerebral cortex?, 30 pages. 1993.
- # 225: H. Kubo, K. Kubota, Asymptotic behaviors of radial solutions to semilinear wave equations in odd space dimensions, 47 pages. 1994.
- # 226: T. Nakazi, K. Takahashi, Two dimensional representations of uniform algebras, 7 pages. 1994.
- # 227: N. Hayashi, T. Ozawa, Global, small radially symmetric solutions to nonlinear Schrödinger equations and a gauge transformation, 16 pages. 1994.
- # 228: S. Izumiya, Characteristic vector fields for first order partial differential equations, 9 pages. 1994.

Lower bounds for the life span of solutions of semilinear wave equations with data of non compact support

Kimitoshi Tsutaya

Department of Mathematics
Hokkaido University
Sapporo 060, Japan

1 Introduction

We consider the life span of classical solutions of the Cauchy problem for the semilinear wave equation

$$\begin{cases} \partial_t^2 u - \Delta u = |u|^{p-1}u, & (x, t) \in R^n \times (0, \infty), \\ u(x, 0) = \varepsilon f(x), \partial_t u(x, 0) = \varepsilon g(x), & x \in R^n, \end{cases} \quad (1.1)$$

where $p > 1$, $n = 2, 3$ and $\varepsilon > 0$ is a small parameter.

Given f, g, p , we define the life span $T^* = T^*(\varepsilon)$ as the supremum of all T such that a C^2 -solution of (1.1) exists for all $x \in R^n$ and $0 \leq t < T$. Existence of global solutions and blow-up of local solutions in finite time correspond to $T^*(\varepsilon) = \infty$, $T^*(\varepsilon) < \infty$, respectively. In the case $T^*(\varepsilon) < \infty$ the basic problem consists in estimating $T^*(\varepsilon)$ in terms of ε both from above and from below, namely, upper and lower bounds for $T^*(\varepsilon)$. Upper and lower bounds are related to the construction of blow-up and local solutions, respectively, and it is expected that both bounds have the same growth rate as $\varepsilon \rightarrow 0$. There is a large literature on this problem [1-16], though there still remains a gap between the upper and lower bounds for $T^*(\varepsilon)$. The purpose of this paper is to improve the previous lower bound estimates and reduce that gap. Before we state our result more precisely, we summarize a number of available results on the life span of solutions of (1.1). In the sequel it is convenient

to introduce the critical power $p_0(n)$ defined by the positive root of the quadratic equation

$$(n-1)p^2 - (n+1)p - 2 = 0 \quad (n = 2, 3).$$

We note that $p_0(3) = 1 + \sqrt{2}$ and $p_0(2) = (3 + \sqrt{17})/2$. It is well understood that in most cases the situation is remarkably different between the super critical case $p > p_0(n)$, the critical case $p = p_0(n)$, and the subcritical case $p < p_0(n)$. Moreover, recent developments reveal that the compactness of the data f, g has a definitive influence on the problem. Therefore we describe the previous results by dividing into two cases, whether f, g have compact support or not. First, we assume that the data $f(x), g(x)$ are smooth and "compactly supported".

John [5, 6], Glassey [3, 4] and Schaeffer [10] have proved in two and three space dimensions that if $p > p_0(n)$ and ε is sufficiently small, then $T^*(\varepsilon) = \infty$, and also that if $1 < p \leq p_0(n)$, $T^*(\varepsilon) < \infty$ for small data.

In the case that blow-up occurs in finite time, that is, $T^*(\varepsilon) < \infty$, there arises the problem of estimating the life span $T^*(\varepsilon)$ in terms of ε . John [5] has shown that

$$A\varepsilon^{-2} \leq T^*(\varepsilon) \leq B\varepsilon^{-2}$$

with certain constants A, B in the case $n = 3$ and $p = 2$. Recently, Lindblad [8] has proved the following results:

(i) if $n = 3$ and $1 < p < p_0(3)$,

$$\lim_{\varepsilon \rightarrow 0} \varepsilon^{p(p-1)/(1+2p-p^2)} T^*(\varepsilon) = T > 0 \quad \text{exists,}$$

(ii) if $n = 2$ and $p = 2$,

$$\lim_{\varepsilon \rightarrow 0} \varepsilon T^*(\varepsilon) = T > 0 \quad \text{exists, provided } \int g(x) dx = 0,$$

and

$$\lim_{\varepsilon \rightarrow 0} a(\varepsilon)^{-1} T^*(\varepsilon) = T > 0 \quad \text{exists, provided } \int g(x) dx \neq 0,$$

where $a(\varepsilon)$ satisfies

$$\varepsilon^2 a(\varepsilon)^2 \ln(1 + a(\varepsilon)) = 1.$$

In [15, 16], Zhou has proved the following:

(iii) when $n = 3$,

$$A\varepsilon^{-p(p-1)/(1+2p-p^2)} \leq T^*(\varepsilon) \leq B\varepsilon^{-p(p-1)/(1+2p-p^2)}, \quad \text{if } 2 \leq p < p_0(3), \quad (1.2)$$

and

$$\exp\{A\varepsilon^{-p(p-1)}\} \leq T^*(\varepsilon) \leq \exp\{B\varepsilon^{-p(p-1)}\}, \quad \text{if } p = p_0(3),$$

(iv) when $n = 2$,

$$\lim_{\varepsilon \rightarrow 0} \varepsilon^{2p(p-1)/(2+3p-p^2)} T^*(\varepsilon) = T > 0 \quad \text{exists,} \quad \text{if } 2 < p < p_0(2),$$

and

$$\exp\{A\varepsilon^{-p(p-1)}\} \leq T^*(\varepsilon) \leq \exp\{B\varepsilon^{-p(p-1)}\}, \quad \text{if } p = p_0(2),$$

where A and B are positive constants. Concerning the above results due to Zhou for the case $p = p_0(n)$ ($n = 2, 3$), the simplified proofs of the upper bound of $T^*(\varepsilon)$ have been recently given by Takamura [11]. He has proved that if $p = p_0(n)$ ($n = 2, 3$) and

$$f(x) = 0, \quad g(x) \geq 0, \quad (1.3)$$

then

$$T^*(\varepsilon) \leq \exp\{B\varepsilon^{-p(p-1)}\}, \quad (1.4)$$

even if the data are not compactly supported. In [1], Agemi and Takamura have proved similar results to Lindblad and Zhou's for the case $n = 2$. Especially, the upper bound of $T^*(\varepsilon)$ in [1]

$$T^*(\varepsilon) \leq B\varepsilon^{-2p(p-1)/(2+3p-p^2)}, \quad \text{if } 1 < p < p_0(2) \quad (1.5)$$

holds even with the data of non compact support, under the condition (1.3). In the case $n = 3$ and $1 < p < p_0(3)$, we can also derive for the data of non compact

support, the upper bound of $T^*(\varepsilon)$ in (1.2),

$$T^*(\varepsilon) \leq B\varepsilon^{-p(p-1)/(1+2p-p^2)} \quad (1.6)$$

under (1.3).

We next assume that the data do not have compact support. Asakura [2], Kubota [7] and the author [12, 13, 14] have proved the following:

Let the data $f(x) \in C^3(R^n)$, $g(x) \in C^2(R^n)$ satisfy

$$D_x^\alpha f(x), D_x^\beta g(x) = O(|x|^{-1-k}) \quad \text{as } |x| \rightarrow \infty, \quad |\alpha| \leq 3, \quad |\beta| \leq 2.$$

If $k \geq 2/(p-1)$, $p > p_0(3)$ for $n = 3$ and $p > p_0(2)$ for $n = 2$, and ε is small, then $T^*(\varepsilon) = \infty$ (see also Pecher [9]).

On the other hand, Asakura [2], Agemi and Takamura [1] and the author [12] have also proved that if the data satisfy

$$f(x) = 0, \quad g(x) \geq \frac{1}{(1+|x|)^{1+k}}$$

and $0 < k < 2/(p-1)$, then $T^*(\varepsilon) < \infty$, even with $p > p_0(n)$ ($n = 2, 3$). In this case, the upper bound for $T^*(\varepsilon)$ is

$$T^*(\varepsilon) \leq B\varepsilon^{-(p-1)/\{2-k(p-1)\}} \quad \text{for } n = 2, 3, \quad (1.7)$$

where B is a constant depending only on p and k .

We note here that when $1 < p < p_0(n)$ ($n = 2, 3$), if $k > 1/2 + 1/p$ (resp. $0 < k < 1/2 + 1/p$) for $n = 2$ and $k > 1 + 1/p$ (resp. $0 < k < 1 + 1/p$) for $n = 3$, then the upper bounds of $T^*(\varepsilon)$ in (1.5), (1.6) (resp. in (1.7)) are sharper than those in (1.7) (resp. in (1.5), (1.6)), respectively.

For the lower bound, it has been shown by Kubota [7] and the author [13, 14] that if $p > p_0(n)$ ($n = 2, 3$) and $0 < k < 2/(p-1)$,

$$T^*(\varepsilon) \geq A\varepsilon^{-(p-1)/\{2-k(p-1)\}}.$$

For $n = 2$, Kubota has also obtained the lower bounds in the other cases:

$$\begin{aligned} p = p_0(2) \quad \text{and} \quad k &\geq \frac{1}{2} + \frac{1}{p}, \\ 2 < p \leq p_0(2) \quad \text{and} \quad 0 < k &< \frac{1}{2} + \frac{1}{p}, \\ 2 < p < p_0(2) \quad \text{and} \quad k &\geq \frac{1}{2} + \frac{1}{p}. \end{aligned}$$

In particular, for the critical case $p = p_0(2)$ and $k \geq 1/2 + 1/p$, the lower bound due to Kubota is the following:

$$T^*(\varepsilon) \geq \exp\{A\varepsilon^{-(p-1)}\}. \quad (1.8)$$

We note that there is a gap between (1.4) and (1.8) about the order of ε in the estimates of life span. In this paper, we shall improve the lower bound (1.8) and give an almost optimal lower bound for $T^*(\varepsilon)$ for the critical cases $p = p_0(2)$ and $k > 1/2 + 1/p$, and $p = p_0(3)$ and $k > 1 + 1/p$ in two and three space dimensions.

The crucial part of the proof of our theorem below is to establish the basic estimate for (1.1). We now state the difference between the proofs of the basic estimates for the data of compact support and for those of non compact support. The weight function in the norm is related with the decay estimate of the solution to the linear problem. In three space dimensions, if the data $f(x)$, $g(x)$ satisfy $\text{supp}\{f, g\} \subset \{x \in R^3 : |x| \leq K\}$, then, as is well known, the solution u^0 of $\partial_t^2 u^0 - \Delta u^0 = 0$ is equal to zero in the inside of the characteristic cone $\Gamma = \{(x, t) : |x| + K \leq t\}$ by Huygens' Principle. Hence we can define the special weight function (see [5], (58b) and also [15], (3.5)). This leads to the sharp basic estimate. In two space dimensions, we need to use another method due to the lack of Huygens' Principle. In fact, Zhou [16] has applied the Friedlander radiation field technique to (1.1) to obtain the sharp estimate. However, the above methods do not seem to be applicable to our problem, since the data are not compactly supported.

The main idea in the proof of the theorem is to introduce new weight functions which are slightly different from those used in [7]. The logarithmic factor in the weight function gives more precise information on the space-time behavior

of solutions and plays an important role to prove our main theorem (see (3.6)).

Accordingly, our method is simpler than those by John and Zhou.

2 Assumptions and Main Theorem

We study the Cauchy problem for the semilinear wave equation

$$\begin{cases} \partial_t^2 u - \Delta u = F(u), & (x, t) \in R^n \times (0, \infty), \\ u(x, 0) = \varepsilon f(x), \partial_t u(x, 0) = \varepsilon g(x), & x \in R^n. \end{cases} \quad (2.1)$$

Before stating the main result, we make the following hypotheses: Let $p = p_0(n)$.

(H1) $F(u) \in C^2(R)$ and there exists $\lambda > 0$ such that

if $n = 3$,

$$|F^{(j)}(u)| \leq \left(\frac{d}{dz} \right)^j (\lambda z^p) \Big|_{z=|u|} \quad \text{for } |u| \leq 1, j = 0, 1, 2,$$

$$|F'''(u) - F'''(v)| \leq \lambda p(p-1)|u-v|^{p-2} \quad \text{for } |u|, |v| \leq 1,$$

and if $n = 2$,

$$|F^{(j)}(u)| \leq \lambda |u|^{p-j} \quad \text{for } |u| \leq 1, j = 0, 1, 2,$$

$$|F'''(u) - F'''(v)| \leq \lambda p(p-1)|\Phi|^{p-3}|u-v| \quad \text{for } |u|, |v| \leq 1,$$

where

$$\Phi = \max\{|u|, |v|\}.$$

(H2) $f(x) \in C^3(R^n)$, $g(x) \in C^2(R^n)$ ($n = 2, 3$) satisfy

$$\sum_{|\alpha| \leq 3} |D_x^\alpha f(x)| + \sum_{|\beta| \leq 2} |D_x^\beta g(x)| \leq \frac{1}{(1+|x|)^{1+k}}$$

with $k > 0$.

Our main theorem is the following:

Theorem 2.1 *Let $n = 2, 3$, $p = p_0(n)$ and $k > 2/(p - 1)$. Assume (H1) and (H2). Then there exist two sufficiently small constants $A, \varepsilon_0 > 0$ depending only on λ, p and k such that for all $0 < \varepsilon \leq \varepsilon_0$, (2.1) has a C^2 -solution for $0 \leq t < T^*(\varepsilon)$, where $T^*(\varepsilon)$ satisfies the following:*

$$\left(\ln(3 + T^*(\varepsilon))\right)^{1/p} \left(\ln(\ln(3 + T^*(\varepsilon)))\right)^{1/(p-1)} \varepsilon^{p-1} \geq A. \quad (2.2)$$

Remark 2.2 *If the left-hand side of (2.2) does not include the factor $\left(\ln(\ln(3 + T^*(\varepsilon)))\right)^{1/(p-1)}$ of slower growth, the estimate is sharp, since the dominant factor $\left(\ln(3 + T^*(\varepsilon))\right)^{1/p}$ with the growth rate $A\varepsilon^{-(p-1)}$ gives precisely (1.4). In this respect (2.2) is almost optimal.*

The plan in the present paper is as follows. In Section 3, we first describe the decay estimates for the homogeneous wave equation (see [2], [7], [12, 13, 14]). Next, we derive the basic estimates needed for the proof of Theorem 2.1. Finally, in Section 4 we prove Theorem 2.1. We denote a constant in the estimates by C , which will change from step to step.

3 Decay Estimates and Basic Estimates

We consider the linear problem

$$\begin{cases} \partial_t^2 u - \Delta u = w(x, t), & (x, t) \in R^n \times (0, \infty), \\ u(x, 0) = \varepsilon f(x), \partial_t u(x, 0) = \varepsilon g(x), & x \in R^n \ (n = 2, 3). \end{cases} \quad (3.1)$$

We see that the solution $u(x, t)$ of (3.1) is given by

$$u(x, t) = u^0(x, t) + Lw(x, t), \quad (3.2)$$

where $u^0(x, t)$ is the solution of $\partial_t^2 u^0 - \Delta u^0 = 0$ with initial data $\varepsilon f(x)$, $\varepsilon g(x)$, and

$$Lw(x, t) = \begin{cases} \frac{1}{4\pi} \int_0^t (t - \tau) \int_{|\omega|=1} w(x + (t - \tau)\omega, \tau) d\omega d\tau & (n = 3), \\ \frac{1}{2\pi} \int_0^t \int_0^{t-s} \frac{\rho}{\sqrt{(t-s)^2 - \rho^2}} \int_{|\omega|=1} w(x + \rho\omega, s) d\omega d\rho ds & (n = 2) \end{cases} \quad (3.3)$$

is the solution of $\partial_t^2 u - \Delta u = w$ with zero data, where $d\omega$ is the surface measure on the unit sphere $|\omega| = 1$. Note that $w \geq 0$ implies $Lw \geq 0$.

The following lemma is proved in [2], [7] and [12, 13, 14].

Lemma 3.1 *Assume that $f(x)$, $g(x)$ satisfy (H2). Then, the solution u^0 of $\partial_t^2 u^0 - \Delta u^0 = 0$ with initial data $\varepsilon f(x)$, $\varepsilon g(x)$ satisfies the following:*

(i) if $n = 3$,

$$\sum_{|\alpha| \leq 2} |D_x^\alpha u^0(x, t)| \leq \begin{cases} \frac{C_k \varepsilon}{(1+t+r)(1+|t-r|)^{k-1}} & (k > 1), \\ \frac{C_k \varepsilon}{1+t+r} \left(1 + \ln \frac{1+t+r}{1+|t-r|} \right) & (k = 1), \\ \frac{C_k \varepsilon}{(1+t+r)^k} & (0 < k < 1), \end{cases} \quad (3.4)$$

(ii) if $n = 2$,

$$\sum_{|\alpha| \leq 2} |D_x^\alpha u^0(x, t)| \leq \begin{cases} \frac{C_k \varepsilon}{\sqrt{(1+t+r)(1+|t-r|)}} & (k > 1), \\ \frac{C_k \varepsilon \ln(2+|t-r|)}{\sqrt{(1+t+r)(1+|t-r|)}} & (k = 1), \\ \frac{C_k \varepsilon}{\sqrt{1+t+r}(1+|t-r|)^{k-1/2}} & (\frac{1}{2} < k < 1), \\ \frac{C_k \varepsilon}{\sqrt{1+t+r}} \left(1 + \ln \frac{1+t+r}{1+|t-r|} \right) & (k = \frac{1}{2}), \\ \frac{C_k \varepsilon}{(1+t+r)^k} & (0 < k < \frac{1}{2}), \end{cases} \quad (3.5)$$

where $r = |x|$, and a constant C_k depends only on k .

We next introduce the function

$$Z_n(|x|, t) = (1+t+|x|)^m (1+|t-|x||)^{1/p} (\ln(3+|t-|x||))^{1/p} (\ln(\ln(3+|t-|x||)))^{1/(p-1)}, \quad (3.6)$$

where

$$m = \begin{cases} 1 & (n = 3), \\ \frac{1}{2} & (n = 2), \end{cases}$$

and define the norm for functions $u(x, t)$ which are continuous on $R^n \times [0, T)$

$$\|u\|_n = \sup_{\substack{x \in R^n \\ 0 \leq t < T}} Z_n(|x|, t) |u(x, t)|. \quad (3.7)$$

The following lemma provides the basic estimate for the existence proof.

Lemma 3.2 *If $p = p_0(n)$ ($n = 2, 3$), then there exists a constant C_p depending only on p such that*

$$\|L|u|^p\|_n \leq C_p (\ln(3+T))^{1/p} (\ln(\ln(3+T)))^{1/(p-1)} \|u\|_n^p. \quad (3.8)$$

Proof) Following [13], we first prove the lemma above in the case $n = 3$. If w is spherically symmetric, i.e., $w(x, t) = w(r, t)$, then from (3.3) we can write the solution of $\partial_t^2 u - \Delta u = w$ with zero data in the form

$$u(r, t) = \frac{1}{2r} \int_0^t \int_{|r-t+\tau|}^{r+t-\tau} \rho w(\rho, \tau) d\rho d\tau. \quad (3.9)$$

We denote (3.9) by $u = Pw$. We define the function

$$\tilde{u}(r, t) = \sup_{|x|=r} |u(x, t)|.$$

Then, we have

$$|(Lw)(x, t)| \leq (P\tilde{w})(r, t) \quad (3.10)$$

(see [2], p.1470 and also [5], Lemma II). Thus, it is sufficient to estimate

$$(P\tilde{u}^p)(r, t) = \frac{1}{2r} \int_0^t \int_{|r-t+\tau|}^{r+t-\tau} \rho \tilde{u}(\rho, \tau)^p d\rho d\tau.$$

We use the following lemma.

Lemma 3.3 *Let $k > 1$. Then,*

$$\frac{1}{2r} \int_{|r-t|}^{r+t} \frac{1}{(1+\rho)^k} d\rho \leq \frac{\max(1, k-1) \min(r, t)}{(k-1)r(1+t+r)(1+|t-r|)^{k-1}}$$

for all $t, r \geq 0$.

Proof) See [2]. ■

To prove (3.8), we distinguish two cases, $r \geq t$ and $t \geq r$.

(a) $r \geq t$.

By (3.6) and (3.7), we have

$$(P\tilde{u}^p)(r, t) \leq \frac{\|u\|_3^p}{2r} \int_0^t \int_{r-t+\tau}^{r+t-\tau} \frac{\rho}{(1+\tau+\rho)^p(1+\rho-\tau)} \\ \times \frac{1}{\ln(3+\rho-\tau)(\ln(\ln(3+\rho-\tau)))^{p/(p-1)}} dp d\tau.$$

Changing variables by

$$\alpha = \rho + \tau, \quad \beta = \rho - \tau, \quad (3.11)$$

we obtain

$$(P\tilde{u}^p)(r, t) \leq \frac{\|u\|_3^p}{4r} \int_{r-t}^{r+t} \frac{1}{(1+\alpha)^{p-1}} \int_{r-t}^{\alpha} \frac{1}{(1+\beta) \ln(3+\beta) (\ln(\ln(3+\beta)))^{p/(p-1)}} d\beta d\alpha \\ \leq \frac{C\|u\|_3^p}{r(\ln(\ln(3+r-t)))^{1/(p-1)}} \int_{r-t}^{r+t} \frac{1}{(1+\alpha)^{p-1}} d\alpha \\ \leq \frac{C\|u\|_3^p (\ln(3+r))^{1/p} t}{r(1+t+r)(1+r-t)^{p-2} (\ln(3+r-t))^{1/p} (\ln(\ln(3+r-t)))^{1/(p-1)}}.$$

We have used here Lemma 3.3. Since the function $\varphi(x) = x/(\ln(3+x))^q$ ($0 < q < 1$) is increasing for $x \geq 0$ and $p-2 = 1/p$, we obtain

$$(P\tilde{u}^p)(r, t) \leq \frac{C\|u\|_3^p (\ln(3+t))^{1/p}}{(1+t+r)(1+r-t)^{p-2} (\ln(3+r-t))^{1/p} (\ln(\ln(3+r-t)))^{1/(p-1)}} \\ \leq \frac{C\|u\|_3^p (\ln(3+t))^{1/p}}{(1+t+r)(1+r-t)^{1/p} (\ln(3+r-t))^{1/p} (\ln(\ln(3+r-t)))^{1/(p-1)}}.$$

(b) $t \geq r$.

We have

$$(P\tilde{u}^p)(r, t) \leq \frac{\|u\|_3^p}{2r} \int_0^t \int_{|r-t+\tau|}^{r+t-\tau} \frac{\rho}{(1+\tau+\rho)^p(1+|\tau-\rho|)} \\ \times \frac{1}{\ln(3+|\rho-\tau|)(\ln(\ln(3+|\rho-\tau|)))^{p/(p-1)}} dp d\tau.$$

By (3.11),

$$\begin{aligned}
& (P\tilde{u}^p)(r, t) \\
& \leq \frac{\|u\|_3^p}{4r} \int_{t-r}^{t+r} \int_{r-t}^{\alpha} \frac{1}{(1+\alpha)^{p-1}(1+|\beta|)\ln(3+|\beta|)(\ln(\ln(3+|\beta|)))^{p/(p-1)}} d\beta d\alpha \\
& = \frac{\|u\|_3^p}{4r} \int_{t-r}^{t+r} \frac{1}{(1+\alpha)^{p-1}} \int_{r-t}^{t-r} \frac{1}{(1+|\beta|)\ln(3+|\beta|)(\ln(\ln(3+|\beta|)))^{p/(p-1)}} d\beta d\alpha \\
& \quad + \frac{\|u\|_3^p}{4r} \int_{t-r}^{t+r} \frac{1}{(1+\alpha)^{p-1}} \int_{t-r}^{\alpha} \frac{1}{(1+\beta)\ln(3+\beta)(\ln(\ln(3+\beta)))^{p/(p-1)}} d\beta d\alpha \\
& = \frac{\|u\|_3^p}{2r} \int_{t-r}^{t+r} \frac{1}{(1+\alpha)^{p-1}} \int_0^{t-r} \frac{1}{(1+\beta)\ln(3+\beta)(\ln(\ln(3+\beta)))^{p/(p-1)}} d\beta d\alpha \\
& \quad + \frac{\|u\|_3^p}{4r} \int_{t-r}^{t+r} \frac{1}{(1+\alpha)^{p-1}} \int_{t-r}^{\alpha} \frac{1}{(1+\beta)\ln(3+\beta)(\ln(\ln(3+\beta)))^{p/(p-1)}} d\beta d\alpha \\
& \leq \frac{C\|u\|_3^p}{r} \int_{t-r}^{t+r} \frac{1}{(1+\alpha)^{p-1}} d\alpha.
\end{aligned}$$

We use Lemma 3.3 to obtain

$$(P\tilde{u}^p)(r, t) \leq \frac{C\|u\|_3^p (\ln(3+t))^{1/p} (\ln(\ln(3+t)))^{1/(p-1)}}{(1+t+r)(1+t-r)^{1/p} (\ln(3+t-r))^{1/p} (\ln(\ln(3+t-r)))^{1/(p-1)}}.$$

Next, the proof of Lemma 3.2 for the case $n = 2$ proceeds in the same way as that of Lemma 4.1 in [14] except that we need to use the following inequality instead of Lemma 4.3 in [14].

$$\int_0^{\alpha} \frac{1}{\sqrt{a+t-\beta}(1+\beta)\ln(3+\beta)(\ln(\ln(3+\beta)))^k} d\beta \leq \frac{C_k}{\sqrt{t+a}} \quad (3.12)$$

for $k > 1$ and $0 \leq \alpha \leq t+a$,

where C_k is a constant depending only on k . Slight modification of the proof of Lemma 4.3 in [14] leads to (3.12).

This completes the proof of Lemma 3.2. ■

4 Proof of Theorem 2.1

Let X_n be the linear space defined by

$$X_n = \{u(x, t) : D_x^\alpha u(x, t) \in C(R^n \times [0, T]), \|D_x^\alpha u\|_n < \infty \text{ for } |\alpha| \leq 2\},$$

$n = 2, 3$.

We can verify easily that X_n is complete with respect to the norm

$$\|u\|_{X_n} = \sum_{|\alpha| \leq 2} \|D_x^\alpha u\|_n.$$

We define the sequence of functions $\{u_n\}$ by

$$u_0 = u^0, \quad u_{n+1} = u^0 + LF(u_n).$$

Here we note that

$$k > \frac{2}{p-1} = \begin{cases} 1 + \frac{1}{p} & (n = 3), \\ \frac{1}{2} + \frac{1}{p} & (n = 2), \end{cases}$$

$$> \begin{cases} 1 & (n = 3), \\ \frac{1}{2} & (n = 2). \end{cases}$$

Then, Lemma 3.1 yields

$$\|u^0\|_{X_n} \leq C_k \varepsilon.$$

Hence, $u^0 \in X_n$.

We now assume that ε is so small that

$$2^p p \lambda C_p (\ln(3+T))^{1/p} (\ln(\ln(3+T)))^{1/(p-1)} (C_k \varepsilon)^{p-1} \leq 1 \quad \text{and} \quad C_k \varepsilon \leq \frac{1}{2}, \quad (4.1)$$

where C_p is a constant given in Lemma 3.2. Then, we have

$$2^p p \lambda C_p (\ln(3+T))^{1/p} (\ln(\ln(3+T)))^{1/(p-1)} \|u^0\|_n^{p-1} \leq 1 \quad \text{and} \quad \|u^0\|_n \leq \frac{1}{2}. \quad (4.2)$$

Therefore, as in [5], [2], we see that if u^0 satisfies (4.2), there exists a unique local solution of (2.1). The lower bound estimate (2.2) follows immediately from (4.1).

This completes the proof of Theorem 2.1. ■

Acknowledgement

The author would like to thank Professor T. Ozawa for his helpful advice.

References

- [1] R. Agemi and H. Takamura, The lifespan of classical solutions to nonlinear wave equations in two space dimensions, *Hokkaido Math. J.* **21**(1992), 517-542.
- [2] F. Asakura, Existence of a global solution to a semi-linear wave equation with slowly decreasing initial data in three space dimensions, *Comm. Partial Differential Equations* **11**, No. 13(1986), 1459-1487.
- [3] R. T. Glassey, Finite-time blow-up for solutions of nonlinear wave equations, *Math. Z.* **177**(1981), 323-340.
- [4] R. T. Glassey, Existence in the large for $\square u = F(u)$ in two space dimensions, *Math. Z.* **178**(1981), 233-261.
- [5] F. John, Blow-up of solutions of nonlinear wave equations in three space dimensions, *Manuscripta Math.* **28**(1979), 235-268.
- [6] F. John, Nonlinear wave equations, Formation of singularities, Lehigh University, University Lecture Series, Amer. Math. Soc. Providence, 1990.
- [7] K. Kubota, Existence of a global solution to a semi-linear wave equation with initial data of non-compact support in low space dimensions, *Hokkaido Math. J.* **22**(1993), 123-180.
- [8] H. Lindblad, Blow-up for solutions of $\square u = |u|^p$ with small initial data, *Comm. Partial Differential Equations* **15**(6) (1990), 757-821.
- [9] H. Pecher, Scattering for semilinear wave equations with small data in three space dimensions, *Math. Z.* **198**(1988), 277-289.
- [10] J. Schaeffer, The equation $u_{tt} - \Delta u = |u|^p$ for the critical value of p , *Proc. Roy. Soc. Edinburgh* **101A**(1985), 31-44.

- [11] H. Takamura, An elementary proof of the exponential blow-up for semilinear wave equations, to appear in Math. Meth. Appl. Sci.
- [12] K. Tsutaya, Global existence theorem for semilinear wave equations with non-compact data in two space dimensions, J. Differential Equations 104(1993), 332-360.
- [13] K. Tsutaya, Global existence and the life span of solutions of semilinear wave equations with data of non compact support in three space dimensions, to appear in Funkcialaj Ekvacioj.
- [14] K. Tsutaya, A global existence theorem for semilinear wave equations with data of non compact support in two space dimensions, Comm. Partial Differential Equations 17, No. 11&12(1992), 1925-1954.
- [15] Y. Zhou, Blow up of classical solutions to $\square u = |u|^{1+\alpha}$ in three space dimensions, J. Partial Differential Equations 5 (1992), 21-32.
- [16] Y. Zhou, Life span of classical solutions to $\square u = |u|^p$ in two space dimensions, to appear in Chin. Ann. of Math.